

MATGEN-Q: Scaling a Two-Stage Generative Quantum Eigensolver to 40 Qubits on NVIDIA CUDA-Q for EUV Photoresist Chemistry

Team EIGENNEXUS — Advanced Materials (Mitsubishi Chemical Group & AIST) — GIC 2026 Phase 3

1. Focus Area and Rationale

Ground-state energy estimation governs reaction thermodynamics, redox behavior, and excited-state properties across materials chemistry, yet classical methods scale exponentially with electron correlation and the gold-standard CCSD(T) costs $O(N^7)$. The Generative Quantum Eigensolver (GQE; Nakaji et al. 2024) replaces variational circuit optimization with a classical generative transformer that proposes quantum circuits, side-stepping the barren-plateau problem of VQE (Tilly et al. 2022; a GQE property established by Nakaji et al. 2024, adopted not re-demonstrated) — but statevector GQE stayed small-scale, and QSCI-paired GQE only recently reached 32 qubits (Kemmoku, Gao et al. 2026). **Scaling this generative approach to the ~40-qubit, industrially relevant regime — with executed, reproducible results — is the problem we address in Phase 3.** Our primary, executed contribution is quantum-accurate energies for the exact chemistry where BOTH classical pillars fail — DFT's spin-state ordering flips sign across functionals (CrO splitting spans 1.9 eV — so a DFT-only screen can assign the wrong spin ground state and rank the wrong candidate before costly synthesis) and single-reference CCSD(T) collapses non-variationally below the exact energy as bonds break — on open-shell, multireference transition-metal oxides (CrO, NiO) and EUV-relevant tin oxides (SnO, SnO₂) at 20 qubits, validated against exact diagonalization and third-party HamLib Hamiltonians. On top we add scalable machinery — tensor-network + selected-CI evaluation, MP2 pool compression, and a size-/chemistry-transferable generator — targeting the 40q+ regime, every claim reproducible and every limitation stated.

Our target is EUV semiconductor photoresist chemistry (tin-oxo clusters; Sn, Hf, Zr oxides), whose open-shell, multi-reference character is where DFT's functional-dependent errors make candidate rankings unreliable, and which is the focus of an active quantum-simulation program by one of this challenge's providers (Kharazi et al., Xanadu & Mitsubishi Chemical, 2026). We quantify the unreliability directly: across six standard functionals the CrO spin-state splitting spans 1.9 eV — PBE0 places the quintet 1.84 eV below the triplet while B3LYP inverts the order (−0.08 eV) — and even the milder NiO gap varies 0.11 eV, against a chemical-accuracy target of 0.044 eV. We resolve the ambiguity directly (Figure 1): CASCI and QSCI both place the quintet 1.89 eV below the triplet — agreeing with the experimental X⁵Π ground term — so B3LYP simply predicts the wrong ground state, and a single quantum-accurate number replaces the entire functional-choice band. The pipeline (AI proposes → DFT filters → GQE refines → Bayesian selects) generalizes to battery electrolytes, catalysts, and polymers central to Mitsubishi Chemical and AIST; in an \$800B+ semiconductor industry, multi-fidelity screening economics (Fare et al., 2022) are the cost lever MATGEN-Q targets via quantum-accurate pre-selection.



Figure 1. CrO spin-state decision. Six DFT functionals span 1.9 eV for the triplet–quintet gap and B3LYP flips the ground state to the (wrong) triplet; CASCI and QSCI in a fixed CAS(10,10) give one consistent quintet (⁵Π) ground state, agreeing with experiment. CAS/def2-SVP is a fixed modest active space, so the multireference value is the single number DFT scatters around — the robust claim is the 1.9 eV spread, the B3LYP sign flip, and the experimental-sign agreement, not the precise gap magnitude.

Stakeholder relevance. For Mitsubishi Chemical the value is a quantum-accurate filter ahead of expensive synthesis and lithographic testing, where DFT mis-ranks tin-oxo redox/bond-cleavage energetics; for AIST/G-QuAT it is a scalable, reproducible GQE workflow on NVIDIA CUDA-Q that extends the jointly-developed program past its size limits and composes with its excited-state work into one materials-informatics loop.

2. Target System and Data Modeling Strategy

Scaling vehicle. The linear hydrogen-chain series H_n (n = 2...20; 4–40 qubits, STO-6G, Jordan–Wigner) — the canonical strong-correlation benchmark; a single bond-length parameter tunes weak (area-law) to strong (volume-law) correlation, directly stressing the simulation layer that underpins any scaling claim.

Target chemistry. The same QSCI engine reaches chemical accuracy on real materials chemistry: Sn-oxide active spaces (Sn ECP-CASCI, validated on H₄ to 0.0000 mHa) — SnO (16q) and SnO₂ (20q) to chemical accuracy (≤0.6 mHa asserted by reproduce.py; observed 0.11–0.45 / 0.18–0.23 across environments), and — to reach the genuine EUV motif rather than a diatomic — a bridged **Sn–O–Sn tin-oxo fragment (Sn₂O₂)**

rhombus) to 0.41 mHa at 16q; and genuine open-shell, multireference transition-metal oxides — **CrO $^5\Pi$ 0.038 mHa and NiO $^3\Sigma^-$ 0.197 mHa at 20 qubits.** We report these executed 20-qubit results; the 38-qubit regime is the GPU scaling target of §5, not a pre-claimed figure — explicitly superseding the aspirational ≤ 0.08 mHa @38q of our Phase 2 draft, which we do not claim.

Data integrity. We adopt HamLib (Sawaya et al., Quantum 2024) and built a generation pipeline replicating its methodology (PySCF $\rightarrow$ OpenFermion $\rightarrow$ Jordan–Wigner, STO-6G). **Validated against the published files: FCI reproduced to 0.00000 mHa (H_2-H_6), and at 28/32/40 qubits our Hamiltonians match HamLib exactly in term count (27,735 / 47,489 / 116,577) and to ~ 15 significant figures in coefficient magnitude** (differing only by a spectrum-invariant orbital-phase convention) — third-party reproducible.

Accuracy anchors / classical references. FCI exactly ($\leq 20q$), CCSD(T) near equilibrium, and DMRG — verified to reproduce FCI to 0.000 mHa at 20q — as the reference under strong correlation where CCSD(T) breaks down.

3. GQE-Based Approach and Algorithmic Innovation

MATGEN-Q uses a two-stage GQE. Stage 1 (generative structure discovery): a decoder-only GPT-style transformer, trained by sequence–energy matching (Nakaji et al.), generates circuits as token sequences over a UCC single/double excitation pool. **Stage 2 (continuous refinement):** adjoint-gradient angle optimization converges to chemical accuracy. Four innovations make it scalable:

(1) Tensor-network (MPS) simulation — the GPU scaling enabler, bond-dimension argument now demonstrated on CPU. CUDA-Q's tensornet-mps memory scales with entanglement, not 2^n ; we verify the enabling claim with block2 DMRG (the classical MPS analogue): **the bond dimension χ for chemical accuracy grows only modestly with size ($\chi \approx 50/100/400$ at 20/28/40 qubits), and entanglement is area-law-bounded near equilibrium ($S_{\max} 0.39, \chi=50$) rising into strong correlation ($S_{\max} 4.43, \chi=200$).** So at 40q, $\chi \approx 400$ puts the MPS footprint ($\sim n \cdot \chi^2 \cdot d$, megabytes) far below the ~ 16 TB exact statevector — quantifying the GPU memory win. The full GPU tensornet-mps run at 40q remains the owed deliverable (§5c [QBRAID-RUN]); the measured χ target above de-risks it.

(2) QSCI energy evaluation — accuracy and noise-aware. Quantum-Selected Configuration Interaction (Kanno et al. 2023): dominant configurations are sampled from the generated circuit and H is classically diagonalized in that subspace — intrinsically noise-robust (the device defines only the subspace; at 20q with 2×10^6 shots the energy degrades gracefully to ≤ 3.3 mHa even with 30% of measurements corrupted; and executed through CUDA-Q's density-matrix backend with a physical depolarizing channel, QSCI on H_4 holds chemical accuracy up to 5% per-gate noise — a selection-principle robustness check on a small determinant space, not a hardware forecast).

(3) Operator-pool compression — efficiency (demonstrated). Ranking the $O(N^4)$ double-excitation pool by active-space MP2 amplitude and keeping the top fraction shrinks the transformer vocabulary with negligible accuracy loss, where random pruning collapses. Deterministic CI-subspace test ($CO/N_2/SiO$, 12q): **N_2 retains full-pool accuracy (2.26 mHa) keeping only 25% of doubles (vocabulary 1170 $\rightarrow$ 430) vs 50.4 mHa for random pruning at the same size ($\sim 22\times$);** CO holds 3.7 mHa at 40% kept vs 29.7 mHa random.

(4) Distributed hybrid workflow. Detailed in §4.

Transfer learning — a generative prior that transfers across system size and chemistry. Using a canonical, frontier-relative tokenization (each excitation encoded as occupied-depth-below-HOMO and virtual-height-above-LUMO, independent of qubit count), a small system's token vocabulary is a provable subset of a larger one ($H_6/12q \subset H_8/16q \subset H_{10}/20q$). One GPT-QE generator trained only on H_4+H_6 (8q+12q) is deployed zero-shot across 16 $\rightarrow$ 56 qubits. **The robust signals are statistical, not a single-number win: $\sim 3.7\times$ tighter selection spread than random (≈ 2.0 vs ≈ 7.4 mHa across-seed SD) and 13/18 size $\times$ seed paired wins (binomial $p \approx 0.05$)** — reliable selection versus a lottery. The per-size mean advantage is large and all-seeds-positive at 16q and stays positive in the mean through 28q ($+8.9/+8.3/+7.1$ mHa), then narrows into run-to-run noise at 40–56q; we report it as a mean trend, not an every-seed guarantee. That the learned policy captures real chemistry — not a per-system fit — is corroborated by interpretability: the generator's learned token distribution (trained on energy alone, blind to perturbation theory) recovers the MP2 double-excitation amplitude hierarchy with a significant rank correlation (Spearman $\rho=0.31$, $p<0.002$; 4 of its top-8 doubles are MP2's top-8). This is the *train-small, deploy-large* idea: GQE training is CPU-bound near ~ 16 qubits, so transferring a small-trained generator toward the 40q+ target — instead of training at scale — speaks directly to the scalability criterion. The enabler is a

determinant-space evaluation (each excitation maps by bitmask to a determinant; H is diagonalized in that subspace via Slater-Condon, validated to 0.0000 mHa against Jordan–Wigner) that needs no 2^n statevector, so 56q runs on CPU. This is a *selected-CI proxy* for circuit-sampled QSCI (deterministic construction, no shots), relative at a small fixed budget ($K=96$), not absolute chemical accuracy.

Cross-chemistry transfer (honest, partial). The same H-chain-trained generator, deployed to oxide active spaces, beats random clearly on 3 of 6 targets (CO-12q/SiO/CO-20q), ties within noise on BeO, and loses on SnO (heavy-element ECP chemistry too dissimilar from training). Target-specific MP2 selected-CI is the strongest selector on every target. Honest conclusion: cross-size transfer is robust; cross-chemistry works only when the target resembles training — a zero-cost prior, not a universal one. A same-size conditional-GQE variant (cf. Minami, Nakaji et al. 2025) gave only a within-noise gain, also reported as a negative.

4. Hybrid Architecture

The classical transformer trains on GPU (PyTorch); circuit evaluations are dispatched across GPUs via CUDA-Q's mcpu in an asynchronous generate → evaluate → update loop — quantum resources do state preparation and energy estimation (MPS/QSCI), classical resources do generation, optimization, and active-space selection. **We validate this execution path on CUDA-Q itself: the UCCSD/QSCI pipeline runs through CUDA-Q's qpp-cpu backend and reproduces exact FCI on H_4 (VQE within 0.02 mHa via `cudaq.observe`; QSCI within chemical accuracy via `cudaq.sample`), confirming the generated circuits compile and sample through the SDK; the 12–40-qubit regime is the owed GPU (`cuStateVec/tensornet-mps`) run.**

5. Phase 3 Execution and Results

Verified results (CPU). Core rows are re-executed end-to-end from a clean checkout by *reproduce.py* (13/13 PASS); every value traces to a committed result file (Table 1).

System	Qubits	Quantum (GQE/QSCI)	Classical ref.	Notes
$H_2/H_4/H_6$	4/8/12	0.000 / 0.009 / 0.297 mHa	FCI (exact)	two-stage GQE (UCCSD)
H_6 GQE → QSCI	12	1.05 mHa (51 → 1.05, $\sim 50\times$)	FCI	measured pipeline
H_{10}/H_{14}	20/28	0.57 / 1.21 mHa	FCI / CCSD(T)	2,401 / 18,201 dets
CrO $^5\Pi$ / NiO $^3\Sigma^-$	20	0.038 / 0.197 mHa	CASCI (exact)	open-shell multiref.
SnO / SnO ₂	16/20	chem. acc. ≤ 0.6 (obs. 0.11–0.45 / 0.18–0.23)	FCI	EUV target; values version/run sensitive
Noise (H_{10})	20	≤ 3.3 mHa @ 30% corrupt (2×10^6 shots)	—	noise-aware bonus

Table 1. Verified quantum results vs classical references on matched instances (CPU).

Classical baseline and the exact wall (matched instances, timed). On identical STO-6G H_n geometries, classical FCI wall-clock grows steeply ($\sim 6\times$ from 20 → 24 qubits, 0.60 s → 3.63 s on a single CPU thread) and is intractable by 32q on CPU; CCSD(T) stays cheap but its error climbs with correlation and breaks down under strong correlation (at $H_{24}/48q$, classical DMRG is itself 6.83 mHa off CCSD(T)). Exact statevector needs ~ 16 TB at 40q vs ~ 0.2 GB for the MPS tier (measured $\chi=400$), and $\sim 10^6$ QSCI determinants vs $\sim 3\times 10^{10}$ for full CI — the memory and determinant walls the MPS + QSCI tiers remove (Table 2; quantified in results/crossover).

System	Qubits	FCI (exact) wall-clock	CCSD(T) err vs FCI	CCSD(T) wall-clock
H_6	12	0.04 s	0.026 mHa	0.08 s
H_{10}	20	0.60 s	0.173 mHa	0.17 s
H_{12}	24	3.63 s	0.282 mHa	0.21 s
H_{14}	28	minutes	—	~ 0.2 s
H_{16}^+	32+	intractable (CPU)	—	—

Table 2. Classical reference cost on the H_n ladder (single-thread CPU; wall-clocks machine-dependent, reproduced in *classical_baselines_evidence.json* — the steep growth and the exact-method wall, not absolute seconds, are the claim; H_{14}^+ /intractable rows are qualitative) — the exact-method wall the quantum approach is measured against.

Scaling results targeted on qBraid GPU (owed — not yet executed, and pre-registered). Each owed run carries a committed quantitative pass/fail prediction derived only from measured scaling

(results/preregistration_v1.json, git-timestamped pre-access): a falsifiable test, not a post-hoc report. **40-qubit MPS GQE/QSCI on H_{20} (primary criterion):** *[QBRAID-RUN: energy err vs DMRG, circuit depth, MPS bond dimension, shot budget, GPU wall-clock]* (CPU baseline today: operational, converging through 39 mHa). **CrO/NiO near-38q on GPU:** *[QBRAID-RUN: accuracy + wall-clock]* (scaling demonstration on real open-shell chemistry, building on the executed 20q results). **Quantum-vs-classical wall-clock:** *[QBRAID-RUN: MPS/QSCI vs exact statevector vs VQE]*. **Hardware validation:** *[QBRAID-RUN: IonQ/IBM at 10–16q]*.

Determinant-budget sweep and a classical baseline. Sweeping the QSCI-proxy budget K: the generator beats random at every budget up to K=96 (at K=128 the two converge, random marginally ahead). Target-specific MP2 selected-CI is the strongest selector and generator+MP2 fusion adds nothing — the transferred generator sits between random and target-specific MP2 at *zero target-specific cost*, a zero-shot prior for high-throughput screening.

What the results establish. Three things are demonstrated, not asserted. (i) Correctness: the GQE/QSCI pipeline reproduces the FCI/CASCI reference to chemical accuracy on every instance we can run, including open-shell multireference oxides (CrO, NiO) and EUV-relevant Sn-oxides. (ii) Scalability: QSCI reaches chemical accuracy using a vanishing fraction of the CI space (3.8% at 20q, 0.15% at 28q), and operator-pool compression holds accuracy at 25–40% of the doubles pool — cost grows far slower than the Hilbert space. (iii) Transferable structure: a generator trained on small systems proposes good circuits for a larger one it never saw (§3). The honest boundary: at sizes classical FCI/DMRG still solve, we show correctness and favourable scaling, not a head-to-head speed win; that win is the 40q+ regime the GPU run targets, where exact classical methods break down (Table 2).

Where it matters most: trust under strong correlation. Stretching H_{10} (20q) to dissociation drives genuine multireference character (Hartree–Fock weight in the exact wavefunction falls $0.93 \rightarrow 0.06$), and the classical gold standard fails qualitatively: CCSD(T) collapses *non-variationally* to 217 mHa BELOW the exact FCI energy at $R=2.4 \text{ \AA}$ — a confidently wrong, unphysical answer. The determinant-subspace method (the QSCI principle, here via iterative selected-CI) instead **reaches chemical accuracy with only ~500 determinants** at every geometry across the dissociation, a rigorous variational bound systematically convergent to the exact answer. **Critically this is not a hydrogen-chain artifact: on a real Cr–O bond stretch (CrO $^5\Pi$, CAS(10,10)=20q), in-active-space CCSD(T) develops large, frequently non-convergent errors (tens to ~140 mHa vs exact CASCI) as the dominant-determinant weight falls $0.87 \rightarrow 0.16$, while selected-CI/QSCI stays variational and within a few mHa throughout (Figure 2).** And we certify the error rather than assert it: an Epstein–Nesbet PT2 correction tightens the rigorous variational upper bound toward FCI — **$E_{\text{var}}+\text{PT2}$ is a second-order estimate (not a bound) that at our budgets converges to FCI from above:** near equilibrium the CIPSI extrapolation reproduces FCI to ~4 mHa ($R^2=0.999$), and even at the strongest correlation ($R=2.4 \text{ \AA}$, 468 determinants) $E_{\text{var}}+\text{PT2}$ sits +10.9 mHa above FCI with the shrinking $E_{\text{var}} \leftrightarrow E_{\text{var}}+\text{PT2}$ gap as the convergence certificate — while CCSD(T) sits 217 mHa BELOW FCI with no error signal at all. (Selected-CI here is classical on CPU; QSCI is its device-driven instantiation, with a quantum sampler as the determinant selector at scales where the selected space is large.) *[QBRAID-RUN: device-sampled selection at scale]*




Figure 2. Trust on a real oxide. As the CrO $^5\Pi$ bond stretches (stronger multireference character, left $\rightarrow$ right), in-active-space CCSD(T) becomes erratic and frequently non-convergent (open markers), a few to ~ 140 mHa vs exact CASCI, while selected-CI/QSCI stays a variational bound at or within the chemical-accuracy band (shaded; ≤ 2.8 mHa even at the strongest correlation). CCSD(T) is run on the identical embedded active-space Hamiltonian as CASCI (apples-to-apples).

Proxy validated against real measurement; cost scaling. The scaling ladder uses a determinant-space selected-CI proxy; we confirm it tracks the real circuit-sampled pipeline by executing measured QSCI (qml.sample, 3,000 shots $\times$ 160 circuits) at 16q and 20q — measured energies match the proxy on the trained generator (16q: 28.1 measured vs 26.6 proxy; 20q: 50.4 vs 49.3 mHa), extending the sampled pipeline past the prior 12q point. One honest wrinkle: at this small shot budget, diffuse random-circuit sampling collects more distinct determinants than the peaked generator and edges it on absolute energy (measured-random 24.0/46.7 vs trained 28.1/50.4 mHa) — a low-shot coverage effect, not a generator-quality reversal: the deterministic proxy and the larger-budget sweeps both favour the generator. Determinants for chemical accuracy grow $\sim 1.38\times$ per qubit vs $2.0\times$ for full FCI (log $R^2=0.99$) — a vanishing fraction (75% at 8q $\rightarrow$ 0.15% at 28q) but still exponential ($\sim 10^6$ dets at 40q+, where device-driven selection becomes valuable); we do not claim polynomial scaling.

6. Platform Use and Resourcing

qBraid provides classical (CPU/GPU) and quantum (QPU) credits. We use **NVIDIA H100/A100 (80 GB)** with CUDA-Q: tensornet-mps for the 24–40-qubit tier and cuStateVec for exact validation to ~ 32 qubits. One high-memory GPU suffices for MPS; 4–8 GPUs (NVLink) enable distributed circuit evaluation (pillar 4) and the >40 -qubit bonus attempt; QPU (IonQ/IBM) for 10–16-qubit validation. The concrete platform value: the ~ 16 TB exact statevector at 40 qubits collapses onto a single 80 GB GPU through CUDA-Q MPS — a $\sim 200\times$ memory reduction that is what makes the 40-qubit tier reachable at all.

Resource estimate. UCC excitations decompose to ≤ 2 -qubit gates; with pool compression a length-8 circuit is shallow (tens of two-qubit gates), within MPS reach near equilibrium. QSCI needs $\sim 10^3$ – 10^4 shots/circuit. We estimate ~ 1 – 2 weeks single-GPU wall-clock for the full 24 $\rightarrow$ 40q sweep plus the oxide runs; backend-agnostic, degrades gracefully. Concrete per-run depth/shot/wall-clock: **[QBRAID-RUN: from §5]**.

7. Limitations and Honest Scope

We state where the approach does and does not yet provide benefit. (i) **Measurement vs proxy:** the integrated GQE $\rightarrow$ QSCI loop is measured at 12q and validated against measured QSCI at 16q/20q; larger QSCI numbers use a hardware-independent determinant-space proxy — quantum-inspired at scale until executed with real sampling on qBraid. (ii) **No classical-beating claim yet:** at ≤ 28 qubits classical FCI/DMRG already solve these instances, so we demonstrate correctness and favourable scaling, not a head-to-head speed advantage; that regime is 40q+ strong correlation where DMRG bond dimension explodes (visible at $H_{24}/48q$, where DMRG is 6.83 mHa off CCSD(T)). (iii) **40q is operational, not converged:** on CPU the 40-qubit run converges through 39 mHa, above chemical accuracy; closing this is the GPU deliverable. (iv) **Transfer scope:** cross-size transfer is a 3-seed-mean trend that narrows into noise beyond $\sim 28q$; cross-chemistry transfer wins clearly on 3 of 6 oxide targets (BeO a within-noise tie) and fails on the heaviest (SnO); target-specific MP2 beats the transferred prior. We claim a zero-cost transferable prior, not a universal or MP2-beating one. (v) **Strong-correlation regime:** where single-reference CCSD(T) collapses (§5: H_{10} and real CrO), the determinant-subspace method stays a rigorous variational bound with an EN-PT2 estimate tightening it from above, but reaching strict accuracy at the strongest correlation needs the large-subspace (device-sampled) regime — motivating the quantum approach. A clear-eyed reading: MATGEN-Q is a working, reproducible pipeline whose scaling mechanisms are demonstrated and whose decisive large-scale quantum-vs-classical comparison is the executed GPU work this phase enables.

8. Conclusion and Reproducibility

MATGEN-Q is a working two-stage GQE whose tensor-network and QSCI tiers, plus MP2 operator-pool compression, target the 40-qubit regime on a single GPU, with every claim third-party reproducible. A one-command driver (*reproduce.py*) and a README with a “Launch on qBraid” button let judges re-run each headline result without modification.

References

- [1] K. Nakaji, et al. “The generative quantum eigensolver (GQE) and its application for ground state search.” arXiv:2401.09253 (2024). [AIST]
- [2] N. P. D. Sawaya, et al. “HamLib: A library of Hamiltonians for benchmarking quantum algorithms and hardware.” Quantum 8, 1559 (2024).
- [3] S. Minami, K. Nakaji, et al. “Generative quantum combinatorial optimization ... conditional generative quantum eigensolver.” Digital Discovery 4(8) (2025).
- [4] K. Kanno, et al. “Quantum-selected configuration interaction ... subspaces selected by quantum computers.” arXiv:2302.11320 (2023).
- [5] NVIDIA Corporation. CUDA-Q and cuQuantum SDK (cuStateVec, tensornet / tensornet-mps backends).
- [6] J. Tilly, et al. “The Variational Quantum Eigensolver: A review of methods and best practices.” Physics Reports 986 (2022).
- [7] C. Fare, et al. “A multi-fidelity machine learning approach to high-throughput materials screening.” npj Computational Materials 8, 257 (2022).
- [8] T. D. Kharazi, et al. “Quantum Simulations for Extreme Ultraviolet Photolithography.” arXiv:2602.20234 (2026). [Xanadu & Mitsubishi Chemical]
- [9] R. Kemmoku, Q. Gao, S. Kanno, et al. “Generative Circuit Design for Quantum-Selected Configuration Interaction.” arXiv:2604.09756 (2026). [Mitsubishi Chemical]