

Changing Views of the Pathophysiology of Parkinsonism **CME**

Thomas Wichmann, MD*

Department of Neurology/School of Medicine and Yerkes National Primate Research Center, Emory University, Atlanta, Georgia, USA

ABSTRACT: Studies of the pathophysiology of parkinsonism (specifically akinesia and bradykinesia) have a long history and primarily model the consequences of dopamine loss in the basal ganglia on the function of the basal ganglia/thalamocortical circuit(s). Changes of firing rates of individual nodes within these circuits were originally considered central to parkinsonism. However, this view has now given way to the belief that changes in firing patterns within the basal ganglia and related nuclei are more important, including the emergence of burst discharges, greater synchrony of firing between neighboring neurons, oscillatory activity patterns, and the excessive

coupling of oscillatory activities at different frequencies. Primarily focusing on studies obtained in nonhuman primates and human patients with Parkinson's disease, this review summarizes the current state of this field and highlights several emerging areas of research, including studies of the impact of the heterogeneity of external pallidal neurons on parkinsonism, the importance of extrastriatal dopamine loss, parkinsonism-associated synaptic and morphologic plasticity, and the potential role(s) of the cerebellum and brainstem in the motor dysfunction of Parkinson's disease. © 2019 International Parkinson and Movement Disorder Society

The term “parkinsonism” encompasses the classical motor signs of Parkinson's disease (PD), that is, the combination of bradykinesia, akinesia, tremor, and rigidity (as well as gait/balance problems). These signs and symptoms are assumed to be triggered by the loss of dopaminergic neurons in the substantia nigra pars compacta (SNc). Discussions of the pathophysiology of parkinsonism are therefore mostly discussions of the immediate and chronic brain circuit effects of the degeneration of dopaminergic neurons in the SNc.

In the following paragraphs, a basic understanding of the anatomical structure of the basal ganglia and related

brain areas (see Fig. 1) is assumed. We will first discuss past and current findings pertaining to a brain circuit-level understanding of the pathophysiology of parkinsonism, coming from experiments in parkinsonian animals and patients with PD. This will be followed by a discussion of recent conceptual changes that are beginning to shape our knowledge in this field. Most of the information presented here will focus on results obtained from studies in non-human primates and patients with PD.

Parkinsonism as Dysfunction of the Basal Ganglia Thalamocortical Network

Most studies of brain dysfunction in parkinsonism assume that the primary brain activity changes that account for the parkinsonian state occur within the basal ganglia/thalamocortical circuitry (Fig. 1). The fact that motor impairments dominate early stages of PD is understandable considering that dopamine loss in (early) PD preferentially affects the dopamine supply to the *motor* circuit of the basal ganglia,¹ which is spatially segregated from nonmotor circuits within the basal ganglia/thalamocortical network of connections.² This section summarizes the current knowledge within this framework.

*Correspondence to: Dr. Thomas Wichmann, Department of Neurology/School of Medicine and Yerkes National Primate Research Center, Emory University, 954 Gatewood Road Northeast, Atlanta, GA 30329, USA; E-mail: twichma@emory.edu

Funding agencies: The writing of this article was supported by NIH/NINDS grant P50-NS098685 (Udall Center grant), and by a grant from the NIH Office of Research Infrastructure Programs OD P51-OD011132 to the Yerkes National Primate Research Center.

Relevant conflicts of interest/financial disclosures: Nothing to report.

Full financial disclosures and author roles may be found in the online version of this article.

Received: 9 April 2019; **Revised:** 15 May 2019; **Accepted:** 20 May 2019

Published online 19 June 2019 in Wiley Online Library (wileyonlinelibrary.com). DOI: 10.1002/mds.27741




FIG. 1. Corticosubcortical motor circuits involved in the pathophysiology of parkinsonism. Dark arrows indicate inhibitory connections; gray arrows indicate excitatory connections. CM/Pf, centromedian and parafascicular nuclei of the thalamus; Cereb. Cortex, cerebellar cortex; CMA, cingulate motor area; DCN, deep cerebellar nuclei; DP, direct pathway; D1, D1-like dopamine receptor subtype; D2, D2-like dopamine receptor subtype; HP, hyperdirect pathway; IP, indirect pathway; MC, motor cortex; PMC, premotor cortex; PN, pontine nuclei; SMA, supplementary motor area; VA/VL, ventral anterior and ventral lateral nuclei of the thalamus. Figure published in a previous work,²⁸⁵ here shown with permission. [Color figure can be viewed at wileyonlinelibrary.com]

Changes of Firing Rates

Thirty years ago, the results of anatomical, metabolic, and electrophysiologic studies were summarized in an influential model of global activity changes in the basal ganglia nuclei that described the emergence of parkinsonism and chorea as opposite ends of a spectrum of activity changes in these nuclei.^{3,4} In this model, the lack of striatal dopamine in the parkinsonian state increases striatal inhibition of neurons in the external pallidal segment (GPe), which would then lead to disinhibition of the glutamatergic STN, and subsequent increased driving of basal ganglia output neurons in the internal pallidal segment (GPi) and the substantia nigra pars reticulata (SNr). In parallel, striatal dopamine loss was also thought to reduce direct striatal inhibition of GPi and SNr. In combination, these changes result in *increased* tonic firing in the GABAergic output neurons in the GPi and SNr, leading to downstream inhibition of thalamocortical projection neurons and reduced activation of dependent areas in the frontal cerebral cortex, triggering the behavioral manifestations of parkinsonism. Events of inverse polarity were predicted to underlie chorea and other hyperkinetic disorders in which *reduced* output of the basal ganglia would lead to release of thalamocortical neurons from tonic inhibition and the emergence of involuntary movements.

Support for this “rate” model of movement disorders came from studies of the effects of inactivation (lesions) of the STN and GPi in parkinsonian animals. By reducing basal ganglia output, this was thought to “normalize” the (rate-model) disturbance underlying parkinsonism. As predicted, these interventions improved parkinsonism

in parkinsonian animals and in patients with PD.⁵⁻²⁹ Recent studies in rodents have further emphasized the importance of rate changes in basal ganglia output for the regulation of movements, by showing that optogenetic or chemogenetic activation of the striatal neurons that give rise to projections to the GPe reduces movement in normal animals.^{30,31}

However, the model would also predict that interventions that lower GPi/SNr output below its normal range would produce involuntary movements, but this is not the case. In fact, lesions of the GPi are used to *treat* hyperkinetic states such as chorea.^{32,33} Furthermore, according to the rate model, inactivation of the thalamic regions that receive basal ganglia input would be expected to worsen parkinsonism, but this is also not observed in humans or monkeys.^{34,35} In addition, recent optogenetic rodent studies have cast doubt on the rate-model notion that parkinsonism arises from an imbalance between the (antikinetic) actions of striatal neurons that project to the GPe and the (prokinetic) actions of neurons that project to the GPi. These studies emphasize that movement initiation and execution is accompanied by combined activation of *both* populations of striatal neurons,³⁶⁻³⁸ inconsistent with the rate model. These and other discrepancies suggest that rate changes in the basal ganglia/thalamocortical circuits may not be important for the pathophysiology of movement disorders than originally thought.

Altered Firing Patterns and Oscillations

Even the earliest reports documenting neuronal activity changes in MPTP-treated parkinsonian monkeys showed that there are prominent changes in firing patterns of basal ganglia neurons.³⁹ In the 1990s, these pattern changes quickly evolved from being regarded as a curiosity to being considered to be a central component of the pathophysiology of parkinsonism. Since then, many researchers have emphasized the role of increased oscillatory fluctuations of firing rates of neurons in the basal ganglia, thalamus, and cortex in the low beta-range of frequencies in humans and in slightly different (neighboring) frequency ranges in animals.⁴⁰⁻⁴⁵ Oscillatory and nonoscillatory bursts of neuronal spiking are also increased throughout the basal ganglia, the ventral motor thalamus, and frontal motor areas of the cerebral cortex in the parkinsonian state.⁴⁶

A related important change of neuronal activities in parkinsonism is a breakdown of the normal separation of the firing of individual neurons in the basal ganglia. The resulting neuronal synchrony has been directly documented in single-cell recordings with multiple electrodes^{44,47,48} and is also implied by the finding of increased amplitudes of local field potentials (LFPs; summed electrical potentials at a brain location) in the beta-band range of frequencies (10–30 Hz) in the basal ganglia and cortex, in recordings from parkinsonian animals,⁴⁹⁻⁵⁷ and from PD patients in whom implanted DBS electrodes were used for

recording purposes.^{44,58-65} The increased beta-band activity is associated with a reduction in gamma-band oscillatory power.^{44,65} In recent studies, it has been emphasized that an increase in the number and duration of beta-bursts in cortex or STN may be even more predictive of parkinsonism than the absolute level of beta-band activity.⁶⁶⁻⁶⁸ The finding of increased low-frequency synchronization not only applies to the individual nodes of the basal ganglia/thalamocortical motor circuit, but similarly affects interactions *between* components of these circuits, as documented, for example, by the finding of substantial cortico-subthalamic coherence in parkinsonian animals (e.g., see a previous work⁵¹) or in patients with PD (e.g., see previous works^{69,70}).

Shifts in the spectral composition of cortical or basal ganglia LFP signals occur also in conjunction with levodopa-induced dyskinesias (LIDs). Recent reports have focused on narrow-band gamma oscillations (60–85 Hz) in the cerebral cortex and the STN during episodes of dyskinesia,⁷³ similar to the narrow-band ~70-Hz oscillations described to occur in L-dopa-treated parkinsonian patients at rest or during voluntary movement,⁷⁴ and, interestingly, to oscillations found in electrocorticography (ECoG) signals in patients with isolated dystonia (described in a dystonia-1 [DYT1]-negative and a DYT1-positive patient in a previous work⁷⁵). This narrow-band gamma-band activity differs from the broad-band cortical gamma-band oscillations in these brain regions that accompany physiological movements, where oscillations in the 75- to 100-Hz range are associated with movement onset and oscillations in the 30- to 50-Hz range are present throughout ongoing movements.⁷⁶

Overall, studies of spectral changes involving single-cell or LFP signals have demonstrated that most elements of the basal ganglia/thalamocortical circuitry are engaged in synchronized low-frequency oscillations in the parkinsonian state. This may work to the detriment of normal processing in the basal ganglia/thalamocortical circuits, which may depend on transient focal synchronization of activity at gamma frequency bands and which is suppressed or temporally/spatially misaligned in the parkinsonian state where beta oscillatory activities prevail.

An interesting recent series of studies has directly focused on these interactions between the beta- and gamma-band oscillations. In studies of ECoG and EEG signals from PD patients,⁷⁷ the amplitude of gamma-band oscillations was found to be abnormally coupled to the phase of ongoing beta-band oscillations in a process called phase-amplitude coupling (PAC). Together with beta-band power, and corticocortical and basal ganglia/cortical coherence, cortical PAC was associated with bradykinesia and rigidity⁷⁸ and lowered by therapeutically beneficial DBS in the STN or GPi.⁷⁹⁻⁸¹ As is true for other oscillatory activities (see above), increased *cortical* PAC is not specific for PD, but also occurs in patients with (isolated) dystonia⁸² or essential tremor.⁸³

Studies of *subcortical* PAC in parkinsonian patients have focused on LFP recordings from the STN and GPi. As in cortical recordings, PAC between beta- and gamma-bands is a sensitive marker of parkinsonism in STN LFPs⁸⁴⁻⁸⁶ and is also found in cases of isolated dystonia.⁵⁹ Interestingly, it is not clear whether the situation is similar in the GPi. Thus, in a study of LFP PAC in GPi, Tsiokos and colleagues reported an inverse relationship between PAC and parkinsonian signs, suggesting that a high level of PAC could be a normal feature of GPi physiology.⁸⁷

The development of parkinsonism in chronically MPTP-treated monkeys is also associated with PAC in LFPs recorded in the basal ganglia^{49,88,89} or frontal cortex,^{49,90} in this case involving the phase of theta-beta-range frequencies and the amplitude of gamma-range frequencies (150–400 Hz). The studies also showed that basal ganglia or cortical PAC is highly dependent on the state of vigilance of the animal.^{49,90} As in patients with PD, beta-gamma PAC in MPTP-treated animals findings appear to be more predictive of parkinsonism than changes of spectral power in specific frequency ranges.^{88,89}

The Question of Causality

Most of the aforementioned abnormalities of neuronal activity patterns have been identified in multiple basal ganglia nuclei, in the thalamus, and are beginning to be identified in the cerebral cortex, suggesting that they may affect processing throughout the entire cortico-basal ganglia/thalamocortical circuits. If true, this may reduce the ability of the basal ganglia/thalamocortical system to normally process information and could interfere or eliminate major functions of the affected basal ganglia, such as their ability to contribute to motor learning, or to facilitate movement. It is important to emphasize again that removal of basal ganglia processing by creation of lesions in the basal ganglia output nuclei reduces rather than increases parkinsonism,^{91,92} suggesting that the oscillatory and abnormally synchronized activity in the cortico-basal ganglia/thalamocortical circuits does not simply disrupt processing of information in the basal ganglia, but that the abnormal basal ganglia activity actively disrupts information processing downstream from the basal ganglia. Specific constellations of rate and pattern changes in the basal ganglia/thalamocortical circuits may affect cortical processing in a manner that triggers bradykinesia or akinesia in the parkinsonian state. Other diseases may be associated with separate (but overlapping) combinations of abnormalities that bias cortical processing in other ways, manifesting, for example, as changes in muscle tone and posture as observed in dystonia.

It remains an open question which of the observed abnormalities in basal ganglia activity are causal for parkinsonism.^{43,51,93,94} This was already mentioned in the discussion of firing rates, and the same uncertainty applies to the synchronized oscillatory activities that

were described above. In fact, several published experiments did not find a strong correlation between spectral changes and the degree of (early) parkinsonism. For example, the described increase in low-frequency synchronization is not always found in animals with behavioral parkinsonism,^{89,90} or in humans with PD (e.g., see a previous work⁹⁵). Some of the observed changes may thus be inconstant (see discussion of beta bursts above) or simply represent epiphenomena of a yet-to-be-discovered underlying disruptive process. Of course, the lack of a causal relationship of spectral (or other) changes to parkinsonism does not diminish their potential value as biomarkers for the severity of parkinsonism if they *can* be detected in a given case.

Beyond Akinesia and Bradykinesia

The pathophysiological considerations mentioned in the previous paragraphs relate primarily to the emergence of akinesia and bradykinesia. Our knowledge concerning the pathophysiology of other principal signs of parkinsonism (rigidity, tremor, and gait impairments) is more limited. A brief summary of the major themes of research pertaining to the emergence of rigidity and gait impairments in PD patients is presented here. Information on the pathophysiology of tremor is found in the next section.

Most studies of the mechanisms underlying *rigidity* come from the human literature. Many studies focus on alterations in spinal and brainstem reflex mechanisms.⁹⁶ A role of the stretch reflex system⁹⁷⁻¹⁰⁰ and its facilitation in PD patients (see earlier works,^{101,102} but see another work¹⁰³) was originally favored, whereas later studies emphasized a role of changes of the latency and amplitude of long-latency reflexes (see earlier works,^{104,105} but see another work¹⁰⁶). Changes in the basal ganglia are likely also involved, given that the severity of rigidity correlates with the degree of striatal dopamine loss,^{107,108} and with the power or stability of beta-band oscillations in the STN.^{109,110} The involvement of the basal ganglia and related area of the ventral motor thalamus is also suggested by the fact that surgical interventions targeting these systems reduce rigidity.¹¹¹⁻¹¹⁴

Although the polarity of changes is not entirely consistent between studies, imaging experiments have shown that frontal and parietal cortical activity are altered in rigid parkinsonian patients.^{115,116} For example, in studies of default-mode network expression in resting-state functional MRI (fMRI) experiments, the connectivity of posterior cortical network components was found to be reduced whereas that of frontal cortical components of the network was pathologically increased.^{117,118} An interpretational issue with these and similar studies is, of course, that rigidity does not develop in isolation. It is difficult to assign brain activity abnormalities specifically to this parkinsonian sign while others, such as bradykinesia or cognitive impairments, also emerge.

The pathophysiological basis of *gait impairments* in parkinsonian patients is complex and likely multifactorial. A specific parkinsonian impairment that has garnered interest in recent years is freezing of gait (FOG). The phrase FOG is applied to sudden and unpredictable blocks in stepping, specifically occurring when the patient initiates gait or turns, and under certain situations, for instance when walking through narrow passages. The presence of FOG is disabling and significantly affects the quality of life.¹¹⁹ Pathophysiologically relevant clinical observations are that FOG typically affects patients with long disease duration.¹¹⁹⁻¹²² is at least partially corrected by L-dopa in many patients,^{119,121} and is typically associated with changes in cognition, such as altered attention,¹²³ cognitive dysfunction,¹²⁴⁻¹²⁷ or psychosis,¹²⁸ suggesting cortical involvement. In experiments in which patients were engaged in cognitive activities while walking, those suffering from FOG typically showed worse walking performance than those without FOG (“dual task interference”),¹²⁹ supporting the notion that FOG tends to arise in susceptible individuals when cognitive resources that are needed for maintenance of gait under parkinsonian conditions have to be shared with other activities. Volumetric and fMRI studies have demonstrated that specific frontal and parietal cortical regions are atrophic in patients with FOG,¹³⁰⁻¹³⁵ and EEG studies have suggest that patients who suffer from FOG have high beta-band oscillatory activity in frontal regions EEG,¹³⁶ specifically increased during conditions that demand increased attention.¹³⁷

Subcortical gait networks, including the striatum, GPi, thalamus, the mesencephalic locomotor region, and the cerebellum, are also involved in FOG.^{126,132,133,136,138} Whereas STN- or GPi-DBS reduces FOG in some patients, others suffer from stimulation-induced freezing,¹³⁹ potentially depending on the choice of stimulation parameters.¹⁴⁰ In an interesting recording study comparing STN-DBS implanted PD patients with and without freezing, Storz and colleagues discovered that pedaling movements that are (by themselves) not specifically affected by freezing bring out increased power in the low beta band in LFPs in the STN.¹⁴¹

It is believed that freezing can also occur in parkinsonian rodents¹⁴²⁻¹⁴⁴ and monkeys.¹⁴⁵ The few available studies point to the involvement of components of the mesencephalic locomotor region/pedunculopontine nucleus (PPN).¹⁴² Recordings in MPTP-treated, parkinsonian monkeys indicated a greater tendency of mesencephalic locomotor region neurons to discharge in bursts, which could be related to FOG.¹⁴⁶

Neurochemically, FOG may be associated with changes in the norepinephrine¹⁴⁷ and/or cholinergic systems.¹⁴⁸ The former is supported by the finding that the catecholaminergic locus coeruleus appears to be damaged in parkinsonian monkeys that show FOG (compared to those without FOG),¹⁴⁹ by neuromelanin MRI findings in

parkinsonian patients with and without FOG,¹⁵⁰ and by the (empirically determined) effectiveness of medications that alters catecholamine release or levels in the brain, such as atomoxetine,^{151,152} or L-threo-DOPS.¹⁵³⁻¹⁵⁶ Cholinergic dysfunction is implicated by imaging studies that show cholinergic loss in cortical regions,^{148,157} and in the striatum¹⁵⁷ of patients with FOG. Studies in rodents and monkeys have supported the notion that severe gait abnormalities may arise in situations where both the dopaminergic nigrostriatal system and cholinergic cells in the pedunculopontine nucleus (area) are destroyed.^{144,158,159}

Evolving Topics

The ever-increasing pace of discovery in basal ganglia research has led to many exciting evolving topics in studies of the pathophysiology of parkinsonism. Only some examples of these can be reviewed here.

Heterogeneity of GPe Neurons

In traditional schemes of the basal ganglia, the GPe is depicted as a homogeneous structure in which neurons relay input from the striatum to STN, GPi, and SNr (Fig. 1). An important recent insight has changed this concept by showing that GPe neurons are, in fact, heterogeneous, belonging to at least two major classes, that is, “arkypallidal” neurons that project to the striatum and “prototypical” neurons, which project to the STN, GPi, and SNr. In rodents, molecular differences among GPe neurons have been used to genetically target and manipulate these neuron subtypes using optogenetic and pharmacogenetic techniques.¹⁶⁰⁻¹⁶⁴ It is not clear whether and how these protein expression patterns translate to the primate GPe.

Prototypical GPe neurons powerfully influence the frequency and firing patterns of STN neurons¹⁶⁵⁻¹⁶⁸ and, perhaps to a lesser degree, of GPi neurons.¹⁶⁹⁻¹⁷¹ Global activation of GPe activity in parkinsonian monkeys¹⁷² induces pathological STN neuron activity and can ameliorate parkinsonian motor signs, but may produce dyskinetic movements.^{169,173,174} Arkypallidal neurons, on the other hand, make abundant connections and strongly inhibit direct and indirect pathway striatal projection neurons and interneurons.^{160,162,163,175-177} Although this may play a role in normal basal ganglia function(s), the role of these cells in parkinsonism is not known.^{175,178} Recent optogenetic and pharmacogenetic studies in parkinsonian rodents have suggested that selective activation of arkypallidal neurons may be detrimental^{160,179} whereas selective activation of prototypical neurons can produce antiparkinsonian effects.¹⁶⁴ It remains to be seen whether similarly specific manipulations can lead to antiparkinsonian effects in nonhuman

primates or patients with PD, without the risk of inducing dyskinesias.

Role of Extrastriatal Dopamine Loss

There is ample evidence that dopaminergic midbrain neurons innervate not only the striatum, but also extrastriatal sites in basal ganglia, thalamus, and cortex.¹⁸⁰ The loss of dopamine at these sites in is also well documented to occur in parkinsonian animals¹⁸¹ and in PD patients.¹⁸²⁻¹⁸⁴

In the primate GPe, D2-like receptors (D2LRs) are expressed on GABAergic striatopallidal inputs. Local activation of these receptors by injections of small amounts of D2LR agonists increases the firing of GPe neurons in monkeys, likely because of reduction of the GABAergic inhibition.¹⁸⁵ In the monkey STN, D1 and D2 receptors are found on preterminal axons and glutamatergic and GABAergic terminals and on postsynaptic D5 receptors.¹⁸⁶ Locally administered agonists at D1LRs lower STN firing rates, but increase bursting in these animals,¹⁸⁶ perhaps through a combination of effects on presynaptic D1- and postsynaptic D5-receptors, respectively.

In GPi and SNr, D1LRs are expressed on axons and terminals of the striatopallidal and striatonigral pathways. D2LRs are also found, on (presumptive) glutamatergic inputs. Local administration of D1LR or D2LR agonists into the GPi and SNr reduces firing in these nuclei.¹⁸⁵ The comparatively maintained pallidal and nigral dopamine levels in early parkinsonism may thus compensate for the striatal dopamine loss in early PD, helping to maintain GPi/SNr firing rates at levels close to normal.¹⁸⁷ In the fully parkinsonian state, the fraction of membrane-bound (active) D1LRs (among all D1LRs) at these sites increases.¹⁸⁸

The basal ganglia-receiving portion of the ventral motor thalamus is also innervated by dopaminergic fibers,¹⁸⁹ arising from several dopaminergic cell groups in the midbrain.¹⁹⁰⁻¹⁹² There appears to be a substantial loss of the thalamic dopaminergic innervation in MPTP-induced parkinsonism in monkeys,¹⁸⁹ which could be related to the emergence of tremor (see also below).^{193,194}

Activation of extrastriatal dopamine receptors may contribute to the antiparkinsonian efficacy of systemically administered dopamine receptor agonists, specifically to that of D2LR agonists such as ropinirole or pramipexole. Conceivably, activation of extrastriatal D2LRs by these agents could lower firing irregularities through actions in the STN, while reducing the overall basal ganglia output through actions in SNr and GPi.¹⁹⁵⁻¹⁹⁷

The expression of extrastriatal dopamine receptors is obviously not restricted to the motor areas of the basal ganglia, thalamus, and cortex. Activation of dopamine receptors at nonmotor sites may contribute to some of the frequent side effects of D2LR agonist therapies. For example, specific associations to changes in extrastriatal

dopamine levels (or receptor occupancy) were recently documented in the context of impulse control problems,^{198,199} depression,²⁰⁰ and dementia.²⁰⁰⁻²⁰²

Parkinsonism-Related Synaptic Plasticity

There is growing awareness of the fact that the effects of dopamine depletion in the basal ganglia/thalamocortical motor circuit does not simply lead to stable changes of neuronal activity or excitability in the affected circuitry. Instead, parkinsonism is associated with considerable morphological and functional plasticity at synapses within the basal ganglia, thalamus, and cortex. Thus, it had been known since the late 1980s that dopamine depletion causes loss or remodeling of glutamatergic synapses within the striatum.²⁰³⁻²⁰⁸ Whereas some of these changes affect cortical inputs to the striatum,²⁰⁹ thalamostriatal inputs are also involved, likely because of the overall loss of neurons in the intralaminar nuclei.²¹⁰⁻²¹² The loss of glutamatergic inputs to the striatum is associated with a coarsening of the remaining synapses and the formation of multisynaptic neuronal terminations.²⁰⁹

More recently, glutamatergic plasticity was also shown to occur in the STN in parkinsonian rodents, nonhuman primates, and patients.^{31,213,214} The remodeling of glutamatergic transmission in the STN involves N-methyl-D-aspartate receptor activation and appears to trigger subsequent remodeling of the GABAergic synapses of the pallidosubthalamic projection.^{31,215,216} The emergence of these changes may set the stage for the presence of dopamine-treatment-resistant features of parkinsonism, triggered by loss of dopamine loss in the striatum.³¹ The specific physiological causes and consequences of these morphological changes are a very active area of research, as is the question whether these changes represent homeostatic or maladaptive plasticity.

Finally, glutamatergic plasticity also occurs in the cerebral cortex.^{217,218} In the motor cortex, there is a prominent dropout of the thalamic (glutamatergic) innervation of layer 5, which could be caused by the loss of neurons in the thalamic intralaminar nucleus and their cortical projections.²¹⁸ The behavioral effects of these (substantial) changes remain uncertain at this time.

Role(s) of the Cerebellum

A role of the cerebellum in the expression of parkinsonism has long been suspected,²¹⁹ and several studies have indeed identified morphological or biochemical changes in the cerebellum, ranging from low levels of alpha-synuclein aggregates^{220,221} in the deep cerebellar nuclei, to a reduction of dopamine D1 and D3 receptor expression,²²² or reduced norepinephrine levels.²²³ However, many researchers interpret the involvement of the cerebellum and its outflow pathways in parkinsonism as an indirect process, triggered through primary pathology in the striatum or midbrain.²²⁴⁻²²⁶ This view

has recently received additional attention because of the identification of anatomical pathways that link the basal ganglia to the cerebellum, so that dopamine loss in the basal ganglia could lead to changes in the function of cerebellar circuits (see Fig. 1 and previous publications²²⁷⁻²³⁰).

According to imaging studies, the cerebellar hemispheres are overactive in PD patients.²³¹⁻²³³ This may indicate compensatory effects of the cerebellum for the “lost” basal ganglia functions,^{234,235} although it remains uncertain what level or type of “compensation” could specifically be provided in this case.

Dysfunction of the cerebellum and/or of its output pathways has likely a more direct role in the generation of parkinsonian tremor. This is supported by the fact that DBS of the thalamic cerebellar-receiving territory effectively treats parkinsonian tremor (e.g., see previous works^{236,237}), while reducing the activity of the ipsilateral cerebellum, as measured by PET.²³⁸ The pathophysiology of parkinsonian tremor thus likely differs from that of bradykinesia or akinesia (which do not respond to such interventions). Imaging and modeling studies have suggested that dopaminergic denervation of the cerebellar receiving territory of the thalamus may contribute to tremor, and that dopaminergic drugs could act directly in the thalamus in such patients.¹⁹³ STN-DBS also affects cerebellar function,^{239,240} perhaps through the pathways shown in Figure 1, which may account for some of its antiparkinsonian effects,²⁴⁰ specifically its effects on parkinsonian tremor.²⁴¹ Although there is mounting evidence that parkinsonian tremor involves the cerebello-thalamocortical system in some way, details of the pathophysiology of parkinsonian tremor remain entirely speculative. For example, the fundamental question regarding the circuit origin of the oscillatory activity that eventually manifests as tremor remains unanswered.

Subtle changes in sensory processing (such as altered responses to sensory uncertainty) in PD may attest to potential additional roles of the cerebellum in other parkinsonian symptoms, such as bradykinesia.²⁴² MRI studies demonstrating an attenuation of the modulation of corticocortical interactions by cerebellar inputs in the parkinsonian state²⁴³ may further support this concept. Finally, the cerebellum has also been implicated in the development of LIDs.²⁴⁴⁻²⁴⁷

Role of Brainstem Nuclei

Traditional pathophysiology schemes strongly emphasize the role of the transthalamic route of transmission of basal ganglia output to the cerebral cortex as the main route for mediating parkinsonism (Fig. 1). However, parkinsonism is associated with relatively minor changes in firing in the basal ganglia receiving portion of the thalamus.²⁴⁸⁻²⁵² Whereas traditional models predict that interventions aimed at the basal ganglia-receiving portion of the ventral motor thalamus “should” impact

bradykinesia or akinesia, these predictions have not been confirmed in parkinsonian animals or human patients with the disease (e.g., see an earlier work).³⁵ These findings suggest that the branching GPi projections to brainstem nuclei, such as the PPN,^{253,254} may be more important for the development of bradykinesia or akinesia than traditionally assumed. PPN dysfunction is usually blamed for dopamine-insensitive gait and balance problems in PD (e.g., see previous works^{159,255}), but experiments in primates and rodents have suggested that lesions of the PPN (area) can elicit at least transient hypokinesia, and that stimulation of this site in parkinsonian animals appears to have beneficial effects.²⁵⁶⁻²⁶⁸ In distinction from the parkinsonism-associated oscillations in the basal ganglia/thalamocortical circuits, parkinsonism is accompanied by increased alpha-band oscillations in the PPN in PD patients, modulated by movement and gait performance,²⁶⁹⁻²⁷¹ and emphasized in the ON-state.²⁷²⁻²⁷⁴ Interestingly, beta-band synchronization, which is proparkinsonian in the basal ganglia and cortex, appears to be associated with reversal of parkinsonism in the PPN.²⁷⁵

It deserves mentioning that many of the recording and interventional studies targeting the PPN area are difficult to interpret because of the fact that this nucleus and the surrounding area are anatomically and functionally highly heterogeneous. To better understand the involvement of this nucleus in parkinsonian pathophysiology, it will be especially important to explore the anatomical connectivity of the specific PPN neurons that are targeted by basal ganglia output.

Conclusion

Studies of the consequences of dopamine loss in the basal ganglia have a long history, and substantial progress has been made. The studies reviewed here have identified some of the major abnormalities by which PD manifests itself at the single-cell or network level within the basal ganglia/thalamocortical circuitry. It is clear, however, that much work needs to be done to better define the anatomical network(s) that are affected by dopamine loss and account for the clinical manifestations of PD, and the mechanisms that bring about these changes.

Simplified models of the basal ganglia, and of the changes in activity that are caused by parkinsonism (such as the “rate model”), have directed our thinking of the pathophysiology of akinesia and bradykinesia. As a guide to therapy development, these traditional models have benefitted many thousands of patients with PD, by having a significant impact on the use of DBS approaches in the treatment of parkinsonism and by providing the backdrop for the development and testing of new medications. An important recent use of the knowledge gained from the studies mentioned in this review has been the generation of biomarkers that scale with the severity of parkinsonism

and can be used to regulate the strength or activity of DBS devices. Thus, studies in monkeys showed that pathologically increased cortical beta-band oscillatory activity can be used as control signal for intermittent “closed loop” regulation of pallidal stimulation in monkeys,²⁷⁶ and subsequent studies in parkinsonian patients demonstrated that beta oscillations²⁷⁷ or beta-burst durations⁶⁷ of signals recorded in the basal ganglia may serve the same purpose, allowing discontinuous DBS, focused on periods of time of significant parkinsonism. Recently, the strength of the dyskinesia-associated cortical gamma-band peaks was used as a signal to limit ongoing DBS,²⁷⁸ and cortical PAC has been discussed as a potentially useful additional control signal for the same purpose.⁷⁷

Developing knowledge of parkinsonism-associated activity changes in the basal ganglia and their output pathways goes hand in hand with developing more detailed and precise anatomic insights. Future experimental exploration of the pathophysiology of PD will therefore continue to focus on parallel lines of studies of functional abnormalities, and of the anatomical connections whose activity and morphology may explain them.

It is also remarkable that most of our knowledge of the pathophysiology of parkinsonism was acquired through studies of neuronal activity at rest—when, strictly speaking, neither bradykinesia nor akinesia are present. Although technically more complicated, it will be important to put increasing emphasis on the study of perimovement activities in parkinsonian subjects, as has already been achieved in some studies in rodents,²⁷⁹ nonhuman primates,^{280,281} and patients with PD (e.g., see previous works²⁸²⁻²⁸⁴).

Models such as the “rate” model are now clearly outdated, but new convincing and testable models that take into account the multitude of oscillatory and non-oscillatory changes in the basal ganglia, the anatomical loci and connections that were newly identified to be pathophysiologically relevant, or the effects of synaptic plasticity remain elusive. To develop such models and carry out experiments designed to identify those abnormalities that are causal rather than secondary continues to have very high priority. ■

Acknowledgment: The author thanks Adriana Galvan, PhD, for critically reading and discussing the content of the final version of the review.

References

1. Rinne JO. Nigral degeneration in Parkinson's disease. *Mov Disord* 1993;8(Suppl 1):S31-S35.
2. Alexander GE, DeLong MR, Strick PL. Parallel organization of functionally segregated circuits linking basal ganglia and cortex. *Annu Rev Neurosci* 1986;9:357-381.
3. Albin RL, Young AB, Penney JB. The functional anatomy of basal ganglia disorders. *Trends Neurosci* 1989;12:366-375.
4. DeLong MR. Primate models of movement disorders of basal ganglia origin. *Trends Neurosci* 1990;13:281-285.

5. Scoditti U, Rustichelli P, Calzetti S. Spontaneous hemiballism and disappearance of parkinsonism following contralateral lenticular lacunar infarct. *Ital J Neurol Sci* 1989;10:575-577.
6. Bergman H, Wichmann T, DeLong MR. Reversal of experimental parkinsonism by lesions of the subthalamic nucleus. *Science* 1990; 249:1436-1438.
7. Aziz TZ, Peggs D, Sambrook MA, Crossman AR. Lesion of the subthalamic nucleus for the alleviation of 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP)-induced parkinsonism in the primate. *Mov Disord* 1991;6:288-292.
8. Aziz TZ, Peggs D, Agarwal E, Sambrook MA, Crossman AR. Subthalamic nucleotomy alleviates parkinsonism in the 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP)-exposed primate. *Br J Neurosurg* 1992;6:575-582.
9. Guridi J, Herrero MT, Luquin R, Guillen J, Obeso JA. Subthalamotomy improves MPTP-induced parkinsonism in monkeys. *Stereotact Funct Neurosurg* 1994;62:98-102.
10. Wichmann T, Bergman H, DeLong MR. The primate subthalamic nucleus. III. Changes in motor behavior and neuronal activity in the internal pallidum induced by subthalamic inactivation in the MPTP model of parkinsonism. *J Neurophys* 1994;72:521-530.
11. Baunez C, Nieoullon A, Amalric M. In a rat model of parkinsonism, lesions of the subthalamic nucleus reverse increases of reaction time but induce a dramatic premature responding deficit. *J Neurosci* 1995;15:6531-6541.
12. Lozano AM, Lang AE, Galvez-Jimenez N, et al. Effect of GPi pallidotomy on motor function in Parkinson's disease. *Lancet* 1995; 346:1383-1387.
13. Guridi J, Herrero MT, Luquin MR, et al. Subthalamotomy in parkinsonian monkeys. Behavioural and biochemical analysis. *Brain* 1996;119(Pt 5):1717-1727.
14. Lang AE, Duff J, Saint-Cyr JA, et al. Posteroventral medial pallidotomy in Parkinson's disease. *J Neurol* 1999;246(Suppl 2): II28-II41.
15. Alvarez L, Macias R, Guridi J, et al. Dorsal subthalamotomy for Parkinson's disease. *Mov Disord* 2001;16:72-78.
16. Parkin S, Nandi D, Giladi N, et al. Lesioning the subthalamic nucleus in the treatment of Parkinson's disease. *Stereotact Funct Neurosurg* 2001;77:68-72.
17. Chen CC, Lee ST, Wu T, Chen CJ, Huang CC, Lu CS. Hemiballism after subthalamotomy in patients with Parkinson's disease: report of 2 cases. *Mov Disord* 2002;17:1367-1371.
18. Su PC, Tseng HM, Liou HH. Postural asymmetries following unilateral subthalamotomy for advanced Parkinson's disease. *Mov Disord* 2002;17:191-194.
19. Su PC, Tseng HM, Liu HM, Yen RF, Liou HH. Subthalamotomy for advanced Parkinson disease. *J Neurosurg* 2002;97:598-606.
20. Vilela F, da Silva DJ. Unilateral subthalamic nucleus lesioning: a safe and effective treatment for Parkinson's disease. *Arq Neuropsiquiatr* 2002;60:935-948.
21. Patel NK, Heywood P, O'Sullivan K, McCarter R, Love S, Gill SS. Unilateral subthalamotomy in the treatment of Parkinson's disease. *Brain* 2003;126(Pt 5):1136-1145.
22. Su PC, Tseng HM, Liu HM, Yen RF, Liou HH. Treatment of advanced Parkinson's disease by subthalamotomy: one-year results. *Mov Disord* 2003;18:531-538.
23. Alvarez L, Macias R, Lopez G, et al. Bilateral subthalamotomy in Parkinson's disease: initial and long-term response. *Brain* 2005;128 (Pt 3):570-583.
24. Cif L, Hariz M. Seventy years of pallidotomy for movement disorders. *Mov Disord* 2017;32:972-982.
25. Rodriguez-Rojas R, Carballo-Barreda M, Alvarez L, et al. Subthalamotomy for Parkinson's disease: clinical outcome and topography of lesions. *J Neurol Neurosurg Psychiatry* 2018;89:572-578.
26. Baron MS, Vitek JL, Turner RS, et al. Treatment of medically intractable Parkinson's disease with micro-electrode guided pallidotomy: one year follow-up [abstract]. *Mov Disord* 1994;112 (Suppl 1).
27. Baron MS, Vitek JL, Bakay RA, et al. Treatment of advanced Parkinson's disease by posterior GPi pallidotomy: 1-year results of a pilot study. *Ann Neurol* 1996;40:355-366.
28. Baron MS, Vitek JL, Bakay RA, et al. Treatment of advanced Parkinson's disease by unilateral posterior GPi pallidotomy: 4-year results of a pilot study. *Mov Disord* 2000;15:230-237.
29. Baron MS, Wichmann T, Ma D, DeLong MR. Effects of transient focal inactivation of the basal ganglia in parkinsonian primates. *J Neurosci* 2002;22:592-599.
30. Kravitz AV, Freeze BS, Parker PR, et al. Regulation of parkinsonian motor behaviours by optogenetic control of basal ganglia circuitry. *Nature* 2010;466:622-626.
31. Chu HY, McIver EL, Kovalski RF, Atherton JF, Bevan MD. Loss of hyperdirect pathway cortico-subthalamic inputs following degeneration of midbrain dopamine neurons. *Neuron* 2017;95: 1306-1318.e5.
32. Cubo E, Shannon KM, Penn RD, Kroin JS. Internal globus pallidotomy in dystonia secondary to Huntington's disease. *Mov Disord* 2000;15:1248-1251.
33. Lenders MW, Buschman HP, Vergouwen MD, Steur EN, Kölling P, Hariz M. Long term results of unilateral posteroventral pallidotomy for antipsychotic drug induced tardive dyskinesia. *J Neurol Neurosurg Psychiatry* 2005;76:1039.
34. Pan L, Dai JZ, Wang BJ, Xu WM, Zhou LF, Chen XR. Stereotactic Gamma thalamotomy for the treatment of parkinsonism. *Stereotact Funct Neurosurg* 1996;66(Suppl 1):329-332.
35. Duval C, Panisset M, Strafella AP, Sadikot AF. The impact of ventrolateral thalamotomy on tremor and voluntary motor behavior in patients with Parkinson's disease. *Exp Brain Res* 2006;170: 160-171.
36. Klaus A, Martins GJ, Paixao VB, Zhou P, Paninski L, Costa RM. The spatiotemporal organization of the striatum encodes action space. *Neuron* 2017;95:1171-1180.e7.
37. Vicente AM, Galvao-Ferreira P, Tecuapetla F, Costa RM. Direct and indirect dorsolateral striatum pathways reinforce different action strategies. *Curr Biol* 2016;26:R267-R269.
38. Tecuapetla F, Jin X, Lima SQ, Costa RM. Complementary contributions of striatal projection pathways to action initiation and execution. *Cell* 2016;166:703-715.
39. Miller WC, DeLong MR. Altered tonic activity of neurons in the globus pallidus and subthalamic nucleus in the primate MPTP model of parkinsonism. In: Carpenter MB, Jayaraman A, eds. *The Basal Ganglia II*. New York, NY: Plenum; 1987:415-427.
40. Bergman H, Wichmann T, Karmon B, DeLong MR. The primate subthalamic nucleus. II. Neuronal activity in the MPTP model of parkinsonism. *J Neurophys* 1994;72:507-520.
41. Rubin JE, McIntyre CC, Turner RS, Wichmann T. Basal ganglia activity patterns in parkinsonism and computational modeling of their downstream effects. *Eur J Neurosci* 2012;36:2213-2228.
42. Wichmann T, Dostrovsky JO. Pathological basal ganglia activity in movement disorders. *Neurosci* 2011;198:232-244.
43. Gatev P, Darbin O, Wichmann T. Oscillations in the basal ganglia under normal conditions and in movement disorders. *Mov Disord* 2006;21:1566-1577.
44. Hammond C, Bergman H, Brown P. Pathological synchronization in Parkinson's disease: networks, models and treatments. *Trends Neurosci* 2007;30:357-364.
45. Salvade A, D'Angelo V, Di Giovanni G, et al. Distinct roles of cortical and pallidal beta and gamma frequencies in hemiparkinsonian and dyskinetic rats. *Exp Neurol* 2016;275(Pt 1):199-208.
46. Wichmann T, Soares J. Neuronal firing before and after burst discharges in the monkey basal ganglia is predictably patterned in the normal state and altered in parkinsonism. *J Neurophys* 2006;95: 2120-2133.
47. Moran A, Bergman H, Israel Z, Bar-Gad I. Subthalamic nucleus functional organization revealed by parkinsonian neuronal oscillations and synchrony. *Brain* 2008;131(Pt 12):3395-3409.
48. Bergman H, Zaidel A, Rosin B, et al. Pathological synchrony of basal ganglia-cortical networks in the systemic MPTP primate model of Parkinson's disease. In: Steiner H, Tseng K, eds. *Handbook of*

- Basal Ganglia Structure and Function, 1st ed. Amsterdam: Academic (Elsevier); 2016:653-658.
49. Escobar Sanabria D, Johnson LA, Nebeck SD, et al. Parkinsonism and vigilance: alteration in neural oscillatory activity and phase-amplitude coupling in the basal ganglia and motor cortex. *J Neurophysiol* 2017;118:2654-2669.
 50. Zhang S, Connolly AT, Madden LR, Vitek JL, Johnson MD. High-resolution local field potentials measured with deep brain stimulation arrays. *J Neural Eng* 2018;15:046019.
 51. Devergnas A, Pittard D, Bliwise D, Wichmann T. Relationship between oscillatory activity in the cortico-basal ganglia network and parkinsonism in MPTP-treated monkeys. *Neurobiol Dis* 2014; 68:156-166.
 52. Wang J, Johnson LA, Jensen AL, et al. Network-wide oscillations in the parkinsonian state: alterations in neuronal activities occur in the premotor cortex in parkinsonian nonhuman primates. *J Neurophysiol* 2017;117:2242-2249.
 53. Connolly AT, Muralidharan A, Hendrix C, et al. Local field potential recordings in a non-human primate model of Parkinson's disease using the Activa PC + S neurostimulator. *J Neural Eng* 2015; 12:066012.
 54. Kammermeier S, Pittard D, Hamada I, Wichmann T. Effects of high-frequency stimulation of the internal pallidal segment on neuronal activity in the thalamus in parkinsonian monkeys. *J Neurophysiol* 2016;116:2869-2881.
 55. Goldberg JA, Rokni U, Boraud T, Vaadia E, Bergman H. Spike synchronization in the cortex/basal-ganglia networks of Parkinsonian primates reflects global dynamics of the local field potentials. *J Neurosci* 2004;24:6003-6010.
 56. Courtemanche R, Fujii N, Graybiel AM. Synchronous, focally modulated beta-band oscillations characterize local field potential activity in the striatum of awake behaving monkeys. *J Neurosci* 2003;23:11741-11752.
 57. Hendrix CM, Campbell BA, Tittle BJ, et al. Predictive encoding of motor behavior in the supplementary motor area is disrupted in parkinsonism. *J Neurophysiol* 2018;120:1247-1255.
 58. Mugge L, Krafcik B, Pontasch G, Alnemari A, Neimat J, Gaudin D. A review of biomarkers use in Parkinson with deep brain stimulation: a successful past promising a bright future. *World Neurosurg* 2019;123:197-207.
 59. Wang DD, de Hemptinne C, Miocinovic S, et al. Subthalamic local field potentials in Parkinson's disease and isolated dystonia: an evaluation of potential biomarkers. *Neurobiol Dis* 2016;89: 213-222.
 60. Priori A, Foffani G, Rossi L, Marceglia S. Adaptive deep brain stimulation (aDBS) controlled by local field potential oscillations. *Exp Neurol* 2013;245:77-86.
 61. Lempka SF, McIntyre CC. Theoretical analysis of the local field potential in deep brain stimulation applications. *PLoS One* 2013;8: e59839.
 62. Giannicola G, Rosa M, Servello D, et al. Subthalamic local field potentials after seven-year deep brain stimulation in Parkinson's disease. *Exp Neurol* 2012;237:312-317.
 63. Brown P, Williams D. Basal ganglia local field potential activity: character and functional significance in the human. *Clin Neurophysiol* 2005;116:2510-2519.
 64. Silberstein P, Kuhn AA, Kupsch A, et al. Patterning of globus pallidus local field potentials differs between Parkinson's disease and dystonia. *Brain* 2003;126(Pt 12):2597-2608.
 65. Brown P, Oliviero A, Mazzone P, Insola A, Tonali P, Di Lazzaro V. Dopamine dependency of oscillations between subthalamic nucleus and pallidum in Parkinson's disease. *J Neurosci* 2001;21:1033-1038.
 66. Tinkhauser G, Torrecillos F, Duclos Y, et al. Beta burst coupling across the motor circuit in Parkinson's disease. *Neurobiol Dis* 2018;117:217-225.
 67. Tinkhauser G, Pogossyan A, Little S, et al. The modulatory effect of adaptive deep brain stimulation on beta bursts in Parkinson's disease. *Brain* 2017;140:1053-1067.
 68. Tinkhauser G, Pogossyan A, Tan H, Herz DM, Kuhn AA, Brown P. Beta burst dynamics in Parkinson's disease OFF and ON dopaminergic medication. *Brain* 2017;140:2968-2981.
 69. Hirschmann J, Hartmann CJ, Butz M, et al. A direct relationship between oscillatory subthalamic nucleus-cortex coupling and rest tremor in Parkinson's disease. *Brain* 2013;136(Pt 12):3659-3670.
 70. Litvak V, Jha A, Eusebio A, et al. Resting oscillatory cortico-subthalamic connectivity in patients with Parkinson's disease. *Brain* 2011;134(Pt 2):359-374.
 71. Alonso-Frech F, Zamarbide I, Alegre M, et al. Slow oscillatory activity and levodopa-induced dyskinesias in Parkinson's disease. *Brain* 2006;129(Pt 7):1748-1757.
 72. Li X, Zhuang P, Hallett M, Zhang Y, Li J, Li Y. Subthalamic oscillatory activity in parkinsonian patients with off-period dystonia. *Acta Neurol Scand* 2016;134:327-338.
 73. Swann NC, de Hemptinne C, Miocinovic S, et al. Gamma oscillations in the hyperkinetic state detected with chronic human brain recordings in Parkinson's disease. *J Neurosci* 2016;36:6445-6458.
 74. Cassidy M, Mazzone P, Oliviero A, et al. Movement-related changes in synchronization in the human basal ganglia. *Brain* 2002;125(Pt 6):1235-1246.
 75. Miocinovic S, Swann NC, de Hemptinne C, Miller A, Ostrem JL, Starr PA. Cortical gamma oscillations in isolated dystonia. *Parkinsonism Relat Disord* 2018;49:104-105.
 76. Crone NE, Miglioretti DL, Gordon B, Lesser RP. Functional mapping of human sensorimotor cortex with electrocorticographic spectral analysis. II. Event-related synchronization in the gamma band. *Brain* 1998;121:2301-2315.
 77. Swann NC, de Hemptinne C, Aron AR, Ostrem JL, Knight RT, Starr PA. Elevated synchrony in Parkinson disease detected with electroencephalography. *Ann Neurol* 2015;78:742-750.
 78. Qasim SE, de Hemptinne C, Swann NC, Miocinovic S, Ostrem JL, Starr PA. Electrocorticography reveals beta desynchronization in the basal ganglia-cortical loop during rest tremor in Parkinson's disease. *Neurobiol Dis* 2016;86:177-186.
 79. de Hemptinne C, Ryapolova-Webb ES, Air EL, et al. Exaggerated phase-amplitude coupling in the primary motor cortex in Parkinson disease. *Proc Natl Acad Sci U S A* 2013;110:4780-4785.
 80. de Hemptinne C, Swann NC, Ostrem JL, et al. Therapeutic deep brain stimulation reduces cortical phase-amplitude coupling in Parkinson's disease. *Nat Neurosci* 2015;18:779-786.
 81. Malekmohammadi M, AuYong N, Ricks-Oddie J, Bordelon Y, Pouratian N. Pallidal deep brain stimulation modulates excessive cortical high β phase amplitude coupling in Parkinson disease. *Brain Stimul* 2018;11:607-617.
 82. Miocinovic S, de Hemptinne C, Qasim S, Ostrem JL, Starr PA. Patterns of cortical synchronization in isolated dystonia compared With Parkinson disease. *JAMA Neurol* 2015;72:1244-1251.
 83. Kondylis ED, Randazzo MJ, Alhourani A, et al. Movement-related dynamics of cortical oscillations in Parkinson's disease and essential tremor. *Brain* 2016;139(Pt 8):2211-2223.
 84. van Wijk BCM, Pogossyan A, Hariz MI, et al. Localization of beta and high-frequency oscillations within the subthalamic nucleus region. *Neuroimage Clin* 2017;16:175-183.
 85. van Wijk BC, Beudel M, Jha A, et al. Subthalamic nucleus phase-amplitude coupling correlates with motor impairment in Parkinson's disease. *Clin Neurophysiol* 2016;127:2010-2019.
 86. Meidahl AC, Moll CKE, van Wijk BCM, et al. Synchronised spiking activity underlies phase amplitude coupling in the subthalamic nucleus of Parkinson's disease patients. *Neurobiol Dis* 2019;127: 101-113.
 87. Tsiokos C, Malekmohammadi M, AuYong N, Pouratian N. Pallidal low beta-low gamma phase-amplitude coupling inversely correlates with Parkinson disease symptoms. *Clin Neurophysiol* 2017; 128:2165-2178.
 88. Connolly AT, Jensen AL, Baker KB, Vitek JL, Johnson MD. Classification of pallidal oscillations with increasing parkinsonian severity. *J Neurophysiol* 2015;114:209-218.

89. Connolly AT, Jensen AL, Bello EM, et al. Modulations in oscillatory frequency and coupling in globus pallidus with increasing parkinsonian severity. *J Neurosci* 2015;35:6231-6240.
90. Devergnas A, Caiola M, Pittard D, Wichmann T. Cortical phase-amplitude coupling in a progressive model of parkinsonism in non-human primates. *Cereb Cortex* 2019;29:167-177.
91. Marsden CD, Obeso JA. The functions of the basal ganglia and the paradox of stereotaxic surgery in Parkinson's disease. *Brain* 1994;117:877-897.
92. Obeso JA, Jahanshahi M, Alvarez L, et al. What can man do without basal ganglia motor output? The effect of combined unilateral subthalamotomy and pallidotomy in a patient with Parkinson's disease. *Exp Neurol* 2009;220:283-292.
93. Leblois A, Meissner W, Bioulac B, Gross CE, Hansel D, Boraud T. Late emergence of synchronized oscillatory activity in the pallidum during progressive Parkinsonism. *Eur J Neurosci* 2007;26:1701-1713.
94. Quiroga-Varela A, Walters JR, Brazhnik E, Marin C, Obeso JA. What basal ganglia changes underlie the parkinsonian state? The significance of neuronal oscillatory activity. *Neurobiol Dis* 2013;58:242-248.
95. Weinberger M, Mahant N, Hutchison WD, et al. Beta oscillatory activity in the subthalamic nucleus and its relation to dopaminergic response in Parkinson's disease. *J Neurophys* 2006;96:3248-3256.
96. Lee RG. Pathophysiology of rigidity and akinesia in Parkinson's disease. *Eur Neurol* 1989;29(Suppl 1):13-18.
97. Dietrichson P. The fusimotor system in relation to spasticity and parkinsonian rigidity. *Scand J Rehabil Med* 1973;5:174-178.
98. Andrews CJ, Burke D. Quantitative study of the effect of L-dopa and phenoxybenzamine on the rigidity of Parkinson's disease. *J Neurol Neurosurg Psychiatry* 1973;36:321-328.
99. Mortimer JA, Webster DD. Evidence for a quantitative association between EMG stretch responses and Parkinsonian rigidity. *Brain Res* 1979;162:169-173.
100. Xia R, Sun J, Threlkeld AJ. Analysis of interactive effect of stretch reflex and shortening reaction on rigidity in Parkinson's disease. *Clin Neurophysiol* 2009;120:1400-1407.
101. McLellan DL. Dynamic spindle reflexes and the rigidity of Parkinsonism. *J Neurol Neurosurg Psychiatry* 1973;36:342-349.
102. Cantello R, Gianelli M, Civardi C, Mutani R. Pathophysiology of Parkinson's disease rigidity. Role of corticospinal motor projections. *Adv Neurol* 1996;69:129-133.
103. Noth J, Schurmann M, Podoll K, Schwarz M. Reconsideration of the concept of enhanced static fusimotor drive in rigidity in patients with Parkinson's disease. *Neurosci Lett* 1988;84:239-243.
104. Berardelli A, Sabra A, Hallett M. Physiological mechanisms of rigidity in Parkinson's disease. *J Neurol Neurosurg Psychiatry* 1983;46:45-53.
105. Delwaide PJ, Sabbatino M, Delwaide C. Some pathophysiological aspects of the parkinsonian rigidity. *J Neural Transm Suppl* 1986;22:129-139.
106. Meara RJ, Cody FW. Stretch reflexes of individual parkinsonian patients studied during changes in clinical rigidity following medication. *Electroencephalogr Clin Neurophysiol* 1993;89:261-268.
107. Winogrodzka A, Wagenaar RC, Bergmans P, et al. Rigidity decreases resting tremor intensity in Parkinson's disease: a [(123)I] beta-CIT SPECT study in early, nonmedicated patients. *Mov Disord* 2001;16:1033-1040.
108. Pikstra ARA, van der Hoorn A, Leenders KL, de Jong BM. Relation of 18-F-Dopa PET with hypokinesia-rigidity, tremor and freezing in Parkinson's disease. *Neuroimage Clin* 2016;11:68-72.
109. Kuhn AA, Tsui A, Aziz T, et al. Pathological synchronisation in the subthalamic nucleus of patients with Parkinson's disease relates to both bradykinesia and rigidity. *Exp Neurol* 2009;215:380-387.
110. Little S, Pogossyan A, Kuhn AA, Brown P. Beta band stability over time correlates with Parkinsonian rigidity and bradykinesia. *Exp Neurol* 2012;236:383-388.
111. Yamada A, Takeuchi H, Miki H. [Unilateral abolition of parkinsonian rigidity after subthalamic nucleus hemorrhage]. [Article in Japanese]. *Rinsho Shinkeigaku* 1992;32:887-889.
112. Cooper IS, Samra K, Bergmann L. The thalamic lesion which abolishes tremor and rigidity of Parkinsonism. A radiologico-clinico-anatomic correlative study. *J Neurol Sci* 1969;8:69-84.
113. Shapiro MB, Vaillancourt DE, Sturman MM, Metman LV, Bakay RA, Corcos DM. Effects of STN DBS on rigidity in Parkinson's disease. *IEEE Trans Neural Syst Rehabil Eng* 2007;15:173-181.
114. Tabbal SD, Ushe M, Mink JW, et al. Unilateral subthalamic nucleus stimulation has a measurable ipsilateral effect on rigidity and bradykinesia in Parkinson disease. *Exp Neurol* 2008;211:234-242.
115. Hirato M, Ishihara J, Horikoshi S, Shibasaki T, Ohye C. Parkinsonian rigidity, dopa-induced dyskinesia and chorea—dynamic studies on the basal ganglia-thalamocortical motor circuit using PET scan and depth microrecording. *Acta Neurochir Suppl* 1995;64:5-8.
116. Karunanayaka PR, Lee EY, Lewis MM, et al. Default mode network differences between rigidity- and tremor-predominant Parkinson's disease. *Cortex* 2016;81:239-250.
117. Hou Y, Luo C, Yang J, et al. Default-mode network connectivity in cognitively unimpaired drug-naïve patients with rigidity-dominant Parkinson's disease. *J Neurol* 2017;264:152-160.
118. Hou Y, Wei Q, Ou R, et al. Impaired topographic organization in cognitively unimpaired drug-naïve patients with rigidity-dominant Parkinson's disease. *Parkinsonism Relat Disord* 2018;56:52-57.
119. Perez-Lloret S, Negre-Pages L, Damier P, et al. Prevalence, determinants, and effect on quality of life of freezing of gait in Parkinson disease. *JAMA Neurol* 2014;71:884-890.
120. Giladi N, McMahon D, Przedborski S, et al. Motor blocks in Parkinson's disease. *Neurology* 1992;42:333-339.
121. Amboni M, Stocchi F, Abbruzzese G, et al. Prevalence and associated features of self-reported freezing of gait in Parkinson disease: the DEEP FOG study. *Parkinsonism Relat Disord* 2015;21:644-649.
122. Giladi N, McDermott MP, Fahn S, et al. Freezing of gait in PD—prospective assessment in the DATATOP cohort. *Neurology* 2001;56:1712-1721.
123. Peterson DS, Fling BW, Mancini M, Cohen RG, Nutt JG, Horak FB. Dual-task interference and brain structural connectivity in people with Parkinson's disease who freeze. *J Neurol Neurosurg Psychiatry* 2015;86:786-792.
124. Myers PS, McNeely ME, Koller JM, Earhart GM, Campbell MC. Cerebellar volume and executive function in Parkinson disease with and without freezing of gait. *J Parkinsons Dis* 2017;7:149-157.
125. Picillo M, Dubbioso R, Iodice R, et al. Short-latency afferent inhibition in patients with Parkinson's disease and freezing of gait. *J Neural Transm (Vienna)* 2015;122:1533-1540.
126. Sunwoo MK, Cho KH, Hong JY, Lee JE, Sohn YH, Lee PH. Thalamic volume and related visual recognition are associated with freezing of gait in non-demented patients with Parkinson's disease. *Parkinsonism Relat Disord* 2013;19:1106-1109.
127. Factor SA, Scullin MK, Sollinger AB, et al. Freezing of gait subtypes have different cognitive correlates in Parkinson's disease. *Parkinsonism Relat Disord* 2014;20:1359-1364.
128. Factor SA, Steenland NK, Higgins DS, et al. Postural instability/gait disturbance in Parkinson's disease has distinct subtypes: an exploratory analysis. *J Neurol Neurosurg Psychiatry* 2011;82:564-568.
129. Peterson DS, Fling BW, Mancini M, Cohen RG, Nutt JG, Horak FB. Dual-task interference and brain structural connectivity in people with Parkinson's disease who freeze. *J Neurol Neurosurg Psychiatry* 2015;86:786-792.
130. Kostic VS, Agosta F, Pievani M, et al. Pattern of brain tissue loss associated with freezing of gait in Parkinson disease. *Neurology* 2012;78:409-416.
131. Tessitore A, Amboni M, Cirillo G, et al. Regional gray matter atrophy in patients with Parkinson disease and freezing of gait. *AJNR Am J Neuroradiol* 2012;33:1804-1809.

132. Shine JM, Matar E, Ward PB, et al. Freezing of gait in Parkinson's disease is associated with functional decoupling between the cognitive control network and the basal ganglia. *Brain* 2013;136(Pt 12):3671-3681.
133. Shine JM, Moustafa AA, Matar E, Frank MJ, Lewis SJ. The role of frontostriatal impairment in freezing of gait in Parkinson's disease. *Front Syst Neurosci* 2013;7:61.
134. Snijders AH, Takakusaki K, Debu B, et al. Physiology of freezing of gait. *Ann Neurol* 2016;80:644-659.
135. Lewis SJG, Shine JM. The next step: a common neural mechanism for freezing of gait. *Neuroscientist* 2016;22:72-82.
136. Toledo JB, Lopez-Azcarate J, Garcia-Garcia D, et al. High beta activity in the subthalamic nucleus and freezing of gait in Parkinson's disease. *Neurobiol Dis* 2014;64:60-65.
137. Scholten M, Govindan RB, Braun C, et al. Cortical correlates of susceptibility to upper limb freezing in Parkinson's disease. *Clin Neurophysiol* 2016;127:2386-2393.
138. Peterson DS, Pickett KA, Duncan R, Perlmutter J, Earhart GM. Gait-related brain activity in people with Parkinson disease with freezing of gait. *PLoS One* 2014;9:e90634.
139. Ferraye MU, Debu B, Fraix V, et al. Effects of subthalamic nucleus stimulation and levodopa on freezing of gait in Parkinson disease. *Neurology* 2008;70(Pt 2):1431-1437.
140. Moreau C, Defebvre L, Destee A, et al. STN-DBS frequency effects on freezing of gait in advanced Parkinson disease. *Neurology* 2008;71:80-84.
141. Storz L, Butz M, Hirschmann J, et al. Bicycling suppresses abnormal beta synchrony in the Parkinsonian basal ganglia. *Ann Neurol* 2017;82:592-601.
142. Gut NK, Winn P. Deep brain stimulation of different pedunculopontine targets in a novel rodent model of parkinsonism. *J Neurosci* 2015;35:4792-4803.
143. Kucinski A, Albin RL, Lustig C, Sarter M. Modeling falls in Parkinson's disease: Slow gait, freezing episodes and falls in rats with extensive striatal dopamine loss. *Behav Brain Res* 2015;282:155-164.
144. Xiao H, Li M, Cai J, et al. Selective cholinergic depletion of pedunculopontine tegmental nucleus aggravates freezing of gait in parkinsonian rats. *Neurosci Lett* 2017;659:92-98.
145. Revuelta GJ, Uthayathas S, Wahlquist AE, Factor SA, Papa SM. Non-human primate FOG develops with advanced parkinsonism induced by MPTP Treatment. *Exp Neurol* 2012;237:464-469.
146. Goetz L, Piallat B, Bhattacharjee M, Mathieu H, David O, Chabardes S. Spike discharge characteristic of the caudal mesencephalic reticular formation and pedunculopontine nucleus in MPTP-induced primate model of Parkinson disease. *Neurobiol Dis* 2018 Aug 4. pii: S0969-9961(18)30394-2. <https://doi.org/10.1016/j.nbd.2018.08.002>. [Epub ahead of print]
147. Ono SA, Sato T, Muramatsu S. Freezing of gait in Parkinson's disease is associated with reduced 6-[(18)F]Fluoro-L-m-tyrosine uptake in the locus coeruleus. *Parkinsons Dis* 2016;2016:5430920.
148. Bohnen NI, Frey KA, Studenski S, et al. Extra-nigral pathological conditions are common in Parkinson's disease with freezing of gait: an in vivo positron emission tomography study. *Mov Disord* 2014;29:1118-1124.
149. Masilamoni G, Beck G, Factor SA, et al. Pathology of brainstem noradrenergic cell groups associated with freezing of gait in MPTP-treated non-human primates. 4th International Workshop on Freezing of Gait 2018;Abstract Volume:11.
150. Huddleston D, Goldstein F, Langley J, et al. Neuromelanin-sensitive MRI correlates of executive function and verbal memory in Parkinson's disease with freezing of gait. 4th International Workshop on Freezing of Gait 2018;Abstract Volume:55-56.
151. Jankovic J. Atomoxetine for freezing of gait in Parkinson disease. *J Neurol Sci* 2009;284:177-178.
152. Revuelta GJ, Embry A, Elm JJ, et al. Pilot study of atomoxetine in patients with Parkinson's disease and dopa-unresponsive Freezing of Gait. *Transl Neurodegener* 2015;4:24.
153. Katsube J, Narabayashi H, Hayashi A, Tanaka C, Suzuki T. Development of L-threo-DOPS, a norepinephrine precursor amino-acid. *Yakugaku Zasshi-J Pharm Soc Jpn* 1994;114:823-846.
154. Narabayashi H, Kondo T, Hayashi A, Suzuki T. L-threo-3,4-dihydroxyphenylserine treatment for akinesia and freezing of parkinsonism. *Neurochem Res* 1982;7:850-851.
155. Narabayashi H, Kondo T, Nagatsu T. L-threo-3,4-dihydroxyphenylserine treatment for freezing and akinesia in parkinsonism. *Neurology* 1982;32:A180.
156. Fukada K, Endo T, Yokoe M, Hamasaki T, Hazama T, Sakoda S. L-threo-3,4-dihydroxyphenylserine (L-DOPS) co-administered with entacapone improves freezing of gait in Parkinson's disease. *Med Hypotheses* 2013;80:209-212.
157. Bohnen NI, Kanel P, Zhou Z, et al. Cholinergic system changes of falls and freezing of gait in Parkinson's disease. *Ann Neurol* 2019;85:538-549.
158. Karachi C, Grabli D, Bernard FA, et al. Cholinergic mesencephalic neurons are involved in gait and postural disorders in Parkinson disease. *J Clin Invest* 2010;120:2745-2754.
159. Grabli D, Karachi C, Folgoas E, et al. Gait disorders in parkinsonian monkeys with pedunculopontine nucleus lesions: a tale of two systems. *J Neurosci* 2013;33:11986-11993.
160. Glajch KE, Kelver DA, Hegeman DJ, et al. Npas1+ pallidal neurons target striatal projection neurons. *J Neurosci* 2016;36:5472-5488.
161. Hernandez VM, Hegeman DJ, Cui Q, et al. Parvalbumin+ neurons and Npas1+ neurons are distinct neuron classes in the mouse external globus pallidus. *J Neurosci* 2015;35:11830-11847.
162. Saunders A, Huang KW, Sabatini BL. Globus pallidus externus neurons expressing parvalbumin interconnect the subthalamic nucleus and striatal interneurons. *PLoS One* 2016;11:e0149798.
163. Mastro KJ, Bouchard RS, Holt HA, Gittis AH. Transgenic mouse lines subdivide external segment of the globus pallidus (GPe) neurons and reveal distinct GPe output pathways. *J Neurosci* 2014;34:2087-2099.
164. Mastro KJ, Zitelli KT, Willard AM, Leblanc KH, Kravitz AV, Gittis AH. Cell-specific pallidal intervention induces long-lasting motor recovery in dopamine-depleted mice. *Nat Neurosci* 2017;20:815-823.
165. Hallworth NE, Bevan MD. Globus pallidus neurons dynamically regulate the activity pattern of subthalamic nucleus neurons through the frequency-dependent activation of postsynaptic GABAA and GABAB receptors. *J Neurosci* 2005;25:6304-6315.
166. Bevan MD, Magill PJ, Hallworth NE, Bolam JP, Wilson CJ. Regulation of the timing and pattern of action potential generation in rat subthalamic neurons in vitro by GABA-A IPSPs. *J Neurophys* 2002;87:1348-1362.
167. Baufreton J, Kirkham E, Atherton JF, et al. Sparse but selective and potent synaptic transmission from the globus pallidus to the subthalamic nucleus. *J Neurophysiol* 2009;102:532-545.
168. Baufreton J, Atherton JF, Surmeier DJ, Bevan MD. Enhancement of excitatory synaptic integration by GABAergic inhibition in the subthalamic nucleus. *J Neurosci* 2005;25:8505-8517.
169. Vitek JL, Zhang J, Hashimoto T, Russo GS, Baker KB. External pallidal stimulation improves parkinsonian motor signs and modulates neuronal activity throughout the basal ganglia thalamic network. *Exp Neurol* 2012;233:581-586.
170. Zhang J, Russo GS, Mewes K, Rye DB, Vitek JL. Lesions in monkey globus pallidus externus exacerbate parkinsonian symptoms. *Exp Neurol* 2006;199:446-453.
171. Kita H, Tachibana Y, Nambu A, Chiken S. Balance of monosynaptic excitatory and disynaptic inhibitory responses of the globus pallidus induced after stimulation of the subthalamic nucleus in the monkey. *J Neurosci* 2005;25:8611-8619.
172. Tachibana Y, Iwamuro H, Kita H, Takada M, Nambu A. Subthalamo-pallidal interactions underlying parkinsonian neuronal oscillations in the primate basal ganglia. *Eur J Neurosci* 2011;34:1470-1484.

173. Vitek JL, Hashimoto T, Peoples J, DeLong MR, Bakay RA. Acute stimulation in the external segment of the globus pallidus improves parkinsonian motor signs. *Mov Disord* 2004;19:907-915.
174. Yelnik J, Damier P, Bejjani BP, et al. Functional mapping of the human globus pallidus: contrasting effect of stimulation in the internal and external pallidum in Parkinson's disease. *Neuroscience* 2000;101:77-87.
175. Mallet N, Micklem BR, Henny P, et al. Dichotomous organization of the external globus pallidus. *Neuron* 2012;74:1075-1086.
176. Bevan MD, Booth PA, Eaton SA, Bolam JP. Selective innervation of neostriatal interneurons by a subclass of neuron in the globus pallidus of the rat. *J Neurosci* 1998;18:9438-9452.
177. Corbit VL, Whalen TC, Zitelli KT, Crilly SY, Rubin JE, Gittis AH. Pallidostriatal projections promote β oscillations in a dopamine-depleted biophysical network model. *J Neurosci* 2016;36:5556-5571.
178. Abdi A, Mallet N, Mohamed FY, et al. Prototypic and arkypallidal neurons in the dopamine-intact external globus pallidus. *J Neurosci* 2015;35:6667-6688.
179. Mastro KJ, Zitelli KT, Willard AM, Leblanc KH, Kravitz AV, Gittis AH. Cell-type specific pallidal stimulation provides long-lasting relief of immobility in a model of Parkinson's disease. *Society for Neuroscience* 2016;San Diego:Program no. 602.624.
180. Rommelfanger KS, Wichmann T. Extrastriatal dopaminergic circuits of the basal ganglia. *Front Neuroanat* 2010;4:139.
181. Pifl C, Bertel O, Schingnitz G, Hornykiewicz O. Extrastriatal dopamine in symptomatic and asymptomatic Rhesus monkeys treated with 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP). *Neurochem Int* 1990;17:263-270.
182. Hornykiewicz O. Biochemical aspects of Parkinson's disease. *Neurology* 1998;51(2 Suppl 2):S2-S9.
183. Leenders KL, Salmon EP, Tyrrell P, et al. The nigrostriatal dopaminergic system assessed in vivo by positron emission tomography in healthy volunteer subjects and patients with Parkinson's disease. *Arch Neurol* 1990;47:1290-1298.
184. Porritt M, Stanic D, Finkelstein D, et al. Dopaminergic innervation of the human striatum in Parkinson's disease. *Mov Disord* 2005;20:810-818.
185. Hadipour-Niktarash A, Rommelfanger KS, Masilamoni GJ, Smith Y, Wichmann T. Extrastriatal D2-like receptors modulate basal ganglia pathways in normal and Parkinsonian monkeys. *J Neurophysiol* 2012;107:1500-1512.
186. Galvan A, Hu X, Rommelfanger KS, et al. Localization and function of dopamine receptors in the subthalamic nucleus of normal and parkinsonian monkeys. *J Neurophysiol* 2014;112:467-479.
187. Whone AL, Moore RY, Piccini PP, Brooks DJ. Plasticity of the nigropallidal pathway in Parkinson's disease. *Ann Neurol* 2003;53:206-213.
188. Kliem MA, Pare JF, Khan ZU, Wichmann T, Smith Y. Ultrastructural localization and function of dopamine D1-like receptors in the substantia nigra pars reticulata and the internal segment of the globus pallidus of parkinsonian monkeys. *Eur J Neurosci* 2010;31:836-851.
189. Freeman A, Ciliax B, Bakay R, et al. Nigrostriatal collaterals to thalamus degenerate in parkinsonian animal models. *Ann Neurol* 2001;50:321-329.
190. Garcia-Cabezas MA, Rico B, Sanchez-Gonzalez MA, Cavada C. Distribution of the dopamine innervation in the macaque and human thalamus. *Neuroimage* 2007;34:965-984.
191. Garcia-Cabezas MA, Martinez-Sanchez P, Sanchez-Gonzalez MA, Garzon M, Cavada C. Dopamine innervation in the thalamus: monkey versus rat. *Cereb Cortex* 2009;19:424-434.
192. Sanchez-Gonzalez MA, Garcia-Cabezas MA, Rico B, Cavada C. The primate thalamus is a key target for brain dopamine. *J Neurosci* 2005;25:6076-6083.
193. Dirks MF, den Ouden HE, Aarts E, et al. Dopamine controls Parkinson's tremor by inhibiting the cerebellar thalamus. *Brain* 2017;140:721-734.
194. Pifl C, Hornykiewicz O, Blesa J, Adanez R, Cavada C, Obeso JA. Reduced noradrenaline, but not dopamine and serotonin in motor thalamus of the MPTP primate: relation to severity of parkinsonism. *J Neurochem* 2013;125:657-662.
195. El-Banoua F, Caraballo I, Flores JA, Galan-Rodriguez B, Fernandez-Espejo E. Effects on turning of microinjections into basal ganglia of D(1) and D(2) dopamine receptors agonists and the cannabinoid CB(1) antagonist SR141716A in a rat Parkinson's model. *Neurobiol Dis* 2004;16:377-385.
196. Mehta A, Theros K, Chesselet MF. Increased behavioral response to dopaminergic stimulation of the subthalamic nucleus after nigrostriatal lesions. *Synapse* 2000;37:298-307.
197. Galvan A, Floran B, Erlij D, Aceves J. Intrapallidal dopamine restores motor deficits induced by 6-hydroxydopamine in the rat. *J Neural Transm* 2001;108:153-166.
198. Ray NJ, Miyasaki JM, Zurowski M, et al. Extrastriatal dopaminergic abnormalities of DA homeostasis in Parkinson's patients with medication-induced pathological gambling: a [11C] FLB-457 and PET study. *Neurobiol Dis* 2012;48:519-525.
199. Lee JY, Seo SH, Kim YK, et al. Extrastriatal dopaminergic changes in Parkinson's disease patients with impulse control disorders. *J Neurol Neurosurg Psychiatry* 2014;85:23-30.
200. Frosini D, Unti E, Guidoccio F, et al. Mesolimbic dopaminergic dysfunction in Parkinson's disease depression: evidence from a 123I-FP-CIT SPECT investigation. *J Neural Transm (Vienna)* 2015;122:1143-1147.
201. Ko JH, Antonelli F, Monchi O, et al. Prefrontal dopaminergic receptor abnormalities and executive functions in Parkinson's disease. *Hum Brain Mapp* 2013;34:1591-1604.
202. Marquie M, Locascio JJ, Rentz DM, et al. Striatal and extrastriatal dopamine transporter levels relate to cognition in Lewy body diseases: an (11)C altropane positron emission tomography study. *Alzheimers Res Ther* 2014;6:52.
203. Ingham CA, Hood SH, Arbuthnott GW. Spine density on neostriatal neurones changes with 6-hydroxydopamine lesions and with age. *Brain Res* 1989;503:334-338.
204. Ingham CA, Hood SH, Taggart P, Arbuthnott GW. Plasticity of synapses in the rat neostriatum after unilateral lesion of the nigrostriatal dopaminergic pathway. *J Neurosci* 1998;18:4732-4743.
205. Smith Y, Villalba RM, Raju DV. Striatal spine plasticity in Parkinson's disease: pathological or not? *Parkinsonism Relat Disord* 2009;15(Suppl 3):S156-S161.
206. Villalba RM, Mathai A, Smith Y. Morphological changes of glutamatergic synapses in animal models of Parkinson's disease. *Front Neuroanat* 2015;9:117.
207. Villalba RM, Smith Y. Differential striatal spine pathology in Parkinson's disease and cocaine addiction: a key role of dopamine? *Neurosci* 2013;251:2-20.
208. Villalba RM, Smith Y. Loss and remodeling of striatal dendritic spines in Parkinson's disease: from homeostasis to maladaptive plasticity? *J Neural Transm (Vienna)* 2018;125:431-447.
209. Villalba RM, Smith Y. Differential structural plasticity of corticostriatal and thalamostriatal axo-spinous synapses in MPTP-treated Parkinsonian monkeys. *J Comp Neurol* 2011;519:989-1005.
210. Villalba RM, Wichmann T, Smith Y. Neuronal loss in the caudal intralaminar thalamic nuclei in a primate model of Parkinson's disease. *Brain Struct Funct* 2014;219:381-394.
211. Smith Y, Galvan A, Ellender TJ, et al. The thalamostriatal system in normal and diseased states. *Front Syst Neurosci* 2014;8:5.
212. Halliday GM. Thalamic changes in Parkinson's disease. *Parkinsonism Relat Disord* 2009;15(Suppl 3):S152-S155.
213. Mathai A, Ma Y, Pare JF, Villalba RM, Wichmann T, Smith Y. Reduced cortical innervation of the subthalamic nucleus in MPTP-treated parkinsonian monkeys. *Brain* 2015;138(Pt 4):946-962.
214. Patriat R, Kaplan J, Niederer J, et al. Fractional anisotropy within the subthalamic nucleus: an imaging biomarker for early Parkinson's disease? A high-resolution 7Tesla magnetic resonance imaging study. *Soc Neurosci Abstr*. 2018: 654.13.
215. Fan KY, Baufreton J, Surmeier DJ, Chan CS, Bevan MD. Proliferation of external globus pallidus-subthalamic nucleus synapses

- following degeneration of midbrain dopamine neurons. *J Neurosci* 2012;32:13718-13728.
216. Chu HY, Atherton JF, Wokosin D, Surmeier DJ, Bevan MD. Heterosynaptic regulation of external globus pallidus inputs to the subthalamic nucleus by the motor cortex. *Neuron* 2015;85:364-376.
 217. Elsworth JD, Leraneth C, Redmond DE, Roth RH. Loss of asymmetric spine synapses in prefrontal cortex of motor-asymptomatic, dopamine-depleted, cognitively impaired MPTP-treated monkeys. *Int J Neuropsychopharmacol* 2013;16:905-912.
 218. Villalba RM, Pare JF, Wichmann T, Smith Y. Loss of glutamatergic thalamic inputs in deep layers of primary motor cortex in MPTP-treated parkinsonian monkeys. *Soc Neurosci Abstr* 2014;248.211.
 219. Narabayashi H. Possible role of cerebellar circuits in parkinsonian symptoms. *Confin Neurol* 1974;36:292-301.
 220. Kakita A, Takahashi H, Homma Y, Ikuta F. Lewy bodies in the cerebellar dentate nucleus of a patient with Parkinson's disease. *Pathol Int* 1994;44:878-880.
 221. Seidel K, Bouzrou M, Heidemann N, et al. Involvement of the cerebellum in Parkinson disease and dementia with Lewy bodies. *Ann Neurol* 2017;81:898-903.
 222. Hurley MJ, Mash DC, Jenner P. Markers for dopaminergic neurotransmission in the cerebellum in normal individuals and patients with Parkinson's disease examined by RT-PCR. *Eur J Neurosci* 2003;18:2668-2672.
 223. Kish SJ, Shannak KS, Rajput AH, Gilbert JJ, Hornykiewicz O. Cerebellar norepinephrine in patients with Parkinson's disease and control subjects. *Arch Neurol* 1984;41:612-614.
 224. Lewis MM, Galley S, Johnson S, Stevenson J, Huang X, McKeown MJ. The role of the cerebellum in the pathophysiology of Parkinson's disease. *Can J Neurol Sci* 2013;40:299-306.
 225. Wu T, Hallett M. The cerebellum in Parkinson's disease. *Brain* 2013;136(Pt 3):696-709.
 226. Martinu K, Monchi O. Cortico-basal ganglia and cortico-cerebellar circuits in Parkinson's disease: pathophysiology or compensation? *Behav Neurosci* 2013;127:222-236.
 227. Bostan AC, Dum RP, Strick PL. The basal ganglia communicate with the cerebellum. *Proc Natl Acad Sci U S A* 2010;107:8452-8456.
 228. Bostan AC, Dum RP, Strick PL. Cerebellar networks with the cerebral cortex and basal ganglia. *Trends Cogn Sci* 2013;17:241-254.
 229. Bostan AC, Dum RP, Strick PL. Functional anatomy of basal ganglia circuits with the cerebral cortex and the cerebellum. *Prog Neurol Surg* 2018;33:50-61.
 230. Bostan AC, Strick PL. The cerebellum and basal ganglia are interconnected. *Neuropsychol Rev* 2010;20:261-270.
 231. Rascol O, Sabatini U, Fabre N, et al. The ipsilateral cerebellar hemisphere is overactive during hand movements in akinetic parkinsonian patients. *Brain* 1997;120(Pt 1):103-110.
 232. Cerasa A, Hagberg GE, Peppe A, et al. Functional changes in the activity of cerebellum and frontostriatal regions during externally and internally timed movement in Parkinson's disease. *Brain Res Bull* 2006;71:259-269.
 233. Yu H, Sternad D, Corcos DM, Vaillancourt DE. Role of hyperactive cerebellum and motor cortex in Parkinson's disease. *Neuroimage* 2007;35:222-233.
 234. Sen S, Kawaguchi A, Truong Y, Lewis MM, Huang X. Dynamic changes in cerebello-thalamo-cortical motor circuitry during progression of Parkinson's disease. *Neurosci* 2010;166:712-719.
 235. Simioni AC, Dagher A, Fellows LK. Compensatory striatal-cerebellar connectivity in mild-moderate Parkinson's disease. *NeuroImage Clinical* 2016;10:54-62.
 236. O'Sullivan D, Pell M. Long-term follow-up of DBS of thalamus for tremor and STN for Parkinson's disease. *Brain Research Bulletin* 2009;78:119-121.
 237. Koller WC, Pahwa PR, Lyons KE, Wilkinson SB. Deep brain stimulation of the Vim nucleus of the thalamus for the treatment of tremor. *Neurology* 2000;55(12 Suppl 6):S29-S33.
 238. Deiber MP, Pollak P, Passingham R, et al. Thalamic stimulation and suppression of parkinsonian tremor. Evidence of a cerebellar deactivation using positron emission tomography. *Brain* 1993;116(Pt 1):267-279.
 239. Hilker R, Voges J, Weisenbach S, et al. Subthalamic nucleus stimulation restores glucose metabolism in associative and limbic cortices and in cerebellum: evidence from a FDG-PET study in advanced Parkinson's disease. *J Cereb Blood Flow Metab* 2004;24:7-16.
 240. Sutton AC, O'Connor KA, Pilitsis JG, Shin DS. Stimulation of the subthalamic nucleus engages the cerebellum for motor function in parkinsonian rats. *Brain Struct Funct* 2015;220:3595-3609.
 241. Helmich RC, Janssen MJ, Oyen WJ, Bloem BR, Toni I. Pallidal dysfunction drives a cerebellothalamic circuit into Parkinson tremor. *Ann Neurol* 2011;69:269-281.
 242. Stevenson JK, Oishi MM, Farajian S, Cretu E, Ty E, McKeown MJ. Response to sensory uncertainty in Parkinson's disease: a marker of cerebellar dysfunction? *Eur J Neurosci* 2011;33:298-305.
 243. Lu MK, Chen JC, Chen CM, Duann JR, Ziemann U, Tsai CH. Impaired cerebellum to primary motor cortex associative plasticity in Parkinson's disease and spinocerebellar ataxia type 3. *Front Neurol* 2017;8:445.
 244. Lewis MM, Slagle CG, Smith AB, et al. Task specific influences of Parkinson's disease on the striato-thalamo-cortical and cerebello-thalamo-cortical motor circuitries. *Neurosci* 2007;147:224-235.
 245. Kishore A, Popa T. Cerebellum in levodopa-induced dyskinesias: the unusual suspect in the motor network. *Front Neurol* 2014;5:157.
 246. Koch G, Brusa L, Carrillo F, et al. Cerebellar magnetic stimulation decreases levodopa-induced dyskinesias in Parkinson disease. *Neurology* 2009;73:113-119.
 247. Brusa L, Ceravolo R, Kiferle L, et al. Metabolic changes induced by theta burst stimulation of the cerebellum in dyskinetic Parkinson's disease patients. *Parkinsonism Relat Disord* 2012;18:59-62.
 248. Pessiglione M, Guehl D, Rolland AS, et al. Thalamic neuronal activity in dopamine-depleted primates: evidence for a loss of functional segregation within basal ganglia circuits. *J Neurosci* 2005;25:1523-1531.
 249. Muralidharan A, Zhang J, Ghosh D, Johnson MD, Baker KB, Vitek JL. Modulation of neuronal activity in the motor thalamus during GPi-DBS in the MPTP nonhuman primate model of Parkinson's disease. *Brain Stimul* 2017;10:126-138.
 250. Elder CM, Vitek JL. The motor thalamus: alteration of neuronal activity in the parkinsonian state. In: Kultas-Ilinsky K, Ilinsky IA, eds. *Basal Ganglia and Thalamus in Health and Movement Disorders*. New York, NY: Kluwer Academic; 2001.
 251. Lenz FA, Jaeger CJ, Seike MS, et al. Thalamic single neuron activity in patients with dystonia: dystonia-related activity and somatic sensory reorganization. *J Neurophys* 1999;82:2372-2392.
 252. Lenz FA, Vitek JL, DeLong MR. Role of the thalamus in parkinsonian tremor: evidence from studies in patients and primate models. *Stereotact Funct Neurosurg* 1993;60:94-103.
 253. Mena-Segovia J, Bolam JP. Rethinking the pedunculopontine nucleus: from cellular organization to function. *Neuron* 2017;94:7-18.
 254. Mena-Segovia J, Bolam JP, Magill PJ. Pedunculopontine nucleus and basal ganglia: distant relatives or part of the same family? *Trends Neurosci* 2004;27:585-588.
 255. Fling BW, Cohen RG, Mancini M, Nutt JG, Fair DA, Horak FB. Asymmetric pedunculopontine network connectivity in parkinsonian patients with freezing of gait. *Brain* 2013;136(Pt 8):2405-2418.
 256. Nandi D, Jenkinson N, Stein J, Aziz T. The pedunculopontine nucleus in Parkinson's disease: primate studies. *Br J Neurosurg* 2008;22(Suppl 1):S4-S8.
 257. Geng X, Xie J, Wang X, et al. Altered neuronal activity in the pedunculopontine nucleus: An electrophysiological study in a rat model of Parkinson's disease. *Behav Brain Res* 2016;305:57-64.

258. Jimenez-Martin J, Blanco-Lezcano L, González-Fraguela ME, et al. Effect of neurotoxic lesion of pedunculopontine nucleus in nigral and striatal redox balance and motor performance in rats. *Neuroscience* 2015;289:300-314.
259. Stein JF. Akinesia, motor oscillations and the pedunculopontine nucleus in rats and men. *Exp Neurol* 2009;215:1-4.
260. Jenkinson N, Nandi D, Muthusamy K, et al. Anatomy, physiology, and pathophysiology of the pedunculopontine nucleus. *Mov Disord* 2009;24:319-328.
261. Winn P. Experimental studies of pedunculopontine functions: are they motor, sensory or integrative? *Parkinsonism Relat Disord* 2008;14(Suppl 2):S194-S198.
262. Breit S, Lessmann L, Benazzouz A, Schulz JB. Unilateral lesion of the pedunculopontine nucleus induces hyperactivity in the subthalamic nucleus and substantia nigra in the rat. *Eur J Neurosci* 2005;22:2283-2294.
263. Jenkinson N, Nandi D, Miall RC, Stein JF, Aziz TZ. Pedunculopontine nucleus stimulation improves akinesia in a Parkinsonian monkey. *Neuroreport* 2004;15:2621-2624.
264. Jeon MF, Ha Y, Cho YH, Lee BH, Park YG, Chang JW. Effect of ipsilateral subthalamic nucleus lesioning in a rat parkinsonian model: study of behavior correlated with neuronal activity in the pedunculopontine nucleus. *J Neurosurg* 2003;99:762-767.
265. Nandi D, Aziz TZ, Giladi N, Winter J, Stein JF. Reversal of akinesia in experimental parkinsonism by GABA antagonist micro-injections in the pedunculopontine nucleus. *Brain* 2002;125(Pt 11):2418-2430.
266. Lee MS, Rinne JO, Marsden CD. The pedunculopontine nucleus: its role in the genesis of movement disorders. *Yonsei Med J* 2000;41:167-184.
267. Kojima J, Yamaji Y, Matsumura M, et al. Excitotoxic lesions of the pedunculopontine tegmental nucleus produce contralateral hemiparkinsonism in the monkey. *Neurosci Lett* 1997;226:111-114.
268. Zweig RM, Jankel WR, Hedreen JC, Mayeux R, Price DL. The pedunculopontine nucleus in Parkinson's disease. *Ann Neurol* 1989;26:41-46.
269. Thevathasan W, Pogoyan A, Hyam JA, et al. Alpha oscillations in the pedunculopontine nucleus correlate with gait performance in parkinsonism. *Brain* 2012;135(Pt 1):148-160.
270. Li M, Zhang W. Oscillations in pedunculopontine nucleus in Parkinson's disease and its relationship with deep brain stimulation. *Front Neural Circuits* 2015;9:47.
271. Tattersall TL, Stratton PG, Coyne TJ, et al. Imagined gait modulates neuronal network dynamics in the human pedunculopontine nucleus. *Nat Neurosci* 2014;17:449-454.
272. Androulidakis AG, Khan S, Litvak V, Pleydell-Pearce CW, Brown P, Gill SS. Local field potential recordings from the pedunculopontine nucleus in a Parkinsonian patient. *Neuroreport* 2008;19:59-62.
273. Androulidakis AG, Mazzone P, Litvak V, et al. Oscillatory activity in the pedunculopontine area of patients with Parkinson's disease. *Exp Neurol* 2008;211:59-66.
274. Fraix V, Bastin J, David O, et al. Pedunculopontine nucleus area oscillations during stance, stepping and freezing in Parkinson's disease. *PLoS One* 2013;8:e83919.
275. Tsang EW, Hamani C, Moro E, et al. Involvement of the human pedunculopontine nucleus region in voluntary movements. *Neurology* 2010;75:950-959.
276. Rosin B, Slovik M, Mitelman R, et al. Closed-loop deep brain stimulation is superior in ameliorating parkinsonism. *Neuron* 2011;72:370-384.
277. Little S, Pogoyan A, Neal S, et al. Adaptive deep brain stimulation in advanced Parkinson disease. *Ann Neurol* 2013;74:449-457.
278. Swann NC, de Hemptinne C, Thompson MC, et al. Adaptive deep brain stimulation for Parkinson's disease using motor cortex sensing. *J Neural Eng* 2018;15:046006.
279. Parker JG, Marshall JD, Ahanonu B, et al. Diametric neural ensemble dynamics in parkinsonian and dyskinetic states. *Nature* 2018;557:177-182.
280. Deffains M, Iskhakova L, Katabi S, Haber SN, Israel Z, Bergman H. Subthalamic, not striatal, activity correlates with basal ganglia downstream activity in normal and parkinsonian monkeys. *eLife* 2016;5:e16443.
281. Pasquereau B, DeLong MR, Turner RS. Primary motor cortex of the parkinsonian monkey: altered encoding of active movement. *Brain* 2016;139(Pt 1):127-143.
282. Kempf F, Kuhn AA, Kupsch A, et al. Premovement activities in the subthalamic area of patients with Parkinson's disease and their dependence on task. *Eur J Neurosci* 2007;25:3137-3145.
283. Rowland NC, De Hemptinne C, Swann NC, et al. Task-related activity in sensorimotor cortex in Parkinson's disease and essential tremor: changes in beta and gamma bands. *Front Hum Neurosci* 2015;9:512.
284. Alegre M, Lopez-Azcarate J, Obeso I, et al. The subthalamic nucleus is involved in successful inhibition in the stop-signal task: a local field potential study in Parkinson's disease. *Exp Neurol* 2013;239:1-12.
285. Wichmann T. Pathophysiologic basis of movement disorders. *Prog Neurol Surg* 2018;33:13-24.