



## Extras for Chapter 2


- A bit more what is likelihood
- Why do we need the normalization term
- Plotting a continuous function
- Why probability density can be larger than 1
- Explicit conditioning on model  $M$

# Binomial: known $\theta$

- **Observation model** (function of  $y$ , discrete)

$$p(y|\theta, n) = \binom{n}{y} \theta^y (1 - \theta)^{n-y}$$

Binomial distribution with  $\theta = 0.5$ ,  $n = 10$




# Binomial: known $\theta$

- Observation model (function of  $y$ , discrete)

$$p(y|\theta, n) = \binom{n}{y} \theta^y (1 - \theta)^{n-y}$$

Binomial distribution with  $\theta = 0.5$ ,  $n = 10$




$p(y|n = 10, \theta = 0.5)$ : 0.00 0.01 0.04 0.12 0.21 0.25 0.21 0.12 0.04 0.01 0.00

# Binomial: known $\theta$

- **Observation model** (function of  $y$ , discrete)

$$p(y|\theta, n) = \binom{n}{y} \theta^y (1 - \theta)^{n-y}$$

Binomial distribution with  $\theta = 0.9$ ,  $n = 10$




$p(y|n = 10, \theta = 0.9)$ : 0.00 0.00 0.00 0.00 0.00 0.00 0.01 0.06 0.19 0.39 0.35

# Binomial: known $\theta$

- **Observation model** (function of  $y$ , discrete)

$$p(y|\theta, n) = \binom{n}{y} \theta^y (1 - \theta)^{n-y}$$

Binomial distribution with  $\theta = 0.5$ ,  $n = 10$




$p(y = 6 | n = 10, \theta = 0.5)$ : 0.00 0.01 0.04 0.12 0.21 0.25 **0.21** 0.12 0.04 0.01 0.00

# Binomial: known $\theta$

- **Observation model** (function of  $y$ , discrete)

$$p(y|\theta, n) = \binom{n}{y} \theta^y (1 - \theta)^{n-y}$$

Binomial distribution with  $\theta = 0.9$ ,  $n = 10$




$p(y = 6 | n = 10, \theta = 0.9)$ : 0.00 0.00 0.00 0.00 0.00 0.00 **0.01** 0.06 0.19 0.39 0.35

# Binomial: unknown $\theta$

- Likelihood (function of  $\theta$ , continuous)

$$p(y|\theta, n) = \binom{n}{y} \theta^y (1 - \theta)^{n-y}$$




$p(y = 6|n = 10, \theta)$ : 0.00 0.00 0.01 0.04 0.11 **0.21** 0.25 0.20 0.09 **0.01** 0.00



# Binomial: unknown $\theta$

- Likelihood (function of  $\theta$ , continuous)


$$p(y|\theta, n) = \binom{n}{y} \theta^y (1 - \theta)^{n-y}$$



# Binomial: unknown $\theta$

- Likelihood (function of  $\theta$ , continuous)


$$p(y|\theta, n) = \binom{n}{y} \theta^y (1 - \theta)^{n-y}$$



# Binomial: unknown $\theta$

- Likelihood (function of  $\theta$ , continuous)


$$p(y|\theta, n) = \binom{n}{y} \theta^y (1 - \theta)^{n-y}$$



# Binomial: unknown $\theta$

- Likelihood (function of  $\theta$ , continuous)

$$p(y|\theta, n) = \binom{n}{y} \theta^y (1 - \theta)^{n-y}$$



`integrate(function(theta) dbinom(6, 10, theta), 0, 1)  $\approx$  0.09`

- Posterior with Bayes rule (function of  $\theta$ , continuous)

$$p(\theta|y, n) = \frac{p(y|\theta, n)p(\theta|n)}{p(y|n)}$$

- Posterior with Bayes rule (function of  $\theta$ , continuous)

$$p(\theta|y, n) = \frac{p(y|\theta, n)p(\theta|n)}{p(y|n)}$$

where  $p(y|n) = \int p(y|\theta, n)p(\theta|n)d\theta$

# Binomial: unknown $\theta$

- Posterior with Bayes rule (function of  $\theta$ , continuous)

$$p(\theta|y, n) = \frac{p(y|\theta, n)p(\theta|n)}{p(y|n)}$$

where  $p(y|n) = \int p(y|\theta, n)p(\theta|n)d\theta$

- Start with uniform prior

$$p(\theta|n) = p(\theta|M) = 1, \text{ when } 0 \leq \theta \leq 1$$

- Posterior with Bayes rule (function of  $\theta$ , continuous)

$$p(\theta|y, n) = \frac{p(y|\theta, n)p(\theta|n)}{p(y|n)}$$

where  $p(y|n) = \int p(y|\theta, n)p(\theta|n)d\theta$

- Start with uniform prior

$$p(\theta|n) = p(\theta|M) = 1, \text{ when } 0 \leq \theta \leq 1$$

- Then

$$\begin{aligned} p(\theta|y, n) &= \frac{p(y|\theta, n)}{p(y|n)} = \frac{\binom{n}{y} \theta^y (1 - \theta)^{n-y}}{\int_0^1 \binom{n}{y} \theta^y (1 - \theta)^{n-y} d\theta} \\ &= \frac{1}{Z} \theta^y (1 - \theta)^{n-y} \end{aligned}$$



- Normalization term  $Z$  (constant given  $y$ )

$$Z = p(y|n) = \int_0^1 \theta^y (1 - \theta)^{n-y} d\theta = \frac{\Gamma(y+1)\Gamma(n-y+1)}{\Gamma(n+2)}$$

- Evaluate with  $y = 6, n = 10$

```
y<-6;n<-10;
```

```
integrate(function(theta) theta^y*(1-theta)^(n-y), 0, 1)
```

```
≈ 0.0004329
```

- Normalization term  $Z$  (constant given  $y$ )

$$Z = p(y|n) = \int_0^1 \theta^y (1 - \theta)^{n-y} d\theta = \frac{\Gamma(y+1)\Gamma(n-y+1)}{\Gamma(n+2)}$$

- Evaluate with  $y = 6, n = 10$

```
y<-6;n<-10;
```

```
integrate(function(theta) theta^y*(1-theta)^(n-y), 0, 1)
```

```
≈ 0.0004329
```

```
gamma(6+1)*gamma(10-6+1)/gamma(10+2) ≈ 0.0004329
```

# Binomial: unknown $\theta$


- Posterior is

$$p(\theta|y, n) = \frac{\Gamma(n+2)}{\Gamma(y+1)\Gamma(n-y+1)}\theta^y(1-\theta)^{n-y},$$

which is called Beta distribution

$$\theta|y, n \sim \text{Beta}(y+1, n-y+1)$$

$p(\theta | y=6, n=10, M=\text{binom}) + \text{unif. prior}$



Sometimes conditioning on the model  $M$  is explicitly shown

- **Posterior** with Bayes rule (function of  $\theta$ , continuous)

$$p(\theta|y, n, M) = \frac{p(y|\theta, n, M)p(\theta|n, M)}{p(y|n, M)}$$

where  $p(y|n, M) = \int p(y|\theta, n, M)p(\theta|n, M)d\theta$

Sometimes conditioning on the model  $M$  is explicitly shown

- **Posterior** with Bayes rule (function of  $\theta$ , continuous)

$$p(\theta|y, n, M) = \frac{p(y|\theta, n, M)p(\theta|n, M)}{p(y|n, M)}$$

where  $p(y|n, M) = \int p(y|\theta, n, M)p(\theta|n, M)d\theta$

- makes it more clear that likelihood and prior are both part of the model
- makes it more clear that there is no absolute probability for  $p(y|n)$ , but it depends on the model  $M$
- in case of two models, we can evaluate marginal likelihoods  $p(y|n, M_1)$  and  $p(y|n, M_2)$  (more in Ch 7)

Sometimes conditioning on the model  $M$  is explicitly shown

- **Posterior** with Bayes rule (function of  $\theta$ , continuous)

$$p(\theta|y, n, M) = \frac{p(y|\theta, n, M)p(\theta|n, M)}{p(y|n, M)}$$

where  $p(y|n, M) = \int p(y|\theta, n, M)p(\theta|n, M)d\theta$

- makes it more clear that likelihood and prior are both part of the model
- makes it more clear that there is no absolute probability for  $p(y|n)$ , but it depends on the model  $M$
- in case of two models, we can evaluate marginal likelihoods  $p(y|n, M_1)$  and  $p(y|n, M_2)$  (more in Ch 7)
- usually dropped to make the notation more concise