## Advanced probabilistic methods - Sketch

Christian Segercrantz 481056 March 24, 2022

$$p(\psi, \mathbf{Z}, \mathbf{W}, \mathbf{X}) = p(\mathbf{X} \mid \psi, \mathbf{Z}, \mathbf{W}) p(\psi) p(\mathbf{Z}) p(\mathbf{W})$$

$$= \mathcal{N}_{\mathcal{D}}(\mathbf{x}_{w} \mid \mathbf{W} \mathbf{z}_{w}, \operatorname{diag}(\psi)^{-1}) \prod_{a=1}^{D} q^{*}(\mathbf{w}_{d}) \prod_{a=1}^{N} q^{*}(\mathbf{z}_{a}) \prod_{d=1}^{D} q^{*}(\psi_{d})$$

$$= \mathcal{N}_{\mathcal{D}}(\mathbf{x}_{w} \mid \mathbf{W} \mathbf{z}_{w}, \operatorname{diag}(\psi)^{-1}) \mathcal{N}_{K}(\mathbf{w}_{d} \mid \mathbf{0}, \operatorname{ol}) \mathcal{N}_{K}(\mathbf{z}_{w} \mid \mathbf{0}, \operatorname{ol}) \mathcal{N}_{G}(\mathbf{z}_{w} \mid \mathbf{0}, \operatorname{ol}) \mathcal{N}_{G}(\mathbf{w}_{d} \mid \mathbf{0}$$

 $C_{u,v} = [\log a^*(y, \mathbf{Z}, \mathbf{W}, \mathbf{X})] \propto \mathcal{N}(y_v, \mathbf{K}_v)$ 

(24)

$$\mathbb{E}_{\psi, \mathbf{z}} \left[ \log q^*(\psi, \mathbf{Z}, \mathbf{W}, \mathbf{X}) \right] = \mathbb{E}_{\psi, \mathbf{z}} \left[ \log \left( \prod_{n=1}^N \mathcal{N}_D(\mathbf{x}_n \mid \mathbf{W} \mathbf{z}_n, \operatorname{diag}(\psi)^{-1}) \right) + \log \left( \mathcal{N}_K(\mathbf{w}_d \mid \mathbf{0}, \alpha \mathbf{I}) \right) \right]$$
(26)

$$+\log\left(\mathcal{N}_{K}(\mathbf{z}_{n}\mid\mathbf{0},\mathbf{I})\right) + \log\left(\operatorname{Gamma}(\psi_{d}\mid a,b)\right)\right] \tag{27}$$

(28)

$$\propto \mathbb{E}_{\psi, \mathbf{z}} \left[ \sum_{n=1}^{N} \log \left( \mathcal{N}_{D}(\mathbf{x}_{n} \mid \mathbf{W} \mathbf{z}_{n}, \operatorname{diag}(\psi)^{-1}) \right) + \log \left( \mathcal{N}_{K}(\mathbf{w}_{d} \mid \mathbf{0}, \alpha \mathbf{I}) \right) \right]$$

$$\log \left( \mathcal{N}_D(\mathbf{x}_n \mid \mathbf{W} \mathbf{z}_n, \operatorname{diag}(\psi)^{-1}) \right) \propto -\frac{1}{2} (\mathbf{x}_n - \mathbf{W} \mathbf{z}_n)^{\top} \operatorname{diag}(\psi) (\mathbf{x}_n - \mathbf{W} \mathbf{z}_n)$$
(29)

$$\log \left( \mathcal{N}_K(\mathbf{z}_n \mid \mathbf{0}, \mathbf{I}) \right) \propto -\frac{1}{2} \mathbf{w_d}^{\top} (\alpha \mathbf{I}) \mathbf{w_d}$$
(30)

$$\mathbb{E}_{\psi, \mathbf{z}} \left[ \log q^*(\psi, \mathbf{Z}, \mathbf{W}, \mathbf{X}) \right] \propto \mathbb{E}_{\psi, \mathbf{z}} \left[ -\frac{1}{2} (\mathbf{x_n} - \mathbf{W} \mathbf{z}_n)^{\top} \operatorname{diag}(\psi) (\mathbf{x_n} - \mathbf{W} \mathbf{z}_n) - \frac{1}{2} \mathbf{w_d}^{\top} (\alpha \mathbf{I}) \mathbf{w_d} \right]$$
(31)