

Advanced probabilistic methods - Sketch

Christian Segercrantz 481056

March 29, 2022

$$-\log p(\mathcal{D} \mid \mathbf{W}) = -\log \prod_{i=0}^1 \mathcal{N}(y_i \mid w_0 + w_1 x_i, \sigma_l^2) \quad (1)$$

$$= \sum_{i=0}^N -\log \mathcal{N}(y_i \mid w_0 + w_1 x_i, \sigma_l^2) \quad (2)$$

$$-\log \mathcal{N}(y_i \mid w_0 + w_1 x_i, \sigma_l^2) = -\log \frac{1}{\sigma_l \sqrt{2\pi}} + \frac{1}{2\sigma_l^2} (y_i - w_0 - w_1 x_i)^2 \quad (3)$$

$$= \log \sigma_l \sqrt{2\pi} + \frac{1}{2\sigma_l^2} (y_i - w_0 - w_1 x_i)^2 \quad (4)$$

$$-\log p(\mathcal{D} \mid \mathbf{W}) = \sum_{i=0}^N \log \sigma_l \sqrt{2\pi} + \frac{1}{2\sigma_l^2} \sum_{i=0}^N (y_i - w_0 - w_1 x_i)^2 \quad (5)$$

$$= N \log \sigma_l \sqrt{2\pi} + \frac{1}{2\sigma_l^2} \sum_{i=0}^N (y_i - w_0 - w_1 x_i)^2, \quad \log \sigma_l \sqrt{2\pi} \approx 0.001 \quad (6)$$

$$\approx \frac{1}{2\sigma_l^2} \sum_{i=0}^N (y_i - w_0 - w_1 x_i)^2 \quad (7)$$

Since the prior $p(\mathbf{w})$ is a MVN with diagonal covariance and that the mean-field approximation of the posterior $q(\mathbf{w})$ is product of gaussians we can write the KL-divergence as a sum of the separate parts

$$\text{KL}[q(\mathbf{w})|p(\mathbf{w})] = \text{KL}_0[q(w_0)|p(w_0)] + \text{KL}_1[q(w_1)|p(w_1)] \quad (8)$$

$$\text{KL}_i[q(w_i) \mid p(w_i)] = \int -q(w_i) \log \frac{p(w_i)}{q(w_i)} dw_i \quad (9)$$

$$= \int q(w_i) \log q(w_i) dw_i - \int q(w_i) \log p(w_i) dw_i \quad (10)$$

$$q(w_i) = \mathcal{N}(w_i \mid \mu_i, \sigma_i^2) \propto \exp\left(-\frac{1}{2\sigma_i^2} (w_i - \mu_i)^2\right) \quad (11)$$

$$\log q(w_i) \propto -\frac{1}{2\sigma_i^2} (w_i - \mu_i)^2 \quad (12)$$

$$p(w_i) = \mathcal{N}(w_i \mid 0, \alpha) \propto \exp\left(-\frac{1}{2\alpha^2} w_i^2\right) \quad (13)$$

$$\log p(w_i) = -\frac{1}{2\alpha^2} w_i^2 \quad (14)$$

$$\text{KL}_i[q(w_i) \mid p(w_i)] \propto \int \exp\left(-\frac{1}{2\sigma_i^2} (w_i - \mu_i)^2\right) - \frac{1}{2\sigma_i^2} (w_i - \mu_i)^2 dw_i \quad (15)$$

$$- \int \exp\left(-\frac{1}{2\sigma_i^2} (w_i - \mu_i)^2\right) - \frac{1}{2\alpha^2} w_i^2 dw_i \quad (16)$$

$$= \int -\frac{1}{2\sigma_i^2} (w_i - \mu_i)^2 dw_i + \int \frac{1}{2\alpha^2} w_i^2 dw_i \quad (17)$$

$$= \int -\frac{1}{2\sigma_i^2} (w_i - \mu_i)^2 + \frac{1}{2\alpha^2} w_i^2 dw_i \quad (18)$$

$$= \frac{1}{2} \int -\frac{1}{\sigma_i^2} (w_i - \mu_i)^2 + \frac{1}{\alpha^2} w_i^2 dw_i \quad (19)$$

$$(20)$$