



Stegosaurus

Stegosaurus (/ˌstɛɡəˈsɔːrəs/^[1] lit. ˈroɒfˈliːzɑːd) is a genus of herbivorous, four-legged, armored dinosaur from the Late Jurassic, characterized by the distinctive kite-shaped upright plates along their backs and spikes on their tails. Fossils of the genus have been found in the western United States and in Portugal, where they are found in Kimmeridgian- to early Tithonian-aged strata, dating to between 155 and 145 million years ago. Of the species that have been classified in the upper Morrison Formation of the western US, only three are universally recognized: *S. stenops*, *S. ungulatus* and *S. sulcatus*. The remains of over 80 individual animals of this genus have been found. *Stegosaurus* would have lived alongside dinosaurs such as *Apatosaurus*, *Diplodocus*, *Brachiosaurus*, ``*Ceratosaurus*`, *Allosaurus*, and *Torvosaurus*; the latter two may have preyed on it.

They were large, heavily built, herbivorous quadrupeds with rounded backs, short fore limbs, long hind limbs, and tails held high in the air. Due to their distinctive combination of broad, upright plates and tail tipped with spikes, *Stegosaurus* is one of the most recognizable kinds of dinosaurs. The function of this array of plates and spikes has been the subject of much speculation among scientists. Today, it is generally agreed that their spiked tails were most likely used for defense against predators, while their plates may have been used primarily for display, and secondarily for thermoregulatory functions. *Stegosaurus* had a relatively low brain-to-body mass ratio. It had a short neck and a small head, meaning it most likely ate low-lying bushes and shrubs. One species, *Stegosaurus ungulatus*, is the largest known of all the stegosaurians (bigger than related dinosaurs such as *Kentrosaurus* and *Huayangosaurus*).

Stegosaurus remains were first identified during the "Bone Wars" by Othniel Charles Marsh at Dinosaur Ridge National Landmark. The first known skeletons were fragmentary and the bones were scattered, and it would be many years before the true appearance of these animals, including their posture and plate arrangement, became well understood. Despite its popularity in books and film, mounted skeletons of *Stegosaurus* did not become a staple of major natural history museums until the mid-20th century, and many museums have had to assemble composite displays from several different specimens due to a lack of complete skeletons. *Stegosaurus* is one of the better-known dinosaurs, and has been featured in film, postal stamps, and many other types of media.

| Stegosaurus |
|--|
| Temporal range: <u>Late Jurassic</u> (<u>Kimmeridgian</u> to <u>Tithonian</u>), |
|  |
| Mounted skeleton of <i>S. stenops</i> (nicknamed "Sophie"), <u>Natural History</u> <u>Museum, London</u> |
| Scientific classification  |
| Kingdom: <u>Animalia</u> |
| Phylum: <u>Chordata</u> |
| Clade: <u>Dinosauria</u> |
| Order: † <u>Ornithischia</u> |
| Clade: † <u>Thyreophora</u> |
| Suborder: † <u>Stegosauria</u> |
| Family: † <u>Stegosauridae</u> |
| Subfamily: † <u>Stegosaurinae</u> |
| Genus: † <u>Stegosaurus</u> <u>Marsh, 1877</u> |
| Type species |
| † <i>Stegosaurus stenops</i> Marsh, 1887 |
| Other species |
| <ul style="list-style-type: none">▪ †<i>S. ungulatus</i> Marsh, 1879▪ †<i>S. sulcatus</i> Marsh, 1887 |
| Synonyms |
| <ul style="list-style-type: none">▪ <u><i>Hypsirhophus</i></u>? Cope, 1878 |

Contents

Discovery and history

[Bone Wars and *Stegosaurus armatus*](#)

[Early skeletal mounts and plate interpretation](#)

[Plate arrangement](#)

[Second Dinosaur Rush](#)

[Resurgent Discoveries](#)

Description

[Skull](#)

[Skeleton](#)

[Plates](#)

Classification and species

[Species](#)

[Doubtful species and junior synonyms](#)

[Reassigned species](#)

Paleobiology

[Posture and movement](#)

[Plate function](#)

[Thagomizer \(tail spikes\)](#)

["Second brain"](#)

[Growth](#)

[Diet](#)

Paleoecology

In popular culture

See also

References


External links

Discovery and history

Bone Wars and *Stegosaurus armatus*

Stegosaurus, one of the many dinosaurs described in the [Bone Wars](#), was first collected by [Arthur Lakes](#) and consisted of several caudal vertebrae, a dermal plate, and several additional postcranial elements that were collected north of [Morrison, Colorado](#) at Lakes' YPM Quarry 5.^[2] These first, fragmented bones (YPM 1850) became the [holotype](#) of *Stegosaurus armatus* when Yale paleontologist Othniel Charles Marsh described them in 1877.^[3] Marsh initially believed the remains were from an aquatic turtle-like animal, and the basis for its [scientific name](#), 'roof(ed) lizard' was due to his early belief that the plates lay flat over the animal's back, overlapping like the [shingles \(tiles\)](#) on a roof.^[3] Though several more complete specimens have been attributed to *Stegosaurus armatus*, preparation of the bones and analysis has discovered that this type specimen is actually dubious, which is not an ideal situation for the type species of a well-known genus like *Stegosaurus*.^[2] Because of this, the [International Code of Zoological](#)

Nomenclature decided to replace the type species with the more well known species *Stegosaurus stenops*.^[4] Marsh also incorrectly referred several fossils to *S. armatus*, including the dentary and teeth of the sauropod *Diplodocus* and putting sauropod limb bones and an *Allosaurus* tibia under YPM 1850.^[5]



Marsh's 1891 illustration of *S. unguatus*. Note the single row of 12 large rounded plates, based on those of *S. stenops*, and eight spikes

On the other side of the Bone Wars, Edward Drinker Cope named *Hypsirhophus discurus* as another stegosaurian based on fragmentary fossils from Cope's Quarry 3 near the "Cope's Nipple" site in Garden Park, Colorado in 1878.^[6] Many later researchers have considered *Hypsirhophus* to be a synonym of *Stegosaurus*,^[7] though Peter Galton (2010) suggested that it is distinct based on differences in the vertebrae.^[2] F. F. Hubbell, a collector for Cope, also found a partial *Stegosaurus* skeleton while digging at Como Bluff in 1877 or '78 that are now part of the *Stegosaurus* mount (AMNH 5752) at the American Museum of Natural History.^[8]

Arthur Lakes made another discovery later in 1879 at Como Bluff in Albany County, Wyoming, the site also dating to the Upper Jurassic of the Morrison Formation, when he found several large *Stegosaurus* fossils in August of that year.^[5] The majority of the fossils came from Quarry 13, including the type specimen of *Stegosaurus unguatus* (YPM 1853), which was collected by Lakes and William Harlow Reed the same year and named by Marsh.^[5] The specimen was one of many found at the quarry, the specimen consisting of a partial skull, several vertebrae, an ischium, partial limbs, several plates, and four thagomizers, though eight thagomizers were referred based on a specimen preserved alongside the type.^[5] The type specimen also preserved the pes, which was the namesake of the species, meaning "hoofed roofed lizard".^{[9][2]} In 1881, he named a third species *Stegosaurus "affinis"*, based only on a hip bone, though the fossil has since been lost and the species declared a nomen nudum.^{[5][2]} Later in 1887, Marsh two more species of *Stegosaurus* from Como Bluff, *Stegosaurus duplex*, based on a partial vertebral column, partial pelvis, and partial left hindlimb (YPM 1858) from Reed's Quarry 11, though the species is now seen as synonymous with *Stegosaurus unguatus*.^[7] The other, *Stegosaurus sulcatus*, was named based on a left forelimb, scapula, left femur, several vertebrae, and several plates and dermal armor elements (USNM V 4937) collected in 1883.^{[7][2]} *Stegosaurus sulcatus* most notably preserves a large spike that has been speculated to have been a shoulder spike that is used to diagnose the species.^[2]



Type specimen of *S. stenops* on display at the National Museum of Natural History.

The greatest *Stegosaurus* discovery came in 1885 with the discovery of a nearly complete, articulated skeleton of a subadult that included previously undiscovered elements like a complete skull, throat ossicles, and articulated plates.^{[9][7]} Marshall P. Felch collected the skeleton throughout 1885 and 1886 from Morrison Formation strata at his quarry in Garden Park, a town near Cañon City, Colorado.^{[10][7]} The skeleton was expertly unearthed by Felch, who first divided the skeleton into labeled blocks and prepared them separately.^[7] The skeleton was shipped to Marsh in 1887, who named it *Stegosaurus stenops* ("narrow-faced roof lizard") that year. Though it had not yet been completely prepared, the nearly complete and articulated type specimen of *Stegosaurus*

stenops allowed Marsh to complete the first attempt at a reconstructed *Stegosaurus* skeleton.^{[9][11]} This first reconstruction, of *S. unguatus* with missing parts filled in from *S. stenops*, was published by Marsh in 1891. (In 1893, Richard Lydekker mistakenly re-published Marsh's drawing under the label *Hypsirhophus*.)^[7]

Early skeletal mounts and plate interpretation

The skeleton of *S. stenops* has since been deposited at the National Museum of Natural History in Washington D. C., where it has been on display since 1915.^[12] Another mount was made for the NMNH in the form of a mounted composite skeleton consisting of several specimens referred to *S. stenops* that were collected at Quarry 13 at Como Bluff in 1887, the most complete being USNM 6531.^[12] The type specimen of *S. ungulatus* (YPM 1853) was incorporated into the first ever mounted skeleton of a stegosaur at the Peabody Museum of Natural History in 1910 by Richard Swann Lull. It was initially mounted with paired plates set wide, above the base of the ribs, but was remounted in 1924 with two staggered rows of plates along the midline of the back.^[13] Additional specimens recovered from the same quarry by the United States National Museum of Natural History, including tail vertebrae and an additional large plate (USNM 7414), belong to the same individual as YPM 1853.^[2]

The next species of *Stegosaurus* to be named was *S. marshi* by Frederick Lucas in 1901. Lucas reclassified this species in the new genus *Hoplitosaurus* later that year. Lucas also re-examined the issue of the life appearance of *Stegosaurus*, coming to the conclusion that the plates were arranged in pairs in two rows along the back, arranged above the bases of the ribs. Lucas commissioned Charles R. Knight to produce a life restoration of *S. ungulatus* based on his new interpretation. However, the following year, Lucas wrote that he now believed the plates were probably attached in staggered rows. In 1910, Richard Swann Lull wrote that the alternating pattern seen in *S. stenops* was probably due to shifting of the skeleton after death. He led the construction of the first ever *Stegosaurus* skeletal mount at the Peabody Museum of Natural History, which was depicted with paired plates. In 1914, Charles Gilmore argued against Lull's interpretation, noting that several specimens of *S. stenops*, including the now-completely prepared holotype, preserved the plates in alternating rows near the peak of the back, and that there was no evidence of the plates having shifted relative to the body during fossilization.^[7] Gilmore and Lucas' interpretation became the generally accepted standard, and Lull's mount at the Peabody Museum was changed to reflect this in 1924.^[13]



1901 life restoration of *S. ungulatus* by Charles R. Knight with paired dorsal plates and eight tail spikes

Plate arrangement

One of the major subjects of books and articles about *Stegosaurus* is the plate arrangement.^[14] The argument has been a major one in the history of dinosaur reconstruction. Four possible plate arrangements have been proposed over the years:

- The plates lie flat along the back, as a shingle-like armor. This was Marsh's initial interpretation, which led to the name 'roof lizard'. As further and complete plates were found, their form showed they stood on edge, rather than lying flat.
- By 1891, Marsh published a more familiar view of *Stegosaurus*,^[11] with a single row of plates. This was dropped fairly early on (apparently because it was poorly understood how the plates were embedded in the skin and they were thought to overlap too much in this arrangement). It was revived, in somewhat modified form, in the 1980s, by Stephen Czerkas,^[15] based on the arrangement of iguana dorsal spines.
- The plates were paired in a double row along the back, such as in Knight's 1901 reconstruction and the 1933 film King Kong.
- Two rows of alternating plates. By the early 1960s, this had become (and remains) the prevalent idea, mainly because some *S. stenops* fossils in which the plates are still partially

articulated show this arrangement. This arrangement is chiral and so demands that a specimen be distinguished from its distinct, hypothetical mirror-image form.^{[16][17]}

Second Dinosaur Rush



Mounted composite skeleton referred to *S. unguulatus*, Carnegie Museum of Natural History

After the end of the Bone Wars, many major institutions in the eastern United States were inspired by the depictions and finds by Marsh and Cope to assemble their own dinosaur fossil collections. The competition was foremost started by the American Museum of Natural History, Carnegie Museum of Natural History, and Field Museum of Natural History which all sent expeditions to the west to make their own dinosaur collections and mount skeletons in their fossil halls.^[18] The American Museum of Natural History was the first to launch an expedition in 1897, finding several assorted, but incomplete, *Stegosaurus* specimens at Bone Cabin Quarry in Como Bluff.^{[18][8]} These remains haven't been described and were mounted in 1932, the mount being a composite primarily of specimens AMNH 650 & 470 from Bone Cabin Quarry.^{[8][19]} The

AMNH mount is cast and on display at the Field Museum, which didn't collect any *Stegosaurus* skeletons during the Second Dinosaur Rush.^[19] The Carnegie Museum in Pittsburgh on the other hand collected many *Stegosaurus* specimens, first at Freezout Hills in Carbon County, Wyoming in 1902–03. The fossils included only a couple postcranial remains, though in the 1900s-1920s Carnegie crews at Dinosaur National Monument discovered dozens of *Stegosaurus* specimens in one of the greatest single sites for the taxon.^[20] CM 11341, the most complete skeleton found at the quarry, was used for the basis of a composite *Stegosaurus* mount in 1940 along with several other specimens to finish the mount. A cranium (CM 12000) was also found by Carnegie crews, one of the few known.^[20] Both the AMNH and CM material has been referred to *Stegosaurus unguulatus*.^{[20][5]}


Resurgent Discoveries

As part of the Dinosaur Renaissance and the resurgent interest in dinosaurs by museums and the public, fossils of *Stegosaurus* were once again being collected, though few have been fully described. An important discovery came in 1937 again at Garden Park by a high school teacher named Frank Kessler in 1937 while leading a nature hike. Kessler contacted the Denver Museum of Nature and Science, who sent paleontologist Robert Landberg.^[21] Landberg excavated the skeleton with the DMNS crews, recovering a 70% complete *Stegosaurus* skeleton along with turtles, crocodiles, and isolated dinosaur fossils at the quarry that would be nicknamed "The Kessler Site".^[21] Phillip Reinheimer, a steel worker, mounted the *Stegosaurus* skeleton at the DMNS in 1938. The skeleton remained mounted until 1989 when the museum curator of the DMNS began a revision of the museum's fossil hall and dispatched an expedition to find additional *Stegosaurus* remains.^[21] The expedition was successful in finding a nearly complete *Stegosaurus* near the Kessler site by Bryan Small, whose name would become the namesake of the new site.^{[21][22]} The "Small Quarry" *Stegosaurus*' articulation and completeness clarified the position of plates and spikes on the back of *Stegosaurus* and the position and size of the throat ossicles found earlier first by Felch with the *Stegosaurus stenops* holotype, though like the *S. stenops* type, the fossils were flattened in a "roadkill" condition.^{[23][22][21]} The *Stegosaurus* skeletons have been mounted alongside an *Allosaurus* skeleton collected in Moffat County, Colorado originally in 1979.^[21]

1987 saw the discovery of a 40% complete *Stegosaurus* skeleton in Rabbit Valley in Mesa County, Colorado by Harold Bollen near the Dinosaur Journey Museum.^[24] The skeleton was nicknamed the "Bollen *Stegosaurus*" and is in the collections of the Dinosaur Journey Museum.^[24] At Jensen-Jensen Quarry, an articulated torso including several dorsal plates from a small individual were collected and briefly described in 2014, though the specimen was collected years before and is still in preparation at Brigham Young University.^[25] 2007 saw the description of a *Stegosaurus* specimen from the Upper Jurassic Lourinha Formation of Portugal, the specimen was placed as cf. *Stegosaurus unguatus* by the describers.^[26] The specimen is one of the few associated *Stegosaurus* skeletons known, though it only contains a tooth, 13 vertebrae, partial limbs, a cervical plate, and several assorted postcranial elements.^{[26][23]}

Sophie the *Stegosaurus* is the best preserved *Stegosaurus* specimen, being 85% intact and containing 360 bones. Sophie was first discovered by Bob Simon in 2003 at the Red Canyon Quarry near Shell, Wyoming and was excavated by crews from the Swiss Sauriermuseum in 2004.^{[23][27]} The skeleton was excavated on private land, so it was interned by US federal authorities who then gave Sophie to the Natural History Museum, London where it was put on display in December of 2014 and later described in 2015.^[23] It is a young adult of undetermined sex, 5.8 m (19 ft) long and 2.9 m (9.5 ft) tall.^[23] The Sauriermuseum found several partial Stegosaurid skeletons throughout their excavations at Howe Quarry, Wyoming in the 1990s, though only Sophie has been described in detail. One skeleton collected at the site known as "Victoria" is very well preserved including many of the vertebrae preserved in semi-articulation and next to an *Allosaurus* skeleton found nicknamed "Big Al II".^[27]

Description



The quadrupedal *Stegosaurus* is one of the most easily identifiable dinosaur genera, due to the distinctive double row of kite-shaped plates rising vertically along the rounded back and the two pairs of long spikes extending horizontally near the end of the tail. Although large individuals (YPM Coll^[28]) could grow up to 9 m (29.5 ft) in length^[29] and 5.3–7 metric tons (5.8–7.7 short tons) in body mass,^{[30][31]} the various species of *Stegosaurus* were dwarfed by contemporaries, the giant sauropods. In 2016, Gregory S. Paul

presented a lower estimate of 6.5 m (21.3 ft) in length and 3.5 metric tons (3.9 short tons) in body mass for *S. stenops*, and 7 m (23.0 ft) in length and 3.8 metric tons (4.2 short tons) in body mass for *S. unguatus*.^[32] Some form of armor appears to have been necessary, as *Stegosaurus* species coexisted with large predatory theropod dinosaurs, such as *Allosaurus*, *Ceratosaurus*, *Saurophaganax* and *Torvosaurus*.

Most of the information known about *Stegosaurus* comes from the remains of mature animals; more recently, though, juvenile remains of *Stegosaurus* have been found. One subadult specimen, discovered in 1994 in Wyoming, is 4.6 m (15.1 ft) long and 2 m (6.6 ft) high, and is estimated to have weighed 1.5–2.2 metric tons (1.6–2.4 short tons)^[33] while alive. It is on display in the University of Wyoming Geological Museum.^[34]

Skull

The long and narrow skull was small in proportion to the body. It had a small antorbital fenestra, the hole between the nose and eye common to most archosaurs, including modern birds, though lost in extant crocodylians. The skull's low position suggests that *Stegosaurus* may have been a browser of low-growing vegetation. This interpretation is supported by the absence of front teeth and their likely replacement by a horny beak or rhamphotheca. The lower jaw had flat downward and upward extensions that would have

completely hidden the teeth when viewed from the side, and these probably supported a turtle-like beak in life. The presence of a beak extended along much of the jaws may have precluded the presence of cheeks in these species.^[35] Such an extensive beak was probably unique to *Stegosaurus* and some other advanced stegosaurids among ornithischians, which usually had beaks restricted to the jaw tips.^{[36][37]} Other researchers have interpreted these ridges as modified versions of similar structures in other ornithischians which might have supported fleshy cheeks, rather than beaks.^[38] Stegosaurian teeth were small, triangular, and flat; wear facets show that they did grind their food.^[39]



S. stenops skull cast, Natural History Museum of Utah

Despite the animal's overall size, the braincase of *Stegosaurus* was small, being no larger than that of a dog. A well-preserved *Stegosaurus* braincase allowed Othniel Charles Marsh to obtain, in the 1880s, a cast of the brain cavity or endocast of the animal, which gave an indication of the brain size. The endocast showed the brain was indeed very small, the smallest proportionally of all dinosaur endocasts then known. The fact that an animal weighing over 4.5 metric tons (5 short tons) could have a brain of no more than 80 g (2.8 oz) contributed to the popular old idea that all dinosaurs were unintelligent, an idea now largely rejected.^[40] Actual brain anatomy in *Stegosaurus* is poorly known, but the brain itself was small even for a dinosaur.^[41]

Skeleton

In *Stegosaurus stenops* there are 27 bones in the vertebral column anterior to the sacrum, a varying number of vertebrae in the sacrum, with four in most subadults, and around 46 caudal (tail) vertebrae. The presacrals are divided into cervical (neck) and dorsal (back) vertebrae, with around 10 cervicals and 17 dorsals, the total number being one greater than in *Hesperosaurus*, two greater than *Huayangosaurus*, although *Miragaia* preserves 17 cervicals and an unknown number of dorsals. The first cervical vertebra is the axis bone, which is connected and often fused to the atlas bone. Farther posteriorly, the proportionately larger the cervicals become, although they do not change greatly in anything other than size. Past the first few dorsals, the centrum of the bones become more elongate front-to-back, and the transverse processes become more elevated dorsal. The sacrum of *S. stenops* includes four sacral vertebrae, but one of the dorsals is also incorporated into the structure. In some specimens of *S. stenops*, a caudal is also incorporated, as a caudosacral. In *Hesperosaurus* there are two dorsosacrals, and only four fused sacrals, but in *Kentrosaurus* there may be as many as seven vertebrae in the sacrum, with both dorsosacrals and caudosacrals. *S. stenops* preserves 46 caudal vertebrae, and up to 49, and along the series both the centra and the neural spines become smaller, until the neural spines disappear at caudal 35. Around the middle of the tail, the neural spines become bifurcated, meaning they are divided near the top.^[23]

With multiple well-preserved skeletons, *S. stenops* preserves all regions of the body, including the limbs. The scapula (shoulder blade) is sub-rectangular, with a robust blade. Though it is not always perfectly preserved, the acromion ridge is slightly larger than in *Kentrosaurus*. The blade is relatively straight, although it curves towards the back. There is a small bump on the back of the blade, that would have served as the base of the triceps muscle. Articulated with the scapula, the coracoid is sub-circular.^[23] The hind feet each had three short toes, while each fore foot had five toes; only the inner two toes had a blunt hoof. The phalangeal formula is 2-2-2-2-1, meaning the innermost finger of the fore limb has two bones, the next has two, etc.^[42] All four limbs were supported by pads behind the toes.^[43] The fore limbs were much shorter than the stocky hind limbs, which resulted in an unusual posture. The tail appears to have been held well clear of the ground, while the head of *Stegosaurus* was positioned relatively low down, probably no higher than 1 m (3.3 ft) above the ground.^[22]

Plates

The most recognizable features of *Stegosaurus* are its dermal plates, which consisted of between 17 and 22 separate plates and flat spines.^[13] These were highly modified osteoderms (bony-cored scales), similar to those seen in crocodiles and many lizards today. They were not directly attached to the animal's skeleton, instead arising from the skin. The largest plates were found over the hips and could measure over 60 cm (24 in) wide and 60 cm (24 in) tall.^[13]




Life restoration of *S. stenops* based on the "Sophie" specimen

In a 2010 review of *Stegosaurus* species, Peter Galton suggested that the arrangement of the plates on the back may have varied between species, and that the pattern of plates as viewed in profile may have been important for species recognition. Galton noted that the plates in *S. stenops* have been found articulated in two staggered rows, rather than paired. Fewer *S. ungulatus* plates have been found, and none articulated, making the arrangement in this species more difficult to determine. However, the type specimen of *S. ungulatus* preserves two flattened spine-like plates from the tail that are nearly identical in shape and size, but are mirror images of each other, suggesting that at least these were arranged in pairs.^[2] Many of the plates are manifestly chiral^{[16][17]} and no two plates of the same size and shape have been found for an individual; however plates have been correlated between individuals. Well preserved integumentary impressions of the plates of *Hesperosaurus* show a smooth surface with long and parallel, shallow grooves. This indicates that the plates were covered in keratinous sheaths.^[44]

Classification and species

Like the spikes and shields of ankylosaurs, the bony plates and spines of stegosaurians evolved from the low-keeled osteoderms characteristic of basal thyreophorans.^[45] Galton (2019) interpreted plates of an armored dinosaur from the Lower Jurassic (Sinemurian-Pliensbachian) Lower Kota Formation of India as fossils of a member of Ankylosauria; the author argued that this finding indicates a probable early Early Jurassic origin for both Ankylosauria and its sister group Stegosauria.^[46]





Restoration of *S. stenops* with alternating plate arrangement.

The vast majority of stegosaurian dinosaurs thus far recovered belong to the Stegosauridae, which lived in the later part of the Jurassic and early Cretaceous, and which were defined by Paul Sereno as all stegosaurians more closely related to *Stegosaurus* than to *Huayangosaurus*.^[47] This group is widespread, with members across the Northern Hemisphere, Africa and possibly South America.^[48]

Stegosaurus frequently is discovered in its own clade in Stegosauridae called Stegosauridae, usually including the taxa Wuerhosaurus and Loricatosaurus,^[49] though Hesperosaurus is sometimes found in the group.^[50] In 2017, Raven and Maidment published a new phylogenetic analysis, including almost every known stegosaurian genus:^{[51][52]}

Thyreophora





Species

Many of the species initially described have since been considered to be invalid or synonymous with earlier named species,^[5] leaving two well-known and one poorly known species. Confirmed *Stegosaurus* remains have been found in the Morrison Formation's stratigraphic zones 2–6, with additional remains possibly referable to *Stegosaurus* recovered from stratigraphic zone 1.^[53]

- *Stegosaurus ungulatus*, meaning "hoofed roof lizard", was named by Marsh in 1879,^[9] from remains recovered at Como Bluff, Wyoming (Quarry 12, near Robber's Roost).^[5] It might be synonymous with *S. stenops*.^[23] At 9 m (29.5 ft), it was the longest species within the genus *Stegosaurus*. A fragmentary *Stegosaurus* specimen discovered in Portugal and dating from the upper Kimmeridgian-lower Tithonian stage has been tentatively assigned to this species.^[26] *Stegosaurus ungulatus* can be distinguished from *S. stenops* by the presence of longer hind limbs, proportionately smaller, more pointed plates with wide bases and narrow tips, and by several small, flat, spine-like plates just before the spikes on the tail. These spine-like plates appear to have been paired, due to the presence of at least one pair that are identical but mirrored. *S. ungulatus* also appears to have had longer legs (femora) and hip bones than other species. The type specimen of *S. ungulatus* was discovered with eight spikes, though they were scattered away from their original positions. These have often been interpreted as indicating that the animal had four pairs of tail spikes. No specimens have been found with complete or articulated sets of tail spikes, but no additional specimens have been found that preserve eight spikes together. It is possible the extra pair of spikes came from a different individual, and though no other extra bones were found with the specimen, these may be found if more digging were done at the original site.^[2] Specimens from other quarries (such as a tail from Quarry 13, now forming part of the composite skeleton AMNH 650 at the American Museum of Natural History), referred to *S. ungulatus* on the basis of their notched tail vertebrae, are preserved with only four tail spikes.^[5] The type specimen of *S. ungulatus* (YPM 1853) was incorporated into the first ever mounted skeleton of a stegosaur at the Peabody Museum of Natural History in 1910 by Richard Swann Lull. It was initially mounted with paired plates set wide, above the base of the ribs, but was remounted in 1924 with two staggered rows of plates along the midline of the back.^[13] Additional specimens recovered from the same quarry by the United States National Museum of Natural History, including tail vertebrae and an additional large plate (USNM 7414), belong to the same individual as YPM 1853.^[2]
- *Stegosaurus stenops*, meaning "narrow-faced roof lizard", was named by Marsh in 1887,^[10] with the holotype having been collected by Marshall Felch at Garden Park, north of Cañon City, Colorado, in 1886. This is the best-known species of *Stegosaurus*, mainly because its remains include at least one complete articulated skeleton. It had proportionately large, broad plates and rounded tail plates. Articulated specimens show that the plates were arranged alternating in a staggered double row. *S. stenops* is known from at least 50 partial skeletons of adults and juveniles, one complete skull, and four partial skulls. It was shorter than other species, at 7 m (23 ft). Found in the Morrison Formation, Colorado, Wyoming, and Utah.^[38]



Articulated holotype of *S. stenops* (USNM 4934), known as the "road kill" specimen

- *Stegosaurus sulcatus*, meaning "furrowed roof lizard", was described by Marsh in 1887 based on a partial skeleton.^[10] It has traditionally been considered a synonym of *S. armatus*,^[38] though more recent studies suggest it is not.^[2] *S. sulcatus* is distinguished mainly by its unusually large, furrowed spikes with very large bases. A spike associated with the type specimen, originally thought to be a tail spike, may in fact come from the shoulder or hip, since its base is much larger than the corresponding tail vertebrae. A review published by Maidment and colleagues in 2008 regarded it as an indeterminate species possibly not even belonging to *Stegosaurus* at all, but to a different genus.^{[19][54]} Peter Galton suggested it should be considered a valid species due to its unique spikes.^[2]

Susannah Maidment and colleagues in 2008 proposed extensive alterations to the taxonomy of *Stegosaurus*. They advocated synonymizing *S. stenops* and *S. unguatus* with *S. armatus*, and sinking *Hesperosaurus* and *Wuerhosaurus* into *Stegosaurus*, with their type species becoming *Stegosaurus mjosi* and *Stegosaurus homheni*, respectively. They regarded *S. longispinus* as dubious. Thus, their conception of *Stegosaurus* would include three valid species (*S. armatus*, *S. homheni*, and *S. mjosi*) and would range from the Late Jurassic of North America and Europe to the Early Cretaceous of Asia.^[19] However, this classification scheme was not followed by other researchers, and a 2017 cladistic analysis co-authored by Maidment with Thomas Raven rejects the synonymy of *Hesperosaurus* with *Stegosaurus*.^{[2][55]} In 2015, Maidment *et al.* revised their suggestion due to the recognition by Galton of *S. armatus* as a *nomen dubium* and its replacement by *S. stenops* as type species.^[23]

Doubtful species and junior synonyms

- *Stegosaurus armatus*, meaning "armored roof lizard", was the first species to be found and the original type species named by O.C. Marsh in 1877.^[3] It is known from a partial skeleton, and more than 30 fragmentary specimens have been referred to it.^[38] However, the type specimen was very fragmentary, consisting only of a partial tail, hips, and leg, parts of some back vertebrae, and a single fragmentary plate (the presence of which was used to give the animal its name). No other plates or spikes were found, and the entire front half of the animal appears not to have been preserved.^[5] Because the type specimen is very fragmentary, it is extremely difficult to compare it with other species based on better specimens, and it is now generally considered to be a *nomen dubium*. Because of this, it was replaced by *S. stenops* as the type species of *Stegosaurus* in a ruling of the ICZN in 2013.^[4]
- *Stegosaurus* "affinis", named by Marsh in 1881, is only known from a pubis which has since been lost. Because Marsh did not provide an adequate description of the bone with which to distinguish a new species, this name is considered a *nomen nudum*.^[7]
- *Diracodon laticeps* was described by Marsh in 1881, from some jawbone fragments.^[56] Bakker resurrected *D. laticeps* in 1986 as a senior synonym of *S. stenops*,^[57] although others note that the material is not diagnostic and is only referable to *Stegosaurus* sp., making it a *nomen dubium*.^[5]
- *Stegosaurus duplex*, meaning "two plexus roof lizard" (in allusion to the greatly enlarged neural canal of the sacrum which Marsh characterized as a "posterior brain case"), was named by Marsh in 1887 (including the holotype specimen). The disarticulated bones were actually collected in 1879 by Edward Ashley at Como Bluff. Marsh initially distinguished it from *S. unguatus* based on the fact that each sacral (hip) vertebra bore its own rib, which he claimed was unlike the anatomy of *S. unguatus*; however, the sacrum of *S. unguatus* had not actually been discovered. Marsh also suggested that *S. duplex* may have lacked armor, since no plates or spikes were found with the specimen, though a single spike may actually have been present nearby, and re-examination of the site maps has shown that the entire specimen was found highly disarticulated and scattered.^[5] It is generally considered a

synonym of *S. ungulatus* today, and parts of the specimen were actually incorporated into the Peabody Museum *S. ungulatus* skeletal mount in 1910.^[2]

Reassigned species

- *Stegosaurus marshi*, which was described by Lucas in 1901, was renamed *Hoplitosaurus* in 1902.^[58]
- *Stegosaurus priscus*, described by Nopcsa in 1911, was reassigned to *Lexovisaurus*,^[38] and is now the type species of *Loricatosaurus*.^[19]
- *Stegosaurus longispinus* was named by Charles W. Gilmore in 1914 based on a fragmentary postcranial skeleton that has largely been lost.^{[59][7]} It is now the type species of the genus *Alcovasaurus*, though it has been referred to *Miragaia*.^{[60][59]}
- *Stegosaurus madagascariensis* from Madagascar is known solely from teeth and was described by Piveteau in 1926. The teeth were variously attributed to a stegosaur, the theropod *Majungasaurus*,^[61] a hadrosaur or even a crocodylian, but is now considered a possible ankylosaur.^[19]
- *Stegosaurus homheni* is an alternative combination for the Chinese Cretaceous stegosaur *Wuerhosaurus homheni*, which was described based on a partial postcranial skeleton in 1973 by Dong Zhiming.^[62] It was referred to *Stegosaurus* in 2008 by Maidment *et al*,^[19] but some still consider the species to be in its own genus.^{[63][64]}
- *Stegosaurus mjosi* was described as *Hesperosaurus mjosi* by Carpenter *et al* in 2001 based on a partial skull and incomplete postcranial skeleton from the Morrison Formation of Johnson County, Wyoming. The species was referred to *Stegosaurus* mostly by Maidment *et al* starting in 2008,^{[65][19]} but *Hesperosaurus* has been the more popular combination since the discovery of more remains.^[66]

Paleobiology

Posture and movement

Soon after its discovery, Marsh considered *Stegosaurus* to have been bipedal, due to its short forelimbs.^[67] He had changed his mind, however, by 1891, after considering the heavy build of the animal.^[11] Although *Stegosaurus* is undoubtedly now considered to have been quadrupedal, some discussion has occurred over whether it could have reared up on its hind legs, using its tail to form a tripod with its hind limbs, to browse for higher foliage.^[38] This has been proposed by Bakker^{[57][68]} and opposed by Carpenter.^[22] A study by Mallison (2010) found support for a rearing up posture in *Kentrosaurus*, though not for ability for the tail to act as a tripod.^[69]

Stegosaurus had short fore limbs in relation to its hind limbs. Furthermore, within the hind limbs, the lower section (comprising the tibia and fibula) was short compared with the femur. This suggests it could not walk very fast, as the stride of the back legs at speed would have overtaken the front legs, giving a maximum speed of 15.3–17.9 km/h (9.5–11.1 mph).^{[70][39]}

Tracks discovered by Matthew Mossbrucker (Morrison Natural History Museum, Colorado) suggest that



Adult and juvenile *S. stenops* mounted as if under attack from an *Allosaurus fragilis*, Denver Museum of Nature and Science

Stegosaurus lived and traveled in multiple-age herds. One group of tracks is interpreted as showing four or five baby stegosaurs moving in the same direction, while another has a juvenile stegosaur track with an adult track overprinting it.^[71]

As the plates would have been obstacles during copulation, it is possible the female stegosaur laid on her side as the male mounted her from behind. Another suggestion is that the female would stand on all fours but squat the fore limbs and raise the tail slightly to the side as the male rested its fore limbs on her broad pelvis. However, this would not explain how their reproductive organs can touch as there is no evidence of muscle attachments for a mobile penis nor a baculum in dinosaurs.^[72]

Plate function

The function of *Stegosaurus*' plates has been much debated. Marsh suggested that they functioned as some form of armor,^[67] though Davitashvili (1961) disputed this, claiming that they were too fragile and ill-placed for defensive purposes, leaving the animal's sides unprotected.^[73] Nevertheless, others have continued to support a defensive function. Bakker suggested in 1986 that the plates were covered in horn comparing the surface of the fossilized plates to the bony cores of horns in other animals known or thought to bear horns.^[74] Christiansen and Tschopp (2010), having studied a well-preserved specimen of *Hesperosaurus* with skin impressions, concluded that the plates were covered in a keratin sheath which would have strengthened the plate as a whole and provided it with sharp cutting edges.^[44] Bakker stated that *Stegosaurus* could flip its osteoderms from one side to another to present a predator with an array of spikes and blades that would impede it from closing sufficiently to attack the *Stegosaurus* effectively. He contends that they had insufficient width for them to stand erect easily in such a manner as to be useful in display without continuous muscular effort.^[74] Mobility of the plates, however, has been disputed by other paleontologists.^[75]



Back plate cast, Museum of the Rockies

Another possible function of the plates is they may have helped to control the body temperature of the animal,^[75] in a similar way to the sails of the pelycosaurs *Dimetrodon* and *Edaphosaurus* (and modern elephant and rabbit ears). The plates had blood vessels running through grooves and air flowing around the plates would have cooled the blood.^[76] Buffrénil, et al. (1986) found "extreme vascularization of the outer layer of bone",^{[77][75]} which was seen as evidence that the plates "acted as thermoregulatory devices".^[77] Likewise, 2010 structural comparisons of *Stegosaurus* plates to *Alligator* osteoderms seem to support the conclusion that the potential for a thermoregulatory role in the plates of *Stegosaurus* definitely exists.^[78]



Plates of *Stegosaurus* specimen "Sophie"

The thermoregulation hypothesis has been seriously questioned, since other stegosaurs such as *Kentrosaurus*, had more low surface area spikes than plates, implying that cooling was not important enough to require specialized structural formations such as plates. However, it has also been suggested that the plates could have helped the animal increase heat absorption from the sun. Since a cooling trend occurred towards the end of the Jurassic, a large ectothermic reptile might have used the increased surface area afforded by the plates to absorb radiation from the sun.^[79] Christiansen and Tschopp (2010) state that the presence of a smooth, insulating keratin covering would have hampered thermoregulation, but such a function cannot be entirely ruled out

as extant cattle and ducks use horns and beaks to dump excess heat despite the keratin covering.^[44] Histological surveys of plate microstructure attributed the vascularization to the need to transport nutrients for rapid plate growth.^{[77][80]}

The vascular system of the plates have been theorized to have played a role in threat displaying as *Stegosaurus* could have pumped blood into them, causing them to "blush" and give a colorful, red warning.^[81] However, Christiansen and Tschopp (2010) consider this unlikely, as stegosaur plates were covered in horn rather than skin.^[44] The plates' large size suggests that they may have served to increase the apparent height of the animal, either to intimidate enemies^[7] or to impress other members of the same species in some form of sexual display.^[73] A 2015 study of the shapes and sizes of *Hesperosaurus* plates suggested that they were sexually dimorphic, with wide plates belonging to males and taller plates belonging to females.^[25] Christiansen and Tschopp (2010) proposed that the display function would have been reinforced by the horny sheath which would have increased the visible surface and such horn structures are often brightly colored.^[44] Some have suggested that plates in stegosaurs were used to allow individuals to identify members of their species.^[80] The use of exaggerated structures in dinosaurs as species identification has been questioned, as no such function exists in modern species.^[82]

Thagomizer (tail spikes)

Debate has been going on about whether the tail spikes were used for display only, as posited by Gilmore in 1914^[7] or used as a weapon. Robert Bakker noted the tail was likely to have been much more flexible than that of other dinosaurs, as it lacked ossified tendons, thus lending credence to the idea of the tail as a weapon. However, as Carpenter^[22] has noted, the plates overlap so many tail vertebrae, movement would be limited. Bakker also observed that *Stegosaurus* could have maneuvered its rear easily, by keeping its large hind limbs stationary and pushing off with its very powerfully muscled but short forelimbs, allowing it to swivel deftly to deal with attack.^[57] More recently, a study of the tail spikes by McWhinney *et al.*,^[83] which showed a high incidence of trauma-related damage, lends more weight to the position that the spikes were indeed used in combat. This study showed that 9.8% of *Stegosaurus* specimens examined had injuries to their tail spikes. Additional support for this idea was a punctured tail vertebra of an *Allosaurus* into which a tail spike fits perfectly.^[84]



Thagomizer on mounted tail

S. stenops had four dermal spikes, each about 60–90 cm (2.0–3.0 ft) long. Discoveries of articulated stegosaur armor show, at least in some species, these spikes protruded horizontally from the tail, not vertically as is often depicted.^[22] Initially, Marsh described *S. ungulatus* as having eight spikes in its tail, unlike *S. stenops*. However, recent research re-examined this and concluded this species also had four.^[5]

"Second brain"

At one time, stegosaurs were described as having a "second brain" in their hips. Soon after describing *Stegosaurus*, Marsh noted a large canal in the hip region of the spinal cord, which could have accommodated a structure up to 20 times larger than the famously small brain. This has led to the influential idea that dinosaurs like *Stegosaurus* had a "second brain" in the tail, which may have been responsible for controlling reflexes in the rear portion of the body. This "brain" was proposed to have given a *Stegosaurus* a temporary boost when it was under threat from predators.^[39]

This space, however, is more likely to have served other purposes. The sacro-lumbar expansion is not unique to stegosaurs, nor even ornithischians. It is also present in birds. In their case, it contains what is called the glycogen body, a structure whose function is not definitely known, but which is postulated to facilitate the supply of glycogen to the animal's nervous system.^[85] It also may function as a balance organ, or reservoir of compounds to support the nervous system.^[86]

Growth

Juveniles of *Stegosaurus* have been preserved, probably showing the growth of the genus. The two juveniles are both relatively small, with the smaller individual being 1.5 m (4.9 ft) long, and the larger having a length of 2.6 m (8.5 ft). The specimens can be identified as not mature because they lack the fusion of the scapula and coracoid, and the lower hind limbs. Also, the pelvic region of the specimens are similar to *Kentrosaurus* juveniles.^[87] One 2009 study of *Stegosaurus* specimens of various sizes found that the plates and spikes had delayed histological growth in comparison to the skeleton and when the dinosaur reached maturity, growth in the osteoderms may have increased.^[88] A 2013 study concluded, based on the rapid deposition of highly vascularised fibrolamellar bone, that *Kentrosaurus* had a quicker growth rate than *Stegosaurus*, contradicting the general rule that larger dinosaurs grew faster than smaller ones.^[89]

Diet

Stegosaurus and related genera were herbivores. However, their teeth and jaws are very different from those of other herbivorous ornithischian dinosaurs, suggesting a different feeding strategy that is not yet well understood. The other ornithischians possessed teeth capable of grinding plant material and a jaw structure capable of movements in planes other than simply orthal (i.e. not only the fused up-down motion to which stegosaur jaws were likely limited). Unlike the sturdy jaws and grinding teeth common to its fellow ornithischians, *Stegosaurus* (and all stegosaurians) had small, peg-shaped teeth that have been observed with horizontal wear facets associated with tooth-food contact^[90] and their unusual jaws were probably capable of only orthal (up-down) movements.^[38] Their teeth were "not tightly pressed together in a block for efficient grinding",^[91] and no evidence in the fossil record of stegosaurians indicates use of gastroliths—the stone(s) some dinosaurs (and some present-day bird species) ingested—to aid the grinding process, so how exactly *Stegosaurus* obtained and processed the amount of plant material required to sustain its size remains "poorly understood".^[91]

The stegosaurians were widely distributed geographically in the late Jurassic.^[38] Palaeontologists believe it would have eaten plants such as mosses, ferns, horsetails, cycads, and conifers or fruits.^[92] One hypothesized feeding behavior strategy considers them to be low-level browsers, eating low-growing fruit of various nonflowering plants, as well as foliage. This scenario has *Stegosaurus* foraging at most 1 m above the ground.^[93] Conversely, if *Stegosaurus* could have raised itself on two legs, as suggested by Bakker, then it could have browsed on vegetation and fruits quite high up, with adults being able to forage up to 6 m (20 ft) above the ground.^[39]



Punctured *Allosaurus* tail vertebra by a *Stegosaurus* thagomizer, the damage shows that the spike entered at an angle from below and displaced a piece of the process upward, remodeled bone on the underside of the process shows that an infection developed



Brain cavity of *S. stenops* marked with red



Tooth crown illustration

A detailed computer analysis of the biomechanics of *Stegosaurus*'s feeding behavior was performed in 2010, using two different three-dimensional models of *Stegosaurus* teeth given realistic physics and properties. Bite force was also calculated using these models and the known skull proportions of the animal, as well as simulated tree branches of different size and hardness. The resultant bite forces calculated for *Stegosaurus* were 140.1 newtons (N), 183.7 N, and 275 N (for anterior, middle and posterior teeth, respectively), which means its bite force was less than half that of a Labrador retriever. *Stegosaurus* could have easily bitten through smaller green branches, but would have had difficulty with anything over 12 mm in diameter. *Stegosaurus*, therefore, probably browsed primarily among smaller twigs and foliage, and would have been unable to

handle larger plant parts unless the animal was capable of biting much more efficiently than predicted in this study.^[94] However, a 2016 study indicates that *Stegosaurus*'s bite strength was stronger than previously believed. Comparisons were made between it (represented by a specimen known as "Sophie" from the United Kingdom's Natural History Museum) and two other herbivorous dinosaurs; *Erlikosaurus* and *Plateosaurus* to determine if all three had similar bite forces and similar niches. Based on the results of the study, it was revealed that the subadult *Stegosaurus* specimen had a bite similar in strength to that of modern herbivorous mammals, in particular, cattle and sheep. Based on this data, it is likely *Stegosaurus* also ate woodier, tougher plants such as cycads, perhaps even acting as a means of spreading cycad seeds.^[95]

Paleoecology

The Morrison Formation is interpreted as a semiarid environment with distinct wet and dry seasons, and flat floodplains. Vegetation varied from river-lining forests of conifers, tree ferns, and ferns (gallery forests), to fern savannas with occasional trees such as the Araucaria-like conifer *Brachyphyllum*. The flora of the period has been revealed by fossils of green algae, fungi, mosses, horsetails, ferns, cycads, ginkoes, and several families of conifers. Animal fossils discovered include bivalves, snails, ray-finned fishes, frogs, salamanders, turtles like *Glyptops*, sphenodonts, lizards, terrestrial and aquatic crocodylomorphans like *Hoplosuchus*, several species of pterosaurs such as *Harpactognathus* and *Mesadactylus*, numerous dinosaur species, and early mammals such as docodonts (like *Docodon*), multituberculates, symmetrodonts, and triconodonts.^[96]



Footprints of an adult and juvenile from the Morrison Formation

Dinosaurs that lived alongside *Stegosaurus* included theropods *Allosaurus*, *Saurophaganax*, *Torvosaurus*, *Ceratosaurus*, *Marshosaurus*, *Stokesosaurus*, *Ornitholestes*, *Coelurus* and *Tanycolagreus*. Sauropods dominated the region, and included *Brachiosaurus*, *Apatosaurus*, *Diplodocus*, *Camarasaurus*, and *Barosaurus*. Other ornithischians included *Camptosaurus*, *Gargoyleosaurus*, *Dryosaurus*, *Othnielosaurus* and *Drinker*.^[97] *Stegosaurus* is commonly found at the same sites as *Allosaurus*, *Apatosaurus*, *Camarasaurus*, and *Diplodocus*. *Stegosaurus* may have preferred drier settings than these other dinosaurs.^[98]

In popular culture



Early restoration of *Stegosaurus* by
A. Tobin, 1884

One of the most recognizable of all dinosaurs,^[39] *Stegosaurus* has been depicted on film, in cartoons and comics and as children's toys. Due to the fragmentary nature of most early *Stegosaurus* fossil finds, it took many years before reasonably accurate restorations of this dinosaur could be produced. The earliest popular image of *Stegosaurus* was an engraving produced by A. Tobin for the November 1884 issue of *Scientific American*, which included the dinosaur amid a speculative Morrison age landscape. Tobin restored the *Stegosaurus* as bipedal and long-necked, with the plates arranged along the tail and the back covered in spikes. This covering of spikes might have been based on a misinterpretation of the teeth, which Marsh had noted were oddly shaped, cylindrical, and found scattered, such that he thought they might turn out to be small dermal spines.^[99]

Marsh published his more accurate skeletal reconstruction of *Stegosaurus* in 1891, and within a decade *Stegosaurus* had become among the most-illustrated types of dinosaur.^[99] Artist Charles R. Knight published his first illustration of *Stegosaurus unguulatus* based on Marsh's skeletal reconstruction in a November 1897 issue of *The Century Magazine*. This illustration would later go on to form the basis of the stop-motion puppet used in the 1933 film *King Kong*. Like Marsh's reconstruction, Knight's first restoration had a single row of large plates, though he next used a double row for his more well-known 1901 painting, produced under the direction of Frederic Lucas. Again under Lucas, Knight revised his version of *Stegosaurus* again two years later, producing a model with a staggered double row of plates. Knight would go on to paint a stegosaur with a staggered double plate row in 1927 for the Field Museum of Natural History, and was followed by Rudolph F. Zallinger, who painted *Stegosaurus* this way in his "Age of Reptiles" mural at the Peabody Museum in 1947.^[100]

Stegosaurus made its major public debut as a paper mache model commissioned by the U.S. National Museum of Natural History for the 1904 Louisiana Purchase Exposition. The model was based on Knight's latest miniature with the double row of staggered plates,^[12] and was exhibited in the United States Government Building at the exposition in St. Louis before being relocated to Portland, Oregon for the Lewis and Clark Centennial Exposition in 1905. The model was moved to the Smithsonian National Museum of Natural History (now the Arts and Industries Building) in Washington, D.C. along with other prehistory displays, and to the current National Museum of Natural History building in 1911. Following renovations to the museum in the 2010s, the model was moved once again for display at the Museum of the Earth in Ithaca, New York.^[101]



Life-sized restoration of *Stegosaurus stenops* in the U.S. National Museum, ca. 1911

The popularity of *Stegosaurus* is owed partly to its prominent display in natural history museums. Though considered one of the most distinctive types of dinosaur, *Stegosaurus* displays were missing from a majority of museums during the first half of the 20th century, due largely to the disarticulated nature of most fossil specimens. Until 1918, the only mounted skeleton of *Stegosaurus* in the world was O. C. Marsh's type specimen of *S. unguulatus* at the Peabody Museum of Natural History, which was put on display in 1910. However, this mount was dismantled in 1917 when the old Peabody Museum building was demolished.^[12] This historically significant specimen was re-mounted ahead of the opening of the new Peabody Museum building in 1925.^[13] 1918 saw the completion of the second *Stegosaurus* mount, and the first depicting *S. stenops*. This mount was created under the direction of Charles Gilmore at the U.S. National Museum of Natural History. It was a composite of several skeletons, primarily USNM 6531, with proportions designed



The first mounted skeleton of a stegosaur (*S. unguatus*), Peabody Museum of Natural History, 1910

to closely follow the *S. stenops* type specimen, which had been on display in relief nearby since 1918.^[12] The aging mount was dismantled in 2003 and replaced with a cast in an updated pose in 2004.^[102] A third mounted skeleton of *Stegosaurus*, referred to *S. stenops*, was put on display at the American Museum of Natural History in 1932. Mounted under the direction of Charles J. Long, the American Museum mount was a composite consisting of partial remains filled in with replicas based on other specimens. In his article about the new mount for the museum's journal, Barnum Brown described (and disputed) the popular misconception that the *Stegosaurus* had a "second brain" in its hips.^[103] Another composite mount, using specimens referred to *S. unguatus* collected from Dinosaur National Monument between 1920 and 1922, was put on display at the Carnegie Museum of Natural History in 1940.^[104]

See also

- Timeline of stegosaur research

References

1. "Stegosaurus" (<https://www.merriam-webster.com/dictionary/Stegosaurus>). *Merriam-Webster Dictionary*.
2. Galton, P.M. (2010). "Species of plated dinosaur *Stegosaurus* (Morrison Formation, Late Jurassic) of western USA: new type species designation needed". *Swiss Journal of Geosciences*. **103** (2): 187–198. doi:10.1007/s00015-010-0022-4 (<https://doi.org/10.1007%2Fs00015-010-0022-4>). S2CID 140688875 (<https://api.semanticscholar.org/CorpusID:140688875>).
3. Marsh, O. C. (1877). "A new order of extinct Reptilia (*Stegosauria*) from the Jurassic of the Rocky Mountains" (<https://zenodo.org/record/1450038>). *American Journal of Science*. **3** (14): 513–514. Bibcode:1877AmJS...14..513M (<https://ui.adsabs.harvard.edu/abs/1877AmJS...14..513M>). doi:10.2475/ajs.s3-14.84.513 (<https://doi.org/10.2475%2Fajs.s3-14.84.513>). S2CID 130078453 (<https://api.semanticscholar.org/CorpusID:130078453>).
4. International Commission on Zoological Nomenclature (2013). "Opinion 2320 (Case 3536): *Stegosaurus* Marsh, 1877 (Dinosauria, Ornithischia): type species replaced with *Stegosaurus stenops* Marsh, 1887". *Bulletin of Zoological Nomenclature*. **70** (2): 129–130. doi:10.21805/bzn.v70i2.a4 (<https://doi.org/10.21805%2Fbzn.v70i2.a4>). S2CID 198149935 (<https://api.semanticscholar.org/CorpusID:198149935>).
5. Carpenter K, Galton PM (2001). "Othniel Charles Marsh and the Myth of the Eight-Spiked *Stegosaurus*". In Carpenter K (ed.). *The Armored Dinosaurs*. Indiana University Press. pp. 76–102. ISBN 978-0-253-33964-5.
6. Carpenter, Kenneth (1998). "Vertebrate biostratigraphy of the Morrison Formation near Cañon City, Colorado". *Modern Geology*. **23**: 407–426.
7. Gilmore CW (1914). "Osteology of the armored Dinosauria in the United States National Museum, with special reference to the genus *Stegosaurus*". *Series: Smithsonian Institution. United States National Museum. Bulletin* 89 (89).
8. "Division of Paleontology" (http://research.amnh.org/paleontology/search.php?action=detail&specimen_id=48559). *research.amnh.org*. Retrieved April 17, 2022.

9. Marsh, O. C. (1879). "Notice of new Jurassic reptiles" (<http://marsh.dinodb.com/marsh/Marsh%201877%20-%20Notice%20of%20new%20dinosaurian%20reptiles%20from%20the%20Jurassic%20formation.pdf>) (PDF). *American Journal of Science*. **3** (18): 501–505. Bibcode:1879AmJS...18..501M (<https://ui.adsabs.harvard.edu/abs/1879AmJS...18..501M>). doi:10.2475/ajs.s3-18.108.501 (<https://doi.org/10.2475%2Fajs.s3-18.108.501>). S2CID 131001110 (<https://api.semanticscholar.org/CorpusID:131001110>).
10. Marsh, O. C. (1887). "Principal characters of American Jurassic dinosaurs, part IX. The skull and dermal armour of *Stegosaurus*" (<https://zenodo.org/record/2387814>). *American Journal of Science*. **3** (34): 413–17. Bibcode:1887AmJS...34..413M (<https://ui.adsabs.harvard.edu/abs/1887AmJS...34..413M>). doi:10.2475/ajs.s3-34.203.413 (<https://doi.org/10.2475%2Fajs.s3-34.203.413>). S2CID 130058870 (<https://api.semanticscholar.org/CorpusID:130058870>).
11. Marsh, Othniel Charles (1891). "Restoration of *Stegosaurus*". *American Journal of Science*. **3** (42): 179–81. Bibcode:1891AmJS...42..179M (<https://ui.adsabs.harvard.edu/abs/1891AmJS...42..179M>). doi:10.2475/ajs.s3-42.248.179 (<https://doi.org/10.2475%2Fajs.s3-42.248.179>). S2CID 129981252 (<https://api.semanticscholar.org/CorpusID:129981252>).
12. Gilmore, Charles W. (1919). "A newly mounted skeleton of the armored dinosaur, *Stegosaurus stenops*, in the United States National Museum" (http://repository.si.edu/bitstream/handle/10088/14684/1/USNMP-54_2241_1918.pdf) (PDF). *Proceedings of the United States National Museum*. **54** (2241): 383–390. doi:10.5479/si.00963801.54-2241.383 (<http://doi.org/10.5479%2Fsi.00963801.54-2241.383>).
13. Revan, A. (2011). *Reconstructing an Icon: Historical Significance of the Peabody's Mounted Skeleton of *Stegosaurus* and the Changes Necessary to Make It Correct Anatomically*. (http://earth.yale.edu/sites/default/files/files/Academics/11-Revan_thesis%2Bappendix.pdf) Doctoral dissertation, faculty of the Department of Geology and Geophysics, Yale University.
14. Colbert, EH (1962). *Dinosaurs: Their Discovery & Their World*. Hutchinson Press, London. pp. 82–99. ISBN 978-1-111-21503-3.
15. Czerkas SA (1987). "A Reevaluation of the Plate Arrangement on *Stegosaurus stenops*". In Czerkas SJ, Olson EC (eds.). *Dinosaurs Past & Present, Vol 2*. University of Washington Press, Seattle. pp. 82–99.
16. Cameron, Robert P.; Cameron, John A.; Barnett, Stephen M. (August 15, 2015). "Were there two forms of *Stegosaurus*?". arXiv:1508.03729 (<https://arxiv.org/abs/1508.03729>) [q-bio.PE (<https://arxiv.org/archive/q-bio.PE>)].
17. Cameron, R. P.; Cameron, J. A.; Barnett, S. M. (November 26, 2016). "*Stegosaurus* chirality". arXiv:1611.08760 (<https://arxiv.org/abs/1611.08760>) [q-bio.PE (<https://arxiv.org/archive/q-bio.PE>)].
18. Brinkman, P. D. (2010). *The second Jurassic dinosaur rush*. University of Chicago Press.
19. Maidment, Susannah C.R.; Norman, David B.; Barrett, Paul M.; Upchurch, Paul (2008). "Systematics and phylogeny of Stegosauria (Dinosauria: Ornithischia)". *Journal of Systematic Palaeontology*. **6** (4): 1. doi:10.1017/S1477201908002459 (<https://doi.org/10.1017%2FS1477201908002459>). S2CID 85673680 (<https://api.semanticscholar.org/CorpusID:85673680>).
20. McIntosh, J. S. (1981). *Annotated catalogue of the dinosaurs (Reptilia, Archosauria) in the collections of Carnegie Museum of Natural History*.
21. "Denver's Fighting Dinosaurs" (<https://extinctmonsters.net/2015/08/14/denvers-fighting-dinosaurs/>). *EXTINCT MONSTERS*. August 14, 2015. Retrieved April 20, 2022.
22. Carpenter, Kenneth (1998). "Armor of *Stegosaurus stenops*, and the taphonomic history of a new specimen from Garden Park, Colorado". *Modern Geology*. **23**: 127–44.

23. Maidment, S. C. R.; Brassey, Charlotte; Barrett, Paul Michael (2015). "The Postcranial Skeleton of an Exceptionally Complete Individual of the Plated Dinosaur *Stegosaurus stenops* (Dinosauria: Thyreophora) from the Upper Jurassic Morrison Formation of Wyoming, U.S.A." (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4605687>) *PLOS ONE*. **10** (10): e0138352. Bibcode:2015PLoSO..1038352M (<https://ui.adsabs.harvard.edu/abs/2015PLoSO..1038352M>). doi:10.1371/journal.pone.0138352 (<https://doi.org/10.1371%2Fjournal.pone.0138352>). PMC 4605687 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4605687>). PMID 26466098 (<https://pubmed.ncbi.nlm.nih.gov/26466098>).
24. "Untitled Document" (<http://www.paleofile.com/Dinosaurs/Armor/Stegosaurus.asp>). *www.paleofile.com*. Retrieved April 20, 2022.
25. Saitta E.T. (2015). "Evidence for Sexual Dimorphism in the Plated Dinosaur *Stegosaurus mjosi* (Ornithischia, Stegosauria) from the Morrison Formation (Upper Jurassic) of Western USA" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4406738>). *PLOS ONE*. **10** (4): e0123503. Bibcode:2015PLoSO..1023503S (<https://ui.adsabs.harvard.edu/abs/2015PLoSO..1023503S>). doi:10.1371/journal.pone.0123503 (<https://doi.org/10.1371%2Fjournal.pone.0123503>). PMC 4406738 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4406738>). PMID 25901727 (<https://pubmed.ncbi.nlm.nih.gov/25901727>).
26. Escaso F, Ortega F, Dantas P, Malafaia E, Pimentel NL, Pereda-Suberbiola X, Sanz JL, Kullberg JC, Kullberg MC, Barriga F (2007). "New Evidence of Shared Dinosaur Across Upper Jurassic Proto-North Atlantic: *Stegosaurus* From Portugal". *Naturwissenschaften*. **94** (5): 367–74. Bibcode:2007NW....94..367E (<https://ui.adsabs.harvard.edu/abs/2007NW....94..367E>). doi:10.1007/s00114-006-0209-8 (<https://doi.org/10.1007%2Fs00114-006-0209-8>). PMID 17187254 (<https://pubmed.ncbi.nlm.nih.gov/17187254>). S2CID 10930309 (<https://api.semanticscholar.org/CorpusID:10930309>).
27. Siber, H. J., & Möckli, U. (2009). The stegosaurs of the Sauriermuseum Aathal. (https://doc.re.ro.ch/record/209444/files/PAL_E4048.pdf)
28. Gauthier, Advisers Jacques; Brinkman, D. (2012). "Reconstructing an Icon : Historical Significance of the Peabody 's Mounted Skeleton of *Stegosaurus* and the Changes Necessary to Make It Correct Anatomically" (<https://www.semanticscholar.org/paper/Reconstructing-an-Icon-%3A-Historical-Significance-of-Gauthier-Brinkman/95044daee27fb47dbdea7ef06c87b534ffa5866>). *www.semanticscholar.org*. S2CID 15111495 (<https://api.semanticscholar.org/CorpusID:15111495>). Retrieved April 17, 2022.
29. Holtz, Thomas R. Jr. (2012) *Dinosaurs: The Most Complete, Up-to-Date Encyclopedia for Dinosaur Lovers of All Ages*, Winter 2011 Appendix. (<http://www.geol.umd.edu/~tholtz/dinosaurappendix/HoltzappendixWinter2011.pdf>)
30. Foster, J.R. (2003). *Paleoecological analysis of the vertebrate fauna of the Morrison Formation (Upper Jurassic), Rocky Mountain region, U.S.A.* New Mexico Museum of Natural History and Science Bulletin, **23**. Albuquerque, New Mexico: New Mexico Museum of Natural History and Science.
31. Benson, RBJ; Campione, NE; Carrano, MT; Mannion, PD; Sullivan, C; Evans, David C.; et al. (2014). "Rates of Dinosaur Body Mass Evolution Indicate 170 Million Years of Sustained Ecological Innovation on the Avian Stem Lineage" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4011683>). *PLOS Biol*. **12** (5): e1001853. doi:10.1371/journal.pbio.1001853 (<https://doi.org/10.1371%2Fjournal.pbio.1001853>). PMC 4011683 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4011683>). PMID 24802911 (<https://pubmed.ncbi.nlm.nih.gov/24802911>).
32. Paul, G. S. (2016). *The Princeton Field Guide to Dinosaurs* (2nd ed.). Princeton, New Jersey: Princeton University Press. pp. 162–168. ISBN 9780691167664.
33. <http://rsbl.royalsocietypublishing.org/content/11/3/20140984>

34. *Stegosaurus*. University of Wyoming Geological Museum. 2006. Retrieved October 6, 2006. University of Wyoming Geological Museum (<http://www.uwyo.edu/geomuseum/tour/stegosaurus.asp>) Archived (<https://web.archive.org/web/20060827043650/http://www.uwyo.edu/geomuseum/tour/stegosaurus.asp>) August 27, 2006, at the [Wayback Machine](#)
35. Czerkas, S (1999). "The beaked jaws of stegosaurs and their implications for other ornithischians". *Miscellaneous Publication of the Utah Geological Survey*. **99–1**: 143–150.
36. Knoll, F (2008). "Buccal soft anatomy in *Lesothosaurus* (Dinosauria: Ornithischia)" (<https://www.researchgate.net/publication/233720216>) (PDF). *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen*. **248** (3): 355–364. doi:10.1127/0077-7749/2008/0248-0355 (<https://doi.org/10.1127/0077-7749/2008/0248-0355>).
37. Barrett, P.M. (2001). Tooth wear and possible jaw action of *Scelidosaurus harrisonii* Owen and a review of feeding mechanisms in other thyreophoran dinosaurs. Pp. 25-52 in Carpenter, K. (ed.): *The Armored Dinosaurs*. Bloomington: Indiana University Press.
38. Galton PM, Upchurch P (2004). "Stegosauria". In Weishampel DB, Dodson P, Osmólska H (eds.). *The Dinosauria* (2nd ed.). University of California Press. ISBN 978-0-520-24209-8.
39. Fastovsky DE, Weishampel DB (2005). "Stegosauria: Hot Plates". In Fastovsky DE, Weishampel DB (eds.). *The Evolution and Extinction of the Dinosaurs* (2nd ed.). Cambridge University Press. pp. 107–30. ISBN 978-0-521-81172-9.
40. Bakker RT (1986). *The Dinosaur Heresies*. William Morrow, New York. pp. 365–74.
41. Buchholtz, Emily; Holtz, Thomas R. Jr.; Farlow, James O.; Walters, Bob (June 27, 2012). Brett-Surman, M.K. (ed.). *The complete dinosaur* (http://www.google.no/books?hl=no&lr=&id=_pX_I24sDARwC&oi=fnd&pg=PA191&dq=stegosaurus+brain+intelligence&ots=W4wYavmVri&sig=L_93Z6f8BWSWIQKWdKpCx8oFhjl&redir_esc=y#v=onepage&q=stegosaurus%20brain%20intelligence&f=false) (2nd ed.). Bloomington, Ind.: Indiana University Press. p. 201. ISBN 978-0-253-00849-7. Retrieved February 27, 2013. "Proportions and anatomy of endocasts of *Stegosaurus* vary little from those of ancestral archosaurs, with an elongated shape, large olfactory lobes, and extremely narrow cerebral hemispheres. Lack of surface detail suggest that the brain did not fill the braincase. EQ estimates are below 0,6 (Hoppson, 1977), agreeing well with predictions of a slow herbivorous lifestyle."
42. Martin, A.J. (2006). Introduction to the Study of Dinosaurs. Second Edition. Oxford, Blackwell Publishing. 560 pp. ISBN 1-4051-3413-5.
43. Lambert, David (1993). *The Ultimate Dinosaur Book* (<https://archive.org/details/ultimatedinosaur00lamb/page/110>). Dorling Kindersley, New York. pp. 110–29. ISBN 978-1-56458-304-8.
44. Christiansen, N. A.; Tschopp, E. (2010). "Exceptional stegosaur integument impressions from the Upper Jurassic Morrison Formation of Wyoming". *Journal of Geosciences*. **103** (2): 163–171. doi:10.1007/s00015-010-0026-0 (<https://doi.org/10.1007/s00015-010-0026-0>). S2CID 129246092 (<https://api.semanticscholar.org/CorpusID:129246092>).
45. Norman, David (2001). "Scelidosaurus, the earliest complete dinosaur" in *The Armored Dinosaurs*, pp 3-24. Bloomington: Indiana University Press. ISBN 0-253-33964-2.
46. Peter M. Galton (2019). "Earliest record of an ankylosaurian dinosaur (Ornithischia: Thyreophora): Dermal armor from Lower Kota Formation (Lower Jurassic) of India". *Neues Jahrbuch für Geologie und Paläontologie - Abhandlungen*. **291** (2): 205–219. doi:10.1127/njgpa/2019/0800 (<https://doi.org/10.1127/njgpa/2019/0800>). S2CID 134302379 (<https://api.semanticscholar.org/CorpusID:134302379>).
47. Sereno, P.C., 1998, "A rationale for phylogenetic definitions, with application to the higher-level taxonomy of Dinosauria", *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen* **210**: 41-83

48. Pereda-Suberbiola, Xabier; Galton, Peter M.; Mallison, Heinrich; Novas, Fernando (2013). "A plated dinosaur (Ornithischia, Stegosauria) from the Early Cretaceous of Argentina, South America: an evaluation". *Alcheringa: An Australasian Journal of Palaeontology*. **37** (1): 65–78. doi:10.1080/03115518.2012.702531 (<https://doi.org/10.1080%2F03115518.2012.702531>). S2CID 128632458 (<https://api.semanticscholar.org/CorpusID:128632458>).
49. Madzia, Daniel; Arbour, Victoria M.; Boyd, Clint A.; Farke, Andrew A.; Cruzado-Caballero, Penélope; Evans, David C. (December 9, 2021). "The phylogenetic nomenclature of ornithischian dinosaurs" (<https://peerj.com/articles/12362>). *PeerJ*. **9**: e12362. doi:10.7717/peerj.12362 (<https://doi.org/10.7717%2Fpeerj.12362>). ISSN 2167-8359 (<https://www.worldcat.org/issn/2167-8359>). PMC 8667728 (<https://www.ncbi.nlm.nih.gov/pmc/article/s/PMC8667728>). PMID 34966571 (<https://pubmed.ncbi.nlm.nih.gov/34966571>).
50. Mateus, Octávio; Maidment, Susannah C.R.; Christiansen, Nicolai A. (May 22, 2009). "A new long-necked 'sauropod-mimic' stegosaur and the evolution of the plated dinosaurs" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2674496>). *Proceedings of the Royal Society B: Biological Sciences*. **276** (1663): 1815–1821. doi:10.1098/rspb.2008.1909 (<https://doi.org/10.1098%2Frspb.2008.1909>). PMC 2674496 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2674496>). PMID 19324778 (<https://pubmed.ncbi.nlm.nih.gov/19324778>).
51. Raven, T.J.; Maidment, S.C.R. (2017). "A new phylogeny of Stegosauria (Dinosauria, Ornithischia)" (http://eprints.brighton.ac.uk/16844/1/Raven%26Maidment_AcceptedPreprint.pdf) (PDF). *Palaeontology*. **2017** (3): 401–408. doi:10.1111/pala.12291 (<https://doi.org/10.1111%2Fpala.12291>). hdl:10044/1/45349 (<https://hdl.handle.net/10044%2F1%2F45349>). S2CID 55613546 (<https://api.semanticscholar.org/CorpusID:55613546>).
52. Madzia, D.; Arbour, V.M.; Boyd, C.A.; Farke, A.A.; Cruzado-Caballero, P.; Evans, D.C. (2021). "The phylogenetic nomenclature of ornithischian dinosaurs" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8667728>). *PeerJ*. **9**: e12362. doi:10.7717/peerj.12362 (<https://doi.org/10.7717%2Fpeerj.12362>). PMC 8667728 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8667728>). PMID 34966571 (<https://pubmed.ncbi.nlm.nih.gov/34966571>).
53. Foster, J. (2007). *Jurassic West: The Dinosaurs of the Morrison Formation and Their World*. Indiana University Press. pp. 327–329. ISBN 978-0-253-34870-8.
54. Senter, Phil (2010). "Evidence for a Sauropod-Like Metacarpal Configuration in Stegosaurian Dinosaurs" (http://doc.rero.ch/record/32124/files/PAL_E2887.pdf) (PDF). *Acta Palaeontologica Polonica*. **55** (3): 427–432. doi:10.4202/app.2009.1105 (<https://doi.org/10.4202%2Fapp.2009.1105>). S2CID 53328847 (<https://api.semanticscholar.org/CorpusID:53328847>).
55. Raven, T. J.; Maidment, S. C. R. (2017). "A new phylogeny of Stegosauria (Dinosauria, Ornithischia)" (http://eprints.brighton.ac.uk/16844/1/Raven%26Maidment_AcceptedPreprint.pdf) (PDF). *Palaeontology*. **60** (3): 401–408. doi:10.1111/pala.12291 (<https://doi.org/10.1111%2Fpala.12291>). hdl:10044/1/45349 (<https://hdl.handle.net/10044%2F1%2F45349>). S2CID 55613546 (<https://api.semanticscholar.org/CorpusID:55613546>).
56. Marsh OC (1881). "Principal characters of American Jurassic dinosaurs, part V". *American Journal of Science*. **3** (21): 417–23. Bibcode:1881AmJS...21..417M (<https://ui.adsabs.harvard.edu/abs/1881AmJS...21..417M>). doi:10.2475/ajs.s3-21.125.417 (<https://doi.org/10.2475%2Fajs.s3-21.125.417>). S2CID 219234316 (<https://api.semanticscholar.org/CorpusID:219234316>).
57. Bakker RT (1986). *The Dinosaur Heresies*. William Morrow, New York. ISBN 978-0-8217-2859-8.
58. *Science* (<https://books.google.com/books?id=sbQ7AQAAMAAJ&q=Scientific+research:+The+art+of+revelation+and+of+prophecy&pg=PA401>). American Association for the Advancement of Science. 1902.

59. Costa, Francisco; Mateus, Octávio (November 13, 2019). "Dacentrurine stegosaurs (Dinosauria): A new specimen of *Miragaia longicollum* from the Late Jurassic of Portugal resolves taxonomical validity and shows the occurrence of the clade in North America" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6853308>). *PLOS ONE*. **14** (11): e0224263. Bibcode:2019PLoSO..1424263C (<https://ui.adsabs.harvard.edu/abs/2019PLoSO..1424263C>). doi:10.1371/journal.pone.0224263 (<https://doi.org/10.1371%2Fjournal.pone.0224263>). ISSN 1932-6203 (<https://www.worldcat.org/issn/1932-6203>). PMC 6853308 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6853308>). PMID 31721771 (<https://pubmed.ncbi.nlm.nih.gov/31721771>).
60. Galton, Peter M.; Carpenter, Kenneth (2016). "The plated dinosaur *Stegosaurus longispinus* Gilmore, 1914 (Dinosauria: Ornithischia; Upper Jurassic, western USA), type species of *Alcovasaurus* n. gen". *Neues Jahrbuch für Geologie und Paläontologie - Abhandlungen*. **279** (2): 185–208. doi:10.1127/njgpa/2016/0551 (<https://doi.org/10.1127%2Fnjgpa%2F2016%2F0551>).
61. Galton, PM (1981). "*Craterosaurus pottonensis* Seeley, a stegosaurian dinosaur from the Lower Cretaceous of England, and a review of Cretaceous stegosaurs". *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen*. **161** (1): 28–46.
62. Dong, Z. M. (1973). Dinosaurs from Wuerho. *Institute of Paleontology and Paleoanthropology Memoir*, 11, 45-52.
63. HAO, B., PENG, G., QIN, G., YE, Y., & JIANG, S. (2018). History and evolution of stegosaurus in China. *Geological Bulletin of China*, 37(10), 1777-1782.
64. Xing, L., Lockley, M. G., PERSONS IV, W. S., Klein, H., Romilio, A., Wang, D., & Wang, M. (2021). Stegosaur track assemblage from Xinjiang, China, featuring the smallest known stegosaur record. *Palaios*, 36(2), 68-76.
65. Maidment, S. C. (2010). Stegosauria: a historical review of the body fossil record and phylogenetic relationships. *Swiss Journal of Geosciences*, 103(2), 199-210.
66. Maidment, Susannah C. R.; Woodruff, D. Cary; Horner, John R. (January 2, 2018). "A new specimen of the ornithischian dinosaur *Hesperosaurus mjosi* from the Upper Jurassic Morrison Formation of Montana, U.S.A., and implications for growth and size in Morrison stegosaurs" (<https://doi.org/10.1080/02724634.2017.1406366>). *Journal of Vertebrate Paleontology*. **38** (1): e1406366. doi:10.1080/02724634.2017.1406366 (<https://doi.org/10.1080%2F02724634.2017.1406366>). hdl:10141/622747 (<https://hdl.handle.net/10141%2F622747>). ISSN 0272-4634 (<https://www.worldcat.org/issn/0272-4634>). S2CID 90752660 (<https://api.semanticscholar.org/CorpusID:90752660>).
67. Marsh, Othniel Charles (1880). "Principal characters of American Jurassic dinosaurs, part III" (<https://zenodo.org/record/2042250>). *American Journal of Science*. **3** (19): 253–59. Bibcode:1880AmJS...19..253M (<https://ui.adsabs.harvard.edu/abs/1880AmJS...19..253M>). doi:10.2475/ajs.s3-19.111.253 (<https://doi.org/10.2475%2Fajs.s3-19.111.253>). S2CID 130249498 (<https://api.semanticscholar.org/CorpusID:130249498>).
68. Bakker RT (1978). "Dinosaur feeding behavior and the origin of flowering plants". *Nature*. **274** (5672): 661–63. Bibcode:1978Natur.274..661B (<https://ui.adsabs.harvard.edu/abs/1978Natur.274..661B>). doi:10.1038/274661a0 (<https://doi.org/10.1038%2F274661a0>). S2CID 4162574 (<https://api.semanticscholar.org/CorpusID:4162574>).
69. Mallison, H. (2010). "CAD assessment of the posture and range of motion of *Kentrosaurus aethiopicus* HENNIG 1915". *Swiss Journal of Geosciences*. **103** (2): 211–233. doi:10.1007/s00015-010-0024-2 (<https://doi.org/10.1007%2Fs00015-010-0024-2>). S2CID 132746786 (<https://api.semanticscholar.org/CorpusID:132746786>).
70. Calculating the speed of Quadrupedal graviportal animals by Ruben Molina-Perez, Asier Larramendi

71. Rajewski, Genevieve (May 2008). "Where Dinosaurs Roamed" (<http://arquivo.pt/wayback/20091015085900/http%3A/www.smithsonianmag.com/history%2Darchaeology/phenom%2Ddino%2D200805.html>). *Smithsonian*: 20–24. Archived from the original (<http://www.smithsonianmag.com/history-archaeology/phenom-dino-200805.html>) on October 15, 2009. Retrieved April 30, 2008.
72. Isles, T. (2009). "The socio-sexual behaviour of extant archosaurs: Implications for understanding dinosaur behaviour" (http://doc.rero.ch/record/232923/files/PAL_E4545.pdf) (PDF). *Historical Biology*. **21** (3–4): 139–214. doi:10.1080/08912960903450505 (<https://doi.org/10.1080%2F08912960903450505>). S2CID 85417112 (<https://api.semanticscholar.org/CorpusID:85417112>).
73. Davitashvili L (1961). *The Theory of sexual selection*. Izdatel'stvo Akademia nauk SSSR. p. 538.
74. Bakker, R (1986). *The Dinosaur Heresies*. Penguin Books. pp. 229–34. ISBN 978-0-14-015792-5.
75. Buffrénil (1986). "Growth and Function of *Stegosaurus* Plates". *Paleobiology*. **12** (4): 459–73. doi:10.1017/S0094837300003171 (<https://doi.org/10.1017%2FS0094837300003171>).
76. Farlow JO, Thompson CV, Rosner DE (1976). "Plates of the dinosaur *Stegosaurus*: Forced convection heat loss fins?". *Science*. **192** (4244): 1123–25. Bibcode:1976Sci...192.1123F (<https://ui.adsabs.harvard.edu/abs/1976Sci...192.1123F>). doi:10.1126/science.192.4244.1123 (<https://doi.org/10.1126%2Fscience.192.4244.1123>). PMID 17748675 (<https://pubmed.ncbi.nlm.nih.gov/17748675>). S2CID 44506996 (<https://api.semanticscholar.org/CorpusID:44506996>).
77. Stephen Brusatte (February 2012). *Dinosaur Paleobiology*. Hoboken, NJ: Wiley-Blackwell. pp. 63–64.
78. Farlow, James O.; Hayashi, Shoji; Tattersall, Glenn J. (2010). "Internal vascularity of the dermal plates of *Stegosaurus* (Ornithischia, Thyreophora)" (https://web.archive.org/web/20131005022905/http://www.brocku.ca/researchers/glenn_tattersall/PDFs/farlow%20et%20al%20stegosaurus.pdf) (PDF). *Swiss J Geoscia*. **103** (2): 173–85. doi:10.1007/s00015-010-0021-5 (<https://doi.org/10.1007%2Fs00015-010-0021-5>). S2CID 129933940 (<https://api.semanticscholar.org/CorpusID:129933940>). Archived from the original (http://www.brocku.ca/researchers/glenn_tattersall/PDFs/farlow%20et%20al%20stegosaurus.pdf) (PDF) on October 5, 2013. Retrieved June 13, 2012.
79. Main RP, Padian K, Horner J (2000). "Comparative histology, growth and evolution of archosaurian osteoderms: why did *Stegosaurus* have such large dorsal plates?". *Journal of Vertebrate Paleontology*. **20**: 56A.
80. Main R, de Ricqlès A, Horner JR, Padian K (2005). "The evolution and function of thyreophoran dinosaur scutes: implications for plate function in stegosaurs". *Paleobiology*. **31** (2): 291–314. doi:10.1666/0094-8373(2005)031[0291:teafot]2.0.co;2 (<https://doi.org/10.1666%2F0094-8373%282005%29031%5B0291%3Ateafot%5D2.0.co%3B2>).
81. Kenneth Carpenter; Dan Chure; James Ian Kirkland; Denver Museum of Natural History (1998). *The Upper Jurassic Morrison Formation: an interdisciplinary study Part 2*. Taylor & Francis. p. 137. ISBN 978-90-5699-183-8.
82. Hone, D. W. E.; Naish, D. (2013). "The 'species recognition hypothesis' does not explain the presence and evolution of exaggerated structures in non-avian dinosaurs" (<https://doi.org/10.1111%2Fjzo.12035>). *Journal of Zoology*. **290** (3): 172–180. doi:10.1111/jzo.12035 (<https://doi.org/10.1111%2Fjzo.12035>).
83. McWhinney LA, Rothschild BM, Carpenter K (2001). "Posttraumatic Chronic Osteomyelitis in *Stegosaurus* dermal spikes". In Carpenter K (ed.). *The Armored Dinosaurs*. Indiana University Press. pp. 141–56. ISBN 978-0-253-33964-5.

84. Carpenter K, Sanders F, McWhinney L, Wood L (2005). "Evidence for predator-prey relationships: Examples for *Allosaurus* and *Stegosaurus*". In Carpenter K (ed.). *The Carnivorous Dinosaurs*. Indiana University Press. pp. 325–50. ISBN 978-0-253-34539-4.
85. Buchholz (née Giffin) EB (1990). "Gross Spinal Anatomy and Limb Use in Living and Fossil Reptiles". *Paleobiology*. **16** (4): 448–58. doi:10.1017/S0094837300010186 (<https://doi.org/10.1017/S0094837300010186>).
86. Wedel, Matt (December 15, 2009). "Lies, damned lies, and Clash of the Dinosaurs" (<https://svpow.com/2009/12/15/lies-damned-lies-and-clash-of-the-dinosaurs/>). *Sauropod Vertebra Picture of the Week*.
87. Galton, P.M. (1982). "Juveniles of the stegosaurian dinosaur *Stegosaurus* from the Upper Jurassic of North America". *Journal of Vertebrate Paleontology*. **2** (1): 47–62. doi:10.1080/02724634.1982.10011917 (<https://doi.org/10.1080/02724634.1982.10011917>).
88. Hayashi, S.; Carpenter, K.; Suzuki, D. (2009). "Different Growth Patterns between the Skeleton and Osteoderms of *Stegosaurus* (Ornithischia: Thyreophora)". *Journal of Vertebrate Paleontology*. **29** (1): 123–131. doi:10.1080/02724634.2009.10010366 (<https://doi.org/10.1080/02724634.2009.10010366>). S2CID 86613451 (<https://api.semanticscholar.org/CorpusID:86613451>).
89. Redelstorff, R.; Hübner, T. R.; Chinsamy, A.; Sander, P. M. (2013). "Bone histology of the stegosaur *Kentrosaurus aethiopicus* (Ornithischia: Thyreophora) from the Upper Jurassic of Tanzania" (<https://doi.org/10.1002/Far.22701>). *The Anatomical Record*. **296** (6): 933–952. doi:10.1002/ar.22701 (<https://doi.org/10.1002/Far.22701>). PMID 23613282 (<https://pubmed.ncbi.nlm.nih.gov/23613282>). S2CID 23433029 (<https://api.semanticscholar.org/CorpusID:23433029>).
90. Barrett, PM (2001). "Tooth wear and possible jaw action of *Scelidosaurus harrisoni*, and a review of feeding mechanisms in other thyreophoran dinosaurs". In Carpenter, Kenneth (ed.). *The Armored Dinosaurs*. Indiana University Press. pp. 25–52. ISBN 978-0-253-33964-5.
91. Fastovsky, David E. & Weishampel, David B. (2009). *Dinosaurs: A Concise Natural History*. Cambridge, GBR: Cambridge University Press. pp. 89–90. ISBN 978-0-511-47789-8.
92. *Stegosaurus ungulatus* (<http://www.carnegiemnh.org/exhibits/jurassic/fctsteg.htm>) Archived (<https://web.archive.org/web/20070312204144/http://www.carnegiemnh.org/exhibits/jurassic/fctsteg.htm>) March 12, 2007, at the [Wayback Machine](#) Carnegie Museum of Natural History. Retrieved October 26, 2006.
93. Weishampel, DB (January 1984). "Interactions between Mesozoic Plants and Vertebrates: Fructifications and seed predation". *N. Jb. Geol. Paläontol. Abhandl.* **167** (2): 224–50. ISSN 0077-7749 (<https://www.worldcat.org/issn/0077-7749>).
94. Reichel, Miriam (2010). "A model for the bite mechanics in the herbivorous dinosaur *Stegosaurus* (Ornithischia, Stegosauridae)". *Swiss Journal of Geosciences*. **103** (2): 235–240. doi:10.1007/s00015-010-0025-1 (<https://doi.org/10.1007/s00015-010-0025-1>). S2CID 84869720 (<https://api.semanticscholar.org/CorpusID:84869720>).
95. Lautenschlager, Stephen; Brassey, Charlotte A.; Button, David J.; Barret, Paul Mm. (2016). "Decoupled form and function in disparate herbivorous dinosaur clades" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4873811>). *Scientific Reports*. **6**: 26495. Bibcode:2016NatSR...626495L (<https://ui.adsabs.harvard.edu/abs/2016NatSR...626495L>). doi:10.1038/srep26495 (<https://doi.org/10.1038/srep26495>). PMC 4873811 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4873811>). PMID 27199098 (<https://pubmed.ncbi.nlm.nih.gov/27199098>).

96. Chure, Daniel J.; Litwin, Ron; Hasiotis, Stephen T.; Evanoff, Emmett; and Carpenter, Kenneth (2006). "The fauna and flora of the Morrison Formation: 2006". In Foster, John R.; and Lucas, Spencer G. (eds.). *Paleontology and Geology of the Upper Jurassic Morrison Formation*. New Mexico Museum of Natural History and Science Bulletin, 36. Albuquerque, New Mexico: New Mexico Museum of Natural History and Science. pp. 233–248.
97. Foster, J. (2007). "Appendix." *Jurassic West: The Dinosaurs of the Morrison Formation and Their World*. Indiana University Press. pp. 327-329.
98. Dodson, Peter; Behrensmeyer, A.K.; Bakker, Robert T.; McIntosh, John S. (Spring 1980). "Taphonomy and paleoecology of the dinosaur beds of the Jurassic Morrison Formation". *Paleobiology*. Cambridge University Press. **6** (2): 208–232.
doi:10.1017/S0094837300025768 (<https://doi.org/10.1017%2FS0094837300025768>).
JSTOR 2400253 (<https://www.jstor.org/stable/2400253>).
99. Debus, A. A. (2009). *Prehistoric Monsters: The Real and Imagined Creatures of the Past That We Love to Fear*. McFarland.
100. Moore, R. (2014). *Dinosaurs by the Decades: A Chronology of the Dinosaur in Science and Popular Culture*. ABC-CLIO.
101. "Discover Our Blogs" (<https://nmnh.typepad.com/100years/discover-other-nmnh-blogs.html>). *National Museum of Natural History Unearthed*. Retrieved January 10, 2022.
102. "Extinct Monsters: The Marsh Dinosaurs, Part II" (<https://extinctmonsters.net/2012/09/23/extinct-monsters-the-marsh-dinosaurs-part-ii/>). *EXTINCT MONSTERS*. September 23, 2012. Retrieved January 10, 2022.
103. Brown, Barnum (November–December 1932). "A Spine-Armored Saurian of the Past". *Natural History*. **32** (6): 493–496.
104. McGinnis, H. J. (1984). *Carnegie's Dinosaurs*. Pittsburgh: Carnegie Institute.

External links

- Europe's First Stegosaurus Boosts Pangaea Theory (<https://web.archive.org/web/20081202151845/http://www.foxnews.com/story/0,2933,248979,00.html>)

Retrieved from "<https://en.wikipedia.org/w/index.php?title=Stegosaurus&oldid=1086119143>"

This page was last edited on 4 May 2022, at 08:16 (UTC).

Text is available under the Creative Commons Attribution-ShareAlike License 3.0; additional terms may apply. By using this site, you agree to the Terms of Use and Privacy Policy. Wikipedia® is a registered trademark of the Wikimedia Foundation, Inc., a non-profit organization.