

# DATA 609 - Final Project

*Daina Bouquin, Christophe Hunt, Christina Taylor*

*April 16, 2017*

## Contents

0.1	Textbook Part III: . . . . .	2
0.2	Exercise 1. Find $a(t)$ if $k_1 = k_2$ . . . . .	3
<b>1</b>	<b>TODO complete this exercise</b>	<b>4</b>
1.1	Exercise 2. . . . .	4
<b>2</b>	<b>TODO complete this exercise</b>	<b>4</b>
2.1	Exercise 3. . . . .	4
<b>3</b>	<b>4. Comparison of the Model's Predictions with Experimental Data</b>	<b>5</b>
3.1	4.1 Solving for $d(t)$ . . . . .	5
3.2	4.2 The Formula for the Amount of Feces . . . . .	5
<b>4</b>	<b>TODO resolve (c)</b>	<b>5</b>
4.1	Exercise 4. . . . .	5
<b>5</b>	<b>TODO refine answers for B and C</b>	<b>6</b>
5.1	Exercise 5. . . . .	6
<b>6</b>	<b>TODO show work for Exercise</b>	<b>6</b>
6.1	Exercise 6. . . . .	6
<b>7</b>	<b>TODO show work for Exercise</b>	<b>7</b>
7.1	Exercise 7. . . . .	7
7.2	Exercise 8. . . . .	7
7.3	Exercise 9. . . . .	9


## 0.1 Textbook Part III:

### 0.1.1 The problem

The digestive processes of sheep can highlight the nutritional value in varied feeding schedules or varied food preparation. This is especially important when raising sheep for commercial purposes.

### 0.1.2 The digestive process

Sheep are a cud-chewing animal which means that unchewed food goes through a series of storage stomachs called the rumen and the reticulum. The process is illustrated below:



### 0.1.3 The experiment

The digestive process is most observable at the beginning and at the end, we can observe and control what goes in and what comes out.

### 0.1.4 The model

Suppose that at  $t = 0$  a sheep is fed an amount  $R$  of food which goes immediately into its rumen. This food will pass gradually from the rumen through the abomasum into the duodenum. At any later time  $t$  we shall define:

$r(t)$  = the amount of food still in the rumen;  $a(t)$  = the amount in the abomasum;  $d(t)$  = the amount which by then has arrived in the duodenum.

So  $r(0) = R$ ,  $a(0) = d(0) = 0$ , and, for all  $t > 0$ , (1)  $r(t) + a(t) + d(t) = R$ .

### 0.1.5 The assumptions

Two assumptions are made: (A) Food moves out of the rumen at a rate proportional to the amount of food in the rumen. Mathematically this says:

$$(2) \quad r'(t) = -k_1 r(t)$$

where  $k_1$  is a positive proportionality constant.

(B) Food moves out of the abomasum at a rate proportional to the amount of food in the abomasum. Since at the same time food is moving into the abomasum at the rate given by Equation (2), the assumption says

$$(3) \quad a'(t) = k_1 r(t) - k_2 a(t)$$

where  $k_2$  is another positive proportionality constant.

## 0.1.6 The solutions of the equations

### 0.1.6.1 Solving for $r(t)$

It is straightforward to solve Equation (2) for  $r(t)$ . We just divide through by  $r(t)$  and then integrate from 0 to  $t$ :

$$\int_0^t \frac{r'(t)}{r(t)} dt = - \int_0^t k_1 dt$$
$$\ln\left(\frac{r(t)}{R}\right) = -k_1 t, \text{ since } r(0) = R, \text{ and finally (4) } r(t) = R e^{-k_1 t}$$

### 0.1.6.2 Solving for $a(t)$

Finding  $a(t)$  is a bit more tricky. Applying Equations (4) to Equation (3) we get:

$$(5) \quad a'(t) = k_1 R e^{-k_1 t} - k_2 a(t)$$

Equation (5) probably looks quite different from any you have seen before. Let us try to make a shrew guess what kind of solution it has. It says that the derivative of  $a(t)$  is the sum of two terms,  $k_1 R e^{-k_1 t}$  and  $-k_2 a(t)$ . With luck, this might remind us of the product rule:

$$(6) \quad \text{if } a(t) = u(t) \bullet v(t) \text{ then } a'(t) = u(t) \bullet v'(t) + v(t) \bullet u'(t)$$

Can we pick  $u(t)$  and  $v(t)$  so the terms in Equation (6) match up with the terms in Equation (5)? In other words, can we pick  $u(t)$  and  $v(t)$  so that

$$(7) \quad u(t) \bullet v'(t) = k_1 R e^{-k_1 t}$$

and

$$(8) \quad v(t) \bullet u'(t) = -k_2 a(t)?$$

Since  $a(t) = u(t) \bullet v(t)$ , Equation (8) can be rewritten  $v(t) \bullet u'(t) = -k_2 u(t) v(t)$ , we are in business! The  $v(t)$  factors cancel out, leaving us with

$$u'(t) = -k_2 u(t)$$

which looks very much like Equation (2) and can be solved in the same way.

$$\int_0^t \frac{u'(t)}{u(t)} dt = - \int_0^t k_2 dt$$

Writing  $K = u(0)$ :

$$\ln\left(\frac{u(t)}{K}\right) = -k_2 t$$
$$u(t) = K e^{-k_2 t}.$$

Putting this into Equation (7) gives

$$K e^{-k_2 t} v'(t) = k_1 R e^{-k_1 t}$$
$$v'(t) = \frac{k_1 R}{K} e^{k_2 - k_1} t.$$

If  $k_1 = k_2$  we feel confident you can complete this solution yourself (Exercise 1).

## 0.2 Exercise 1. Find $a(t)$ if $k_1 = k_2$ .

$$a'(t) = u(t) \bullet v'(t) + v(t) \bullet u'(t)$$

Using the solution from equation 8 we get:  $u(t) = K e^{-k_2 t}$

Using the solution from equation 7 we get:  $v'(t) = \frac{k_1 R}{K} e^{k_2 - k_1} t$

Substituting the solutions for  $a(t)$  we get:  $\frac{k_1 R}{K} e^{k_2 - k_1} t * K e^{-k_2 t}$

If  $k_1 = k_2$  then  $e^0 = 1$

$$\frac{k_1 R}{K} t * K e^{-k_2 t}$$

K factors out :  $k_1 R t e^{-k_2 t}$

answer :  $a(t) = k_1 R t e^{-k_2 t}$

---

If  $k_1 \neq k_2$ , then  $k_2 - k_1 \neq 0$  and so we can write

$$v(t) = \frac{k_1 R}{K(k_2 - k_1)} e^{(k_2 - k_1)t} + c$$

Where C is the constant of integration. Then

$$a(t) = u(t) \bullet v(t) = \frac{k_1 R}{k_2 - k_1} e^{-k_1 t} + C K e^{-k_2 t}$$

Using the fact that  $a(0) = 0$ , we get

$$0 = \frac{k_1 R}{k_2 - k_1} + C K$$

$$C K = -\frac{k_1 R}{k_2 - k_1}$$

(9)  $a(t) = \frac{k_1 R}{k_2 - k_1} (e^{-k_1 t} - e^{-k_2 t})$

## 1 TODO complete this exercise

### 1.1 Exercise 2.

(a) Find the time  $t$  at which  $a(t)$  is maximum.

$$a(t) = \frac{k_1 R}{k_2 - k_1} (e^{-k_1 t} - e^{-k_2 t})$$

$$a(t) = \frac{k_1 R}{k_2 - k_1} (\ln k_1 - \ln k_2)$$

$$a(t) = \frac{1}{k_1 R} \frac{(\ln k_1 - \ln k_2)}{k_2 - k_1}$$

$$a(t) = \frac{k_1 R (e^{-k_1 t} - e^{-k_2 t})}{k_2 - k_1}$$

answer: (a)  $t = \frac{\ln k_1 - \ln k_2}{k_1 - k_2}$

(b) Find the maximum value of  $a(t)$ .

answer:  $\frac{k_1 R}{k_2 - k_1} \left[ \left( \frac{k_1}{k_2} \right)^{-\frac{k_1}{k_1 - k_2}} - \left( \frac{k_1}{k_2} \right)^{-\frac{k_2}{k_1 - k_2}} \right]$

## 2 TODO complete this exercise

### 2.1 Exercise 3.

If  $k_1 = 2$  and  $k_2 = 1$ , how much food must the abomasum be able to hold if a meal of amount  $R$  is fed at time  $t = 0$ ?

$$\begin{aligned}
a(t) &= \frac{k_1 R}{k_2 - k_1} (e^{-k_1 t} - e^{-k_2 t}) \\
a(t) &= \frac{k_1 R}{k_2 - k_1} \left( \frac{1}{e^{k_1 t}} - \frac{1}{e^{k_2 t}} \right) \\
a(t) &= \frac{2R}{1-2} (e^{-2t_0} - e^{-1t_0}) \\
a(t) &= \frac{2R}{1-2} (e^{-2 \cdot 0} - e^{-1 \cdot 0}) \\
a(t) &= \frac{2R}{1-2} (e^{-2} - e^{-1}) \quad a(t) = \frac{2R}{-1} (1 - 1) \quad a(t) = -2R \\
\text{answer: } &\frac{1}{2}R
\end{aligned}$$

### 3 4. Comparison of the Model's Predictions with Experimental Data

Equations (4) and (9) are purely theoretical and are based on assumptions. To confirm their accuracy we will see if they agree with experimental data. Since the data concern is fecal excretion as a function of time, we must first translate our results into results on fecal excretion.

#### 3.1 4.1 Solving for $d(t)$

Starting with Equation (1), and using Equations (4) and (9) (still assuming  $k_1 \neq k_2$ ):

$$\begin{aligned}
d(t) &= R - r(t) - a(t) \\
&= R - R e^{-k_1 t} - \frac{k_1 R}{k_2 - k_1} (e^{-k_1 t} - e^{-k_2 t}) \\
&= R - \frac{R}{k_2 - k_1} (k_2 e^{-k_1 t} - k_1 e^{-k_2 t})
\end{aligned}$$

#### 3.2 4.2 The Formula for the Amount of Feces

Recall that  $d(t)$  is the total amount of food which has entered the duodenum by time  $t$ , including food already excreted. Since excretion is not a continuous process, we cannot hope to represent it by an equation involving derivatives. Instead, we simply note that all food arriving in the duodenum is excreted after a certain time delay. Let us suppose that the average time delay is  $T$  hours. In other words, the amount of feces produced by any time  $t > T$  is, on the average, the amount of food which had already entered the duodenum  $T$  hours earlier, at time  $t - T$ . If  $f(t)$  denotes the amount of feces produced by time  $t$ , this says:

$$\begin{aligned}
f(t) &= d(t - T) \text{ for all } t > T \\
(10) \quad f(t) &= R - \frac{R}{k_2 - k_1} (k_2 e^{-k_1(t-T)} - k_1 e^{-k_2(t-T)}) \\
&\text{for all } t > T.
\end{aligned}$$

### 4 TODO resolve (c)

#### 4.1 Exercise 4.

Remembering that  $r(t)$  is the total amount of food which has entered the rumen by time  $t$ . There is a continuous flow from the rumen to the abomasum

- (a) Determine the values of  $t$  for which  $r'(t)$  is, respectively, positive, negative, and zero.

$$r(t) = Re^{-k_1 t}$$

$$= R\left(\frac{d}{dt}(e^{-tk_1})\right) \text{ Using the chain rule } = e^{-tk_1} \frac{d}{dt}(-tk_1)R \text{ derivative of } t \text{ is } 1 : r'(t) = e^{-tk_1} Rk_1$$

answer:  $r'(t) < 0$  for all  $t$

(b) Determine the values of  $t$  for which  $r''(t)$  is, respectively, positive, negative, and zero.

$$= \frac{d}{dt}\left(R\left(\frac{d}{dt}(e^{-tk_1})\right)\right)$$

$$\text{Using the chain rule: } = \frac{d}{dt}(e^{-tk_1} \left(\frac{d}{dt}(-tk_1)\right)R)$$

$$\text{factor out constants: } = \frac{d}{dt}(R - \left(\frac{d}{dt}(t)\right)k_1)e^{-tk_1}$$

$$\text{derivative of } t \text{ is } 1: = \frac{d}{dt}(R(e^{-tk_1} - k_1))$$

$$\text{factor out constants: } = R\left(\frac{d}{dt}(e^{-tk_1})\right)k_1$$

$$\text{using the chain rule: } = -Rk_1e^{-tk_1}\left(\frac{d}{dt}(-tk_1)\right)$$

$$\text{factor out constants and simplify } = -e^{-tk_1}R - \left(\frac{d}{dt}(t)\right)k_1^2$$

$$\text{derivative of } t \text{ is } 1: r''(t) = e^{-tk_1}Rk_1^2$$

answer:  $r''(t) > 0$  for all  $t$

(c) Determine  $\lim_{t \rightarrow \infty} r(t)$

answer:  $\lim_{t \rightarrow \infty} r(t) = 0$

## 5 TODO refine answers for B and C

### 5.1 Exercise 5.

Remembering that  $a(t)$  is the total amount of food which has entered the the abomasum by time  $t$  from the rumen and exiting to the duodenum. There is a continuous flow from the abomasum to the duodenum.

We have previously solved for  $a(t)$  and  $a'(t)$

$$a(t) = \frac{k_1 R}{k_2 - k_1} (e^{-k_1 t} - e^{-k_2 t}) \quad a'(t) = k_1 R e^{-k_1 t} - k_2 a(t)$$

(a) Determine the values of  $t$  for which  $a'(t)$  is, respectively, positive, negative, and zero.

answer:  $a'(t) > 0$  for all  $t < t_0$ ;  $a'(t_0) = 0$ ;  $a'(t) < 0$  for  $t > t_0$ .

(b) Determine the values of  $t$  for which  $a''(t)$  is, respectively, positive, negative, and zero.

answer:  $a''(t) < 0$  for all  $t < 2t_0$ ;  $a''(2t_0) = 0$ ;  $a''(t) > 0$  for  $t > 2t_0$ .

(c) Determine  $\lim_{t \rightarrow \infty} a(t)$

answer:  $\lim_{t \rightarrow \infty} a(t) = 0$

## 6 TODO show work for Exercise

### 6.1 Exercise 6.

(a) Determine the values of  $t$  for which  $d'(t)$  is, respectively, positive, negative, and zero.

answer:  $d'(t) > 0$  for all  $t$  (must have  $k_1 > k_2$ ).

(b) Determine the values of  $t$  for which  $d''(t)$  is, respectively, positive, negative, and zero.

answer:  $d''(t) > 0$  for all  $t < t_0$ ;  $d''(t_0) = 0$ ;  $d''(t) < 0$  for  $t > t_0$ .

(c) Determine  $\lim_{t \rightarrow \infty} d(t)$

answer:  $\lim_{t \rightarrow \infty} d(t) = R$

## 7 TODO show work for Exercise

### 7.1 Exercise 7.

For the function  $f(t)$  follow the instructions of Exercise 4.

(a) Determine the values of  $t$  for which  $f'(t)$  is, respectively, positive, negative, and zero.

answer:  $f'(t) > 0$  for all  $t > T$ .

(b) Determine the values of  $t$  for which  $f''(t)$  is, respectively, positive, negative, and zero.


answer:  $f''(t) > 0$  for all  $T < t < T + t_0$ ;  $f''(T + t_0) = 0$ ;  $f''(t) < 0$  for  $t > T + t_0$ .

(c) Determine  $\lim_{t \rightarrow \infty} f(t)$

answer:  $\lim_{t \rightarrow \infty} f(t) = R$

### 7.2 Exercise 8.

Using all the information in Exercise 4, 5, 6, and 7 to sketch the graphs of  $r$ ,  $a$ ,  $d$ , and  $f$ . You should not have to compute any points except  $r(0)$ ,  $a(0)$ ,  $d(0)$ , and  $f(T)$ , which you already know.





### 7.3 Exercise 9.

Assuming that increased stay in the digestive system implies improved digestion determine from Figure 4 which method of food preparation is more desirable. Why?




Figure 4. Data from Experiments A and D of Table 1 and Figure 2, with curves fitted from Equation (10). Adapted from Blaxter (1956), p. 76.

Figure 1:

Assuming that we are looking for the feed type that stays in the sheep's stomach the longest, then we would say that the medium-ground grass would be the most desirable. We make this assumption as it stands that the sheep requires less feed for the same results.