




Analyzing the Impacts of Stellar and Planetary Parameters to Exoplanet Habitability through Machine Learning

Christina X. Liu¹, Jonathan H. Jiang²

¹Lakeside School, Seattle, WA, ²Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA



INTRODUCTION

Are there any planets other than Earth potentially habitable by humans? What factors affect habitability and how do they impact habitability? Seeking answers to these questions has been an active research field in astrophysics in the past decade.

In this study, we identified the most influential stellar and planetary parameters to exoplanet habitability and analyzed how they impact habitability through the lens of machine learning.

OBJECTIVES

Build high-quality ML models (**Random Forest**, **XGBoost**) for exoplanet habitability prediction.

Perform analysis on the ML models through **SHAP (SHapley Additive exPlanations)** techniques to understand how stellar and planetary parameters impact exoplanet habitability.

DATASET

The primary data sources for this study:


- Planetary Systems Composite Data @ **NASA Exoplanet Archive**: **5,834** confirmed exoplanets.
- Habitable World Catalog (**HWC**), PHL @ UPR Arecibo: **5,599** exoplanets.

The NASA dataset was joined with the HWC dataset. Then the **P_HABITABLE** data field from the HWC dataset to label the training data.

MODEL SELECTION

Research in machine learning shows that tree-based models could be very effective for tabular dataset (on par with deep learning models). In this study, we chose to build **Random Forest** and **XGBoost** classifiers for exoplanet habitability prediction.

FEATURE ENGINEERING & MODEL TRAINING





Random Forest Model Evaluation

Class	Random Forest		
	Precision	Recall	F1 score
1 (Habitable)	0.96	0.94	0.95
0 (Non-Habitable)	0.94	0.96	0.95


XGBoost Model Evaluation

Class	XGBoost		
	Precision	Recall	F1 score
1 (Habitable)	0.94	0.97	0.95
0 (Non-Habitable)	0.97	0.93	0.95


SHAP ANALYSIS



Deep Dive on Random Forest through SHAP




This SHAP beeswarm plot shows that relatively higher stellar radius, planet radius, stellar mass, and stellar effective temperatures contribute to habitability negatively, while relatively higher planet orbit semi-major axis contributes to habitability positively.




This SHAP heatmap plot groups samples (exoplanets) that have the same model predictions and the similar feature impacts together.

SHAP force plot for a positive prediction sample (exoplanet)




SHAP force plot for a negative prediction sample (exoplanet)




MORE SHAP ANALYSIS


SHAP force plot for all samples (exoplanets) - Random Forest



SHAP waterfall plot for a positive prediction sample (exoplanet) - Random Forest



SHAP waterfall plot for a negative prediction sample (exoplanet) - Random Forest



CONCLUSIONS

Random Forest and **XGBoost** models were trained to predict exoplanet habitability with high F1 scores at **0.95**.

Feature importance analysis through **SHAP** identified influential stellar and planetary parameters to habitability. Further **SHAP** analysis on those parameters showed that relatively higher stellar radius, stellar mass, stellar effective temperature, and planet radius have negative impacts on habitability, while relatively higher planet orbit semi-major has a positive impact.

This study demonstrated that machine learning techniques could be effective for both predicting exoplanet habitability and understanding how stellar and planetary parameters impact exoplanet habitability.

FUTURE WORK

Train a **Neural Network** model for habitability prediction and SHAP analysis. Compare with tree-based models.

Study **Planetary Systems** as a whole to understand what planetary systems might be more likely to host habitable planets.