

Metabolism across brain cell types, disease and the *C4b*⁺ oligodendrocyte state

OligoMetab project · in-house Xenium / Visium / Falcão data plus sixteen public datasets · 2026-09-28

The question. Do for metabolism (ketone bodies, glycolysis and lactate, mitochondrial oxidation, fatty acids, lipids and their regulators) what OligoC4b did for complement: which cell types run which pathway, how that changes with disease, age, demyelination and lesion distance in every annotated cell type, and how it relates to the *C4b*⁺ disease-associated oligodendrocyte state.

The short answer. Metabolic identity of the cell types is stable across platforms and species. Neuroinflammation (EAE) shifts the whole tissue, not one cell type: cholesterol synthesis and the TCA cycle fall and lipid uptake and storage rise in nearly every cell type, steepest near lesions, where OPCs and newly formed oligodendrocytes lose cholesterol synthesis most. The disease-associated oligodendrocyte state, both in situ (Xenium, 107 samples) and in sorted single cells (Falcão), is less cholesterol and myelin-lipid synthesis, more lipid uptake and storage (*Plin2* / *Plin4*, *Abca1*, *Lpl*, *Cd36*), a *Hk2* / *Pdk4* / *Ldha* glycolytic-switch signature, *Bdh1* induction and less MCT1 (*Slc16a1*). In human MS oligodendrocytes cholesterol synthesis is, if anything, higher, and the hypoxia / nutrient-stress genes *Hif1a*, *Txnip* and *Ddit4* are up. Across all mouse datasets the one metabolic feature that travels with *C4b* is lipid handling (*Apod*, *Plin4*, *Abca1*), not energy metabolism and not ketone bodies: the ketone enzymes belong to astrocytes and OPCs, the ketone receptor *Hcar2* to microglia, and *Hcar2*⁺ and lactate-exporting *Slc16a3*⁺ cells are enriched around *C4b*-high oligodendrocytes in EAE as *C5aR1*⁺ cells were in OligoC4b.


1. Data and gene panel

Datasets are those of OligoC4b: Xenium mouse AD (347-gene panel, one section per genotype × age), Xenium mouse EAE (5K panel, 107 samples, lesion-distance annotation), Visium aging mouse brain (6 sections), Falcão et al. 2018 EAE scRNA-seq (sorted oligodendrocyte-lineage cells, microglia, VLMC) and sixteen public whole-transcriptome datasets (mouse AD, aging, demyelination, spatial AD; human AD and MS single-nucleus and Visium). The gene panel has 183 mouse genes in 21 pathway groups (ketone-body synthesis and utilisation, ketone / SCFA receptors, monocarboxylate and glucose transporters, glycolysis, pyruvate fate, pentose phosphate, glycogen, TCA, anaplerosis, OXPHOS, mitochondrial biogenesis, fatty-acid uptake and β-oxidation, lipid, cholesterol and myelin-lipid synthesis, cholesterol / lipid transport and storage, nutrient-sensing regulators). Pathway scores are `scanpy.tl.score_genes` values (set mean minus a size-matched control set, log scale) and are compared by difference; single genes by log2 fold change of pseudobulk means; groups by Mann–Whitney across samples when both have ≥3 samples.


Platform	Panel genes present
Xenium mouse AD	3 of 183 (<i>Apoe</i> , <i>Apod</i> , <i>Acsbg1</i>): metabolism is not measurable
Xenium mouse EAE (5K)	100 of 183; no ketogenesis / ketolysis enzymes, 2 of 15 OXPHOS, 3 of 12 cholesterol-synthesis genes
Visium, Falcão, public	177–183 of 183; metabolic genes are quantifiable in human data, unlike <i>C4A/C4B</i>

2. Division of labour between cell types

Astrocytes and neurons carry the highest glycolysis, TCA and OXPHOS scores; homeostatic oligodendrocytes the highest lipid-synthesis and newly formed oligodendrocytes the highest cholesterol-synthesis score; microglia, macrophages and foamy myeloid cells the highest lipid-transport / storage score with the lowest cholesterol synthesis. In sorted cells OPC / COP / NFOL have the highest ketone-utilisation and TCA scores of the lineage. The same profile holds across the seven public mouse single-cell datasets. *Hcar2* is microglial everywhere; *Slc16a1* (MCT1) is in 39 % of EAE oligodendrocytes in situ and 29–57 % of whole oligodendrocytes in droplet data.




Xenium EAE: mean pathway score by cell type (27 types). Scores are relative to size-matched control genes; 0 means no enrichment.




Public mouse datasets: pathway score by coarse cell type, z-scored within each dataset and averaged.

3. EAE is a tissue-wide metabolic shift

Comparing 92 EAE with 15 control samples per cell type, cholesterol synthesis falls and lipid transport / storage rises in essentially every cell type with enough control samples, and the TCA score falls in most; glycolysis rises only in microglia and DA oligodendrocytes. Gene-level, *Hcar2*, *Cd36*, *Hk2*, *Plin2*, *Plin4*, *Abca1*, *Lpl*, *Slc16a3* and *Txnip* rise across many cell types. Toward lesions, glycolysis and TCA fall in neurons, oligodendrocytes, endothelial and vascular cells, lipid storage rises almost everywhere, and cholesterol synthesis falls most in OPCs (-0.33), newly formed oligodendrocytes (-0.29) and DA oligodendrocytes (-0.16).




Xenium EAE: pathway-score difference EAE – CONTROL per cell type, pseudobulk per sample; * Mann–Whitney $p < 0.05$.




Xenium EAE: cholesterol-synthesis score by cell type and lesion distance, relative to each cell type's value beyond 500 μm .

4. The disease-associated oligodendrocyte state

Within each of 107 samples, DA oligodendrocytes score lower than homeostatic ones on glycolysis (−0.10), TCA (−0.12), lipid synthesis (−0.15) and cholesterol synthesis (−0.16) and higher on lipid transport / storage (+0.15; 99 % of samples). Genes up: *Hk2* (3.5×), *Mlxipl*, *Hcar2*, *Cd36*, *Lpl*, *Slc16a3*, *Slc2a4*, *Txnip*, *Fabp7*, *Plin2*, *Abca1*, *Ldha*, *Pdk1*; down: *Slc2a3*, *Eno2*, *Ppargc1a*, *Got2*, *Mpc2*, *Pcx*, *Fh1*, *Ogdh*, *Mfn2*, *Dnm1l*, *Elovl1*, *Pfkfb*. Sorted cells (Falcão, EAE-MOL vs control-MOL) confirm the core intrinsically: cholesterol synthesis −1.0 (*Hmgcs1*, *Fdps*, *Cyp51*, *Dhcr7*, *Dhcr24*, *Sqle* each 20–30 % lower), myelin-lipid synthesis −0.30, lipid transport / storage +0.95 (*Plin4* 5×, *Plin2* 3×, *Abca1* 2.5×), *Pdk4* 4×, *Ldha* 2.2×, *Cpt1a* 2.5×, *Bdh1* 2.3×, *Slc16a1* −0.24 log2. The direction of the glycolysis / TCA scores differs between the two datasets (down in situ, up in sorted cells) because the 15-gene Xenium set is weighted toward *Eno2*, *Pfkfb* and *Slc2a3*, which fall, and the whole-transcriptome set toward *Aldoa*, *Tpi1*, *Gapdh* and *Ldha*, which rise; the gene tables are the safer read-out.




Xenium EAE: median paired difference in pathway score between DA and homeostatic oligodendrocytes (left) and between *C4b*-high and *C4b*-negative lineage cells (right) across samples; red = Wilcoxon p < 0.05.




Falcão 2018 sorted cells: mean pathway score by cluster (OPC → COP → NFOL → MOL, EAE-MOL clusters, microglia, VLMC).

5. Public datasets: small pathway shifts, consistent genes


Pathway-score differences in public oligodendrocytes are mostly within ± 0.05 . Clear ones: Schirmer MS oligodendrocytes more glycolytic (+0.16) and oxidative (+0.11) with more lipid storage; aged oligodendrocytes (Ximerakis) less OXPHOS, cholesterol and myelin-lipid synthesis; slightly lower ketone synthesis / utilisation in MS oligodendrocytes in three human sets; less lipid synthesis at Braak 6. At the gene level the same genes move in most of the 14 oligodendrocyte contrasts: *Lpl* and *Hk2* up in 10, *Apoe* / *Plin2* / *Fabp5* up in 6–7, *Slc16a1* down in 5, *Gal3st1* in 6, *Acsf6*, *Ppara*, *Lss*, *Srebf1*, *Cpt1c*, *Scd1* in 4–5. Microglia show the most reproducible signature of all: *Lpl* up in 12 of 13 contrasts (median log2FC 2.8), *Apoe* and *Plin2* in 8, *Pparg* and *Cd36* in 6–8, *Hcar2* in 6 of 10. Astrocytes gain *Cd36* and *Fabp7* and lose ketone utilisation in every MS dataset. Human MS oligodendrocytes have modestly higher cholesterol-synthesis scores than controls and more *Hif1a*, *Txnip* and *Ddit4*.



Public datasets, oligodendrocytes: pathway-score difference per contrast (disease / aged / demyelinated vs reference); * Mann-Whitney $p < 0.05$ across samples.




Public datasets, microglia: the same. Lipid transport / storage rises in nearly every disease contrast.




Public datasets, oligodendrocytes: pseudobulk log2 fold change of the lipid / regulator focus genes.

6. The *C4b* axis

Inside DA oligodendrocytes (Xenium EAE) the metabolic genes that track *C4b* are lipid-droplet / lipid-handling genes (*Plin4* rank 2 of 3,673 panel genes, *Apod* 42, *Cers2* 72, *Fasn* 86); no metabolic pathway is enriched as a set, and across the lineage glycolysis, TCA and mitochondrial-biogenesis genes are anti-correlated with *C4b*. In sorted MOL lipid transport / storage is the top pathway (*Apod* p 0.46, *Plin4* 0.40, *Abca1* 0.32) and cholesterol synthesis is significantly anti-correlated. In the public mouse datasets the correlations are weak ($|\text{mean } \rho| < 0.1$), *Apod* recurs among the top ranks (8–42), lipid transport / storage is the enriched pathway in LPC / cuprizone, *Serpina3n*-cKO and 5XFAD, and *C4b*-high oligodendrocytes have a higher lipid-storage score in all seven datasets. Ketone-body genes are not part of the program.




Public mouse datasets: mean Spearman correlation with *C4b* of each pathway's genes inside oligodendrocytes; * Mann-Whitney vs all other genes $p < 0.05$.




Public mouse datasets: pathway score in *C4b*-high minus *C4b*-negative oligodendrocytes; * paired Wilcoxon across samples $p < 0.05$.

7. Spatial partners of C4b-high oligodendrocytes (Xenium EAE)


Hcar2⁺ cells (median ratio 1.7), *Hcar2*⁺ myeloid cells (1.8), *Slc16a3*⁺ astrocytes (2.3), *Hk2*⁺ (1.8), *Pdk1*⁺ (1.3) and *Plin2*⁺ (1.4) cells are enriched within 30 μm of C4b-high compared with C4b-negative oligodendrocytes in 87 samples (paired Wilcoxon $p < 10^{-5}$); *Slc2a1*⁺ cells are not. The enrichment survives matching for lesion distance (1.2–1.75) and restriction to non-lesion tissue (1.3–2.2).



Per-sample ratio of the fraction of target-positive neighbours around *C4b*-high versus *C4b*-negative oligodendrocytes (30 μm).




The same ratio after stratifying source oligodendrocytes by lesion distance.



Xenium EAE, one peak-EAE sample: *C4b*, *Slc16a1*, *Ldha*, *Hk2*, *Hcar2*, *Cpt1a* and *Plin2*.

8. Aging (Visium, 6 sections)

Whole-section glycolysis, TCA and OXPHOS scores rise slightly with age ($p \approx 0.03$ – 0.04 , slopes of a few hundredths over 15 months). In white-matter-rich spots the lipid-synthesis score falls, *Fa2h*, *Ugt8a*, *Cyp51* and *Hmgcs1* decline by 10–19 %, *Apoe* and *Plin2* rise and *Ddit4*, *Slc2a1* and *Sirt1* fall: a mild version of the EAE pattern.



Visium aging, white-matter-rich spots: pathway score per section by age group.

9. Caveats

1. Xenium AD has one section per genotype $\times$ age and only three metabolic genes.
2. The EAE 5K panel misses ketogenesis, ketolysis, most OXPHOS and most cholesterol-synthesis genes.

3. In situ transcripts inside a segment can come from a neighbouring cell (*Hcar2*, *Cd36*, *Slc16a3* in "oligodendrocytes"); the sorted Falcão cells are the control, and cells are the replicates there.
4. Pathway scores depend on set composition; compare by difference and check the gene tables.
5. Pseudobulk tests need ≥ 3 samples per group; Park, LPC / cuprizone, *Serpina3n*-cKO and several human contrasts fall short.
6. Nuclei detect cytoplasmic and mitochondrial metabolic transcripts far less than whole cells; microglial ambient RNA in demyelinating-lesion nuclei inflates *Apoe*, *Lpl* and *Hcar2* in "oligodendrocytes".
7. Human *C4B* is not quantifiable, so the *C4b* axis is mouse-only.

10. Suggested next steps

- Lipid-droplet staining (PLIN2 / PLIN4, BODIPY) and cholesterol-synthesis readouts (HMGCS1, SQLE) in *C4b*⁺ oligodendrocytes in EAE and aged white matter, the one metabolic feature that travels with the state in every dataset.
- MCT1 protein in DA oligodendrocytes and the lactate supply to axons near lesions (*Slc16a1* down, *Slc16a3*⁺ astrocytes and *Hcar2*⁺ myeloid cells enriched in the niche).
- Re-quantify human *C4A/C4B* (OligoC4b) so the *C4b* axis can be tested in the human MS sets where cholesterol synthesis moves the other way.
- Metabolic flux (Seahorse or ¹³C tracing) on sorted DA versus homeostatic oligodendrocytes to resolve the glycolysis / TCA direction that the two RNA read-outs disagree on.

11. Where everything lives

Repository `oligoMetab` (GitHub, [christoffermattssonlangseth/oligoMetab](https://github.com/christoffermattssonlangseth/oligoMetab)): executed notebooks under `notebooks/analysis/`, sources under `notebooks/src/`, the gene panel and helpers in `scripts/oligometab.py`, summary tables in `results/`, and plain-language documentation in `docs/` (`metabolism_findings.md`, `gene_panel.md`, `notebooks.md`, `public_datasets.md`). Data are the OligoC4b objects on the group's analysis machine.