

Boxford Open Land Trust's

Property Map and Ecological Management Plan

for the

Hoveys Pond Conservation Area

(POTTER'S FARM)

West Boxford, MA

Property Description

The Hoveys Pond Conservation Area was acquired by the Boxford Open Land Trust in the spring of 1994. The area is approximately 70 acres with a diverse assemblage of habitat types. Wetlands cover just less than half of the property with oak and hemlock dominating the uplands. Both wooded and shrub swamp wetlands occur as well as five state certified vernal pools with additional pools along the property boundary. Powerline corridors bisect the property in two places. The larger of the two corridors bisects the extreme northern part of the property while the smaller corridor bisects the wetlands along the western edge.

Little is known of the property historically. The land is either too wet or too rocky to have been used for agriculture. However, evidence of pasturing remains. Stonewalls outline the property and bisect it. Also remnant juniper and red cedar persist, although many are dead or dying, in the upland portions of the property. It appears there were attempts to drain the wetland to the west of the property. Very steep banks with almost 90 degree angles where the stream flows from the wetland, clearly suggests the stream had been excavated.

This property and the surrounding area is rich in Amphibian and Reptile species. Three Massachusetts state listed rare species have been documented from the property with reports of a third rare species on adjacent properties. The diverse variety of wetland and upland habitats is no doubt responsible. Hoveys Pond and 50 acres of undeveloped land lie just to the north. A tree farm and Sperrys Pond, with its open marshes, lies just to the south. To the west lies houses and route 133. To the east lies the subdivision of Belvedere and Plummer Roads, and 67 acres of undeveloped lands. Efforts to protect much of the undeveloped lands adjacent to the property are underway.

Passive management, including that for wildlife, is the primary management goal. No forestry management is planned because wetlands, rare species, vernal pools,

powerlines, hemlock stands, ledge outcroppings and poor access make the property poorly suited for forestry. More than half of the property has little to no timber value. Even if access of the kind necessary for heavy equipment was negotiated, buffer areas would need to be established around the wetlands and vernal pools, restricting additional acreage from consideration for timber.

However, wildlife management on a scale the land trust could afford and manage is desirable. Consideration for the rare species and those species which are present now or could be present based on current habitat, are the primary species for management. The land trust does not wish to impact the rare species in any negative way thus care should be taken to identify the needs of these species and safeguard these needs.

An additional management consideration, which relates to wildlife, is the control and eradication of exotic and invasive plant species. Several exotic species do occur on the property and include: European buckthorn, purple loosestrife, Japanese honeysuckle and Japanese barberry. The honeysuckle and barberry occur in the uplands and are uncommon. These species could easily be removed. The buckthorn and loosestrife have already become extensively established in the wetlands and pose a removal problem. The few buckthorns that do occur in the uplands should be removed immediately to prevent spreading. Allowing these species to persist and spread will likely degrade habitat for wildlife by competing with native species.

Adjacent property: description and current use.

For planning and land management purposes, the Trust should identify the ownership and land use patterns of adjacent lands. Collectively these properties influence the types of wildlife found in the area and on the Trust's property.

To the south of the property lies Sperrys Pond. This is a small, shallow pond with extensive marshes, especially to the west of the pond. Sperrys Pond and its associated wetlands are likely the reason why there is a diversity of reptiles and amphibians in the area. Much of the "buildable" land around the pond has been developed. The exception is King's Tree Farm. This is a 28 acre property which lies between Sperrys Pond and the Hoveys Pond Conservation Area. Approximately 10 acres of this property is upland. This upland is being managed as a tree nursery and apparently this use is compatible with at least some of the rare species. The remaining acreage is marsh associated with Sperrys Pond.

Most of the lands to the west have been developed as well. Three small parcels totaling approximately 9 acres, which abut the Conservation Area, remain undeveloped. Two of these properties are promising turtle nesting areas. The soils are well drained sand and gravel. The current owner of these lots are unknown but it is likely they will end up as building lots, although topography and wetlands suggest these lots are less than ideal for development.

To the north and northwest lies the Nason's property and Hoveys Pond. The Nason's property is approximately 50 acres of mixed woodlands with powerlines bisecting the property roughly in half. This property has extensive frontage on Washington Street and Hoveys Pond. Two small fields also occur on this property. Hoveys Pond is a 45 acre pond with marshes. Three quarters of the shoreline is undeveloped.

To the east lies the Bergstrom lot and New England Power Company land. The Bergstrom lot is a narrow 17 acres which parallels the Conservation Area. This lot is

mostly forested except where the powerlines bisect it. This lot also has frontage on Hoveys Pond. Because this lot is land locked, it is doubtful the property will be developed. However, access is possible through the Nason's property. Finally, the New England Power Company owns 50 acres beyond the Bergstrom lot. This property abuts the Conservation Area as well and also has frontage on Hoveys Pond. This property is mostly upland mixed forest except for the powerlines and several forested and shrub swamps.

Beyond these properties along the remaining undeveloped shoreline of Hoveys Pond lies 60+ acres of farmland. Both fields and powerline habitat are common in the area, however, little is permanently protected. Efforts should be made to identify those habitats and lands which are utilized by the rare turtle species so protection efforts can then concentrate on these areas.

Scale

1" = 400'

70 Acres

Boxford Open Land Trust
West Boxford

BASE MAP

Scale

1" = 400'

70 Acres

SCALE
1" = 400'

--- TRAIL/DIRT ROAD


--- FENCE

--- STREAM

--- FIELDS

--- FENCE

Boxford Open Land Trust
West Boxford



SCALE
1" = 400'

----- TRAIL / DIRT ROAD


●●●● STONEWALL

..... STREAM


↓ WETLAND

☼ LEDGE

COVER TYPES



COMPARTMENTS



Doxford Open Land Trust
West Doxford

COVER TYPE

Scale

1" = 400'

SCALE

1" = 400'

TRAIL / DIRT ROAD


STONE WALL

STREAM

WETLAND

WOOD

Boxford Open Land Trust West Boxford



SCALE
1" = 400'

----- TRAIL / DIRT ROAD

●●●● STONEWALL

..... STREAM

↓ WETLAND


⊙ LEDGE

COMPARTMENTS

COMPARTMENTS

Scale

1" = 400'



Wildlife Management Goals

TYPE

- WS wooded swamp
- SS shrub swamp
- M marsh
- OW open water
- V vernal pond
- H hemlock stand
- HD hemlock dominated mixed forest
- RMD red maple / deciduous forest
- ROD red oak / deciduous forest
- WPD white pine / deciduous forest
- P powerlines, old field

Wildlife Management Goals

Currently there are no plans to remove any timber from the property. This is likely to remain the case even in the future. Thus management goals will concentrate on wildlife. Tree cutting will be limited to those trees needed in the creation of brush piles, to provide cover for small mammals. Additional tree cutting may be necessary around selected specimen trees to maintain or enhance their wildlife and aesthetic values. Cutting equipment will be limited to what can be hand carried.

Since management will be carried out by volunteers, management goals will reflect this fact. The land trust will rely on volunteers for most of the equipment and materials as well. The property has been divided into compartments to outline management goals and increase efficiency in implementing these goals. Compartments vary in size and shape and take advantage of existing natural and human made boundaries which are easily identifiable. Specific management goals are discussed within the following compartment descriptions. However, all management must either be compatible with, or benefit, the rare species known to occur on the property.

Arctic Pika to be undertaken

1. Build 3 brush piles along the powerline easement edge of the compartment, see Decker and Kelly (1992) start and finish in 1995.
2. Create crickets, see Decker and Kelly (1992), start in 1995.

Specific wildlife habitat information: A small semi-permanent pond exists just over the property boundary in the south east corner of the compartment. Spotted salamanders have been observed in this pond in the spring. Wood ducks have been observed feeding in this pond in the fall.

Compartment A

Location: Compartment A is the most northern part of the property. It is a triangular piece with stonewalls marking the eastern and northern boundaries. The southern boundary is marked by the powerlines.

Description: Compartment A is approximately 2 acres in area. Drainage is good and there is a slight southern aspect. Red oak and hemlock are the dominant species but pignut hickory also occurs. Hop hornbeam represents the scarce understory. There are very few shrubs or herbaceous plants. A few low blueberry and maple-leaved viburnum are scattered throughout the compartment. Areas of exposed bedrock occur. A dirt road loops through this compartment from the powerlines. An additional dirt road enters this compartment from the adjacent property.

Management Objectives and Priorities: Management will consist of enhancement of cover for small mammals and habitat for cavity nesters.

Activities to be undertaken:

1. Build 3 brush piles along the powerline easement edge of the compartment, see Decker and Kelly (1982) start and finish in 1995.
2. Create cavities, see Decker and Kelly (1982), start in 1995.

Specific wildlife habitat information: A small semi-permanent pool exists just over the property boundary in the south east corner of the compartment. Spotted salamander egg clusters have been observed in this pool in the spring. Wood ducks have been observed feeding in this pool in the fall.

Compartment B

Location: Compartment B consists of the powerline easement that bisects the northern end of the property. The boundaries to the north and south are the forest edges. Boundaries to the east and west are broken stonewalls.

Description: The area is approximately 4 acres and drainage is good. The easement is 200' wide with three sets of powerlines. Habitat is typical "old field" with grasses, shrubs and herbaceous plants dominating. Topography is both dramatic and gently sloping. The highest point on the property is near the center of the powerlines and provides excellent views. This high point has exposed ledge and is covered with bearberry, juniper, bear oak and little bluestem. A small shrub swamp lies at the base of the western slope. Other species include but are not limited to: red oak, birch, sassafras, low blueberry, sweet fern and goldenrod. A dirt road meanders under the powerlines.

Management Objectives and Priorities: Enhancement of small mammal cover and habitat for cavity nesting birds of open habitat.

Activities to be undertaken:

1. Build 3 brush piles along the middle of the compartment, see Decker and Kelly (1982), start and finish in 1996.
2. Construct and place bluebird / tree swallow nesting boxes along powerlines. Boxes should be placed back to back and secured on poles with predator guards. To minimize or avoid vandalism, boxes should be removed immediately after the nesting season and not installed until late March / early April, start in 1995.
3. Construct and place 1 kestrel nesting box along powerlines, start and finish in 1996.
4. Talk to power company regarding "soft edges" for powerline easement.

Compartment C

Specific wildlife habitat information: Bluebirds and kestrels have been observed using this area. Besides placing nesting boxes on the Land Trust's portion of the powerlines, efforts should be made to establish a "bluebird trail" along the adjacent properties as well. Red-tailed hawks frequent this area and "improvements" for small mammals will benefit these raptors. Also, prairie warblers and field sparrows occur along the powerlines as well. Managing for soft edges along the powerline cut will benefit these and other early successional species. If power company is unwilling to oversee this "habitat improvement" then perhaps the land trust through its volunteers could take on this responsibility.

Topography is gently rolling with a ravine bisecting the compartment. An intermittent stream flows through this ravine and into a small shrub swamp along the north side of the compartment which extends into the powerline easement. An intermittent stream drains this wetland to the west. Red oak is dominant but white oak, pignut hickory and red maple also occur. Hop hornbeam is common throughout understory. The majority of trees are larger pole sized. The multiple trunking of many of these trees and their steep age suggests the compartment was cut over in the recent past. Shrubs consist of purple-leaved viburnum, low blueberry and highbush blueberry. Ferns and clubmosses dominate herbaceous layer, but these are sparse. Rock outcroppings occur throughout.

Management Objectives and Priorities: Enhance mast production, small mammal cover and habitat for cavity nesting birds.

Activities to be undertaken:

1. Selectively remove smaller trees from around specimen trees as 1 use for construction of brush piles.
2. Build 3 brush piles along forest edge with powerlines, see Decker and Kelly (1997), start in 1997.
3. Create snags $\geq 6"$ dbh, see Decker and Kelly (1997), start in 1997.

Compartment C

Location: Compartment C is located to the north of the property. The northern boundary is the powerline easement. A stonewall runs the length of the western boundary. A stonewall also runs the length of the southern boundary where it intercepts a swamp. The eastern boundary is marked by the shrub swamp and edge of hemlock forest.

Description: Compartment C is approximately 6 acres in size and drainage is good.

Topography is gently rolling with a ravine bisecting the compartment. An intermittent stream flows through this ravine and into a small shrub swamp along the north side of the compartment which extends into the powerline easement. An intermittent stream drains this wetland to the west. Red oak is dominant but white oak, pignut hickory and red maple also occur. Hop hornbeam is common throughout understory. The majority of trees are larger poll sized. The multiple trunking of many of these trees and their even age suggests the compartment was cut over in the recent past. Shrubs consist of maple-leaved viburnum, low blueberry and highbush blueberry. Ferns and clubmoss dominate herbaceous layer, but these are sparse. Rock outcroppings occur throughout.

Management Objectives and Priorities: Enhance mast production, small mammal cover and habitat for cavity nesting birds.

Activities to be undertaken:

1. Selectively remove smaller trees from around specimen trees and use for construction of brush piles.
2. Build 3 brush piles along forest edge with powerlines, see Decker and Kelly (1982), start in 1997.
3. Create snags > 6" dbh, see Decker and Kelly (1982), start in 1997.

4. Maintain owl nesting platform.

Compartment D1

Specific wildlife habitat information: A steel drum platform consisting of the bottom third of a 55 gallon drum container, was discovered mounted in the crotch of a tree when the land trust purchased the property. Inquiry into the platform lead the land trust to believe that an owl "researcher" well known to the area had constructed the platform for owls. It is not known if the platform is being utilized by owls. A great-horned owl has been seen in the area.

Description: Compartment D1 is approximately 3 acres in size and is well drained.

Hemlock is the dominant species. Shagbark hickory, pignut hickory, white ash and hop hornbeam occur as well. Very little shrub and herbaceous layer exists. However, some maple-leaved viburnum and low blueberry does occur and some ferns are clustered around the rock outcroppings. The topography is gently rolling with some abrupt depressions.

Management Objectives and Priorities: None

Activities to be undertaken:

1. create several hemlock snags for woodpecker feeding trees, see Decker and Kelly (1982), start in 1993.

Specific wildlife habitat information: A vernal pool occurs along the property line in the northeast corner on the adjacent property. It is likely that amphibians which use this pool for breeding in the spring, utilize the uplands of this compartment throughout the rest of the year. A great-horned owl has been seen in this hemlock forest. This is a good location for song birds which require coniferous woodlands for breeding. A hemlock snag in-between compartments D1 and D2 shows evidence of pileated woodpecker feeding.

Compartment D1

Location: Compartment D1 is located roughly in the center of the property along the eastern boundary. The deciduous forest of compartment C establishes the northern boundary. The western boundary is the edge of the central swamp and a stonewall marks both the southern and eastern boundaries.

Description: Compartment D1 is approximately 5 acres in size and is well drained. Hemlock is the dominant species. Shagbark hickory, pignut hickory, white ash and hop hornbeam occur as well. Very little shrub and herbaceous layer exists. However, some maple-leaved viburnum and low blueberry does occur and some ferns are clustered around the rock outcroppings. The topography is gently rolling with some abrupt depressions.

Management Objectives and Priorities: None

Activities to be undertaken:

1. create several hemlock snags for woodpecker feeding trees, see Decker and Kelly (1982), start in 1998.

Specific wildlife habitat information: A vernal pool occurs along the property line in the northeast corner on the adjacent property. It is likely that amphibians which use this pool for breeding in the spring, utilize the uplands of this compartment throughout the rest of the year. A great-horned owl has been seen in this hemlock forest. This is a good location for song birds which require coniferous woodlands for breeding. A hemlock snag in-between compartments D1 and D2 shows evidence of pileated woodpecker feeding.

Compartment D2

Location: Compartment D2 is located south of compartment D1 along the eastern boundary of the property. These two compartments are separated by a narrow band of deciduous forest. The compartment runs north and south, tapering at its ends. A stonewall runs the length of the eastern boundary. The limits of the hemlock forest mark the western boundary. Compartment D2 is approximately 3 acres with good to fair drainage.

Description: Compartment D2 is approximately 3 acres and drainage is good. Hemlock is the dominant species with red oak, scarlet oak, white oak and red maple occurring as well. Clubmoss, partridgeberry and ferns are sparse. No shrubs occur. Topography is gently sloping to the east with a ravine along the eastern edge. An intermittent stream flows through this ravine in the spring and wet periods. Rock outcroppings occur throughout compartment.

Management Objectives and Priorities: None

Activities to be undertaken:

1. create several hemlock snags for woodpecker feeding trees, see Decker and Kelly (1982), start in 1999.

Specific wildlife habitat information: A vernal pool lies near the north center of this compartment. Spotted salamander and wood frog egg masses have been found in this pool in the spring.

Compartment E

Location: Compartment E is located at the center of the property. A stonewall marks the western boundary. The eastern boundary is along the central swamp. A stonewall delineates both the northern and southeastern boundaries.

Description: Compartment E is approximately 8 acres with good to fair drainage.

Topography is gently sloping. The compartment slopes to the east and west. Red oak dominates with pignut hickory, white oak and shagbark hickory less common. Hop hornbeam is dominant understory species. Shrubs are represented by maple-leaved viburnum, huckleberry, low blueberry and highbush blueberry. The low areas along the wetland edge to the west include red maple, white ash, white birch, beech and yellow birch. The compartment runs north and south and roughly tapers at its ends. It is fairly narrow with wetlands to both sides.

Management Objectives and Priorities: Enhance habitat for cavity nesters.

Activities to be undertaken:

1. create snags, see Decker and Kelly (1982), start and finish in 2000.

Specific wildlife habitat information: Three low spots have standing water in the spring and may serve as vernal pools. Because this compartment is virtually surrounded by wetlands, opportunities for wildlife are great. Several woodpecker species including red-bellied are common in this area. Additionally there is much evidence of deer and gray squirrels in this area as well, especially along the wetland edges. Oaks are abundant and provide food for many species in the fall. Snags of various sizes do occur except snags with a dbh of 12 inches or greater are scarce.

Compartment F1

Location: Compartment F1 is the majority of the eastern portion of the property. Its eastern boundary is the central stonewall along the wetland. The remaining boundaries are obscure and are not well defined and can be difficult to locate. Broken stonewalls define parts of the northern boundary. Parts of the southern boundary follow the wetland edge. The remaining boundaries are irregular in shape with a narrow portion abutting route 133, (Washington St.).

Description: Compartment F1 is approximately 15 acres. Topography is flat. The majority of the compartment is wetland with three small upland areas along its edges. Red oak is dominant with beech, red maple and white ash also occurring. Along the edge of the upland yellow birch and bitternut hickory with a few spicebush also occur. White pine is rare. There are several types of wetland communities. The forested wetlands are dominated by red maple and black alder swamps. The shrub swamps include black alder, arrowwood, European buckthorn, and dogwoods with a few willows scattered throughout. Sensitive fern, purple loosestrife, cattail and jewelweed represent the herbaceous species. Powerlines bisect the compartment down the middle, running approximately north to south. The outflow of this wetland, at the southeast corner of the compartment, appears to have been excavated in the past in an attempt to drain the wetland.

Management Objectives and Priorities: This compartment, being mostly wetland, is subject to wetland regulations thus any management which alters wetland would need to be approved by the Boxford Conservation Commission, Department of Environmental Protection and likely the Division of Fisheries and Wildlife. However, management of the wetland may prove beneficial to wetland wildlife, particularly rare turtle species. Water level manipulation may assist efforts to control and eradicate the invasive wetland plant

Compartment F2

species. In addition, the increase in water level at times may reduce the shrub cover and increase the emergent vegetation benefiting waterfowl, rails, and other marsh nesting species.

Activities to be undertaken:

1. construct dam at outflow of wetland which will allow manipulation of water levels.
2. establish water level regime that simulates the rise and fall of naturally occurring beaver ponds.

Specific wildlife habitat information: This area provides habitat for many species of wildlife. Bluebirds have been seen along the powerlines. Swamp sparrows, common yellowthroats, gray catbirds and many other songbirds have been seen in this area. The vast number of shrubs provide cover and a source of food for many species. During the fall many species and flocks of birds were observed feeding in and along the wetland edge. Deer tracks and scrapes were common in this area as well. A small open water body along the property boundary provides turtle habitat.

Compartment F2

Compartment G

Location: Compartment F2 is located in the center of the property. Boundaries are edge of wetland.

Description: Compartment F2 is approximately 2 acres and is poorly drained.

Compartment F2 is a wooded shrub swamp dominated by black alder, speckled alder, sweet pepperbush, highbush blueberry and red maple.

Management Objectives and Priorities: None

Activities to be undertaken: None

Specific wildlife habitat information: This swamp could serve as breeding habitat for amphibians but there is no documentation to date.

Compartment G

Location: Compartment G is located along the western boundary of the property.

Compartment F is to the north and the wetland edge marks the boundary. A broken stonewall delineates the southern boundary. The powerlines are to the west and a stonewall to the east.

Description: Compartment G is approximately 1.5 acres in area. It is a narrow compartment which gently slopes to the north. Drainage is fair to poor. Red maple is dominant with some white ash. Understory consists of red maple, white ash, hickory, red oak. Hemlock and hop hornbeam are rare. Spicebush is the dominant shrub with maple-leaved viburnum, highbush blueberry and arrowwood present. Ferns are common. One very large and old white oak occurs along the boundary. This oak is probably two to three hundred years old. This tree is a good landmark.

Management Objectives and Priorities: Maintain mast producing trees and enhance habitat for cavity nesters and small mammals.

Activities to be undertaken:

1. clear competing trees away from the few mast producing trees like oak and hickory.
2. using material from #1 above, create two small brush piles along powerline edge, see Decker and Kelly (1982), start and finish in 2001.
3. girdle trees with >6" dbh, see Decker and Kelly (1982), start and finish in 2001.

Specific wildlife habitat information: None

Compartment H

Location: Compartment H comprises most of the southern portion of the property. A stonewall runs along its northeastern and western boundaries. A stonewall and white pine/deciduous forest delineate the southern boundary. The edge of the hemlock forest and stonewall mark the eastern boundary.

Description: Compartment H is approximately 22 acres and drainage is good. The compartment is very rolling with several ledge outcroppings and kettleholes. Several of the kettleholes fill with water in the spring. Red oak and white oak are dominant. Hop hornbeam is dominant understory species. Red cedar, white pine, pignut hickory, shagbark hickory, beech, swamp white oak and red maple are uncommon. American chestnut and hemlock are rare. Yellow birch and muscledwood occur along low points adjacent to wetland edge. The shrubs consist of highbush blueberry, low blueberry, huckleberry, witch hazel and maple-leaved viburnum. Little to no evidence of herbaceous layer.

Management Objectives and Priorities: Protection of rare species habitat (vernal pools) and the enhancement of habitat for cavity nesters.

Activities to be undertaken:

1. create snags with large dbh, see Decker and Kelly (1982), start in 2002.

Specific wildlife habitat information: Four vernal pools occur in this compartment.

Several rare state listed species have been documented from this area. Both Blanding's and spotted turtles have been found in the vernal pools as well as blue-spotted salamanders. In addition to these species, spotted salamanders, wood frogs, spring peepers and red-spotted newts also occur in the pools. Gray squirrels and chipmunks are abundant and wild turkey

have been seen in this area. Deer rubs have been observed on small trees in fall. The abundance of mast producing trees is no doubt important to the wildlife of the area. Many snags and cavities exist throughout the compartment although there does seem to be a void of snags with a larger dbh.

eastern boundaries. Deciduous forest defines the northern and western boundaries.

Description: Compartment 1 is approximately 1 acre with good drainage. White pine and red oak are dominant with white oak also occurring. Hop hornbeam is the dominant understory species. Shrubs consist of highbush blueberry, low blueberry, maple-leaved viburnum, juniper and huckleberry. Club moss and partridgeberry with a few ferns represent the herbaceous layer. Topography is gently rolling with several small rock outcrops and boulders. Snags of all dbh occur throughout compartment.

Management Objectives and Priorities: None

Activities: None

Specific Wildlife Habitat Information: Gray squirrels are common in this compartment. Also blue jays, chickadees and white-breasted nuthatches are common. This compartment is close to the road and several homes. This may explain the common occurrence of these species since all of these species are common visitors to feeding stations.

Compartment I


Location: Compartment I is located in the southeast corner of the property. Stonewalls delineate the southern and eastern boundaries. Deciduous forest defines the northern and western boundaries.

Description: Compartment I is approximately 2 acres with good drainage. White pine and red oak are dominant with white oak also occurring. Hop hornbeam is the dominant understory species. Shrubs consist of highbush blueberry, low blueberry, maple-leaved viburnum, juniper and huckleberry. Club moss and partridgeberry with a few ferns represent the herbaceous layer. Topography is gently rolling with several small rock outcroppings and boulders. Snags of all dbh occur throughout compartment.

Management Objectives and Priorities: None

Activities: None

Specific wildlife habitat information: Gray squirrels are common in this compartment. Also blue jays, chickadees and white-breasted nuthatch are common. This compartment is close to the road and several homes. This may explain the common occurrence of these species since all of these species are common visitors to feeding stations.




TOPOGRAPHY

Scale

1" = 400'


Sperry



COLOR INFRARED AERIAL PHOTOGRAPH

Scale

1" = 400'



SPECIES LIST (incomplete)

BIRDS

red-tailed hawk (powerlines)
broad-winged hawk
American kestrel (powerlines)
great horned owl
wild turkey
morning dove
northern flicker
red-bellied woodpecker
pileated woodpecker
hairy woodpecker
downy woodpecker
eastern phoebe
blue jay
American crow
tufted titmouse
black-capped chickadee
white-breasted nuthatch
eastern bluebird (powerlines)
American robin
gray catbird
yellow-rumped warbler
prairie warbler (powerlines)
yellow warbler
common yellowthroat
song sparrow (powerlines)
field sparrow (powerlines)
white-throated sparrow
swamp sparrow (powerlines)
rufous-sided towhee (powerlines)
red-winged blackbird
common grackle
scarlet tanager
northern oriole
American goldfinch

MAMMALS

white-tailed deer
gray squirrel
red squirrel
eastern chipmunk

AMPHIBIANS

green frog
wood frog
spring peeper
red-backed salamander
spotted salamander
blue-spotted salamander
red-spotted newt

REPTILES

northern water snake
Blanding's turtle
spotted turtle

References

Beattie, M., C. Thompson and L. Levine. 1993. *Working with Your Woodland: A Landowner's Guide*. University Press of New England, Hanover. 279 pg.

Decker, D. J. and J. W. Kelly. 1982. *Enhancement of Wildlife Habitat on Private Lands*. Information Bulletin 181. Cornell Cooperative Extension Publication.

Decker, D. J., J. W. Kelly, T. W. Seamans, and R. R. Roth. 1990. *Wildlife and Timber from Private Lands: A Landowner's Guide to Planning*. Information Bulletin 193. Cornell Cooperative Extension Publication.

1994