

# Thesis Proposal

Christophe Rouleau-Desrochers

November 25, 2025

## 1 Introduction

### 1.1 Climate change impacts on tree phenology

Research from the past decades has shown convincing evidence that human activity is increasingly affecting many worldwide environmental processes (Ceballos *et al.*, 2017; Intergovernmental Panel On Climate Change (Ipcc), 2023; Laurance, 2007; Parmesan & Yohe, 2003). This can be through land use change and destruction, pollution, invasive species, resource overexploitation and climate change (Driscoll *et al.*, 2018; Parmesan *et al.*, 1999; Wu, 2013). That alone raises major concern, and actions have been deployed to mitigate these impacts, with varying success (e.g. (Campbell *et al.*, 2014)). Even though immediate actions can have positive impacts and potentially reduce some threats to biodiversity, reversing 150 years of human-induced greenhouse gas emissions is harder. These emissions have affected Earth's climate and are projected to keep changing it for many centuries (Intergovernmental Panel On Climate Change (Ipcc), 2023). While there is a scientific consensus that observed climate change is human-caused (Change, 2014; Lynas *et al.*, 2021; Oreskes, 2004), the magnitude and the extent of the consequences that a warming climate will have on biological processes are still debatable (Huey *et al.*, 2012). Historically, the first case of attribution of a biological change to climate change was on poleward shifts of European butterflies in Europe in response to regional warming (Parmesan *et al.*, 1999).

#### 1.1.2. Trends of spring and autumn phenological events and their drivers

The most frequently observed biological impact of climate change over the past decades is major changes in spring and autumn phenology—the timing of recurring life history events (Parmesan & Yohe, 2003; Cleland *et al.*, 2007; Lieth *et al.*, 1974; Woolway *et al.*, 2021; Menzel *et al.*, 2006). Understanding the consequences of these shifts on ecosystems requires understanding how much the growing season has changed (Duputié *et al.*, 2015). Spring phenological events (e.g. budburst and leafout) have been advancing from 0.5 (Wolfe *et al.*, 2005) to 4.2 days/decade (Chmielewski & Rötzer, 2001; Fu *et al.*, 2014) and are mainly driven by temperature (Chuine, 2010; Cleland *et al.*, 2007; Peñuelas & Filella, 2001). In contrast, autumn phenology (e.g. budset and leaf senescence) is delayed, though to a much lesser extent than spring (Gallinat *et al.*, 2015; Jeong & Medvigy, 2014). The drivers regulating autumn phenology are far less understood than those of spring for many reasons. First, autumn phenology has attracted much less attention compared to spring (Piao *et al.*, 2019). Second, the data is often much noisier, since meteorological conditions in the fall can drastically influence phenological phenomena. To illustrate this, trees going through leaf senescence are subjected to a gradual leaf abscission that follows nutrient reabsorption, and the leaves within the same individual might be at different senescence stages, but a strong wind spell may trigger leaf drop for all leaves, thus affecting the temporal resolution of these data (Wu *et al.*, 2024). However, there is a general belief that autumn phenophases are driven by shortening photoperiod and colder temperatures (Cooke *et al.*, 2012; Flynn & Wolkovich, 2018; Körner & Basler, 2010; Delpierre *et al.*, 2016). Different hypotheses are proposed to explain delayed autumn phenophases. First, warmer autumn temperatures may extend the activity of photosynthetic enzymes, which could be maintained at a higher level. Thus, the degradation rate of chlorophyll would decrease and the timing of senescence would be delayed (Yan *et al.*, 2021). Second, summer droughts could make trees pause their activity schedule and delay senescence to increase carbon assimilation (Dox *et al.*, 2022). Third, there could be an antagonistic effect of warming and brightening—caused by reductions in atmospheric pollution and cloud cover (Sanchez-Lorenzo *et al.*, 2015)—on leaf senescence (Wu

*et al.*, 2021). Brightening accelerates the leaf senescence processes and reduces the temperature sensitivity during that period, counteracting the expected warming-induced delays in leaf senescence (Wu *et al.*, 2021). The photo-protection and sink limitation hypotheses provide plausible explanations for the negative effect of radiation on leaf senescence and the declining effect of temperature sensitivity of leaf senescence in response to brightening (Wu *et al.*, 2021; Zani *et al.*, 2020).

### 1.1.3. Misleading evidence of declining sensitivity to warming

While we have convincing proof that spring events advanced in the past decades, there is evidence that this might decelerate because of declining sensitivity to warming (Fu *et al.*, 2015; Meng *et al.*, 2020). The proposed mechanism is through the action of warmer winters on tree dormancy. In the fall, trees in boreal and temperate forests slowly enter dormancy, which is initiated with budset. During this phase, cold hardiness increases, which prepares the tree for the upcoming cold temperatures and prevents tissue damage. Then, the tree enters dormancy, during which a certain duration of cold temperatures (chilling)—with some interaction with photoperiod for some species—is necessary for the trees to be ready to accumulate forcing (Vitasse *et al.*, 2014). In the late winter and early spring, they go through two forms of deacclimation before budburst (Vitasse *et al.*, 2014). When deacclimation is reached, a certain amount of heat (forcing) is required to initiate budburst (Fu *et al.*, 2015). The argument of declining sensitivity appears here: heat requirement is met sooner in warm springs, but it's also negatively correlated with chilling (Fu *et al.*, 2015, 2013; Laube *et al.*, 2014). However, it is this interaction between chilling and forcing requirements that determines the timing of leaf unfolding. In other words, a decrease in chilling accumulation is proposed to explain the observed weaker spring temperature sensitivities, where spring forcing loses its relative importance (Fu *et al.*, 2015; Meng *et al.*, 2020; Wolkovich *et al.*, 2021). However, a meta-analysis compiling 72 studies of 203 species suggests that declining sensitivities observed in Europe may be a statistical artifact of how these responses are calculated, thus casting doubt on this proposed trend (Ettinger *et al.*, 2020). This statistical artifact is explained to be caused by using linear models for calculating non-linear processes (Wolkovich *et al.*, 2021).

### 1.1.4. Mechanisms that could limit growth despite having a longer growing season

Plants' seasonal activity has internal and external controls, both determined by environmental conditions (Wolkovich *et al.*, 2025). Internal controls operate via autonomous clocks, activating genes and releasing hormones which often rely on chilling and/or photoperiod (Körner *et al.*, 2023). The external controls, often referred to as forcing, act directly on the developmental rate, meristem activity, tissue differentiation and metabolism (Körner *et al.*, 2023). Understanding how these controls operate is critical to our comprehension of plants' capacity to adjust their activity schedule in response to changing conditions (Chuine & Régnière, 2017). In light of this, I hypothesize two possible drivers that could explain why a longer growing season might not lead to increased growth: external (environmental) (Kolár *et al.*, 2016) and internal (Zohner *et al.*, 2023) constraints to growth (Wolkovich *et al.*, 2025).

#### 1.1.4.1. External constraints

The complex nature of climate change makes predicting the external limits to growth hard to quantify at the individual level, as these drivers affect communities as a whole. However, drought, spring frost and heat waves are commonly mentioned as the main extreme events that could limit tree growth under climate change (Tyree & Zimmermann, 2002; Choat *et al.*, 2018; Li *et al.*, 2023; Trenberth *et al.*, 2014; Change, 2014; Chiang *et al.*, 2021; Polgar & Primack, 2011; Reinmann *et al.*, 2023). Their definitions, respective mechanisms, global trend of occurrence, consequences and differences among species are described in Tables 3.1. (Spring frosts), 3.2. (Droughts) and 3.3. (Heat waves).

#### 1.1.4.2. Internal constraints

As for the internal constraints to growth, recent hypotheses propose that broadleaf deciduous tree species may be sink-saturated, such that longer growing seasons with more carbon fixation do not necessarily increase growth (Dow *et al.*, 2022). This one pathway is directly linked to the internal controls of plant growth, which are under strong genetic control. In addition to height and radial growth varying across species, these growth responses also vary by site (Silvestro *et al.*, 2023a; Aitken & Bemmels, 2016). Populations from higher altitude or latitude grow less under the same conditions than individuals from lower altitude or latitude (Way & Montgomery, 2015). This is further supported by phenological studies showing that growth cessation arrives

earlier from populations of higher latitude, demonstrating local adaptation to potentially avoid fall frost, before nutrient uptake has finished (Aitken & Bemmels, 2016; Vitasse, 2013; Way & Montgomery, 2015). These trees rely on photoperiod cues for setting buds (stopping height growth) (Way & Montgomery, 2015; Soolanayakanahally *et al.*, 2013).

Growth seasonality has huge consequences on annual growth, but this seasonality is also closely tied to the environmental conditions experienced by an individual. For instance, warming spring temperatures seem to positively affect growth, but recent evidence suggests a shift in this net positive effect near the summer solstice. Indeed Zohner *et al.* (2023) found that warmer temperatures before the solstice advanced the onset of senescence, but it slowed the progression of leaf colouring when temperatures post-solstice were warmer. This further reinforces the importance of understanding how the timing of warmer conditions affects tree species.

#### **1.1.5. How these shifts translate into effects on trees/forests is not clear - Pros and cons of early/late start/end of season**

Shifts in spring and autumn phenology support a long-lasting and intuitive assumption that earlier spring and delayed autumn events lead to longer seasons—and thus increased growth (Keenan *et al.*, 2014; Stridbeck *et al.*, 2022). However, research from the past three years has cast doubt on this hypothesis (Dow *et al.*, 2022; Green & Keenan, 2022; Silvestro *et al.*, 2023b). For instance, (Dow *et al.*, 2022) showed that despite an earlier growth onset, neither growth rate nor overall annual increment was increased by longer seasons. This could substantially affect carbon-cycle model projections and thus feedbacks to future climate (Richardson *et al.*, 2013; Swidrak *et al.*, 2013). Starting to grow earlier and stopping later both have pros and cons (Figure 1).

Understanding these findings requires answering why trees do not grow more despite longer growing seasons. While carbon allocation to above-ground biomass is one of the largest carbon sinks, how this carbon is allocated into wood is poorly understood. Indeed, the linear relationship assumption between wood growth and carbon assimilation is not well supported mechanistically and represents an important limitation of vegetation models (Cabon *et al.*, 2022). Net primary production represents the difference between photosynthesis and plant respiration, but this commonly used metric completely omits the representation of growth processes. This is perhaps because of a long-lasting paradigm of source-limited photosynthesis. This has now been debunked with carbon fertilization experiments, supporting evidence that direct environmental constraints on growth may be stronger than those on photosynthesis (Friend *et al.*, 2019; Parent *et al.*, 2010). This opens up a new—poorly tested—temperature sensitivity relationship between cambial activity and photosynthesis. Growing evidence suggests that cambial activity may be more sensitive than photosynthesis to a range of environmental conditions, such as water and nutrient availability, and temperature (Cabon *et al.*, 2022, 2020; Muller *et al.*, 2011; Peters *et al.*, 2021). This demonstrates that carbon projection models that heavily rely on vegetation alone may mislead the amount of carbon sequestered in our forests.

#### **1.1.6. Growing season shifts and consequences on forest ecosystems and services**

Spring and fall phenological events are shifting with debatable consequences on tree growth. The sensitivity of cambial activity to water, temperature and nutrients has the potential to have far-reaching consequences given the hard-to-predict nature of future climate change, where any of these variables could vary from low to high amplitude (Almagro *et al.*, 2025; Cabon *et al.*, 2022). This expected asymmetry of future environmental changes makes understanding the internal and external limits to growth critical. Especially, the capacity to tease apart different biomes—as for example boreal vs tropical forests are expected to react differently (REF)—is critical and empirical data coming from experiments, but also from observations are paramount to predict the changes of forest carbon offset from human GHG emissions (Wolkovich *et al.*, 2025).

## **1.2 Nature of the problem, and how to address it**

### **1.2.1. Past phenological trends can help (or not) predict future phenological changes**

Observed phenological trends in the last decades cannot be used directly to extrapolate future phenological changes because: (1) the mechanisms guiding them aren't clear, and (2) phenological responses of trees to warming are very likely to be non-linear (Ettinger *et al.*, 2020; Fu *et al.*, 2013). Indeed, accurate pre-

dictions require an in-depth, accurate mechanistic understanding of phenophases and their sensitivities to environmental drivers, especially to temperature and photoperiod (Fu *et al.*, 2013). Therefore, the very foundation of the assumption that longer seasons increase growth may shift with future climate change. The well-observed advance in spring phenology may decelerate, and delayed fall phenology may shift towards earlier leaf senescence (through summer drought-induced growth cessation).

### 1.2.2. The assumption that longer seasons lead to increased growth is called into question

Recent work shows an absence of increased growth despite apparent better environmental conditions, thus casting doubt on a simple and intuitive positive relationship between growing season length and growth. This paradox further emphasizes the need to better understand the drivers regulating growth across biomes, but also how these drivers vary across the species within these biomes. Phenology varies greatly across species (e.g. closely related species tend to budburst at similar times under similar conditions) (Wolkovich *et al.*, 2014), but so does the relationship between growth and season length, which may explain the wide variation of this relationship within communities (Buckley & Kingsolver, 2012). This highlights another weakness of current carbon sequestration models that pool species together, which likely overpass important nuances in the growth responses that could be explained by species differences. This may mislead the future of forest carbon source-sink ratios (Green & Keenan, 2022; Cabon *et al.*, 2022). To better understand how different species respond to warming, different strategies can be used.

First, experiments are paramount to robustly tease apart the external vs internal drivers (e.g. warmer springs from severe drought later in the season—a common co-occurring reality in natural environments) (Morin *et al.*, 2010; Primack *et al.*, 2015). However, experiments are most often performed on juvenile trees, which are critical for their role in forest regeneration projections, but their responses can hardly be translated to mature trees, which hold the overwhelming carbon biomass proportion of forests (Augsburger & Bartlett, 2003; Silvestro *et al.*, 2023b; Vitasse, 2013).

SLeaf phenology through ground-based observations can provide valuable insights into the growth onset and end of trees that are not suitable for experimental trials, since cambial and leaf phenology are closely linked to the other. It is to say that knowing when leaves elongate and colour can guide when trees start and stop growing—fundamental metrics to determine the growing season length. Ground observations have the advantage of providing accurate data on phenological events for specific sites and species. Recently, the widespread use of smartphones has opened a whole new world of possible phenological monitoring through citizen scientists’ records of data over much larger areas and for a wider range of species (Dickinson *et al.*, 2012; Hufkens *et al.*, 2019; Piao *et al.*, 2019). While there are drawbacks to these observations (e.g. non-standard protocols, highly uneven spatiotemporal distribution of these observations), these methods have a huge potential to diversify the phenology data.

### 1.2.3. Goals of my thesis

Using citizen science data, a common garden study and a large-scale experiment, I aim to better understand how different tree species, at different lifespan stages, vary in their growth responses to different season length.

## 1.3 Complexity of measuring growth and defining growing season length

### 1.3.1. What is a growing season?

To understand how trees respond to growing season conditions, it is important to clarify what a growing season and growth are.

First, a problem that arises when one tries to quantify how shifting growing seasons affect growth comes from the definition of the growing season itself. Recently, Körner *et al.* (2023) proposed four definitions addressing this issue: (1) true growing season, based on measurable growth; (2) phenological season, based on visible phenological markers; (3) the productive season, based on primary production and (4) meteorological season, based on environmental conditions.

Here, I will focus on how definition (2), incorporating (4) affects definition (1) as the data collected for this thesis can’t address (3). I will use definition (2) to infer a “window of opportunity” during which

meteorological conditions will be used to calculate growing degree days (GDD). I am using the meteorological season within a constrained window, instead of simply using it irrespective of phenology because of the illusion that an absolute increase in GDD over the last decades—irrespective of the timing of phenophases—also increases growth. Springs are warmer, falls are also sometimes warmer, and summers are warmer, which together increase the number of GDD, which may appear to be a reliable proxy for better environmental conditions. However, models that accumulate GDD before and/or after trees grow could mislead about how this variable drives growth.

Models using degree-days are still widely used even if they have been employed for decades in agriculture (e.g. (Gilmore & Rogers, 1958; McMaster, 1997)). These rely on developmental patterns that are based on temperature dependence to estimate a particular ecological process, in my case, tree-ring width. These models describe a particular response variable as a composite of time and temperature as opposed to time alone. This is a partimonious method that requires three variables: daily minimum and maximum temperatures and the base temperature at which the process of interest cannot occur (cambial activation in this case) (McMaster, 1997; Moore & Remais, 2014). However, this simplicity comes with a drawback of oversimplifying potentially complex developmental processes in response to varying environmental conditions within a season (Bonhomme, 2000).

#### *What is growth?*

Wood formation (xylogenesis) is the major biological process by which carbon is allocated and long-term stored in woody plants. Radial growth is determined by the production of xylem and phloem cells that begins with cambial activation and cell production by cambial initials, followed by cell differentiation through the following events: (1) Cell enlargement (2) Secondary-wall formation and lignification and (3) programmed cell death (Etzold *et al.*, 2022; Silvestro *et al.*, 2025). The rate and duration of these phases lead to irreversible radial growth increments usually represented through tree rings. In these, secondary xylem cells account disproportionately to the number of cells produced because they divide more than phloem cells (Plomion *et al.*, 2001; Rathgeber *et al.*, 2016).

#### **1.3.2. Traditional diameter measurements miss the resolution of annual growth increment**

Foresters have measured tree diameter and height for decades to infer allometries that could give them a good estimate of how much wood they could harvest in a forest (e.g. (Meyer, 1940; Saunders & Wagner, 2008)). The widely used method in forestry is to measure diameter at breast height at punctual time intervals (Yuancai & Parresol, 2001). However, these measurements don't provide short-term indicators of growth—especially if taken at multiple-year intervals. Therefore, extreme events affecting growth are likely to be missed. In addition, many forest inventories only report diameter measurements and exclude height, because of logistical constraints ((Saunders & Wagner, 2008)), which reduces biomass estimation quality. This growth data lacks the temporal resolution necessary to properly infer a relationship between growth and environmental conditions.

Alongside the diameter-height allometric relationship, dendrochronology and its related fields also have long-standing traditions in forestry and silviculture. Indeed, tree-ring research was developed in the early 1900s to perform archeological dating and climate reconstruction (Douglass, 1928; Pearl *et al.*, 2020). Since then, different dendrochronology methods have been developed for different purposes, such as the calibration of the terrestrial radiocarbon curve (Helama, 2023; Reimer *et al.*, 2020) and modelling Earth's past climate and ecological change (Pearl *et al.*, 2020). More recently, dendroecology—applications of dendrochronological techniques to problems in ecology (Fritts & Swetnam, 1989)—emerged to answer ecological problems as well as to hindcast (e.g. (Bergeron *et al.*, 2004)) and forecast ecological processes both at the regional (Gazol *et al.*, 2018) and global scale (Manzanedo & Pederson, 2019; Büntgen *et al.*, 2018). Now, these methods can be used to understand more precise growth patterns and their relationship with different environmental factors.

#### **1.3.3. Growth increment needs to incorporate wood density in order to evaluate how much structural carbohydrates were stored within a single year.**

While tree-ring width is a reliable proxy to how much trees grow in each year, at relatively low cost and time, the inclusion of wood density in the analysis may provide additional information. For example, Čufar

*et al.* (2008) showed that intra- and inter-annual density variations in beech provided more information and at a higher temporal resolution than tree-ring widths alone.

In addition to densitometry, an increasing number of studies are going beyond traditional ring width by performing analysis using wood anatomy data. For instance, low-cost techniques (e.g. "blue intensity" proxy for latewood density (Babst *et al.*, 2016; Campbell *et al.*, 2007) and high resolution imaging (Griffin *et al.*, 2021; Levanič, 2007; Von Arx & Carrer, 2014)) give rise to a whole new world of possibilities regarding the microscopic components of wood anatomical features. Micro-anatomical analyses within tree rings could be applied to a wide range of applications (Pearl *et al.*, 2020). The character of annual rings, cell structure, timing of growth and markers for trauma can assist in answering a variety of ecological and physiological questions previously unanswered with ring width or density alone, such as how growth is affected by growing season length under anthropogenic forcing.

#### 1.3.4. Asynchrony between primary and secondary growth (internal growth control?)

I argue that internal growth control in trees may shape their growth responses to growing season length, and these can be split into two categories: primary and secondary growth synchronicity and growth determinancy.

Primary (shoot elongation) and secondary (xylem and phloem formation) growth both contribute to how much carbon is stored in plant tissue, but how much they differ in their responses to environmental drivers is poorly understood. After a dormancy period, trees will start growing, both vertically (primary) and horizontally (secondary), but there is high variation among species as to when each growth starts, for how long it lasts and when it stops. For instance, ring-porous species initiate primary growth early in the season, sometimes even before budburst (e.g. oaks) (Stridbeck *et al.*, 2022), whereas xylogenesis in diffuse-porous species is usually more synchronized with budburst. These two examples highlight how more complicated it might be to infer general conclusions as to how growing season shifts may also shift growth, where some species may extend their primary growth, but restrict their secondary growth and vice versa.

In addition to differences in primary and secondary growth synchronicity, the role of internal growth control—often overlooked—may misshape our understanding of growth responses to growing season length (Baumgarten *et al.*, 2025). In perennial plants, two dichotomous growth strategies are commonly mentioned: determinate and indeterminate growth, though it appears that species exist along a gradient of these (Baumgarten *et al.*, 2025). Indeterminate growth is usually associated with short-lived and fast growth species, where these life-history strategies may give them a competing advantage as tissue growth can be produced quickly in response to changing environmental conditions, but this comes with a higher risk of late spring and early fall frost as well as late droughts (Baumgarten *et al.*, 2025; Brien *et al.*, 2020). At the opposite side of the spectrum, determinate species are usually long-lived and slow-growing and are mainly constrained by conditions during bud formation; this may increase bud survival at the detriment of opportunistic growth in face of better-than-expected conditions (Baumgarten *et al.*, 2025).

Thus, primary versus secondary growth allocation, coupled with growth determinancy, are variables often overlooked, but which greatly vary across species and that are likely to affect how trees respond to shifting season length.

#### 1.3.5. Getting growth temporal resolution is labor-intensive and expensive (e.g. dendrometer costs)

To know when trees start and stop growing within a single growing season (data not extractable through tree rings), there are two methods, one being labour-intensive (1) and the other being expensive (2).

1. Trephor is a tool that allows recurrent sampling of mature trees where a 2mm microcore is extracted from the tree at different moments during the growing season. By sampling multiple times a single tree, growth temporality can be inferred by counting the ring cell increment between each sample. While this non-destructive tool can be extremely valuable, getting a large sample size can hardly be feasible (Rossi *et al.*, 2006).

2. Dendrometers allow for monitoring stem radius variation, measuring irreversible secondary growth, but also stem water fluctuations and thermal expansion, often leading to biased estimates of growth increment temporality (Camarero *et al.*, 2010).

308 **1.4 Objectives**

309 My objectives for Fuelinex are to assess tree species' potential to prolong or stretch their activity schedule by  
310 artificially manipulating growing season length and analyze how this translates (or not) into growth, during  
311 the current year (2024) and in the following year (2025). I will also conduct a secondary experiment to  
312 examine whether trees can absorb nutrients late in the season and if that translates into growth during the  
313 following season. For the CookieSpotters projects, I will investigate how the timing of phenological events  
314 affects growth across years for juvenile and mature trees, using observational phenology data and tree ring.  
315 Together, my two chapters will allow me to address the paradox of the absence of increased growth  
316 despite longer growing seasons.

317 **1.5 Research questions**

318 *necessary after the objectives? It feels redundant...*  
319 Fuelinex: How do extended growing seasons affect tree growth across different species, both immediately  
320 (in the same year as the extended season) and in subsequent years? CookieSpotters: How does phenology  
321 regulate tree growth in urban ecosystems?  
322  
323

324 **3.1. Spring frosts**

<b>Definition:</b>	Late spring frosts are below-freezing temperatures in late spring (Zohner <i>et al.</i> , 2020)
<b>Mechanisms</b>	Early warm spells → early leaf out → hard frost (<-2Celsius) → tissue death = loss of photosynthetic capacity (Polgar & Primack, 2011); Response: second cohort of leaves are more efficient and mitigate carbon sequestration loss (Reinmann <i>et al.</i> , 2023)
<b>Global trend of occurrence</b>	Most vulnerable regions are the ones with no past risk of occurrence (); ↑ in Europe and East Asia, but ↓ North America; Global trend is controversial (Reinmann <i>et al.</i> , 2023)
<b>Consequences (Individual and Ecosystem level consequences)</b>	Loss of vegetative tissue = ↓ photosynthesis = ↓ and remobilization of NSC to repair damaged tissues = ↓ secondary growth (Meyer <i>et al.</i> , 2024); Loss of reproductive tissue (higher flower mortality) (Sgubin <i>et al.</i> , 2018); Costs for orchards and stuff (Reinmann <i>et al.</i> , 2023)
<b>Differences across species/provenance</b>	

325

### 3.2. Drought

326

<b>Definition:</b>	"Drought is a prolonged absence or marked deficiency of precipitation that results in water shortage or a period of abnormally dry weather sufficiently prolonged for the lack of precipitation to cause a serious hydrological imbalance (Trenberth <i>et al.</i> , 2014; Intergovernmental panel on climate change, 2007).
<b>Mechanisms</b>	<ul style="list-style-type: none"> <li>— Hot temperature + low precipitation (aka global-change-type drought (Tyree &amp; Zimmermann, 2002)) = <math>\uparrow</math> evapotranspiration <math>\rightarrow</math> less water in soil <math>\rightarrow</math> cavitation <math>\rightarrow</math> embolism <math>\rightarrow</math> hydraulic failure (Tyree &amp; Zimmermann, 2002) = tissue death (Choat <i>et al.</i>, 2018);</li> <li>— Earlier spring phenology = longer GS <math>\rightarrow</math> increases vegetative growth <math>\rightarrow</math> increases evapotranspiration <math>\rightarrow</math> increases drawdown of soil moisture = progressive water stress (Li <i>et al.</i>, 2023)</li> <li>— Long-term vs short-term stomatal responses and consequences on tissue death (Choat <i>et al.</i>, 2018);</li> <li>— Recovery and its determinants (Choat <i>et al.</i>, 2018; Li <i>et al.</i>, 2023)</li> </ul>
<b>Global trend of occurrence</b>	<ul style="list-style-type: none"> <li>— <math>\uparrow</math> precipitation anomalies since 1990 (Trenberth <i>et al.</i>, 2014);</li> <li>— Models often exclude PDO/ENSO which limit the capacity to attribute increasing droughts to CC (Trenberth <i>et al.</i>, 2014);</li> <li>— Weak evidence of detection and attribution of changes in meteorological drought since the mid-20th century (Change, 2014);</li> <li>— Using a spacial, model-based perspective, anthropogenic forcing increased the frequency, duration and intensity of SPI-based droughts for North America (Hidalgo <i>et al.</i>, 2009), Europe and the Mediterranean (Spinoni <i>et al.</i>, 2018), (Kurnik <i>et al.</i>, 2011) and E Asia (Chiang <i>et al.</i>, 2021; Marvel <i>et al.</i>, 2019; Spinoni <i>et al.</i>, 2014)</li> </ul>
<b>Consequences (Individual and Ecosystem level consequences)</b>	<ul style="list-style-type: none"> <li>— Recurring droughts may limit trees' ability to recover from other types of stress.</li> <li>— Tree mortality (e.g. Texas and California extreme droughts are estimated to have killed 300 and 102 million trees (Li <i>et al.</i>, 2023))</li> </ul>
<b>Differences across species/provenance</b>	

327

328

### 3.3. Heat waves

<b>Definition:</b>	Heat wave is a period of excessively hot weather (5 or more consecutive days of prolonged heat in which the daily maximum temperature is higher than the average maximum temperature by 5 °C ), which may be accompanied by high humidity (Marx <i>et al.</i> , 2021).
<b>Mechanisms</b>	<p>↑ atmospheric CO<sub>2</sub> = ↑ temperature → ↑ heat waves... More specifically: A mechanism for the increase occurrence of heat waves is a weakening of the polar jet stream (important weather factor for middle latitude regions of North America, Europe and Asia) caused by global warming which increases the occurrence of stationary weather, resulting in heavy rain falls or heat waves (Marx <i>et al.</i>, 2021). Extreme heat → growth either through (1) Directly via cell processes disruption or (2) indirectly via effects of rising leaf-to-air vapour deficit (VPD) (Gagne <i>et al.</i>, 2020).</p> <p>Increased temperature leads to reduced photosynthesis which can be attributed to: 1. Damage to photosynthetic machinery 2. Inactivation of RUBISCO 3. Reduction to RuBP regeneration 4. Membrane stability 5. Increased mitochondrial respiration and photorespiration (Hauck <i>et al.</i>, 2025)</p>
<b>Global trend of occurrence</b>	Heat waves have increased (Gagne <i>et al.</i> , 2020; Meehl & Tebaldi, 2004; Teskey <i>et al.</i> , 2015) and are expected to increase under future climate change (Dosio <i>et al.</i> , 2018; Change, 2014; Teskey <i>et al.</i> , 2015). Summertime extreme temperatures associated with prolonged heat waves, lasting for several weeks, now impact approximately 10% of land surfaces, up from only 1% in the 1960s. (Teskey <i>et al.</i> , 2015). The more intense and more frequently occurring heat waves cannot be explained solely by natural climate variations and without human-made climate change (Marx <i>et al.</i> , 2021).
<b>Consequences (Individual and Ecosystem level consequences)</b>	- Reduced photosynthesis - Increased mortality - Photosynthetic tissue loss (Gagne <i>et al.</i> , 2020).
<b>Differences across species/provenance</b>	Some species have thermal photosynthetic/respiratory acclimation while others don't. Growth and survival will change depending on species to thermally acclimate to both photosynthesis and respiration - This is explained by growth strategies of gymnosperms vs angiosperms (which are usually better)

## References

- Aitken, S.N. & Bemmels, J.B. (2016). Time to get moving: assisted gene flow of forest trees. *Evolutionary Applications*, 9, 271–290.
- Almagro, D., Martin-Benito, D., Rossi, S., Conde, M., Fernández-de-Uña, L. & Gea-Izquierdo, G. (2025). Long-Term Cambial Phenology Reveals Diverging Growth Responses of Two Tree Species in a Mixed Forest Under Climate Change. *Global Change Biology*, 31, e70503.
- Augspurger, C.K. & Bartlett, E.A. (2003). Differences in leaf phenology between juvenile and adult trees in a temperate deciduous forest. *Tree Physiology*, 23, 517–525.
- Babst, F., Wright, W.E., Szejner, P., Wells, L., Belmecheri, S. & Monson, R.K. (2016). Blue intensity parameters derived from Ponderosa pine tree rings characterize intra-annual density fluctuations and reveal seasonally divergent water limitations. *Trees*, 30, 1403–1415.
- Baumgarten, F., Aitken, S., Vitasse, Y., Guy, R.D. & Wolkovich, E. (2025). Invest now, get paid later? Limits and opportunities of woody plants to time growth in a future climate. *In review*.
- Bergeron, Y., Gauthier, S., Flannigan, M. & Kafka, V. (2004). Fire regimes at the transition between mixedwood and coniferous boreal forest in northwestern Quebec. *Ecology*, 85, 1916–1932.
- Bonhomme, R. (2000). Bases and limits to using ‘degree.day’ units. *European Journal of Agronomy*, 13, 1–10.




Figure 1: Pros and cons of early start and late end of growing season.

- 347 Brien, R.J.W., Caldwell, L., Duchesne, L., Voelker, S., Barichivich, J., Baliva, M., Ceccantini, G., Di Fil-  
348 ippo, A., Helama, S., Locosselli, G.M., Lopez, L., Piovesan, G., Schöngart, J., Villalba, R. & Gloor, E.  
349 (2020). Forest carbon sink neutralized by pervasive growth-lifespan trade-offs. *Nature Communications*,  
350 11, 4241.
- 351 Buckley, L.B. & Kingsolver, J.G. (2012). Functional and Phylogenetic Approaches to Forecasting Species'  
352 Responses to Climate Change. *Annual Review of Ecology, Evolution, and Systematics*, 43, 205–226.
- 353 Buntgen, U., Wacker, L., Galván, J.D., Arnold, S., Arseneault, D., Baillie, M., Beer, J., Bernabei, M.,  
354 Bleicher, N., Boswijk, G., Bräuning, A., Carrer, M., Ljungqvist, F.C., Cherubini, P., Christl, M., Christie,  
355 D.A., Clark, P.W., Cook, E.R., Esper, J., Fowler, A.M., Gennaretti, F., GieBinger, J., Grissino-Mayer,  
356 H., Grudd, H., Gunnarson, B.E., Hantemirov, R., Herzig, F., Hessler, A., Heussner, K.U., Jull, A.J.T.,  
357 Kukarskih, V., Kirdyanov, A., Krusic, P.J., Kyncl, T., Lara, A., LeQuesne, C., Linderholm, H.W., Loader,  
358 N.J., Luckman, B., Miyake, F., Myglan, V.S., Nicolussi, K., Oppenheimer, C., Palmer, J., Panyushkina, I.,  
359 Pederson, N., Rybníček, M., Schweingruber, F.H., Seim, A., Sigl, M., Churakova, O., Speer, J.H., Synal,  
360 H.A., Tegel, W., Treydte, K., Villalba, R., Wiles, G., Wilson, R., Winship, L.J., Wunder, J., Yang, B. &  
361 Young, G.H.F. (2018). Tree rings reveal globally coherent signature of cosmogenic radiocarbon events in  
362 774 and 993 CE. *Nature Communications*, 9, 3605.
- 363 Cabon, A., Fernández-de-Uña, L., Gea-Izquierdo, G., Meinzer, F.C., Woodruff, D.R., Martínez-Vilalta, J. &  
364 De Cáceres, M. (2020). Water potential control of turgor-driven tracheid enlargement in Scots pine at its  
365 xeric distribution edge. *New Phytologist*, 225, 209–221.
- 366 Cabon, A., Kannenberg, S.A., Arain, A., Babst, F., Baldocchi, D., Belmecheri, S., Delpierre, N., Guerrieri,  
367 R., Maxwell, J.T., McKenzie, S., Meinzer, F.C., Moore, D.J.P., Pappas, C., Rocha, A.V., Szejner, P.,  
368 Ueyama, M., Ulrich, D., Vincke, C., Voelker, S.L., Wei, J., Woodruff, D. & Anderegg, W.R.L. (2022).  
369 Cross-biome synthesis of source versus sink limits to tree growth. *Science*, 376, 758–761.
- 370 Camarero, J.J., Olano, J.M. & Parras, A. (2010). Plastic bimodal xylogenesis in conifers from continental  
371 Mediterranean climates. *New Phytologist*, 185, 471–480.
- 372 Campbell, L.M., Hagerman, S. & Gray, N.J. (2014). Producing Targets for Conservation: Science and Politics  
373 at the Tenth Conference of the Parties to the Convention on Biological Diversity. *Global Environmental*  
374 *Politics*, 14, 41–63.
- 375 Campbell, R., McCarroll, D., Loader, N.J., Grudd, H., Robertson, I. & Jalkanen, R. (2007). Blue intensity  
376 in *Pinus sylvestris* tree-rings: developing a new palaeoclimate proxy. *The Holocene*, 17, 821–828.
- 377 Ceballos, G., Ehrlich, P.R. & Dirzo, R. (2017). Biological annihilation via the ongoing sixth mass extinction  
378 signaled by vertebrate population losses and declines. *Proceedings of the National Academy of Sciences*,  
379 114.
- 380 Change, I.P.O.C. (2014). Detection and Attribution of Climate Change: from Global to Regional. In:  
381 *Climate Change 2013 – The Physical Science Basis*. Cambridge University Press, pp. 867–952. 1st edn.
- 382 Chiang, F., Mazdiyasn, O. & AghaKouchak, A. (2021). Evidence of anthropogenic impacts on global drought  
383 frequency, duration, and intensity. *Nature Communications*, 12, 2754.
- 384 Chmielewski, F.M. & Rötzer, T. (2001). Response of tree phenology to climate change across Europe.  
385 *Agricultural and Forest Meteorology*, 108, 101–112.
- 386 Choat, B., Brodribb, T.J., Brodersen, C.R., Duursma, R.A., López, R. & Medlyn, B.E. (2018). Triggers of  
387 tree mortality under drought. *Nature*, 558, 531–539.
- 388 Chuine, I. (2010). Why does phenology drive species distribution? *Philosophical Transactions of the Royal*  
389 *Society B: Biological Sciences*, 365, 3149–3160.
- 390 Chuine, I. & Régnière, J. (2017). Process-Based Models of Phenology for Plants and Animals. *Annual*  
391 *Review of Ecology, Evolution, and Systematics*, 48, 159–182.

- Cleland, E., Chuine, I., Menzel, A., Mooney, H. & Schwartz, M. (2007). Shifting plant phenology in response to global change. *Trends in Ecology & Evolution*, 22, 357–365.
- Cooke, J.E.K., Eriksson, M.E. & Junttila, O. (2012). The dynamic nature of bud dormancy in trees: environmental control and molecular mechanisms. *Plant, Cell & Environment*, 35, 1707–1728.
- Delpierre, N., Vitasse, Y., Chuine, I., Guillemot, J., Bazot, S., Rutishauser, T. & Rathgeber, C.B.K. (2016). Temperate and boreal forest tree phenology: from organ-scale processes to terrestrial ecosystem models. *Annals of Forest Science*, 73, 5–25.
- Dickinson, J.L., Shirk, J., Bonter, D., Bonney, R., Crain, R.L., Martin, J., Phillips, T. & Purcell, K. (2012). The current state of citizen science as a tool for ecological research and public engagement. *Frontiers in Ecology and the Environment*, 10, 291–297.
- Dosio, A., Mentaschi, L., Fischer, E.M. & Wyser, K. (2018). Extreme heat waves under 1.5 °C and 2 °C global warming. *Environmental Research Letters*, 13, 054006.
- Douglass, A. (1928). Climate and trees. *Nature Magazine*, 12, 51–53.
- Dow, C., Kim, A.Y., D’Orangeville, L., Gonzalez-Akre, E.B., Helcoski, R., Herrmann, V., Harley, G.L., Maxwell, J.T., McGregor, I.R., McShea, W.J., McMahon, S.M., Pederson, N., Tepley, A.J. & Anderson-Teixeira, K.J. (2022). Warm springs alter timing but not total growth of temperate deciduous trees. *Nature*, 608, 552–557.
- Dox, I., Skrøppa, T., Decoster, M., Prislan, P., Gascó, A., Gričar, J., Lange, H. & Campioli, M. (2022). Severe drought can delay autumn senescence of silver birch in the current year but advance it in the next year. *Agricultural and Forest Meteorology*, 316, 108879.
- Driscoll, D.A., Bland, L.M., Bryan, B.A., Newsome, T.M., Nicholson, E., Ritchie, E.G. & Doherty, T.S. (2018). A biodiversity-crisis hierarchy to evaluate and refine conservation indicators. *Nature Ecology & Evolution*, 2, 775–781.
- Duputié, A., Rutschmann, A., Ronce, O. & Chuine, I. (2015). Phenological plasticity will not help all species adapt to climate change. *Global Change Biology*, 21, 3062–3073.
- Ettinger, A.K., Chamberlain, C.J., Morales-Castilla, I., Buonaiuto, D.M., Flynn, D.F.B., Savas, T., Samaha, J.A. & Wolkovich, E.M. (2020). Winter temperatures predominate in spring phenological responses to warming. *Nature Climate Change*, 10, 1137–1142.
- Etzold, S., Sterck, F., Bose, A.K., Braun, S., Buchmann, N., Eugster, W., Gessler, A., Kahmen, A., Peters, R.L., Vitasse, Y., Walthert, L., Ziemińska, K. & Zweifel, R. (2022). Number of growth days and not length of the growth period determines radial stem growth of temperate trees. *Ecology Letters*, 25, 427–439.
- Flynn, D.F.B. & Wolkovich, E.M. (2018). Temperature and photoperiod drive spring phenology across all species in a temperate forest community. *New Phytologist*, 219, 1353–1362.
- Friend, A.D., Eckes-Shephard, A.H., Fonti, P., Rademacher, T.T., Rathgeber, C.B.K., Richardson, A.D. & Turton, R.H. (2019). On the need to consider wood formation processes in global vegetation models and a suggested approach. *Annals of Forest Science*, 76, 49.
- Fritts, H. & Swetnam, T. (1989). Dendroecology: A Tool for Evaluating Variations in Past and Present Forest Environments. In: *Advances in Ecological Research*. Elsevier, vol. 19, pp. 111–188.
- Fu, Y.H., Campioli, M., Deckmyn, G. & Janssens, I.A. (2013). Sensitivity of leaf unfolding to experimental warming in three temperate tree species. *Agricultural and Forest Meteorology*, 181, 125–132.
- Fu, Y.H., Piao, S., Op De Beeck, M., Cong, N., Zhao, H., Zhang, Y., Menzel, A. & Janssens, I.A. (2014). Recent spring phenology shifts in western Central Europe based on multiscale observations. *Global Ecology and Biogeography*, 23, 1255–1263.

- Fu, Y.H., Zhao, H., Piao, S., Peaucelle, M., Peng, S., Zhou, G., Ciais, P., Huang, M., Menzel, A., Peñuelas, J., Song, Y., Vitasse, Y., Zeng, Z. & Janssens, I.A. (2015). Declining global warming effects on the phenology of spring leaf unfolding. *Nature*, 526, 104–107. Publisher: Nature Publishing Group.
- Gagne, M.A., Smith, D.D. & McCulloh, K.A. (2020). Limited physiological acclimation to recurrent heat-waves in two boreal tree species. *Tree Physiology*, 40, 1680–1696.
- Gallinat, A.S., Primack, R.B. & Wagner, D.L. (2015). Autumn, the neglected season in climate change research. *Trends in Ecology & Evolution*, 30, 169–176.
- Gazol, A., Camarero, J.J., Vicente-Serrano, S.M., Sánchez-Salguero, R., Gutiérrez, E., De Luis, M., Sangüesa-Barreda, G., Novak, K., Rozas, V., Tíscar, P.A., Linares, J.C., Martín-Hernández, N., Martínez Del Castillo, E., Ribas, M., García-González, I., Silla, F., Camisón, A., Génova, M., Olano, J.M., Longares, L.A., Hevia, A., Tomás-Burguera, M. & Galván, J.D. (2018). Forest resilience to drought varies across biomes. *Global Change Biology*, 24, 2143–2158.
- Gilmore, E.C. & Rogers, J.S. (1958). Heat Units as a Method of Measuring Maturity in Corn. *Agronomy Journal*, 50, 611–615.
- Green, J.K. & Keenan, T.F. (2022). The limits of forest carbon sequestration. *Science*, 376, 692–693.
- Griffin, D., Porter, S.T., Trummer, M.L., Carlson, K.E., Crawford, D.J., Schwalen, D. & McFadden, C.H. (2021). Gigapixel Macro Photography of Tree Rings. *Tree-Ring Research*, 77.
- Hauck, M., Schneider, T., Bahlinger, S., Fischbach, J., Oswald, G., Csapek, G. & Dulamsuren, C. (2025). Heat tolerance of temperate tree species from Central Europe. *Forest Ecology and Management*, 580, 122541.
- Helama, S. (2023). Distinguishing Type I and II errors in statistical tree-ring dating. *Quaternary Geochronology*, 78, 101470.
- Hidalgo, H.G., Das, T., Dettinger, M.D., Cayan, D.R., Pierce, D.W., Barnett, T.P., Bala, G., Mirin, A., Wood, A.W., Bonfils, C., Santer, B.D. & Nozawa, T. (2009). Detection and Attribution of Streamflow Timing Changes to Climate Change in the Western United States. *Journal of Climate*, 22, 3838–3855.
- Huey, R.B., Kearney, M.R., Krockenberger, A., Holtum, J.A.M., Jess, M. & Williams, S.E. (2012). Predicting organismal vulnerability to climate warming: roles of behaviour, physiology and adaptation. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 367, 1665–1679.
- Hufkens, K., Melaas, E.K., Mann, M.L., Foster, T., Ceballos, F., Robles, M. & Kramer, B. (2019). Monitoring crop phenology using a smartphone based near-surface remote sensing approach. *Agricultural and Forest Meteorology*, 265, 327–337.
- Intergovernmental panel on climate change (ed.) (2007). *Climate change 2007: the physical science basis*. Cambridge university press, Cambridge.
- Intergovernmental Panel On Climate Change (Ipcc) (2023). *Climate Change 2022 – Impacts, Adaptation and Vulnerability: Working Group II Contribution to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*. 1st edn. Cambridge University Press.
- Jeong, S. & Medvigy, D. (2014). Macroscale prediction of autumn leaf coloration throughout the continental United States. *Global Ecology and Biogeography*, 23, 1245–1254.
- Keenan, T.F., Gray, J., Friedl, M.A., Toomey, M., Bohrer, G., Hollinger, D.Y., Munger, J.W., O’Keefe, J., Schmid, H.P., Wing, I.S., Yang, B. & Richardson, A.D. (2014). Net carbon uptake has increased through warming-induced changes in temperate forest phenology. *Nature Climate Change*, 4, 598–604.
- Kolář, T., Giagli, K., Trnka, M., Bednářová, E., Vavřík, H. & Rybníček, M. (2016). Response of the leaf phenology and tree-ring width of European beech to climate variability. *Silva Fennica*, 50.

478 Kurnik, B., Barbosa, P. & Vogt, J. (2011). Testing two different precipitation datasets to compute the  
479 standardized precipitation index over the Horn of Africa. *International Journal of Remote Sensing*, 32,  
480 5947–5964.

481 Körner, C. & Basler, D. (2010). Phenology Under Global Warming. *Science*, 327, 1461–1462. Publisher:  
482 American Association for the Advancement of Science.

483 Körner, C., Möhl, P. & Hiltbrunner, E. (2023). Four ways to define the growing season. *Ecology Letters*, 26,  
484 1277–1292.

485 Laube, J., Sparks, T.H., Estrella, N., Höfler, J., Ankerst, D.P. & Menzel, A. (2014). Chilling outweighs  
486 photoperiod in preventing precocious spring development. *Global Change Biology*, 20, 170–182.

487 Laurance, W.F. (2007). Have we overstated the tropical biodiversity crisis? *Trends in Ecology & Evolution*,  
488 22, 65–70.

489 Levanič, T. (2007). Atrics – A New System for Image Acquisition in Dendrochronology. *Tree-Ring Research*,  
490 63, 117–122.

491 Li, Y., Zhang, W., Schwalm, C.R., Gentine, P., Smith, W.K., Ciais, P., Kimball, J.S., Gazol, A., Kannenberg,  
492 S.A., Chen, A., Piao, S., Liu, H., Chen, D. & Wu, X. (2023). Widespread spring phenology effects on  
493 drought recovery of Northern Hemisphere ecosystems. *Nature Climate Change*, 13, 182–188.

494 Lieth, H., Jacobs, J., Lange, O.L., Olson, J.S. & Wieser, W. (eds.) (1974). *Phenology and Seasonality*  
495 *Modeling*. vol. 8 of *Ecological Studies*. Springer Berlin Heidelberg, Berlin, Heidelberg.

496 Lynas, M., Houlton, B.Z. & Perry, S. (2021). Greater than 99% consensus on human caused climate change  
497 in the peer-reviewed scientific literature. *Environmental Research Letters*, 16, 114005.

498 Manzanedo, R.D. & Pederson, N. (2019). Towards a More Ecological Dendroecology. *Tree-Ring Research*,  
499 75, 152.

500 Marvel, K., Cook, B.I., Bonfils, C.J.W., Durack, P.J., Smerdon, J.E. & Williams, A.P. (2019). Twentieth-  
501 century hydroclimate changes consistent with human influence. *Nature*, 569, 59–65.

502 Marx, W., Haunschild, R. & Bornmann, L. (2021). Heat waves: a hot topic in climate change research.  
503 *Theoretical and Applied Climatology*, 146, 781–800.

504 McMaster, G. (1997). Growing degree-days: one equation, two interpretations. *Agricultural and Forest*  
505 *Meteorology*, 87, 291–300.

506 Meehl, G.A. & Tebaldi, C. (2004). More Intense, More Frequent, and Longer Lasting Heat Waves in the  
507 21st Century. *Science*, 305, 994–997.

508 Meng, L., Mao, J., Zhou, Y., Richardson, A.D., Lee, X., Thornton, P.E., Ricciuto, D.M., Li, X., Dai, Y., Shi,  
509 X. & Jia, G. (2020). Urban warming advances spring phenology but reduces the response of phenology  
510 to temperature in the conterminous United States. *Proceedings of the National Academy of Sciences*, 117,  
511 4228–4233.

512 Menzel, A., Sparks, T.H., Estrella, N., Koch, E., Aasa, A., Ahas, R., Alm-Kübler, K., Bissolli, P., Braslavská,  
513 O., Briede, A., Chmielewski, F.M., Crepinsek, Z., Curnel, Y., Defila, C., Donnelly, A., Filella, Y., Jatczak,  
514 K., Mestre, A., Peñuelas, J., Pirinen, P., Scheifinger, H., Striz, M., Susnik, A., Van Vliet, A.J.H., Wiel-  
515 golaski, F., Zach, S. & Zust, A. (2006). European phenological response to climate change matches the  
516 warming pattern. *Global Change Biology*, 12, 1969–1976.

517 Meyer, B.F., Buras, A., Gregor, K., Layritz, L.S., Principe, A., Kreyling, J., Rammig, A. & Zang, C.S.  
518 (2024). Frost matters: incorporating late-spring frost into a dynamic vegetation model regulates regional  
519 productivity dynamics in European beech forests. *Biogeosciences*, 21, 1355–1370.

520 Meyer, H.A. (1940). A Mathematical Expression for Height Curves. *Journal of Forestry*, 38, 415–420.

- Moore, J.L. & Remais, J.V. (2014). Developmental Models for Estimating Ecological Responses to Environmental Variability: Structural, Parametric, and Experimental Issues. *Acta Biotheoretica*, 62, 69–90.
- Morin, X., Roy, J., Sonié, L. & Chuine, I. (2010). Changes in leaf phenology of three European oak species in response to experimental climate change. *New Phytologist*, 186, 900–910.
- Muller, B., Pantin, F., Génard, M., Turc, O., Freixes, S., Piques, M. & Gibon, Y. (2011). Water deficits uncouple growth from photosynthesis, increase C content, and modify the relationships between C and growth in sink organs. *Journal of Experimental Botany*, 62, 1715–1729.
- Oreskes, N. (2004). The Scientific Consensus on Climate Change. *Science*, 306, 1686–1686.
- Parent, B., Turc, O., Gibon, Y., Stitt, M. & Tardieu, F. (2010). Modelling temperature-compensated physiological rates, based on the co-ordination of responses to temperature of developmental processes. *Journal of Experimental Botany*, 61, 2057–2069.
- Parmesan, C., Ryrholm, N., Stefanescu, C., Hill, J.K., Thomas, C.D., Descimon, H., Huntley, B., Kaila, L., Kullberg, J., Tammara, T., Tennent, W.J., Thomas, J.A. & Warren, M. (1999). Poleward shifts in geographical ranges of butterfly species associated with regional warming. *Nature*, 399, 579–583.
- Parmesan, C. & Yohe, G. (2003). A globally coherent fingerprint of climate change impacts across natural systems. *Nature*, 421, 37–42.
- Pearl, J.K., Keck, J.R., Tintor, W., Siekacz, L., Herrick, H.M., Meko, M.D. & Pearson, C.L. (2020). New frontiers in tree-ring research. *The Holocene*, 30, 923–941.
- Peters, R.L., Steppe, K., Cuny, H.E., De Pauw, D.J., Frank, D.C., Schaub, M., Rathgeber, C.B., Cabon, A. & Fonti, P. (2021). Turgor – a limiting factor for radial growth in mature conifers along an elevational gradient. *New Phytologist*, 229, 213–229.
- Peñuelas, J. & Filella, I. (2001). Responses to a Warming World. *Science*, 294, 793–795. Publisher: American Association for the Advancement of Science.
- Piao, S., Liu, Q., Chen, A., Janssens, I.A., Fu, Y., Dai, J., Liu, L., Lian, X., Shen, M. & Zhu, X. (2019). Plant phenology and global climate change: Current progresses and challenges. *Global Change Biology*, 25, 1922–1940.
- Plomion, C., Leprovost, G. & Stokes, A. (2001). Wood Formation in Trees. *Plant Physiology*, 127, 1513–1523.
- Polgar, C.A. & Primack, R.B. (2011). Leaf-out phenology of temperate woody plants: from trees to ecosystems. *New Phytologist*, 191, 926–941.
- Primack, R.B., Laube, J., Gallinat, A.S. & Menzel, A. (2015). From observations to experiments in phenology research: investigating climate change impacts on trees and shrubs using dormant twigs. *Annals of Botany*, 116, 889–897.
- Rathgeber, C.B.K., Cuny, H.E. & Fonti, P. (2016). Biological Basis of Tree-Ring Formation: A Crash Course. *Frontiers in Plant Science*, 7.
- Reimer, P.J., Austin, W.E.N., Bard, E., Bayliss, A., Blackwell, P.G., Bronk Ramsey, C., Butzin, M., Cheng, H., Edwards, R.L., Friedrich, M., Grootes, P.M., Guilderson, T.P., Hajdas, I., Heaton, T.J., Hogg, A.G., Hughen, K.A., Kromer, B., Manning, S.W., Muscheler, R., Palmer, J.G., Pearson, C., Van Der Plicht, J., Reimer, R.W., Richards, D.A., Scott, E.M., Southon, J.R., Turney, C.S.M., Wacker, L., Adolphi, F., Büntgen, U., Capano, M., Fahrni, S.M., Fogtmann-Schulz, A., Friedrich, R., Köhler, P., Kudsk, S., Miyake, F., Olsen, J., Reinig, F., Sakamoto, M., Sookdeo, A. & Talamo, S. (2020). The IntCal20 Northern Hemisphere Radiocarbon Age Calibration Curve (0–55 cal kBP). *Radiocarbon*, 62, 725–757.
- Reinmann, A.B., Bowers, J.T., Kaur, P. & Kohler, C. (2023). Compensatory responses of leaf physiology reduce effects of spring frost defoliation on temperate forest tree carbon uptake. *Frontiers in Forests and Global Change*, 6, 988233.

- Richardson, A.D., Keenan, T.F., Migliavacca, M., Ryu, Y., Sonnentag, O. & Toomey, M. (2013). Climate change, phenology, and phenological control of vegetation feedbacks to the climate system. *Agricultural and Forest Meteorology*, 169, 156–173.
- Rossi, S., Anfodillo, T. & Menardi, R. (2006). Trephor: A New Tool for Sampling Microcores from tree stems. *IAWA Journal*, 27, 89–97.
- Sanchez-Lorenzo, A., Wild, M., Brunetti, M., Guijarro, J.A., Hakuba, M.Z., Calbó, J., Mystakidis, S. & Bartok, B. (2015). Reassessment and update of long-term trends in downward surface shortwave radiation over Europe (1939–2012). *Journal of Geophysical Research: Atmospheres*, 120, 9555–9569.
- Saunders, M.R. & Wagner, R.G. (2008). Height-diameter models with random coefficients and site variables for tree species of Central Maine. *Annals of Forest Science*, 65, 203–203.
- Sgubin, G., Swingedouw, D., Dayon, G., García De Cortázar-Atauri, I., Ollat, N., Pagé, C. & Van Leeuwen, C. (2018). The risk of tardive frost damage in French vineyards in a changing climate. *Agricultural and Forest Meteorology*, 250–251, 226–242.
- Silvestro, R., Deslauriers, A., Prislan, P., Rademacher, T., Rezaie, N., Richardson, A.D., Vitasse, Y. & Rossi, S. (2025). From Roots to Leaves: Tree Growth Phenology in Forest Ecosystems. *Current Forestry Reports*, 11, 12.
- Silvestro, R., Mura, C., Alano Bonacini, D., De Lafontaine, G., Faubert, P., Mencuccini, M. & Rossi, S. (2023a). Local adaptation shapes functional traits and resource allocation in black spruce. *Scientific Reports*, 13, 21257.
- Silvestro, R., Zeng, Q., Buttò, V., Sylvain, J.D., Drolet, G., Mencuccini, M., Thiffault, N., Yuan, S. & Rossi, S. (2023b). A longer wood growing season does not lead to higher carbon sequestration. *Scientific Reports*, 13, 4059.
- Soolanayakanahally, R.Y., Guy, R.D., Silim, S.N. & Song, M. (2013). Timing of photoperiodic competency causes phenological mismatch in balsam poplar ( *Populus balsamifera* L.). *Plant, Cell & Environment*, 36, 116–127.
- Spinoni, J., Naumann, G., Carrao, H., Barbosa, P. & Vogt, J. (2014). World drought frequency, duration, and severity for 1951–2010. *International Journal of Climatology*, 34, 2792–2804.
- Spinoni, J., Vogt, J.V., Naumann, G., Barbosa, P. & Dosio, A. (2018). Will drought events become more frequent and severe in Europe? *International Journal of Climatology*, 38, 1718–1736.
- Stridbeck, P., Björklund, J., Fuentes, M., Gunnarson, B.E., Jönsson, A.M., Linderholm, H.W., Ljungqvist, F.C., Olsson, C., Rayner, D., Rocha, E., Zhang, P. & Seftigen, K. (2022). Partly decoupled tree-ring width and leaf phenology response to 20th century temperature change in Sweden. *Dendrochronologia*, 75, 125993.
- Swidrak, I., Schuster, R. & Oberhuber, W. (2013). Comparing growth phenology of co-occurring deciduous and evergreen conifers exposed to drought. *Flora: Morphology, Distribution, Functional Ecology of Plants*, 208, 609–617.
- Teskey, R., Wertin, T., Bauweraerts, I., Ameye, M., McGuire, M.A. & Steppe, K. (2015). Responses of tree species to heat waves and extreme heat events. *Plant, Cell & Environment*, 38, 1699–1712.
- Trenberth, K.E., Dai, A., Van Der Schrier, G., Jones, P.D., Barichivich, J., Briffa, K.R. & Sheffield, J. (2014). Global warming and changes in drought. *Nature Climate Change*, 4, 17–22.
- Tyree, M.T. & Zimmermann, M.H. (2002). *Xylem Structure and the Ascent of Sap*. Springer Series in Wood Science. Springer Berlin Heidelberg, Berlin, Heidelberg.
- Vitasse, Y. (2013). Ontogenic changes rather than difference in temperature cause understory trees to leaf out earlier. *New Phytologist*, 198, 149–155.

- Vitasse, Y., eLenz, A. & eKoerner, C. (2014). The interaction between freezing tolerance and phenology in temperate deciduous trees. *Frontiers in Plant Science*, 5. Publisher: Frontiers Media S.A.
- Von Arx, G. & Carrer, M. (2014). ROXAS – A new tool to build centuries-long tracheid-lumen chronologies in conifers. *Dendrochronologia*, 32, 290–293.
- Way, D.A. & Montgomery, R.A. (2015). Photoperiod constraints on tree phenology, performance and migration in a warming world. *Plant, Cell & Environment*, 38, 1725–1736. \_eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1111/pce.12431>.
- Wolfe, D.W., Schwartz, M.D., Lakso, A.N., Otsuki, Y., Pool, R.M. & Shaulis, N.J. (2005). Climate change and shifts in spring phenology of three horticultural woody perennials in northeastern USA. *International Journal of Biometeorology*, 49, 303–309.
- Wolkovich, E.M., Auerbach, J., Chamberlain, C.J., Buonaiuto, D.M., Ettinger, A.K., Morales-Castilla, I. & Gelman, A. (2021). A simple explanation for declining temperature sensitivity with warming. *Global Change Biology*, 27, 4947–4949.
- Wolkovich, E.M., Cook, B.I. & Davies, T.J. (2014). Progress towards an interdisciplinary science of plant phenology: building predictions across space, time and species diversity. *New Phytologist*, 201, 1156–1162. \_eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1111/nph.12599>.
- Wolkovich, E.M., Ettinger, A.K., Chin, A., Chamberlain, C.J., Baumgarten, F., Pradhan, K., Manzanedo, R.D. & Lambers, J.H.R. (2025). Why longer seasons with climate change may not increase tree growth. *In review*.
- Woolway, R.I., Sharma, S., Weyhenmeyer, G.A., Debolskiy, A., Golub, M., Mercado-Bettín, D., Perroud, M., Stepanenko, V., Tan, Z., Grant, L., Ladwig, R., Mesman, J., Moore, T.N., Shatwell, T., Vanderkelen, I., Austin, J.A., DeGasperi, C.L., Dokulil, M., La Fuente, S., Mackay, E.B., Schladow, S.G., Watanabe, S., Marcé, R., Pierson, D.C., Thiery, W. & Jennings, E. (2021). Phenological shifts in lake stratification under climate change. *Nature Communications*, 12, 2318.
- Wu, J. (2013). Key concepts and research topics in landscape ecology revisited: 30 years after the Allerton Park workshop. *Landscape Ecology*, 28, 1–11.
- Wu, X., Niu, C., Liu, X., Hu, T., Feng, Y., Zhao, Y., Liu, S., Liu, Z., Dai, G., Zhang, Y., Van Meerbeek, K., Wu, J., Liu, L., Guo, Q. & Su, Y. (2024). Canopy structure regulates autumn phenology by mediating the microclimate in temperate forests. *Nature Climate Change*, 14, 1299–1305.
- Wu, Z., Chen, S., De Boeck, H.J., Stenseth, N.C., Tang, J., Vitasse, Y., Wang, S., Zohner, C. & Fu, Y.H. (2021). Atmospheric brightening counteracts warming-induced delays in autumn phenology of temperate trees in Europe. *Global Ecology and Biogeography*, 30, 2477–2487.
- Yan, T., Fu, Y.H., Campioli, M., Peñuelas, J. & Wang, X. (2021). Divergent responses of phenology and growth to summer and autumnal warming. *Global Change Biology*, 27, null.
- Yuancai, L. & Parresol, B.R. (2001). Remarks on Height-Diameter Modeling. Tech. Rep. SRS-RN-10, U.S. Department of Agriculture, Forest Service, Southern Research Station, Asheville, NC.
- Zani, D., Crowther, T.W., Mo, L., Renner, S.S. & Zohner, C.M. (2020). Increased growing-season productivity drives earlier autumn leaf senescence in temperate trees. *Science*, 370, 1066–1071. Publisher: American Association for the Advancement of Science.
- Zohner, C.M., Mirzaghali, L., Renner, S.S., Mo, L., Rebindaine, D., Bucher, R., Palouš, D., Vitasse, Y., Fu, Y.H., Stocker, B.D. & Crowther, T.W. (2023). Effect of climate warming on the timing of autumn leaf senescence reverses after the summer solstice. *Science*, 381.

- Zohner, C.M., Mo, L., Renner, S.S., Svenning, J.C., Vitasse, Y., Benito, B.M., Ordonez, A., Baumgarten, F., Bastin, J.F., Sebold, V., Reich, P.B., Liang, J., Nabuurs, G.J., de Miguel, S., Alberti, G., Antón-Fernández, C., Balazy, R., Brändli, U.B., Chen, H.Y.H., Chisholm, C., Cienciala, E., Dayanandan, S., Fayle, T.M., Frizzera, L., Gianelle, D., Jagodzinski, A.M., Jaroszewicz, B., Jucker, T., Kepfer-Rojas, S., Khan, M.L., Kim, H.S., Korjus, H., Johannsen, V.K., Laarmann, D., Lang, M., Zawila-Niedzwiecki, T., Niklaus, P.A., Paquette, A., Pretzsch, H., Saikia, P., Schall, P., Šebeň, V., Svoboda, M., Tikhonova, E., Viana, H., Zhang, C., Zhao, X. & Crowther, T.W. (2020). Late-spring frost risk between 1959 and 2017 decreased in North America but increased in Europe and Asia. *Proceedings of the National Academy of Sciences*, 117, 12192–12200.
- Čufar, K., Prislan, P., De Luis, M. & Gričar, J. (2008). Tree-ring variation, wood formation and phenology of beech (*Fagus sylvatica*) from a representative site in Slovenia, SE Central Europe. *Trees*, 22, 749–758.