

title tbd

Context: Anthropogenic climate change is attributed to human emitted greenhouse gas, but the consequences on a biological systems are subject of active debate. While this is true, changing climate over the past decades have induced important changes on spring and fall phenological events — but the consequences on species fitness remain unclear. More precisely, spring phenological events have been advancing from 0.5 (Wolfe2005Climate) to 4.2 days/decade (Chmielewski and Rotzer2001;Fu2014Recent) and are driven by mainly driven by temperature (Chuine2010, Cleland2007, Penuelas et Fiella2009). In contrast, autumn phenophases (e.g. budset and leaf colouring) are delayed, though at a much lesser extent than spring's (Gallinat2015Autumn; Medvigy2014Macroscale, Menzel2006European) and are driven by shortening photoperiod (Flynn and Wolkovich2018; Korner and Basler2010, Cooke2010TheDynamic) and colder temperatures (Cooke2012, Delpierre2016, Lang1087). These shifts support a long-lasting —and intuitive— assumption that earlier spring and delayed fall events lead to longer seasons —which translate into increased growth. However, research from the past three years have casted doubt on this hypothesis. For instance Dow 2022, showed that despite an earlier growth onset, growth rate nor overall annual increment were increased by a theroritcal longer season.

This work led to my research questions. What are the mechanisms pushing down trees from growing more despite theoritcal longer growing seasons. Overall, there are two possible explanations: internal (via physiological constraints)(zohner2023) or external (environmental) limits to growth. As climate change has a complex and rather hard-to-predict nature, the external limits to growth are hard to quantify at the individual level as they affect communities as a whole. Drought, spring frost and heat waves are commonly known to be the main mechanisms that could limit tree growth under climate change (Drobyshev2008influence). To better comprehend these mechanisms, experiments are paramount because they can robustly tease out the internal vs external constraints to growth (e.g. warmer spring spring from severe drought later in the season —a common reality in natural environments). This is essential to quantify impacts on carbon sequestration of these constraints where in nature, is merely impossible. However, experiments are most often performed on saplings, which are critical for their role in forest regeneration projections, but they can hardly be translated to mature trees which hold for the overwhelming carbon biomass proportion of forests. For this, drone imagery paired with machine learning hold the promise of acquiring large sample sizes beyond what was achievable using traditional ground work and at a better resolution than sattelites. Therefore, I propose to use a combination or experiments to test internal (Chapter 1), along with external (Chapter 2) limits to growth along with observational data of a mixed-forest community using UAVs imagery and AI (Chapter 3) to address the paradox of the absence of increased growth despite apparent better conditions.

Chapter 1: extended growing season experiment (Fuelinex) continuation Shifts in phenophases have consequences on growth during the current growing season, but experiments

to date have yet to quantify if longer seasons have lagging effects over the following years (REFS). Therefore, for the first two years of my proposed award tenure, I will expand my Master's for an additional two growing seasons: 2025 and 2026, following the treatment year of 2024. Along with the continuation of phenological phase monitoring over the whole growing season, I will also collect the biomass at the end of the 2026 growing season and collect tree cross-sections. I will perform standard dendrochronological methods on them, scan them under a high-resolution scanner and perform ring width analysis. By using growth increment through tree rings, along with shoot elongation data collected for 3 consecutive seasons, I will leverage how the treatments of the first year affected growth over this year and the two following the treatments. With the expansion of my Master's work, I aim to assess tree species' potential to prolong or stretch their activity schedule and if that results in growth increment over the course of the growing seasons.

Chapter 2: drought and spring frost experiment With climate change, not only the growing season length shifts, but trees will experience shifts in the timing of moisture deficits from lower precipitation and higher evapotranspiration that may lead to drought stress (dox2022severe). Tree-ring (not sure) research shows that summer droughts advances growth cessation—leading to an earlier end of season (Kang2023An earlier; REF). In addition, warming springs lead to earlier budburst but come with increased the frequency and severity of late spring frosts which leads to tissue loss. Trees can recover by reinvesting in a second cohort of leaves. However, the lost time that trees cannot photosynthesize along with the increased investment in the second cohort may lead to significant disadvantages, but it's unclear whether trees going through spring frosts grow less. To investigate these processes, I will conduct an experiment consisting of three drought treatments, occurring at different timings during the growing season and an additional two treatments testing for spring frost events both at the start of the growing season when the buds have just started to burst and later, once the leaves have fully leaf out. I will use twelve deciduous tree species (six congeneric pairs to avoid potential confounding effects of shared evolutionary history) native to North America. The spring frost treatments will consist of placing the trees in growth chambers at a temperature 5 degrees warmer than ambient condition to trigger budburst. When the trees started to burst, I will place the first treatment for one hour in freezing growth chambers at a budkilling temperature set to the specific species LT50 values. For the second spring frost treatment, I will wait for the leaves to be elongated and then place the trees under the same freezing condition as the first treatment (Zohner2018Increased autumn).

For the drought treatments, the climate conditions will be set at 5 degrees warmer than ambient condition, at very low air humidity to maximize evapotranspiration rate. Once all the trees of a specific species have reached their respective wilting point (values at which soil water is not extractable by the plant), the trees will be removed from the chambers and put back to ambient condition at constant irrigation. The three drought treatments won't differ in their conditions, but instead in their timing since little attention has been paid to the importance of the moment of the drought. Therefore, the first treatment will be conducted just after leaf-out and before summer solstice. The second will be start one week before solstice — timing of peak growth for a lot of species***. The last drought treatment will occur just at the beginning of budset which is the theoretical end of early wood growth and onset of late wood growth. Phenological phases, shoot

elongation will be monitored weekly throughout the growing season. Biomass will be estimated using allometric equations at the start and end of the growing season. In order to grasp a high temporal resolution of growth responses to treatments, 5 replicates per species/treatments will be equipped with micro magnetic dendrometers that will provide valuable previously impossible to get onto how trees respond to growth at an extremely high temporal resolution.

Chapter 3 : cambial phenology X drone imagery phenological observations Getting a better understanding of accross species differences in growth synchrony with leaf phenology is paramount to refine carbon sequestration models in the face of Anthropogenic climate change. Thus, I aim to launch a large-scale project using cutting-edge technology to gatter a large amount of data of trees' growth onset and end from a mixed-forest community located at Station bilogique des Laurentides (St-Hypollyte (Qc)), during three growing seasons. Using this site will allow me to follow up work previously done by my lab as well as creating a partnership with Dr. Etienne Laliberté from the Plant Functional Ecology Laboratory (PFEL) who curently uses this site for his research. To monitor leaf phenology from budburst to leaf drop, I will use high-frequency repeat overflights using Unmaned Aerian Vehicles (UAVs) over the canopy of these for communities to monitor every single trees over the course of the growing season. Partnering with the PFEL will allow me to automatically acquire hundreds of close-up photos of individual tree crowns per day to monitor critical phenophases. Along with this imagery, I will use 200 DC3 Perimeter Dendrometer placed randomly throughtout the site on 40 trees per species. I believe that using high resolution data accross space and time will allow me to infer a strong relationship between leaf phenophases and growth seasonality

1 References