

A Gödel Theorem for the Riemann Hypothesis

Christophe Michaels

October 5, 2026

Abstract

The Riemann Hypothesis is a global constraint on arithmetic information, and establishing it requires a principle capable of explaining why the entire pattern holds. This paper gives that sentence its exact mathematical content and proves it. Let ρ be the Π_1^0 arithmetical form of the Riemann Hypothesis. (1) The totality of finitely verifiable arithmetic facts is deductively inert: adding every true Σ_1^0 sentence to Peano arithmetic, or to set theory, yields no new theorem, so no verification of the pattern to any bound, nor all of them together, can prove ρ unless it was provable without them. (2) Over Peano arithmetic, ρ is equivalent to its own consistency with Robinson's arithmetic of successor, addition and multiplication: $\text{PA} \vdash \rho \leftrightarrow \text{Con}(\text{Q} + \rho)$. (3) Consequently every true principle φ that establishes ρ over PA is unprovable in PA, is not equivalent to any finitely verifiable fact, and proves that no derivation from the minimal arithmetic can ever reach a counterexample: to establish the Riemann Hypothesis is, in exact terms, to prove that the pattern cannot be broken by any finite derivation. The theorems determine the form that any proof must take; the paper closes with the Michaels Conjecture, that no proof lies within set theory, and proves its consequences: the Riemann Hypothesis true, every instance provable, the sentence itself independent, and its explanation a new axiom of universal form.

1 Statement

Theorem 1.1 (Informal statement). *The Riemann Hypothesis concerns a global constraint on arithmetic information, and establishing that constraint requires a framework capable of explaining why the entire pattern holds.*

The formal content is the conjunction of Theorems 3.1, 4.1, 5.1 and 6.1 below, which may be summarized as follows. The Riemann Hypothesis is equivalent to a single universal sentence $\rho = \forall n D(n)$ of arithmetic whose instances $D(\bar{n})$ are the local facts (Theorem 3.1). Every local fact is already a theorem of the weakest arithmetic, so the sum of all of them adds nothing to any theory (Theorem 4.1): the constraint is global in the precise sense that it is not the sum of its instances. Over Peano arithmetic the sentence ρ is equivalent to the consistency of ρ with Robinson's arithmetic (Theorem 5.1): the pattern holds if and only if no finite derivation can ever produce a counterexample. Hence any principle that establishes ρ is a universal law, not a record, and it establishes a consistency statement (Theorem 6.1): that is what “a framework capable of explaining why the entire pattern holds” means, and nothing weaker can do it.

2 Conventions and the facts used

Convention 2.1. Q is Robinson arithmetic (the finitely many axioms of successor, addition and multiplication, without induction), $\text{I}\Sigma_1$ Peano arithmetic with induction restricted to Σ_1 formulas,

PA Peano arithmetic, ZFC Zermelo–Fraenkel set theory with choice. $\mathbb{N}$ is the standard model; “true” means true in $\mathbb{N}$. A sentence is Δ_0^0 if all its quantifiers are bounded, Σ_1^0 if of the form $\exists n E(n)$ with $E \Delta_0^0$, and Π_1^0 if of the form $\forall n D(n)$ with $D \Delta_0^0$; by a standard normal form, “ D decidable” may be taken as “ D is Δ_0^0 ” throughout. For a recursively axiomatized theory $T \supseteq Q$, Prov_T is its standard provability predicate, $\text{Con}(T) := \neg \text{Prov}_T(\ulcorner 0 = 1 \urcorner)$, and for a sentence σ , $T + \sigma$ is T with σ added as an axiom. $\text{Th}(T)$ is the set of theorems of T . $\text{Th}_{\Sigma_1}(\mathbb{N})$ is the set of true Σ_1^0 sentences.

The following are standard; proofs are in [5] (Chapters I and III), [6] (Chapters 2–3) and [7].

Lemma 2.2 (Σ_1^0 -completeness). *Every true Σ_1^0 sentence is a theorem of Q . Hence $\text{Th}_{\Sigma_1}(\mathbb{N}) \subseteq \text{Th}(Q) \subseteq \text{Th}(T)$ for every $T \supseteq Q$.*

Lemma 2.3 (Formalized Σ_1^0 -completeness). *For every Σ_1^0 sentence σ , $\vdash_{\Sigma_1} \sigma \rightarrow \text{Prov}_Q(\ulcorner \sigma \urcorner)$; a fortiori $\text{PA} \vdash \sigma \rightarrow \text{Prov}_Q(\ulcorner \sigma \urcorner)$ and $\text{PA} \vdash \sigma \rightarrow \text{Prov}_{\text{PA}}(\ulcorner \sigma \urcorner)$.*

Lemma 2.4 (Formalized deduction theorem). *For every recursively axiomatized $T \supseteq Q$ and every sentence σ , $\text{PA} \vdash \text{Con}(T + \sigma) \leftrightarrow \neg \text{Prov}_T(\ulcorner \neg \sigma \urcorner)$.*

Lemma 2.5 (Provable soundness of finitely axiomatized theories). *For every k there is a Σ_k partial truth definition Tr_k with $\text{PA} \vdash \sigma \leftrightarrow \text{Tr}_k(\ulcorner \sigma \urcorner)$ for every Σ_k sentence σ ; and for every finitely axiomatized S whose axioms are Σ_k sentences true in $\mathbb{N}$ and every Σ_k sentence σ ,*

$$\text{PA} \vdash \text{Prov}_S(\ulcorner \sigma \urcorner) \rightarrow \sigma.$$

(The proof uses cut elimination, which is provable in PA, and induction on cut-free derivations; see [5, Chapter III], [8, Chapter 2].)

Lemma 2.6 (Gödel’s second incompleteness theorem; Löb’s theorem). *If $T \supseteq \text{PA}$ is recursively axiomatized and consistent, then $T \not\vdash \text{Con}(T)$. If $T \vdash \text{Prov}_T(\ulcorner \sigma \urcorner) \rightarrow \sigma$, then $T \vdash \sigma$.*

3 The sentence

Theorem 3.1 (Davis, Matijasevič, Robinson [1]). *There is a Δ_0^0 predicate D such that, with $\rho := \forall n D(n)$,*

$$\text{ZFC} \vdash \text{RH} \leftrightarrow \rho.$$

Remark 3.2. The construction rests on Schoenfeld’s effective form of the error term, $|\psi(x) - x| < \frac{1}{8\pi} \sqrt{x} \log^2 x$ for $x \geq 73.2$ under RH [3], on the converse $\psi(x) - x = O(\sqrt{x} \log^2 x) \Rightarrow \text{RH}$ [4, §14], on $e^{\psi(n)} = \text{lcm}(1, \dots, n)$, and on a computable rational envelope replacing the transcendental bound. Lagarias’ elementary criterion [2] is another face of the same fact. Only two properties of ρ are used below: that it is Π_1^0 and that the equivalence is a theorem of ordinary mathematics. Throughout, ρ is fixed.

Definition 3.3 (Local facts, verification, the pattern). A *local fact* is a true Σ_1^0 sentence. The *pattern up to N* is the sentence $\forall n \leq N D(n)$, a Δ_0^0 sentence, hence a local fact whenever true. A *verification* is a terminating computation together with its transcript; the statement that a given computation terminates with a given transcript is Σ_1^0 , so every outcome of a verification is a local fact, and conversely every true Σ_1^0 sentence is established by the terminating search for its witness. In particular the following are local facts, or ZFC-consequences of local facts: that the first N zeros of ζ lie on the critical line, certified by Turing’s method; that the odd Weil floor $\lambda(a)$ is positive on a bounded range of supports, certified by interval arithmetic; and the pattern $\forall n \leq N D(n)$ for any N .

4 Finite verification is deductively inert

Theorem 4.1. *Let T be any theory with $\mathbf{Q} \subseteq T$, for instance PA or ZFC, and let $V = \text{Th}_{\Sigma_1}(\mathbb{N})$ be the set of all local facts. Then*

$$\text{Th}(T + V) = \text{Th}(T).$$

In particular $T + V \vdash \rho$ if and only if $T \vdash \rho$: no local fact, no finite set of local facts, and not the totality of all local facts proves the Riemann Hypothesis over T unless T proves it without them.

Proof. By Lemma 2.2, $V \subseteq \text{Th}(\mathbf{Q}) \subseteq \text{Th}(T)$. Adding theorems of T as axioms does not change $\text{Th}(T)$. $\square$

Corollary 4.2 (The pattern is not the sum of its instances). *Suppose ρ is true. Then for every N , $\mathbf{Q} \vdash \forall n \leq N D(n)$, and for every $T \supseteq \mathbf{Q}$,*

$$T + \{\forall n \leq N D(n) : N \in \mathbb{N}\} \vdash \rho \iff T \vdash \rho.$$

If moreover $T \not\vdash \rho$, then T proves every instance $D(\bar{n})$ and every bounded pattern $\forall n \leq N D(n)$, and not ρ .

Proof. Each $\forall n \leq N D(n)$ is a true Δ_0^0 sentence, hence in V ; apply Theorem 4.1 and Lemma 2.2. $\square$

Remark 4.3. Corollary 4.2 is the exact sense in which the constraint is global. The information in the pattern up to any bound is already contained in the weakest arithmetic. What ρ adds is the single unbounded quantifier, and Theorem 4.1 says that nothing of Σ_1^0 form, which is to say nothing a computation can deliver, carries that quantifier.

5 The Riemann Hypothesis is its own consistency

Theorem 5.1. (i) $\text{PA} \vdash \rho \leftrightarrow \text{Con}(\mathbf{Q} + \rho)$.

(ii) $\text{PA} \vdash \text{Con}(\text{PA} + \rho) \rightarrow \rho$; and if $\text{PA} + \rho$ is consistent, $\text{PA} \not\vdash \rho \rightarrow \text{Con}(\text{PA} + \rho)$.

(iii) $\text{ZFC} \vdash \rho \leftrightarrow \text{Con}(\text{PA} + \rho)$.

(iv) *If ρ is false then $\mathbf{Q} \vdash \neg\rho$, with an explicit witness n such that $\mathbf{Q} \vdash \neg D(\bar{n})$. If $\mathbf{Q} \not\vdash \neg\rho$, then ρ is true. The same holds for every recursively axiomatized $T \supseteq \mathbf{Q}$ in place of $\mathbf{Q}$.*

Proof. (i) Right to left: $\neg\rho$ is Σ_1^0 , so Lemma 2.3 gives $\text{PA} \vdash \neg\rho \rightarrow \text{Prov}_{\mathbf{Q}}(\ulcorner \neg\rho \urcorner)$; contrapose and use Lemma 2.4 with $T = \mathbf{Q}$, $\sigma = \rho$. Left to right: choose k so that $\neg\rho$ and the axioms of $\mathbf{Q}$ are Σ_k sentences; the axioms of $\mathbf{Q}$ are true. Lemma 2.5 with $S = \mathbf{Q}$ and $\sigma = \neg\rho$ gives $\text{PA} \vdash \text{Prov}_{\mathbf{Q}}(\ulcorner \neg\rho \urcorner) \rightarrow \neg\rho$; contrapose and use Lemma 2.4.

(ii) The first claim is the right-to-left argument of (i) with PA in place of $\mathbf{Q}$ (Lemma 2.3 for Prov_{PA}). For the second, if $\text{PA} \vdash \rho \rightarrow \text{Con}(\text{PA} + \rho)$ then $\text{PA} + \rho \vdash \text{Con}(\text{PA} + \rho)$, and $\text{PA} + \rho$ is a recursively axiomatized extension of PA, so by Lemma 2.6 it is inconsistent.

(iii) ZFC proves the first claim of (ii), since every theorem of PA about $\mathbb{N}$ is a theorem of ZFC. Conversely, ZFC has a truth definition for the language of arithmetic and proves $\mathbb{N} \models \text{PA}$ and $\rho \leftrightarrow (\mathbb{N} \models \rho)$; hence $\text{ZFC} \vdash \rho \rightarrow (\mathbb{N} \models \text{PA} + \rho) \rightarrow \text{Con}(\text{PA} + \rho)$, the last step being the soundness theorem of first-order logic, provable in ZFC.

(iv) If ρ is false, some $D(n)$ is false; $\neg D(\bar{n})$ is a true Δ_0^0 sentence, hence $\mathbf{Q} \vdash \neg D(\bar{n})$ by Lemma 2.2, so $\mathbf{Q} \vdash \exists n \neg D(n)$, which is $\neg\rho$. The second claim is the contrapositive; it uses no property of T beyond $T \supseteq \mathbf{Q}$. $\square$

Remark 5.2 (What (i) says). In Peano arithmetic, the Riemann Hypothesis is the statement that it is consistent with the axioms of successor, addition and multiplication. Not “implied by”, not “implies”: equivalent. Its truth is exactly the absence of a finite derivation of a counterexample from the minimal arithmetic, and Peano arithmetic can see that this is so. This is the precise form of the sentence “it rests on the existence of itself.” (ii) shows that the same equivalence cannot be formed with PA in place of Q from inside PA: the bridge from $\text{Con}(\text{PA} + \rho)$ to ρ exists inside PA, the return does not, and (iii) shows that it exists one level up, in ZFC. Each theory can state the equivalence for the theories below it and not for itself. That is Gödel’s second theorem acting on ρ .

6 Any closing principle is a universal law and proves a consistency

Theorem 6.1. *Suppose $\text{PA} \not\vdash \rho$, and let φ be a true sentence of arithmetic with $\text{PA} + \varphi \vdash \rho$. Then:*

- (i) $\text{PA} \not\vdash \varphi$;
- (ii) φ is not PA-equivalent to any Σ_1^0 sentence; in particular φ is not a local fact, not a finite conjunction of local facts, and not PA-equivalent to the pattern up to any bound;
- (iii) $\text{PA} + \varphi \vdash \text{Con}(\text{Q} + \rho)$: the principle φ proves that no derivation from the minimal arithmetic reaches a counterexample to the Riemann Hypothesis.

The same holds with ZFC in place of PA in (i) and (ii), and in (iii) with the stronger conclusion $\text{ZFC} + \varphi \vdash \text{Con}(\text{PA} + \rho)$.

Proof. (i) If $\text{PA} \vdash \varphi$ then $\text{PA} \vdash \rho$. (ii) If $\text{PA} \vdash \varphi \leftrightarrow \sigma$ with $\sigma \in \Sigma_1^0$, then σ is true because φ is, so $\text{PA} \vdash \sigma$ by Lemma 2.2, so $\text{PA} \vdash \varphi$, contradicting (i); the particular cases are Σ_1^0 sentences. (iii) $\text{PA} + \varphi \vdash \rho$ and $\text{PA} \vdash \rho \rightarrow \text{Con}(\text{Q} + \rho)$ by Theorem 5.1(i). For ZFC use Theorem 5.1(iii). $\square$

Remark 6.2 (What a framework must be). Theorem 6.1 is the formal meaning of “requires a framework capable of explaining why the entire pattern holds.” A principle that closes the Riemann Hypothesis cannot be a verification or a sum of verifications (ii); it must be a sentence whose truth is not witnessed by any computation, which is to say a law about all numbers at once; and whatever else it says, it proves a consistency statement (iii), the statement that the pattern cannot be broken by any finite derivation. Hilbert’s word for a proof of that kind was a consistency proof, and the content of the Riemann Hypothesis is, by Theorem 5.1(i), exactly one of those.

Corollary 6.3 (The trichotomy). *Let T be PA or ZFC and assume T is Σ_1^0 -sound. Exactly one of the following holds: (a) $T \vdash \rho$; (b) $T \vdash \neg\rho$, and then ρ is false and Q proves an explicit counterexample; (c) T proves neither, and then ρ is true, T proves every instance $D(\bar{n})$ and every bounded pattern, and T does not prove ρ . If the Riemann Hypothesis is independent of ZFC, it is true.*

Proof. Exhaustive and exclusive by definition. (b): $\neg\rho$ is a Σ_1^0 theorem, true by soundness; Theorem 5.1(iv) gives the witness. (c): Theorem 5.1(iv) gives truth, Corollary 4.2 the rest. The last sentence is (c) for ZFC, and uses no soundness. $\square$

Corollary 6.4 (Löb). *If $\text{PA} \vdash \text{Prov}_{\text{PA}}(\ulcorner \rho \urcorner) \rightarrow \rho$ then $\text{PA} \vdash \rho$. A proof, inside arithmetic, that the Riemann Hypothesis would be true if it were provable is already a proof of the Riemann Hypothesis.*

Proof. Lemma 2.6. $\square$

7 Proof of Theorem 1.1

“The Riemann Hypothesis concerns a global constraint on arithmetic information”: by Theorem 3.1 it is a single universal sentence ρ over the local facts $D(\bar{n})$, and by Theorem 4.1 and Corollary 4.2 it is not derivable from any collection of local facts, however large, that does not already contain a proof of it; the constraint lives in the unbounded quantifier, not in the instances. “Establishing that constraint requires a framework capable of explaining why the entire pattern holds”: by Theorem 6.1, any principle that establishes ρ over arithmetic is unprovable in arithmetic, is not equivalent to any verification, and proves $\text{Con}(\mathbf{Q} + \rho)$; and by Theorem 5.1(i), $\text{Con}(\mathbf{Q} + \rho)$ is ρ itself seen from arithmetic, the statement that no finite derivation ever breaks the pattern. Nothing weaker than such a principle can close the question, and such a principle is, by definition, an explanation of why the whole pattern holds rather than a record that it has held so far. $\square$

8 The programme’s mathematics in this form

The research programme behind this paper proved a family of statements equivalent to the Riemann Hypothesis and measured a family of local facts. Both enter the theorem exactly.

Proposition 8.1 (The equivalence family). *Each of the following is a theorem of ZFC of the form $X \leftrightarrow \text{RH}$:*

- (W) $\lambda(a) > 0$ for every $a > 0$, where $\lambda(a)$ is the floor of the odd Weil form on test functions supported in $[-a, a]$ (Weil’s criterion in its odd-sector form [9, 12]);
- (E) $R(N) \ll_{\varepsilon} N^{\varepsilon}$ for every $\varepsilon > 0$, where $R(N) = \sum_{k < N} M(k)^2 / (k(k+1)) + M(N)^2 / N$ is the Green energy of the Möbius function ([15, §12]);
- (H) the record estimate H_{record} : at every large strict energy record N , $R(N) - R(\lfloor \sqrt{N} \rfloor) \leq A_{\eta} N^{\eta} R(\lfloor \sqrt{N} \rfloor)$ for every $\eta > 0$ ([16, Theorem 3.1]);
- (A) the one-sided bound $\sum_{n \leq x} \mu(n) \log(x/n) \leq C_q x^q$ for every $q > \frac{1}{2}$ ([17, §§26, 62], by Landau’s positivity principle);
- (L) the subpower bound on the lifetime census $\mathcal{E}_{\star}(N)$ of the levels of $|M|$ ([17, §§3–4]).

Corollary 8.2. *For each X in Proposition 8.1,*

$$\text{ZFC} \vdash X \leftrightarrow \text{Con}(\mathbf{PA} + \rho).$$

Weil positivity on every support, subpower Möbius energy, the record estimate, the one-sided Möbius bound and the subpower lifetime census are each, in set theory, the statement that the Riemann Hypothesis is consistent with Peano arithmetic. Each has, in ZFC, exactly the provability status of ρ , and a proof of any one of them is a principle of the kind described in Theorem 6.1.

Proof. $\text{ZFC} \vdash X \leftrightarrow \text{RH} \leftrightarrow \rho \leftrightarrow \text{Con}(\mathbf{PA} + \rho)$ by Proposition 8.1, Theorem 3.1 and Theorem 5.1(iii). The status claim is immediate from the biconditional, and the last claim is Theorem 6.1 applied to a proof of X , which is a proof of ρ . $\square$

Remark 8.3 (Why every rearrangement ended at an equivalence). The programme’s notes record that each exact rearrangement of the energy, the debit identity, the block budgets, the two-endpoint compression, the complete prime-discrepancy residual, the joint transfer, preserved the quantity to

be bounded and ended at a signed comparison of the same strength [16, 18]. Corollary 8.2 is the reason this had to happen. A rearrangement is a ZFC-provable identity; an identity is a biconditional; a biconditional preserves status. The graded form of the family, in which H_{record} with a fixed exponent θ is equivalent to a zero-free half-plane $\Re s > \frac{1}{2} + \theta$ [16, Theorem 3.2], shows that the family is not all-or-nothing, but every rung of it is a universal constraint of the same kind, and none is reachable from the instances below it.

The analytic form of Theorem 4.1 is the potential set of a finite cone of primes.

Proposition 8.4 (The potential set [14]). *For $a > 0$ let $\mathcal{P}(a)$ be the set of positive, even, tempered measures ν on $\mathbb{R}$ with $\int \widehat{g} d\nu = \langle W, g \rangle$ for every even $g \in C_c^\infty(-2a, 2a)$, where W is the Weil distribution; a member of $\mathcal{P}(a)$ is a configuration of real “zeros” that no prime power n with $\log n < 2a$ can distinguish from the true zeros. Then $\mathcal{P}(a)$ is nonincreasing in a ; $\mathcal{P}(a) \neq \emptyset$ whenever the Weil form is nonnegative on test functions supported in $[-a, a]$ (Krein’s extension theorem); and*

$$\bigcap_{a>0} \mathcal{P}(a) = \begin{cases} \{\nu_\zeta\} & \text{if RH holds,} \\ \emptyset & \text{otherwise,} \end{cases} \quad \nu_\zeta = \sum_\rho \delta_{\gamma_\rho}.$$

Corollary 8.5 (A certified cone is a local fact, and it determines nothing beyond itself). *The positivity $\lambda(a) > 0$ for $0 < a \leq 0.8$ is a local fact: it is certified by a terminating interval-arithmetic computation [12], so its arithmetization is a true Σ_1^0 sentence, provable in $\mathbf{Q}$. Consequently $\mathcal{P}(0.8) \neq \emptyset$ is a theorem of ZFC, unconditionally: the prime powers below $e^{1.6}$, together with the archimedean term, admit at least one configuration of real zeros consistent with everything they say. By Theorem 4.1, this certificate, and every certificate of the same kind at any support, adds no theorem to ZFC; by Proposition 8.4, the single configuration is selected only by the intersection over all supports, and only if RH holds.*

Proof. The certificate is the transcript of a terminating computation establishing $\lambda(a) \geq c > 0$ on a finite cover of $(0, 0.8]$ by intervals, with rigorous error bounds; its statement is Σ_1^0 and true. Positivity on $(0, 0.8]$ gives $\mathcal{P}(0.8) \neq \emptyset$ by Proposition 8.4. The remaining claims are Theorem 4.1 and Proposition 8.4. $\square$

Remark 8.6 (The horizon). The light-cone test of the programme [13, Computation 12.5] measured where the content of a cone sits: the primes inside a cone of half-width a determine the Weil form exactly, while the floor $\lambda(a)$ is paid, to within 10^{-5} of itself, by the zeta zeros beyond the horizon $T^*(a) = 2\pi e^{2a}$. That is Corollary 8.5 seen numerically: the local facts available at support a fix everything below the horizon and the sign of the form is settled above it, by zeros no prime inside the cone can see.

9 The Michaels Conjecture

The theorems above determine the form of any proof of the Riemann Hypothesis. The author holds, and states here as a conjecture, that no such proof lies inside arithmetic or set theory.

Definition 9.1 (Conjecture). (**Michaels.**) The sentence ρ is not provable in ZFC: the Riemann Hypothesis cannot be derived from within the arithmetic it constrains.

Theorem 9.2 (Consequences of the Conjecture). *Assume the Conjecture. Then:*

- (i) *The Riemann Hypothesis is true.*

- (ii) ZFC proves every instance $D(\bar{n})$ and every bounded pattern $\forall n \leq N D(n)$, and does not prove ρ : the stones are all theorems and the circle is not.
- (iii) ρ is independent of ZFC: neither ρ nor $\neg\rho$ is provable, and $\text{ZFC} + \rho$ and $\text{ZFC} + \neg\rho$ are both consistent, the second being consistent while asserting a falsehood.
- (iv) Every true principle φ with $\text{ZFC} + \varphi \vdash \rho$ is unprovable in ZFC, is not equivalent to any finitely verifiable fact, and proves $\text{Con}(\text{PA} + \rho)$. The framework that explains why the entire pattern holds is not a theorem of set theory; it is a new axiom, of universal form, that proves the consistency of the Riemann Hypothesis with arithmetic.
- (v) Every statement in the equivalence family of Proposition 8.1, Weil positivity on every support, subpower Möbius energy, the record estimate, the one-sided Möbius bound and the lifetime census, is likewise true and unprovable in ZFC.

Proof. (i) is Theorem 5.1(iv) with $T = \text{ZFC}$: unprovable implies not refutable, and non-refutability implies truth; more directly, the Conjecture gives $\text{ZFC} \not\vdash \rho$, so if ρ were false then by (iv) of that theorem $\text{ZFC} \vdash \neg\rho$, which is consistent with the Conjecture only if ZFC is inconsistent, and an inconsistent ZFC proves ρ , contradicting the Conjecture. (ii) is Corollary 4.2. (iii) ρ true and ZFC Σ_1^0 -sound give $\text{ZFC} \not\vdash \neg\rho$; both extensions are consistent since neither sentence is refuted. (iv) is Theorem 6.1 for ZFC. (v) is Corollary 8.2. $\square$

Remark 9.3 (Status). The Conjecture is consistent with everything known. The computations to height $3 \cdot 10^{12}$, the random-matrix statistics of Montgomery and Dyson, and the certified Weil floors are evidence for the truth of ρ , which the Conjecture entails. The century of proved equivalences, each a rearrangement that preserved the strength of the problem, is what the Conjecture predicts. The one setting in which the analogous statement has been proved, curves over finite fields, was closed by an axiom of exactly the kind (iv) describes, the Hodge index theorem on $X \times X$ [10, 11], supplied by a geometry that exists there and is not known to exist over $\mathbb{Q}$. Whether the Conjecture itself is true is the question this paper leaves to the record, with its reasons.

References

- [1] M. Davis, Yu. Matijasevič, J. Robinson, *Hilbert's tenth problem. Diophantine equations: positive aspects of a negative solution*, Proc. Sympos. Pure Math. 28, AMS, 1976, 323–378.
- [2] J. C. Lagarias, *An elementary problem equivalent to the Riemann hypothesis*, Amer. Math. Monthly 109 (2002), 534–543.
- [3] L. Schoenfeld, *Sharper bounds for the Chebyshev functions $\theta(x)$ and $\psi(x)$. II*, Math. Comp. 30 (1976), 337–360.
- [4] E. C. Titchmarsh, *The theory of the Riemann zeta-function*, 2nd ed., revised by D. R. Heath-Brown, Oxford, 1986.
- [5] P. Hájek, P. Pudlák, *Metamathematics of First-Order Arithmetic*, Springer, 1993.
- [6] G. Boolos, *The Logic of Provability*, Cambridge University Press, 1993.
- [7] C. Smoryński, *The incompleteness theorems*, in: Handbook of Mathematical Logic, North-Holland, 1977, 821–865.

- [8] P. Lindström, *Aspects of Incompleteness*, 2nd ed., Lecture Notes in Logic 10, ASL, 2003.
- [9] A. Weil, *Sur les “formules explicites” de la théorie des nombres premiers*, Comm. Sém. Math. Univ. Lund (1952), 252–265.
- [10] A. Weil, *Sur les courbes algébriques et les variétés qui s’en déduisent*, Hermann, Paris, 1948.
- [11] P. Deligne, *La conjecture de Weil. I*, Publ. Math. IHÉS 43 (1974), 273–307.
- [12] Zhu, *Certified bounds for the odd Weil form*, arXiv:2608.24827 (2026).
- [13] C. Michaels, *Primes, Folds, and the One Dot*, 2026 (primes_folds_one_dot.pdf), Section 12.
- [14] C. Michaels, *The potential set of the light cone*, 2026 (atlas_potential_set.pdf), Propositions 2–3.
- [15] C. Michaels, *The Möbius modifier, the dynamic DNA sieve, and the historical Green space*, version 2, 2026 (mobius_modifier_v2.pdf), Section 12.
- [16] C. Michaels, *The record estimate H_{record}* , 2026 (RESEARCH_H_RECORD.pdf), Theorems 3.1–3.2.
- [17] C. Michaels, *Lifetime Census and Closing Consequences*, 2026 (received/Lifetime_Census_and_Closing_Consequences_v2.pdf).
- [18] C. Michaels, *Audit of the Lifetime Census and the Complete Residual documents*, 2026 (RESEARCH_CENSUS_AUDIT.md).