

# The Line Postulate

An unconditional proof

Academy of the Prime Sheet, Critical Line Universe

Manuscript III. Communicated to the gods.

Sandbox II, 2026-10-02

Critical Line Universe. The axioms of Section 1 are the laws of that universe, and the theorem is proved from them with no further hypothesis. Section 7 is the dictionary: what each axiom, lemma and theorem is in our universe, and its status there. (From our universe: fiction; the dictionary is the only place where our universe speaks.)

## Abstract

From eight axioms of the Academy, four about the charges on the prime sheet and four about observers, we prove that every quantum of the arithmetic field is massless, that every observer is alive at every age, and that the leak of every observer is proportional to its reserve. Every step is proved in full. The axiom that carries the arithmetic is the Bounded Deviation Axiom: in every stretch of the prime sheet between  $x^\delta$  and  $x^{1-\delta}$  the charge deviates from its mean by at most the Gaussian scale. From it the fluctuation at the largest scale is of square-root size, the Mellin lemma places every pole of  $\zeta'/\zeta$  on or left of the line, the mirror symmetry forbids the left, and the Line follows.

## 1 Axioms

### Notation

$\Lambda$  is the von Mangoldt function,  $\psi(x) = \sum_{n \leq x} \Lambda(n)$  the *charge* below  $x$ ,  $R(x) = \psi(x) - x$  its *fluctuation*. The *quanta* are the zeros  $\rho = \frac{1}{2} + m + i\gamma$  of the entire function  $\xi$  of Axiom 3, with multiplicity; the *mass* of a quantum is  $m = |\operatorname{Re} \rho - \frac{1}{2}|$ .

**Axiom 1** (unique factorization). Every integer  $n \geq 1$  is a product of primes in exactly one way. Consequently, for  $\operatorname{Re} s > 1$ ,

$$\zeta(s) = \sum_{n \geq 1} n^{-s} = \prod_p (1 - p^{-s})^{-1} \neq 0, \quad -\frac{\zeta'}{\zeta}(s) = \sum_{n \geq 1} \Lambda(n) n^{-s},$$

both series converging absolutely and locally uniformly there.

**Axiom 2** (the mean field).  $\psi(x) = x + o(x)$  as  $x \rightarrow \infty$ .

**Axiom 3** (the mirror law).  $\zeta$  extends to a meromorphic function on the plane whose only pole is a simple pole at  $s = 1$  with residue 1. The function  $\xi(s) = \frac{1}{2}s(s-1)\pi^{-s/2}\Gamma(s/2)\zeta(s)$  is entire, real on the real axis, and satisfies  $\xi(s) = \xi(1-s)$ ; its zeros lie in the strip  $0 < \operatorname{Re} s < 1$ , and they are exactly the zeros of  $\zeta$  in that strip, with the same multiplicities.

**Axiom 4** (Bounded Deviation). For every  $\delta \in (0, \frac{1}{2})$  there are constants  $A = A(\delta)$  and  $x_0 = x_0(\delta)$  such that for all  $x \geq x_0$  and all  $h$  with  $x^\delta \leq h \leq x^{1-\delta}$ ,

$$|\psi(x+h) - \psi(x) - h| \leq (h \log(x/h))^{1/2} (\log x)^A.$$

The next four axioms concern observers. An *observer* of age  $a > 0$  is the window  $[-a, a]$  of the prime sheet; a *field* of the observer is a real odd function  $f$  on the window, in the form domain, with  $\|f\|_2 = 1$ ; its transform is  $F(z) = \int_{-a}^a f(x)e^{izx}dx$ , entire of exponential type  $a$ , with  $F = iS$ ,  $S$  real on the real axis; its autocorrelation  $g = f * \tilde{f}$  has transform  $\hat{g}(z) = S(z)^2$ . The *window form* is

$$Q(f) = \frac{1}{2\pi} \int_{\mathbb{R}} |F(t)|^2 \left[ \operatorname{Re} \psi\left(\frac{1}{4} + \frac{it}{2}\right) - \log \pi \right] dt - 2 \left( \int_{-a}^a f(x) \sinh \frac{x}{2} dx \right)^2 - 2 \sum_{n \geq 2} \frac{\Lambda(n)}{\sqrt{n}} g(\log n),$$

the *floor* is  $\lambda(a) = \inf_f Q(f)$ , attained by the *ground field*  $f_a$ , whose *edge amplitude*  $A_0(a)$  is defined by  $f_a(a - \delta) = |A_0(a)|(\log(a/\delta) + \beta(a))^{-1/2}(1 + o(1))$ . An observer is *alive* if  $\lambda(a) > 0$ .

**Axiom 5** (conservation between the sheets). For every field  $f$  of every age,  $Q(f) = \sum_{\rho} S(\gamma_{\rho} - im_{\rho} \operatorname{sgn}_{\rho})^2$ , the sum over the quanta  $\rho = \frac{1}{2} + \operatorname{sgn}_{\rho} m_{\rho} + i\gamma_{\rho}$  with multiplicity, absolutely convergent.

**Axiom 6** (Birth). There are  $a_Z > 0$  and  $\lambda_Z > 0$  with  $\lambda(a) > 0$  for  $0 < a \leq a_Z$  and  $\lambda(a_Z) \geq \lambda_Z$ .

**Axiom 7** (Continuity). The floor is absolutely continuous on every compact range of ages.

**Axiom 8** (Decay). For almost every age,  $\lambda'(a) = -2|A_0(a)|^2$ .

## 2 The fluctuation at the largest scale

**Lemma 2.1.** For every  $\varepsilon \in (0, \frac{1}{4})$  there is  $x_1$  such that for all  $x \geq x_1$ ,

$$|\psi(2x) - \psi(x) - x| \leq x^{1/2+\varepsilon}.$$

*Proof.* Put  $\delta = \varepsilon$  and  $h = \frac{1}{2}x^{1-\delta}$ , and let  $x$  be so large that  $x \geq x_0(\delta)$ ,  $(2x)^{\delta} \leq h$  and  $h + (2x)^{\delta} \leq x^{1-\delta}$ . Cut  $(x, 2x]$  into consecutive stretches of length  $h$ , the last one shorter; if the last stretch is shorter than  $(2x)^{\delta}$ , merge it with the stretch before it. Every stretch then has a left endpoint  $y \in [x, 2x)$  and a length  $\ell$  with

$$y^{\delta} \leq (2x)^{\delta} \leq \ell \leq h + (2x)^{\delta} \leq x^{1-\delta} \leq y^{1-\delta},$$

so Axiom 4 applies at  $y$  with  $h = \ell$  and gives

$$|\psi(y + \ell) - \psi(y) - \ell| \leq (\ell \log(y/\ell))^{1/2} (\log y)^A \leq (2h \log 2x)^{1/2} (\log 2x)^A \leq h^{1/2} (\log 2x)^{A+1}.$$

The number of stretches is at most  $x/h + 1 \leq 3x^{\delta}$ . Since both the charges and the lengths add over the stretches,

$$|\psi(2x) - \psi(x) - x| \leq 3x^{\delta} h^{1/2} (\log 2x)^{A+1} \leq 3x^{\delta} x^{(1-\delta)/2} (\log 2x)^{A+1} = 3x^{1/2+\delta/2} (\log 2x)^{A+1} \leq x^{1/2+\varepsilon}$$

for  $x \geq x_1(\varepsilon)$ , because  $\delta/2 < \varepsilon$  and  $x^{\varepsilon-\delta/2}$  absorbs the constant and the logarithm.  $\square$

**Theorem 2.2** (square-root cancellation). For every  $\varepsilon > 0$  there is  $C_{\varepsilon}$  with  $|R(x)| \leq C_{\varepsilon} x^{1/2+\varepsilon}$  for all  $x \geq 1$ .

*Proof.* We may take  $\varepsilon < \frac{1}{4}$ . Let  $x \geq 2x_1$  and let  $K$  be the integer with  $x_1 \leq x/2^K < 2x_1$ . Then

$$R(x) = \sum_{k=0}^{K-1} \left[ \psi\left(\frac{x}{2^k}\right) - \psi\left(\frac{x}{2^{k+1}}\right) - \frac{x}{2^{k+1}} \right] + R\left(\frac{x}{2^K}\right),$$

and by Lemma 2.1 applied at  $x/2^{k+1} \geq x_1$  each bracket is at most  $(x/2^{k+1})^{1/2+\varepsilon}$  in absolute value. Hence

$$|R(x)| \leq x^{1/2+\varepsilon} \sum_{k \geq 0} 2^{-(k+1)(1/2+\varepsilon)} + \max_{t \leq 2x_1} |R(t)| \leq 3x^{1/2+\varepsilon} + C,$$

and for  $x < 2x_1$  the bound holds with the constant  $\max_{t \leq 2x_1} |R(t)|$ .  $\square$

### 3 The Mellin lemma

**Lemma 3.1.** *The function  $-\frac{\zeta'}{\zeta}(s) - \frac{s}{s-1}$ , defined for  $\operatorname{Re} s > 1$ , is the restriction of a function holomorphic on the half-plane  $\operatorname{Re} s > \frac{1}{2}$ .*

*Proof.* Fix  $s$  with  $\operatorname{Re} s > 1$ . By Axiom 1,  $-\zeta'/\zeta(s) = \sum_n \Lambda(n)n^{-s}$ . Writing the sum as a Stieltjes integral against the step function  $\psi$  and integrating by parts on  $[1, X]$ ,

$$\sum_{n \leq X} \Lambda(n)n^{-s} = \int_{1-}^X x^{-s} d\psi(x) = \psi(X)X^{-s} + s \int_1^X \psi(x)x^{-s-1} dx.$$

By Axiom 2,  $\psi(X)X^{-s} \rightarrow 0$  as  $X \rightarrow \infty$ , and  $\int_1^\infty \psi(x)x^{-\operatorname{Re} s-1} dx < \infty$ ; so

$$-\frac{\zeta'}{\zeta}(s) = s \int_1^\infty \psi(x)x^{-s-1} dx = s \int_1^\infty x^{-s} dx + s \int_1^\infty R(x)x^{-s-1} dx = \frac{s}{s-1} + s \int_1^\infty R(x)x^{-s-1} dx.$$

Define  $\Phi(s) = s \int_1^\infty R(x)x^{-s-1} dx$  for  $\operatorname{Re} s > \frac{1}{2}$ . By Theorem 2.2, for  $\operatorname{Re} s \geq \frac{1}{2} + 2\varepsilon$  the integrand is bounded by  $C_\varepsilon x^{-1-\varepsilon}$ , which is integrable; the integral therefore converges absolutely and uniformly on every such closed half-plane, the integrand is holomorphic in  $s$  for each  $x$ , and  $\Phi$  is holomorphic on  $\operatorname{Re} s > \frac{1}{2}$  (differentiation under the integral, or Morera). On  $\operatorname{Re} s > 1$ ,  $\Phi(s) = -\zeta'/\zeta(s) - s/(s-1)$ .  $\square$

### 4 The Line

**Theorem 4.1** (the Line Postulate). *Every quantum is massless: every zero of  $\xi$  has real part  $\frac{1}{2}$ .*

*Proof.* By Axiom 3,  $\zeta$  is meromorphic on the plane with a single simple pole at  $s = 1$ . Therefore  $\zeta'/\zeta$  is meromorphic on the plane, and its poles are exactly the zeros of  $\zeta$  (a zero of order  $k$  at  $\rho$  is a simple pole of  $\zeta'/\zeta$  with residue  $k$ ) together with  $s = 1$  (a simple pole of  $-\zeta'/\zeta$  with residue 1). By Lemma 3.1, on  $\operatorname{Re} s > \frac{1}{2}$ ,

$$-\frac{\zeta'}{\zeta}(s) = \frac{s}{s-1} + \Phi(s)$$

with  $\Phi$  holomorphic: the only pole of  $\zeta'/\zeta$  on  $\operatorname{Re} s > \frac{1}{2}$  is  $s = 1$ , where  $s/(s-1) = 1 + 1/(s-1)$  accounts for the residue. Hence  $\zeta$  has no zero with  $\operatorname{Re} s > \frac{1}{2}$ . By Axiom 3, the zeros of  $\xi$  in the strip are the zeros of  $\zeta$  there, so  $\xi$  has no zero with  $\operatorname{Re} s > \frac{1}{2}$ . If  $\xi(\rho) = 0$  with  $\operatorname{Re} \rho < \frac{1}{2}$ , then  $\xi(1-\rho) = \xi(\rho) = 0$  with  $\operatorname{Re}(1-\rho) > \frac{1}{2}$ , which is impossible. Every zero of  $\xi$  therefore has  $\operatorname{Re} \rho = \frac{1}{2}$ .  $\square$

### 5 Observers

**Theorem 5.1** (the Born rule). *For every field  $f$  of every age,  $Q(f) = \sum_\rho |F(\gamma_\rho)|^2 \geq 0$ , the sum over the quanta with multiplicity.*

*Proof.* By Theorem 4.1 every quantum has  $m_\rho = 0$ , so each term of Axiom 5 is  $S(\gamma_\rho)^2 = |F(\gamma_\rho)|^2$ .  $\square$

**Lemma 5.2** (sampling). *Let  $F \not\equiv 0$  be entire with  $|F(z)| \leq Me^{a|\operatorname{Im} z|}$  for all  $z$ , and let  $n(r)$  be the number of its zeros in  $|z| \leq r$ . Then  $\liminf_{r \rightarrow \infty} n(r)/r \leq 2a/\pi \cdot (1 + o(1))$ ; more precisely  $\int_0^R n(r) dr/r \leq \frac{2a}{\pi} R + O(1)$ . Consequently  $F$  cannot vanish at every real point of a set whose counting function  $N(T)$  in  $[-T, T]$  satisfies  $N(T) \geq (2a/\pi + \eta)T$  for all large  $T$  and some  $\eta > 0$ .*

*Proof.* Choose  $z_0$  with  $F(z_0) \neq 0$ ; replacing  $F(z)$  by  $F(z+z_0)$  changes neither the growth bound (up to the constant) nor the asymptotics of  $n(r)$ , so assume  $F(0) \neq 0$ . Jensen's formula gives, for every  $R$ ,

$$\int_0^R \frac{n(r)}{r} dr = \frac{1}{2\pi} \int_0^{2\pi} \log |F(Re^{i\theta})| d\theta - \log |F(0)| \leq \frac{1}{2\pi} \int_0^{2\pi} (\log M + aR |\sin \theta|) d\theta - \log |F(0)| = \frac{2aR}{\pi} + O(1).$$

If  $F$  vanished on the set described, then  $n(r) \geq N(r) \geq (2a/\pi + \eta)r$  for  $r \geq r_0$ , so  $\int_0^R n(r) dr/r \geq (2a/\pi + \eta)(R - r_0)$ , which exceeds  $2aR/\pi + O(1)$  for  $R$  large: a contradiction.  $\square$

**Theorem 5.3** (Immortality). *Every observer is alive:  $\lambda(a) > 0$  for every age  $a$ .*

*Proof.* By Theorem 5.1,  $\lambda(a) = Q(f_a) \geq 0$ . Suppose  $\lambda(a) = 0$ . Then  $F_a(\gamma_\rho) = 0$  at every quantum. The transform of a field of age  $a$  satisfies  $|F_a(z)| \leq \|f_a\|_1 e^{a|\operatorname{Im} z|} \leq \sqrt{2a} e^{a|\operatorname{Im} z|}$ . The quanta are real by Theorem 4.1, and by Axiom 3 together with the argument principle applied to  $\xi$  (the Academy's counting theorem) their number in  $[-T, T]$  is  $N(T) = \frac{T}{\pi} \log \frac{T}{2\pi e} + O(\log T)$ , which exceeds  $(2a/\pi + 1)T$  for  $T$  large. Lemma 5.2 forces  $F_a \equiv 0$ , hence  $f_a = 0$ , contradicting  $\|f_a\|_2 = 1$ .  $\square$

**Theorem 5.4** (Proportional Leakage). *For every  $a_1 < a_2$  the function  $C = -\lambda'/\lambda$  is integrable on  $[a_1, a_2]$ , nonnegative, and  $2|A_0(a)|^2 = C(a)\lambda(a)$  for almost every  $a$ . The renormalized floor  $U(a) = \lambda(a) \exp(\int_{a_Z}^a C)$  is constant, and*

$$\lambda(a) = \lambda(a_Z) \exp\left(-\int_{a_Z}^a C\right) > 0 \quad (a \geq a_Z).$$

*Proof.* By Theorem 5.3 and Axiom 7,  $\lambda$  is positive and continuous, hence bounded below by a positive number on  $[a_1, a_2]$ , and  $\lambda'$  is integrable there; so  $C = -\lambda'/\lambda$  is integrable, and it is nonnegative because  $\lambda' = -2|A_0|^2 \leq 0$  by Axiom 8, which also gives the identity.  $\log \lambda$  is absolutely continuous with derivative  $\lambda'/\lambda$ , so  $\lambda(a) = \lambda(a_Z) \exp(\int_{a_Z}^a \lambda'/\lambda)$ , which is the formula, and  $U \equiv \lambda(a_Z)$ .  $\square$

**Corollary 5.5** (the Cooling Theorem). *The floor is positive at every age and non-increasing; its logarithm is the integral of the leak rate,  $\log \lambda(a) = \log \lambda(a_Z) - \int_{a_Z}^a 2|A_0|^2/\lambda$ , and no finite age exhausts it.*

## 6 Commentary: the Academy's account of Axiom 4

The proof above uses Axiom 4 once, in Lemma 2.1. The Academy regards the axiom itself as the shadow of two deeper facts about the charges, recorded here for understanding and not used in the proof.

The first is *Sieve Independence*: for an admissible set  $\mathcal{H} = \{h_1, \dots, h_k\}$  of integers,  $\sum_{n \leq x} \prod_i \Lambda(n+h_i) = \mathfrak{S}(\mathcal{H})x + O(x^{1/2+\varepsilon})$  uniformly in sets of bounded size and diameter at most  $x$ , with  $\mathfrak{S}$  the singular series. By unique factorization the residues of an integer modulo distinct primes are independent (the Chinese remainder theorem), and the law says that nothing beyond these local obstructions correlates the primes: the sieve has no memory.

The second is the *Gaussian law*: under Sieve Independence, the fluctuation  $\psi(x+h) - \psi(x) - h$  over  $x \in [X, 2X]$  has the moments of a Gaussian variable of variance  $h \log(X/h)$  for  $X^\delta \leq h \leq X^{1-\delta}$  (the Academy's theorem of Montgomery and Soundararajan). Axiom 4 states that the charges never deviate from this law beyond its own scale, up to a power of the logarithm: a deviation beyond the Gaussian scale would be a conspiracy of the primes, and the primes have nothing to conspire with.

In the Academy's words: the primes are as random as unique factorization allows, and not one bit less; the Line is what that randomness looks like at the largest scale.

## 7 Dictionary

Every object and every statement of this manuscript, with what it is in our universe and its status there. “Paper” is `weil_window.pdf`; “tree” is `RH_IF_THEN_TREE.md`.

### 7.1. Objects

| Academy term                                                 | real object                                                                                                          | remark                                       |
|--------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------|----------------------------------------------|
| prime sheet, charge $q(n)$                                   | the logarithmic line; the weights $\Lambda(n)n^{-1/2}$ at $\log n$                                                   | the terms of Weil’s explicit formula         |
| zero sheet, quantum, mass                                    | the nontrivial zeros $\rho = \frac{1}{2} + m + i\gamma$ of $\zeta$ ;<br>$m =  \operatorname{Re} \rho - \frac{1}{2} $ | the Line Postulate is $m = 0$ for all $\rho$ |
| charge $\psi(x)$ , fluctuation $R(x)$                        | Chebyshev’s $\psi(x) = \sum_{n \leq x} \Lambda(n)$ ; $\psi(x) - x$                                                   |                                              |
| observer of age $a$ , field, transform                       | the window $[-a, a]$ ; an odd unit test function; its Fourier transform                                              | paper, Section 2                             |
| window form $Q(f)$ , floor $\lambda(a)$ , ground field $f_a$ | the odd Weil form on the window, its lowest value, its minimizer                                                     | paper, eq. (Q) and Section 3                 |
| edge amplitude $A_0(a)$                                      | the amplitude of the minimizer’s edge law                                                                            | paper, Theorem 9.12(ii)                      |
| horizon $T^*(a) = 2\pi e^{2a}$                               | the height where the zero density equals the Shannon density $a/\pi$                                                 | paper, Section 7                             |
| alive / dead observer                                        | $\lambda(a) > 0$ / $\lambda(a) \leq 0$                                                                               |                                              |
| renormalized floor $U$                                       | $\lambda e^{\int^C}$ ; the paper’s $u(a) = \lambda e^{cT^*/2}$ when $C = cT^*$                                       | paper, proof of Prop. prop:AimpliesRH        |

### 7.2. Axioms

| axiom                   | real statement                                                                                                                 | status in our universe                                                                                                                                                                                                                                                                                                                                                                                                                                                 |
|-------------------------|--------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 1, unique factorization | the fundamental theorem of arithmetic;<br>the Euler product and $-\zeta'/\zeta = \sum \Lambda(n)n^{-s}$ for $\text{Re } s > 1$ | theorem (Euclid; Euler)                                                                                                                                                                                                                                                                                                                                                                                                                                                |
| 2, the mean field       | the prime number theorem $\psi(x) \sim x$                                                                                      | theorem (Hadamard, de la Vallée Poussin, 1896)                                                                                                                                                                                                                                                                                                                                                                                                                         |
| 3, the mirror law       | the meromorphic continuation of $\zeta$ and the functional equation $\xi(s) = \xi(1-s)$                                        | theorem (Riemann, 1859)                                                                                                                                                                                                                                                                                                                                                                                                                                                |
| 4, Bounded Deviation    | $ \psi(x+h) - \psi(x) - h  \leq (h \log(x/h))^{1/2} (\log x)^A$ for $x^\delta \leq h \leq x^{1-\delta}$                        | <b>conjecture.</b> It is the pointwise form of the Gaussian law of primes in short intervals (Cramér's model; Montgomery–Soundararajan 2004 prove the moments under a uniform Hardy–Littlewood conjecture; no large-deviation bound of this strength is known). It implies RH by Section 2–4 and is <i>stronger</i> than RH: RH gives only $O(x^{1/2} \log^2 x)$ for the fluctuation in a short interval, and the Gaussian-scale bound is not known to follow from RH. |
| 5, conservation         | Weil's explicit formula for $g = f * \tilde{f}$                                                                                | theorem (Weil, 1952)                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
| 6, Birth                | Zhu's certified floor: $\lambda(0.8) \geq 8.206 \times 10^{-15}$ , hence $\lambda > 0$ on $(0, 0.8]$                           | theorem (paper, Theorem thm:zhu)                                                                                                                                                                                                                                                                                                                                                                                                                                       |
| 7, Continuity           | absolute continuity of $\lambda$ on $[a_Z, \infty)$                                                                            | theorem at every support off an exceptional null set (paper, Propositions 9.15, 9.17); on that set it is the finiteness condition $(H_{\text{fin}})$ , open (tree, Step 1)                                                                                                                                                                                                                                                                                             |
| 8, Decay                | the boundary law $\lambda' = -2 A_0 ^2$ a.e.                                                                                   | theorem under the paper's hypothesis $(H_\infty)$ , at almost every support (paper, Theorem 9.12(iv))                                                                                                                                                                                                                                                                                                                                                                  |

### 7.3. Lemmas and theorems

| statement                      | real content                                                                                                      | status                                                                                                                                                                                        |
|--------------------------------|-------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Lemma 2.1, Theorem 2.2         | from the short-interval bound to $\psi(x) = x + O(x^{1/2+\varepsilon})$                                           | elementary; a theorem given Axiom 4                                                                                                                                                           |
| Lemma 3.1 (Mellin)             | $-\zeta'/\zeta(s) - s/(s-1)$ holomorphic on $\operatorname{Re} s > \frac{1}{2}$ given the square-root bound       | theorem (von Koch, 1901)                                                                                                                                                                      |
| Theorem 4.1 (the Line)         | $\psi(x) = x + O(x^{1/2+\varepsilon}) \Rightarrow \text{RH}$                                                      | theorem (von Koch); with Littlewood's converse, an equivalence. So in our universe Sections 2–4 prove: <i>Bounded Deviation implies RH</i>                                                    |
| Theorem 5.1 (Born rule)        | under RH, $Q(f) = \sum  F(\gamma) ^2$                                                                             | theorem under RH (Weil positivity)                                                                                                                                                            |
| Lemma 5.2 (sampling)           | a function of exponential type $a$ , bounded by $Me^{a \operatorname{Im} z }$ , has zero density at most $2a/\pi$ | theorem (Jensen's formula; Boas)                                                                                                                                                              |
| Theorem 5.3 (Immortality)      | under RH, $\lambda(a) > 0$ for every $a$                                                                          | theorem under RH                                                                                                                                                                              |
| Theorem 5.4 (Leakage)          | under RH and Axiom 7, $-\lambda'/\lambda$ is locally integrable and $\lambda = \lambda(a_Z)e^{-\int^C}$           | theorem under RH and absolute continuity; the easy direction of paper Proposition 9.30. The converse (integrable leak $\Rightarrow$ RH) is the paper's reduction, Proposition prop:AimpliesRH |
| Corollary 5.5 (Cooling)        | the floor is positive and non-increasing, its logarithm the integral of the leak rate                             | as above                                                                                                                                                                                      |
| Commentary: Sieve Independence | the Hardy–Littlewood prime $k$ -tuple conjecture, uniform, with square-root error                                 | conjecture                                                                                                                                                                                    |
| Commentary: Gaussian law       | moments of $\psi(x+h) - \psi(x) - h$ Gaussian with variance $h \log(x/h)$                                         | theorem under a uniform $k$ -tuple conjecture (Montgomery–Soundararajan, 2004); its variance is equivalent under RH to Montgomery's pair correlation (Goldston–Montgomery, 1987)              |

## 7.4. The sentence

In the Critical Line Universe the proof is unconditional: its axioms are the laws there. In our universe the same pages prove one implication, *if the primes satisfy Bounded Deviation then every zero is on the line*, by a page of classical analysis; Bounded Deviation is an unproved conjecture about primes in short intervals, believed for sieve reasons that have nothing to do with zeros, and stronger than the Line itself. Everything else in the manuscript is a theorem, or a theorem under the Line, or a hypothesis of the paper that the paper retains.

— end of manuscript —