

# Part IA — Markov Chains Example Sheet 1

Supervised by Prof Weber (rrw1@cam.ac.uk)

Examples worked through by Christopher Turnbull

Michaelmas 2017

**QUESTION 1**

$X = (X_n)$  is a Markov Chain, and so satisfies the Markov Property

$$\mathbb{P}(X_{n+1} = i_{n+1} \mid X_0 = i_0, \dots, X_n = i_n) = \mathbb{P}(X_{n+1} = i_{n+1} \mid X_n = i_n)$$

Given that  $Z_k = X_{m+k}$ , with  $k \geq 0$ , want to show that  $Z_k$  is a Markov Chain, ie.

$$\mathbb{P}(Z_{n+1} \mid Z_0, \dots, Z_n) = \mathbb{P}(Z_{n+1} \mid Z_n) \quad (*)$$

Substituting in our definition of  $Z_k$ ,

$$\begin{aligned} \mathbb{P}(Z_{n+1} \mid Z_0, \dots, Z_n) &= \mathbb{P}(X_{m+n+1} \mid X_m, \dots, X_{m+n}) \\ &= \mathbb{P}(X_{m+n+1} \mid X_{m+n}) \quad \text{as } X \text{ is Markov} \\ &= \mathbb{P}(Z_{n+1} \mid Z_n) \end{aligned}$$

Hence  $Z_k$  satisfies the Markov property, and is thus a Markov Chain.  
 $Z_0 = X_m = i$ , so  $Z_k$  has starting state  $i$ .

**QUESTION 2**

Let  $X_1, \dots, X_n$  be independent random variables, so  $\mathbb{P}(X_{n+1} = i_{n+1} \mid X_n = i_n) = \mathbb{P}(X_{n+1} = i_{n+1})$ . But then

$$\mathbb{P}(X_{n+1} = i_{n+1} \mid X_0 = i_0, \dots, X_n = i_n) = \mathbb{P}(X_{n+1} = i_{n+1})$$

also by independence, so Markov property is satisfied.

Homogeneous if  $\mathbb{P}(X_{n+1} = i_{n+1} \mid X_n = i_n)$  does not depend on the value of  $n$ , so must have  $X_n$  identically distributed.

### QUESTION 3

As  $X_n$  is the *maximum* reading obtained after  $n$  throws, this depends only on the last maximum, so  $X_n$  is a Markov chain. In particular, the state transition matrix is

$$\begin{pmatrix} 1/6 & 1/6 & 1/6 & 1/6 & 1/6 & 1/6 \\ 0 & 2/6 & 1/6 & 1/6 & 1/6 & 1/6 \\ 0 & 0 & 3/6 & 1/6 & 1/6 & 1/6 \\ 0 & 0 & 0 & 4/6 & 1/6 & 1/6 \\ 0 & 0 & 0 & 0 & 5/6 & 1/6 \\ 0 & 0 & 0 & 0 & 0 & 6/6 \end{pmatrix}$$

Now note that

$$p_{i,j}(n) = \begin{cases} 0 & \text{if } j < i \\ \left(\frac{1}{6}\right)^n & \text{if } j = i \\ \left(\frac{j}{6}\right)^n - \left(\frac{j-1}{6}\right)^n & \text{if } j > i \end{cases}$$

## QUESTION 4

We have

$$S_{n+1} = \begin{cases} S_n + 1 & \text{with probability } p \\ S_n - 1 & \text{with probability } q \end{cases}$$

and  $S_0 = 0$ .

To show that  $X_n$  is a Markov chain, consider the probability

$$\mathbb{P}(|S_{n+1}| = i + 1 \mid |S_n| = i, H)$$

for some past event  $H$ . Conditioning, this is equal to

$$\begin{aligned} &= \underbrace{\mathbb{P}(|S_{n+1}| = i + 1 \mid S_n = i)}_p \times \underbrace{\mathbb{P}(S_n = i \mid |S_n| = i, H)}_{\pi_1} \\ &\quad + \underbrace{\mathbb{P}(|S_{n+1}| = i + 1 \mid S_n = -i)}_q \times \underbrace{\mathbb{P}(S_n = -i \mid |S_n| = i, H)}_{\pi_2} \end{aligned}$$

Note that

$$\begin{aligned} \frac{\pi_1}{\pi_2} &= \frac{p^{n/2+i/2} q^{n/2-i/2}}{q^{n/2+i/2} p^{n/2-i/2}} \\ &= \frac{p^i}{q^i} \end{aligned}$$

Hence  $\pi_1 = \frac{p^i}{p^i + q^i}$

Define  $M_k := \max\{S_k \mid 0 \leq k \leq n\}$ , and  $Y_n = M_n - S_n$

---


$$\begin{array}{cc} S_n & M_n \end{array}$$

with the distance between  $S_n$  and  $M_n$  being  $i$ , we have:

$$p_{ij} = \begin{cases} i - 1 & \text{with probability } p \\ i + 1 & \text{with probability } q \end{cases}$$

$$p_{0j} = \begin{cases} 0 & \text{with probability } p \\ 1 & \text{with probability } q \end{cases}$$

## QUESTION 5

$X$  is a Markov chain, thus

$$\mathbb{P}(X_{n+1} = i_{n+1} \mid X_0 = i_0, \dots, X_n = i_n) = \mathbb{P}(X_{n+1} = i_{n+1} \mid X_n = i_n)$$

And so

$$\mathbb{P}(Y_{r+1} \mid Y_0, \dots, Y_r) = \mathbb{P}(X_{n_{r+1}} \mid X_{n_0}, \dots, X_{n_r})$$

If  $n_r = 2r$  and  $X$  is a simple random walk on  $\mathbb{Z}$  st.

$$X_{n+1} = \begin{cases} X_n + 1 & \text{with probability } p \\ X_n - 1 & \text{with probability } q \end{cases}$$

the allowed states are  $\{\dots, -2, -1, 0, 1, 2, \dots\}$ . Of course, the probability of finding oneself in an odd state is always zero (we are always taking an even number of states).

It is difficult to visualise the state transition matrix as the state space is infinite, but a general row has the form

$$(\dots \quad 0 \quad q^2 \quad 0 \quad 2pq \quad 0 \quad p^2 \quad \dots)$$

With probability  $2pq$  of returning to the current state,  $p^2$  moving two to the right and  $q^2$  moving two to the left. (Thus rows summing to  $(p+q)^2 = 1$ )

**QUESTION 6**

I think the answer should be no.  $X$  and  $Y$  are both Markov chains, so the value of  $X_{n+1} + Y_{n+1}$  only depends on  $X_n$ , and  $Y_n$ . But we only have the value of  $X_n + Y_n$  (loss of information).

For an explicit counterexample, try to smuggle information about  $X_1$  into  $Y$ , so that  $X_n + Y_n$  depends on  $X_1$  (contradicting the Markov property) but  $X_n$  and  $Y_n$  are both Markov.

Idea: Let  $X$  be a random walk on the integers, so  $X_1 = \pm 1$  with some probabilities. Define  $Y_n$  as a symmetric ‘boosted’ random walk:

$$Y_n = \begin{cases} Y_{n-1} + 1 + 3X_1 & \text{with probability } 1/2 \\ Y_{n-1} - 1 + 3X_1 & \text{with probability } 1/2 \end{cases}$$

Then  $X_n + Y_n$  depends on  $X_1$ , not just  $X_{n-1}, Y_{n-1}$

Note that we can also make use of  $M_n$  as defined in question 4.

$$\mathbb{P}(M_3 = 2 \mid M_2 = 1, M_1 = 1) = 0$$

$$\mathbb{P}(M_3 = 2 \mid M_2 = 1, M_1 = 0) = \frac{1}{2}$$

So  $M_n$  not M.C.

## QUESTION 7

- (i) Let the probability of the flea being on the original vertex after  $n$  hops be  $p_n$ . Then

$$p_n = \begin{cases} 0 & \text{if flea on first vertex after } n-1 \text{ hops} \\ \frac{1}{2} & \text{otherwise} \end{cases}$$

Hence we set up the recurrence, with  $p_0 = 1$

$$\begin{aligned} p_n &= 0 \cdot p_{n-1} + \frac{1}{2}(1 - p_{n-1}) \\ &= \frac{1}{2}(1 - p_{n-1}) \end{aligned}$$

Solving this difference equation gives general solution  $p_n = \frac{1}{3} + \frac{2}{3} \left(-\frac{1}{2}\right)^n$ .

- (ii) Let  $p_n$  be defined as before. This can be thought of as a Markov chain on the state space of triangle vertices  $\{1, 2, 3\}$ , with transition matrix

$$P = \begin{pmatrix} 0 & \frac{2}{3} & \frac{1}{3} \\ \frac{1}{3} & 0 & \frac{2}{3} \\ \frac{2}{3} & \frac{1}{3} & 0 \end{pmatrix}$$

Given  $p_{1,1}(0) = 1$  (the flea starts on vertex 1) we want to find  $p_{1,1}(n)$ . The eigenvalues  $\kappa$  of  $P$  are the roots of the equation  $\det(P - \kappa I)$ , with solutions

$$\kappa_1 = 1, \quad \kappa_2, \kappa_3 = -\frac{1}{2} \pm \frac{i}{2\sqrt{3}}$$

Therefore

$$P = U^{-1} \begin{pmatrix} 1 & 0 & 0 \\ 0 & \kappa_2 & 0 \\ 0 & 0 & \kappa_3 \end{pmatrix} U$$

for some invertible matrix  $U$ . It follows that

$$P^n = U^{-1} \begin{pmatrix} 1 & 0 & 0 \\ 0 & \kappa_2^n & 0 \\ 0 & 0 & \kappa_3^n \end{pmatrix} U$$

Now note that  $\kappa_2, \kappa_3 = \frac{1}{\sqrt{3}} e^{\pm i\pi/6}$ , so

$$\begin{aligned} p_{1,1}(n) &= A + B e^{i\pi n/6} + C e^{-i\pi n/6} \\ &= A + B' \left(3^{-1/2}\right)^n \cos(\pi n/6) + C' \left(3^{-1/2}\right)^n \sin(\pi n/6) \end{aligned}$$



Using the b.c.s,

$p_{1,1}(0) = 1, p_{1,1}(1) = 0, p_{1,1}(2) = 4/9$  and hence

$$p_{1,1}(n) = \frac{1}{3} + \frac{2}{3} \left(3^{-1/2}\right)^n \cos(\pi n/6)$$

**QUESTION 8**

Note that

$$p_{66}(n-1) = \frac{1}{5}(1 - p_{66}(n-2))$$

Hence,

$$p = \frac{1}{6} + \frac{5}{6} \left(-\frac{1}{2}\right)^{n-1}$$

For the next part, let  $Y_n$  denote the die now, and this is related to the previous die by

$$Y_n - Y_{n-1} = X_n - X_{n-1} \pmod{6}$$

Thus

$$Y_n = X_n + (n-1) \pmod{6}$$

Thus  $n-1 = 0 \pmod{6} \Rightarrow p_{66}(n-1) = p$ , and  $n-1 \neq 0 \pmod{6} \Rightarrow \frac{1}{5}(1-p)$

**QUESTION 9**

- (a) For  $p = 1/16$ , the characteristic equation  $\det(P - \kappa I) = 0$  gives roots  $\kappa_1 = 1, \kappa_2 = -1/4, \kappa_3 = -1/12$ .

Thus

$$P^n = U^{-1} \begin{pmatrix} 1 & 0 & 0 \\ 0 & (-1/4)^n & 0 \\ 0 & 0 & (-1/12)^n \end{pmatrix} U$$

for some invertible matrix  $U$ , and so

$$p_{1,1}(n) = A + B \left(-\frac{1}{4}\right)^n + C \left(-\frac{1}{12}\right)^n$$

Using  $p_{1,1}(0) = 1, p_{1,1}(1) = 0, p_{1,1}(2) = 0$

This gives  $A = 1/65, B = -2/5$  and  $C = 18/13$

- (b) (Check, this is incorrect) For  $p = 1/6$ , the characteristic equation  $\det(P - \kappa I) = 0$  gives roots  $\kappa_1 = 1, \kappa_2, \kappa_3 = -1/6 \pm i/6 = \frac{1}{\sqrt{18}} e^{\pm i\pi/4}$ , so similarly,

$$p_{1,1}(n) = A + \left(18^{-1/2}\right)^n B \cos(\pi n/4) + \left(18^{-1/2}\right)^n C \sin(\pi n/4)$$

yielding  $A = 1/13, B = 12/13, C = -18/13$

- (c)

$$P^n = U^{-1} \begin{pmatrix} 1 & 0 & 0 \\ 0 & (-1/6)^n & 0 \\ 0 & 0 & (-1/6)^n \end{pmatrix} U$$


Hence,  $p_{1,1}(n) = A + B \left(-\frac{1}{6}\right)^n + Cn \left(-\frac{1}{6}\right)^n$

Check, should get

$$\frac{1}{49} + \left(-\frac{1}{6}\right)^n \left(\frac{48}{49} - \frac{6}{7}n\right)$$

**QUESTION 10**

Possible transitions of the chain are illustrated below:



The communicating classes are  $C_1 = \{1, 5\}$ ,  $C_2 = \{3\}$  and  $C_3 = \{2, 4\}$ . The classes  $C_1$  and  $C_3$  are not closed, but  $C_2$  is closed.

**QUESTION 11**

Suppose  $S$  contains no closed communicating classes.

Pick some  $i_0 \in S$ . If  $i$  is a singleton communicating class, then  $\{i_0\}$  is closed. So  $\exists i_1 \in S$  st.  $i_0 \rightarrow i_1$ . Similarly,  $\exists i_2 \in S$  st.  $i_1 \rightarrow i_2$ . ( $i_2 \neq i_0$ , or  $\{i_0, i_1\}$  is a closed communicating class). Continuing in this fashion, we eventually hit the ‘last’ element in the state space  $i_n$ , as it is finite. Then  $S = \{i_0, \dots, i_n\}$  itself is a closed communicating class.

For a transition matrix with no communicating classes a simple example is:

$$\begin{pmatrix} 0 & 1 & 0 & 0 & \cdots \\ 0 & 0 & 1 & 0 & \cdots \\ 0 & 0 & 0 & 1 & \ddots \\ \vdots & \vdots & & & \ddots \end{pmatrix}$$

ie.  $i_0 \rightarrow i_1 \rightarrow i_2 \rightarrow \cdots$

**QUESTION 12**

See Weber's Right Hand Equations

$$\alpha = P\alpha, k = 1 + Pk$$

$$\begin{aligned}\alpha_2 &= \frac{1}{2} \cdot 0 + \frac{1}{2} \alpha_4 \\ \alpha_4 &= \frac{1}{2} \cdot 0 + \frac{1}{2} \alpha_8 \\ \alpha_6 &= \frac{1}{2} \alpha_6 + \frac{1}{2} 1 \\ \alpha_8 &= \frac{1}{2} \alpha_2 + \frac{1}{2} 1 \\ \alpha_2 &= \frac{1}{2} \left( \frac{1}{2} \left( \frac{1}{2} + \frac{1}{2} \left( \frac{1}{2} + \frac{1}{2} \alpha_2 \right) \right) \right) \\ &= \frac{1}{8} + \frac{1}{16} + \frac{1}{16} \alpha_2\end{aligned}$$

Hence

$$\frac{15}{16} \alpha_2 = \frac{3}{16} \quad \alpha_2 = \frac{1}{5}$$

Could also argue it has to be  $1/5$  since it's a fair game, and the placement of 2 between 0 and 10.

Similarly,

$$\begin{aligned}k_2 &= 1 + \frac{1}{2} \cdot 0 + \frac{1}{2} k_4 \\ k_4 &= 1 + \frac{1}{2} \cdot 0 + \frac{1}{2} k_8\end{aligned}$$

But since  $k_6 = k_4$ , and  $k_8 = k_2$ , we have

$$k_2 = 1 + \frac{1}{2} \left( 1 + \frac{1}{2} k_2 \right)$$

and we recover  $k_2 = 2$ , as expected.

**QUESTION 13**

$$\alpha_0 = 1$$

$$\alpha_i = \frac{(i+1)^2}{i^2 + (i+1)^2} \alpha_{i+1} + \frac{i^2}{i^2 + (i+1)^2} \alpha_{i-1}$$

This is equivalent to

$$(i+1)^2(\alpha_{i+1} - \alpha_i) = i^2(\alpha_i - \alpha_{i-1})$$

Iterating down,

$$\dots = 1^2(\alpha_1 - \alpha_0) = \alpha_1 - 1$$

ie.

$$\alpha_{i+1} - \alpha_i = \frac{1}{(i+1)^2}(\alpha_1 - 1)$$

Summing,

$$\alpha_{k+1} - \alpha_0 = \left[ \frac{1}{1^2} + \dots + \frac{1}{(k+1)^2} \right] (\alpha_1 - 1)$$

The result follows, taking  $k \rightarrow \infty$ .

## QUESTION 14

Again, expand on the right side,

$$\begin{aligned}\phi(s) &= \mathbb{E}s^{H_0} \\ &= \frac{1}{2}s^1 + \frac{1}{2}s^1 \underbrace{\mathbb{E}s^{H'_0 + H''_0}}_{=\phi(s)^2}\end{aligned}$$

where  $H'_0, H''_0$  are independent variables.

$$\phi(s) = \frac{1 \pm \sqrt{1 - s^2}}{s}$$

where we pick the  $-$  from the  $\pm$ , as terms in the expansion need to be positive.

Thus

$$\phi(s) = \frac{s}{2} + \frac{s^3}{8} + \frac{s^5}{16} + O(s^6)$$

Next we have the same idea,

$$\phi(s) = \frac{1}{2}s^1 + \frac{1}{2}s^1\phi(s)^3$$

Can use mathematica to solve this. The smallest root of this equation, interestingly, is the golden mean!