

Oral Care

Quick Tips – Care Guide



From City of Hope Cancer Center, Dr. Visha Ahuja

Cancer treatments, such as chemotherapy and radiation, target rapidly dividing cells, including the healthy cells lining the mouth. This collateral damage alters the oral environment, causing tissue to become tender, inflamed, and prone to injury. Standard toothpastes are often too aggressive for this sensitized tissue. Selecting a specialized oral hygiene product is necessary to maintain comfort, prevent severe complications, and support the overall success of cancer therapy.

Irritating Ingredients to Avoid

The first step in selecting a gentle toothpaste is eliminating common additives that can exacerbate existing oral irritation. Sodium Lauryl Sulfate (SLS) is a problematic ingredient; this detergent and foaming agent can attack the protective mucosal membrane. SLS causes mucosal sloughing, stripping away the superficial layer of soft tissue, which significantly worsens the pain and recurrence of mouth sores.

Strong flavorings, particularly sharp mint oils and cinnamon, should also be avoided because they can produce a painful stinging or burning sensation on inflamed tissues. These intense flavors may also trigger or enhance nausea, a common side effect of chemotherapy. Patients should look for mild flavors, such as peach or bubblegum, or opt for an unflavored product.

Many commercial toothpastes contain harsh agents that are too aggressive for a sensitive mouth. Whitening agents (often containing peroxides) and tartar control ingredients (such as pyrophosphates) can irritate sore oral tissues. Additionally, any toothpaste or rinse containing alcohol should be avoided, as alcohol has a drying effect that intensifies xerostomia and increases discomfort.

Essential Ingredients for Comfort and Protection

The most beneficial toothpastes for cancer patients are those specifically formulated to be gentle, moisturizing, and protective. Glycerin and moisturizing agents like betaine and olive oil help lubricate and soothe the dry, tender tissues. These ingredients provide a protective coating over the mucosal lining, offering relief from the sticky feeling of xerostomia.

Xylitol, a natural sugar alcohol, is particularly helpful for patients experiencing dry mouth because it stimulates the salivary glands to produce more saliva. Xylitol cannot be metabolized by the harmful bacteria that cause tooth decay, reducing the elevated risk of cavities associated with dry mouth. A toothpaste with a neutral pH (typically between 6.5 and 7.2) is also important to prevent the product from burning or stinging sensitive tissues.

Gentle abrasives are necessary to clean the teeth without causing friction injuries to the gums or enamel. Specialized toothpastes often use very low-abrasivity silica or other mild polishing agents to ensure effective cleaning. Low-Relative Dentin Abrasivity (RDA) formulations are preferred to minimize mechanical trauma during brushing.

Fluoride remains a significant component in oral care for most cancer patients due to the high risk of rapid tooth decay associated with dry mouth. For those at high risk of caries, a prescription-strength, high-fluoride toothpaste may be recommended by a dentist or oncologist for maximum remineralization and protection. These high-concentration formulas are generally gentle and often SLS-free, designed to protect enamel without irritating sensitive mouth tissues.

Choosing and Using Specialized Toothpaste

Selecting the right product should always begin with a consultation with the oncology care team or dental oncologist, who provides personalized recommendations based on the treatment plan and oral side effects. Patients should look for products labeled as “gentle,” “sensitive,” or designed for “dry mouth” care. They must confirm the absence of irritating ingredients like SLS and strong flavors. Many patients find a smooth, creamy gel more comfortable than a thick paste.

The method of application must be adapted to the mouth's sensitive condition. Patients should use an extra-soft toothbrush with small, round-headed bristles to minimize trauma to the gums and cheek lining. Brushing should be done with minimal pressure, using small, gentle circular strokes to clean the teeth effectively without causing injury.

In cases of severe oral mucositis flare-ups, when brushing with any toothpaste is excruciating, the patient may need to temporarily switch to alternative methods. This may involve using a simple, lukewarm rinse made of salt water or a baking soda solution to gently cleanse the mouth and provide comfort. The goal is to maintain cleanliness to prevent infection, while prioritizing comfort and avoiding further injury to the fragile oral tissues.

Resources

[Cancer Therapies and Dental Considerations](#)

[Guidance on Toothpaste Choices for Chemotherapy Patients](#)

[Cancer Treatment and Your Teeth](#)



Disclaimer: The content is provided for general informational purposes only and is not intended as, nor should it be considered a substitute for, professional medical advice. Do not use the information on this website for diagnosing or treating any medical or health condition. If you have or suspect you have a medical problem, promptly contact your professional healthcare provider.



Sarcoma & LMS Pathways

Care Guide - Quick Tips for Patients & Families

Trusted guidance for navigating diagnosis, treatment, and leiomyosarcoma

www.NLMSF.org