3.1 The Chain Rule

Name _____ Period ____ Date _____

AP Calculus 3.1 (30-minute plan) · Student sheet

I can…

- identify the outer and inner functions of a composite function.
- use the Chain Rule, including the Power Rule for functions.
- use the Chain Rule with e^u, trig functions, and tables of values.

Vocabulary · fill in the missing word

Composite function — $(f \circ g)(x) = f(g(x))$: the _____ of g is the input of f

Chain Rule — $\frac{d}{dx} f(g(x)) =$ _____

Power Rule for functions — $\frac{d}{dx} [u(x)]^n =$ _____

Part 1 · Notes and examples (fill in each note, then work the example)

Chain Rule: if $y = f(u)$ and $u = g(x)$, then $dy/dx = (dy/du) \cdot$ _____.

Example 1

$y = u^5$ and $u = x^2 + 1$. Find $\frac{dy}{dx}$.

Chain Rule (function form): $d/dx\ f(g(x)) = f'(g(x)) \cdot$ _____.

Example 2

$f(2) = 5$, $f'(2) = 3$, $g(1) = 2$, $g'(1) = -4$, $f'(5) = 6$. $h(x) = f(g(x))$. Find $h'(1)$.

Power Rule for functions: $d/dx\ [u(x)]^n = n[u(x)]^{n-1} \cdot$ _____.

Exponential and trig: $d/dx\ e^u = e^u \cdot u'$, $d/dx \sin u =$ _____ $\cdot u'$.

Example 3

Find $f'(x)$ for $f(x) = (3x^2 - 5)^4$, then find $f'(1)$.

Part 2 · Partner practice

Partner practice

(AP® Practice) If $f(x) = (x^2 - 3)^4$, then $f'(2) =$

(A) 4 (B) 16 (C) 32 (D) 64