4.6 Analyzing Lines of Fit

Name _____ Period ____ Date _____

Algebra I 4.6 · Student sheet

I can…

- use residuals to tell whether a line fits data well.
- use technology to find a line of best fit and the correlation coefficient.
- tell the difference between correlation and causation.

Part 1 · Check the fit

Line: y = 2.3x + 1.9

x	1	2	3	4	5
Actual y	4	7	8	12	13
Predicted y					
Residual					

Residual plot:

Good fit? _____ Desmos line of best fit: _____ r = _____

Part 2 · Notes: fill in the blanks

Residual = _____ − _____.

A good fit has residuals scattered _____ above and below 0.

r near 1: _____; r near −1: _____; r near 0: _____.

Part 3 · Examples (work them with the class)

Example 1

Use y = 2.3x + 1.9 and the data x = 1, 2, 3, 4, 5; y = 4, 7, 8, 12, 13. Find the residuals.

Example 2

The residuals from Example 1 are −0.2, 0.5, −0.8, 0.9, and −0.4. Is the line a good fit?

Example 3

Desmos gives r ≈ 0.98 for the data. Describe the correlation.

Example 4

Cities with more ice cream sales have more sunburns. Does eating ice cream cause sunburns?

Part 4 · Partner practice

Partner 1

A model predicts 50, and the actual value is 54. Find the residual.

Partner 2

Which r shows the strongest relationship: 0.45, −0.91, or 0.80?