# 5.4 Solving Radical Equations and Inequalities

Name _____ Period ____ Date _____

Algebra II 5.4 · Student sheet

**I can…**

- solve equations that contain radicals or rational exponents.
- identify extraneous solutions.
- solve radical inequalities.

## Part 1 · Find the impostor

Solve √(x + 2) = x by squaring both sides. Show your work.

| Answer | Check it in √(x + 2) = x | Does it work? |
|---|---|---|
| x = | | |
| x = | | |

Graph y = √(x + 2) and y = x in Desmos. How many times do they cross? _____

Why did one answer fail?

---

## Part 2 · Notes: fill in the blanks

Step 1: _____ the radical. Step 2: raise both sides to the same _____ as the index. Step 3: solve. Step 4:

_____ every answer in the _____ equation.

An **extraneous solution** is an answer that _____.

To solve x^(m/n) = b, raise both sides to the _____ power.

For an inequality with an even root, the radicand must also be _____.

## Part 3 · Examples (work them with the class)

### Example 1

Solve $2\sqrt{x} + 5 = 11$.

**Example 2**

Solve $\sqrt[3]{(2x - 9)} - 1 = 2$.

**Example 3**

Solve $x + 1 = \sqrt{7x + 15}$.

**Example 4**

Solve $\sqrt{2x + 3} = \sqrt{x + 7}$.

**Example 5**

Solve $x^{(\frac{3}{2})} = 8$.

**Example 6**

Solve $3\sqrt{x - 1} \leq 12$.

## Part 4 · Partner practice

### Partner 1

Solve $\sqrt{4x - 3} + 2 = 7$.

### Partner 2

Solve $x - 6 = \sqrt{3x}$.