

Insomnia Disorder

Code: 307.42 (F51.01)

Insomnia Disorder
307.42 (F51.01)
Insomnia Disorder
Diagnostic Criteria 307.42 (F51.01)
A. A predominant complaint of dissatisfaction with sleep quantity or quality, associated with one (or more) of the following symptoms:
1. Difficulty initiating sleep. (In children, this may manifest as difficulty initiating sleep without caregiver intervention.)
2. Difficulty maintaining sleep, characterized by frequent awakenings or problems returning to sleep after awakenings. (In children, this may manifest as difficulty returning to sleep without caregiver intervention.)
3. Early-morning awakening with inability to return to sleep.
B. The sleep disturbance causes clinically significant distress or impairment in social, occupational, educational, academic, behavioral, or other important areas of functioning.
C. The sleep difficulty occurs at least 3 nights per week.
D. The sleep difficulty is present for at least 3 months.
E. The sleep difficulty occurs despite adequate opportunity for sleep.
F. The insomnia is not better explained by and does not occur exclusively during the course of another sleep-wake disorder (e.g., narcolepsy, a breathing-related sleep disorder, a circadian rhythm sleep-wake disorder, a parasomnia).
G. The insomnia is not attributable to the physiological effects of a substance (e.g., a drug of abuse, a medication).
H. Coexisting mental disorders and medical conditions do not adequately explain the predominant complaint of insomnia.
Specify if:
With non-sleep disorder mental comorbidity, including substance use disorders
With other medical comorbidity
With other sleep disorder
Coding note: The code 307.42 (F51.01) applies to all three specifiers. Code also the relevant associated mental disorder, medical condition, or other sleep disorder immediately after the code for insomnia disorder in order to indicate the association.
Insomnia Disorder 365

Specify if:
Epidemic: Symptoms last at least 1 month but less than 3 months.
Persistent: Symptoms last 3 months or longer.
Recurrent: Two (or more) episodes within the space of 1 year.
Note: Acute and short-term insomnia (i.e., symptoms lasting less than 3 months but otherwise meeting all criteria with regard to frequency, intensity, distress, and/or impairment) should be coded as an other specified insomnia disorder.
Note: The diagnosis of insomnia disorder is given whether it occurs as an independent condition or is comorbid with another mental disorder (e.g., major depressive disorder), medical condition (e.g., pain), or another sleep disorder (e.g., a breathing-related sleep disorder). For instance, insomnia may develop in one course with some anxiety and depressive features but in the absence of criteria being met for any one mental disorder. Insomnia may also manifest as a clinical feature of a more predominant mental disorder. Persistent insomnia may even be a risk factor for depression and is a common residual symptom of treatment for this condition. With comorbid insomnia and a mental disorder, treatment may also need to target both conditions. Given these different courses, it is often impossible to establish the precise nature of the relationship between these clinical entities, and this relationship may change over time. Therefore, in the presence of insomnia and a comorbid disorder, it is not necessary to make a causal attribution between the two conditions. Rather, the diagnosis of insomnia disorder is made with concurrent specification of the clinically comorbid conditions. A concurrent insomnia diagnosis should only be considered when the insomnia is sufficiently severe to warrant independent clinical attention, phenotypic, or separate diagnosis is necessary.

Diagnostic Features
The essential feature of insomnia disorder is dissatisfaction with sleep quantity or quality with complaints of difficulty initiating or maintaining sleep. The sleep complaints are often accompanied by clinically significant distress or impairment in social, occupational, or other important areas of functioning. The sleep disturbance may occur during the course of any other mental disorder or medical condition, or it may occur independently.
Other manifestations of insomnia include one or different lines of the sleep period. Sleep-onset insomnia (or initial insomnia) involves difficulty initiating sleep at bedtime. Sleep-maintenance insomnia (or middle insomnia) involves frequent or prolonged awakenings throughout the night. Late insomnia involves early-morning awakening with an inability to return to sleep. The specific type of sleep complaint often varies over time. Individuals may complain of difficulty falling asleep, while a combination of these symptoms is the most common presentation overall. The specific type of sleep complaint often varies over time. Individuals may complain of difficulty falling asleep at one time may later complain of difficulty maintaining sleep, and vice versa. Symptoms of difficulty falling asleep and difficulty maintaining sleep can be quantified by the individual's retrospective self-report, sleep diaries, or other methods, such as actigraphy or polysomnography, but the diagnosis of insomnia disorder is based on the individual's subjective perception of sleep or a caretaker's report.
Nonrestorative sleep, a complaint of poor sleep quality that does not have the individual rested upon awakening despite adequate duration, is a common sleep complaint usually occurring in association with difficulty initiating or maintaining sleep, or too frequently in isolation. This complaint can also be reported in association with other sleep disorders (e.g., breathing-related sleep disorder). When a complaint of nonrestorative sleep occurs in isolation (i.e., in the absence of difficulty initiating and/or maintaining sleep) but all diagnostic criteria with regard to frequency, duration, and daytime distress and impairment are otherwise met, a diagnosis of other specified insomnia disorder or unspecified insomnia disorder is made.

364 Sleep-Wake Disorders
Adequate time for the frequency and duration criteria required to make the diagnosis, additional criteria are useful to quantify insomnia severity. These quantitative criteria, while arbitrary, are provided for descriptive purposes only. For instance, difficulty falling asleep defined by a subjective sleep latency greater than 20–30 minutes, and difficulty maintaining sleep defined by a subjective time awake after sleep onset greater than 20–30 minutes. Although there is no standard definition of early-morning awakening, this symptom involves awakening at least 30 minutes before the scheduled wake time and before last sleep time reaches 6 hours. It is essential to take into account not only the final awakening time but also bedtime or the previous evening. Awakening at 6:00 A.M. does not have the same clinical significance in those who go to bed at 9:00 P.M. as in those who go to bed at 11:00 P.M. Such a symptom may also reflect an age-dependent decrease in the ability to sustain sleep or an age-dependent shift in the timing of the main sleep period.
Insomnia disorder involves daytime impairment as well as nighttime sleep difficulties. These include fatigue or, less commonly, daytime sleepiness; the latter is more common among older individuals and when insomnia is comorbid with another medical condition (e.g., chronic pain) or sleep disorder (e.g., sleep apnea). Impairment in cognitive performance may include difficulties with attention, concentration and memory, as well as performing simple manual skills. Associated mood disturbances are typically described as irritability or mood lability and less commonly as depressive or anxiety symptoms, but all individuals with nighttime sleep disturbances are distressed or have functional impairment. For example, sleep continuity is often impaired in healthy older adults who nevertheless identify themselves as good sleepers. A diagnosis of insomnia disorder should be reserved for those individuals with insomnia after requiring more effort to maintain cognitive performance.

Associated Features Supporting Diagnosis
Insomnia is often associated with physiological and cognitive arousal and conditioning factors that interfere with sleep. A preoccupation with sleep and distress due to the inability to sleep may lead to a vicious cycle: the more the individual strives to sleep, the more frustration builds and further impairs sleep. Thus, excessive attention and efforts to sleep, which override normal sleep-onset mechanisms, may contribute to the development of insomnia. Individuals with persistent insomnia may also acquire maladaptive sleep habits (e.g., spending excessive time in bed, following an erratic sleep schedule, napping) and cognitions (e.g., fear of sleeplessness, apprehensions of daytime impairments, clock monitoring during the course of the disorder). Engaging in such activities in an environment in which the individual has frequently spent sleepless nights may further compound the conditioned arousal and perpetuate sleep difficulties. Conversely, the individual may fall asleep more easily when not trying to do so. Some individuals also report better sleep when away from their own bedrooms and their usual routines.
Insomnia may be accompanied by a variety of daytime complaints and symptoms, including fatigue, decreased energy, and mood disturbances. Symptoms of anxiety or depression that do not meet criteria for a specific mental disorder may be present, as well as excessive focus on the perceived effects of sleep or self-report psychological or personality inventories with profiles indicating mild depression and anxiety, a worrisome cognitive style, or emotion focused and internalizing style of conflict resolution, and a somatic focus. Patterns of neurocognitive impairment among individuals with insomnia disorder are inconsistent, although there may be impairments in performance tasks of higher complexity and those requiring frequent changes in performance strategy. Individuals with insomnia often require more effort to maintain cognitive performance.

Prevalence
Population-based estimates indicate that about one-third of adults report insomnia symptoms, 10%–15% experience associated daytime impairments, and 6%–10% have symptoms of insomnia disorder. 365
that meet criteria for insomnia disorder. Insomnia disorder is the most prevalent of all sleep disorders. In primary care settings, approximately 10%–20% of individuals complain of significant insomnia symptoms. Insomnia is a more prevalent complaint among females than among males, with gender odds of about 1.4:1. Although insomnia can be a symptom or an independent disorder, it is most frequently observed as a comorbid condition with another medical condition or mental disorder. For instance, 40%–50% of individuals with insomnia also present with a comorbid mental disorder.

Development and Course
The onset of insomnia symptoms can occur at any time during life, but the first episode is more common in young adulthood. Less frequently, insomnia begins in childhood or adolescence. In women, non-onset insomnia may occur during menopause and persist even after other symptoms (e.g., hot flashes) have resolved. Insomnia may have a late-life onset, which is often associated with the onset of other health-related conditions.
Insomnia can be situational, persistent, or recurrent. Situational or acute insomnia usually lasts a few days or a few weeks and is often associated with life events or rapid changes in sleep schedules or environment. It usually resolves once the initial precipitating event subsides. For some individuals, perhaps those more vulnerable to sleep disturbances, insomnia may persist long after the initial triggering event, possibly because of conditioning factors and heightened arousal. The factors that precipitate insomnia may differ from those that perpetuate it. For example, an individual who is bedridden with a painful injury and has difficulty sleeping may then develop negative associations for sleep. Conditioned arousal may then persist and lead to persistent insomnia. A similar course may develop in the context of an acute psychological stress or a mental disorder. For instance, insomnia that occurs during an episode of major depressive disorder can become a focus of attention, with consequent negative conditioning, and persist even after resolution of the depressive episode. In some cases, insomnia may also have an insidious onset without any identifiable precipitating factor.
The course of insomnia may also be episodic, with recurrent episodes of sleep difficulties associated with the occurrence of stressful events. Chronicity rates range from 45% to 75% for intervals of 1–7 years. Even when the course of the insomnia has become chronic, there is night-to-night variability in sleep patterns, with an occasional restful night's sleep interspersed with several nights of poor sleep. The characteristics of insomnia may also change over time. Many individuals with insomnia have a history of "light" or easily disturbed sleep prior to onset of more persistent sleep problems.
Insomnia complaints are most prevalent among middle-age and older adults. The type of insomnia symptom changes as a function of age, with difficulties initiating sleep being more common among young adults and problems maintaining sleep occurring more frequently among middle-age and older individuals.
Difficulties initiating and maintaining sleep can also occur in children and adolescents, but there are more limited data on prevalence, risk factors, and comorbidity during these developmental phases of the lifespan. Sleep difficulties in childhood can result from conditioning factors (e.g., a child who does not learn to fall asleep or return to sleep without the presence of a parent) or from the absence of consistent sleep schedules and bedtime routines. Insomnia in adolescence is often triggered or exacerbated by irregular sleep schedules (e.g., phase delay). In both children and adolescents, psychological and medical factors contribute to insomnia.
The increased prevalence of insomnia in older adults is partly explained by the higher incidence of physical health problems with aging. Changes in sleep patterns associated with the normal developmental process must be differentiated from those exceeding age-related changes. Although polysomnography is of limited value in the routine evaluation of insomnia, it may be more useful in the differential diagnosis among older adults because the etiologies of insomnia (e.g., sleep apnea) are more often identifiable in older individuals.

Risk and Prognostic Factors
While the risk and prognostic factors discussed in this section increase vulnerability to insomnia, sleep disturbances are more likely to occur when predisposed individuals are exposed to precipitating events with a major life events (e.g., illness, separation, divorce) are severe but more chronic daily stress. Most individuals resume normal sleep patterns after the initial triggering event disappears, but others—perhaps those more vulnerable to insomnia—continue experiencing persistent sleep difficulties. Predisposing factors such as poor sleep habits, irregular sleep scheduling, and the fear of sleep loss may increase the insomnia problem and may contribute to a vicious cycle that may induce persistent insomnia. Temperament, anxiety or worry-prone personality or cognitive style, increased arousal predisposition, and tendency to repress emotions can increase vulnerability to insomnia. Environmental factors, light, uncontrollably high or low temperature, and high arousal may also increase vulnerability to insomnia.
Genetic and physiological. Female gender and advancing age are associated with increased vulnerability to insomnia. Disrupted sleep and insomnia display a familial disposition. The prevalence of insomnia is higher among monozygotic twins than among dizygotic twins. It is also higher in first-degree family members compared with the general population. The extent to which this risk is inherited through genetic predisposition, learned by observations of parental models, or established as a by-product of another psychopathology remains undetermined.
Course modifiers. Deliberate course modifiers include poor sleep hygiene practices (excessive caffeine use, irregular sleep schedules).

Gender-Related Diagnostic Issues
Insomnia is a more prevalent complaint among females than among males, with first onset often associated with the birth of a new child or with menopause. Despite higher prevalence among older females, polysomnographic studies suggest better preservation of sleep continuity and slow-wave sleep in older females than in older males.
Diagnostic Markers
Polysomnography usually shows impairments of sleep continuity (e.g., increased sleep latency and time awake after sleep onset and decreased sleep efficiency [percentage of time in bed asleep] and may show increased stage 1 sleep and decreased stages 3 and 4 sleep. The severity of these sleep impairments does not always match the individual's clinical complaint or subjective complaint of poor sleep, as individuals with insomnia often underestimate sleep duration and overestimate wakefulness relative to polysomnography. Quantitative electroencephalographic analysis may indicate that individuals with insomnia have greater high-frequency electroencephalography power relative to good sleepers both during the sleep onset period and during non-rapid-eye movement sleep. A feature suggestive of increased cortical arousal. Individuals with insomnia disorder may have a sleep latency that is longer and typically do not show increased daytime sleepiness in objective sleep laboratory measures compared with individuals without sleep disorders. Other laboratory measures show evidence, although not consistently, of increased arousal and a generalized activation of the hypothalamic-pituitary-adrenal axis (e.g., increased cortisol levels, heart rate variability, reactivity to stress, metabolic rate). In general, findings are consistent with the hypothesis that increased physiological and cognitive arousal plays a significant role in insomnia disorder.
Individuals with insomnia disorder may appear either fatigued or haggard or, conversely, irritable and "wired." However, there are no consistently characteristic abnormalities on physical examination. There may be an increased incidence of stress-related psychophysiological symptoms (e.g., tension headaches, muscle tension or pain).

Functional Consequences of Insomnia Disorder