

MATH266: UNSTABLE MOTIVIC HOMOTOPY THEORY  
Fall 2024

Thomas Brazelton

November 8, 2024

# Contents

0.0	Introduction . . . . .	3
<b>1</b>	<b>Torsors</b>	<b>7</b>
1.1	Torsors . . . . .	7
1.2	Torsors II: Simplicial methods . . . . .	15
1.3	Torsors III: Sites, sheaves, and Hilbert 90 . . . . .	22
<b>2</b>	<b>Higher categories</b>	<b>25</b>
2.1	Infinity categories . . . . .	25
2.2	Descent . . . . .	32
2.3	Bonus: more category theory . . . . .	40
<b>3</b>	<b>Motivic spaces</b>	<b>44</b>
3.1	Motivic spaces . . . . .	45
3.2	Jouanolou devices and motivic equivalences . . . . .	50
3.3	Pointed motivic spaces . . . . .	52
3.4	$\mathbb{A}^1$ -homotopy classes of maps . . . . .	54
3.5	Strong and strict $\mathbb{A}^1$ -invariance . . . . .	55
3.6	Milnor $K$ -theory . . . . .	59
3.7	Milnor–Witt $K$ -theory . . . . .	61
3.8	The Milnor conjecture . . . . .	65
3.9	Eilenberg–MacLane spaces . . . . .	67
3.10	Contraction . . . . .	70
3.11	Homotopy modules . . . . .	73
3.12	Gersten complexes . . . . .	76
3.13	The fundamental group of $\mathbb{P}^1$ . . . . .	79
3.14	The Brouwer degree . . . . .	82
3.15	Naive homotopies . . . . .	85
3.16	Unstable connectivity . . . . .	87
3.17	Simplicial Freudenthal . . . . .	89
3.18	Motivic homotopy groups of spheres . . . . .	90

## About these notes

These are notes for a topics class taught at Harvard in Fall 2024. Notes are being updated twice a week with new lectures. Please send any and all changes to

<https://github.com/tbrazel/math266-motivic>

Thank you to Merrick Cai, Juan Fernando Valdes, Jake Levinson, Isabel Longbottom, Sofía Marlasca Aparicio, Matthew Niemi, and Natalie Stewart for valuable changes and suggestions.

Thank you to Aravind Asok, Tom Bachmann, Elden Elmanto, Niels Feld, Mike Hopkins, and Matthias Wendt for answering random motivic questions.

Thank you to Maxine Calle, Connor Cassady, David Chan, Liam Keenan, and Maxmilien Péroux for letting be bother them about technical non-motivic questions.

## 0.0 Introduction

### 0.0.1 Overview

What sorts of things about a ring  $R$  are still true when we move to the polynomial ring  $R[t]$ ? In other words, what sorts of things about  $R$  can't be varied in a 1-parameter family?

Let's give a ton of examples! Don't stress if not all of the words are familiar, we'll break down what's happening here over the course of the semester, this is just motivation.

**Example 0.0.1.** Let  $R$  be a reduced ring. Then the inclusion  $R \rightarrow R[t]$  induces an isomorphism after taking units<sup>1</sup>

$$R^\times \xrightarrow{\sim} (R[t])^\times. \quad (2)$$

Recall that the functor sending a commutative ring to its group of units is corepresented by  $\mathbb{Z}[u, u^{-1}]$ , so **Equation 2** is equivalent to saying that the following map is a bijection

$$\mathrm{Hom}_{\mathrm{Ring}}(\mathbb{Z}[u, u^{-1}], R) \rightarrow \mathrm{Hom}_{\mathrm{Ring}}(\mathbb{Z}[u, u^{-1}], R[t]).$$

After taking  $\mathrm{Spec}$  everywhere, this becomes

$$\mathrm{Hom}_{\mathrm{Sch}}(\mathrm{Spec}(R), \mathbb{G}_m) \rightarrow \mathrm{Hom}_{\mathrm{Sch}}(\mathrm{Spec}(R[t]), \mathbb{G}_m).$$

We therefore might rephrase **Equation 2** as saying that  $\mathrm{Hom}_{\mathrm{Sch}}(-, \mathbb{G}_m)$  is  $\mathbb{A}^1$ -invariant, at least when we plug in something reduced.

**Example 0.0.3.** Let  $k$  be a field. Then the functor  $k \rightarrow k[t_1, \dots, t_n]$  induces an extension of scalars map

$$\begin{aligned} \mathrm{Mod}_k &\rightarrow \mathrm{Mod}_{k[t_1, \dots, t_n]} \\ M &\mapsto M \otimes_k k[t_1, \dots, t_n]. \end{aligned}$$

**Serre's Problem:** Is every finitely generated  $k[t_1, \dots, t_n]$ -module free?

Recall finitely generated projective  $R$ -modules are the same as “algebraic vector bundles” over  $\mathrm{Spec}(R)$ , so we're asking whether every algebraic vector bundle on  $\mathbb{A}_k^n$  is trivial.

**Answer:** Yes (Quillen–Suslin, 1974). Quillen actually proved more— for  $R$  a PID, he proved that the every finitely generated projective  $R[t]$ -module is extended from an  $R$ -module.<sup>2</sup> Lindel proved shortly thereafter that every finitely generated projective  $A[t]$ -module is extended from an  $A$ -algebra, where  $A$  is a smooth algebra containing a field  $k$ . We could read this as saying that the stack of algebraic vector bundles is  $\mathbb{A}^1$ -invariant over the category of smooth affine  $k$ -schemes.

<sup>1</sup>If  $R$  is not reduced, say there is some  $r \in R$  so that  $r^2 = 0$ , then  $(1 + rt)(1 - rt) = 1$ , so  $1 + rt \in R[t]^\times$ .

<sup>2</sup>Quillen's proof involves leveraging some previous work of Horrocks, flat descent for vector bundles, and a very clever technique he invented called *patching*. Suslin's proof, which appeared in the same year, is almost completely linear algebraic, leveraging the theory of *unimodular rows*.

**More general:** (Bass–Quillen conjecture) is it true that for every  $R$  regular Noetherian, the map

$$\begin{aligned} \mathrm{Mod}_R^{\mathrm{f.g., proj}} &\rightarrow \mathrm{Mod}_{R[t]}^{\mathrm{f.g., proj}} \\ M &\mapsto M \otimes_R R[t] \end{aligned}$$

is essentially surjective? *Still open.*

**Fundamental Theorem of Algebraic  $K$ -Theory** (Quillen): For  $R$  regular Noetherian, we have that  $R \rightarrow R[t]$  induces an equivalence<sup>34</sup>

$$K(R) \xrightarrow{\sim} K(R[t]).$$

In other words  $K$ -theory is  $\mathbb{A}^1$ -invariant for regular Noetherian rings (regular Noetherian schemes, more generally).

**Example 0.0.4.** If  $X = \mathrm{Spec}(R)$  or more generally  $X$  is a scheme, then the map  $\pi: X \times \mathbb{A}^1 \rightarrow X$  induces an isomorphism on Chow groups (see for instance [Ful98, 3.3])

$$\pi^*: \mathrm{CH}_*(X) \xrightarrow{\sim} \mathrm{CH}_{*+1}(X \times \mathbb{A}^1).$$

**Example 0.0.5.** Let  $X = \mathrm{Spec}(R)$  where  $R$  is normal and Noetherian.<sup>5</sup> Then every line bundle on  $X \times \mathbb{A}^1$  is extended from a line bundle on  $X$ , in other words  $X \times \mathbb{A}^1 \rightarrow X$  induces an isomorphism

$$\mathrm{Pic}(X) \xrightarrow{\sim} \mathrm{Pic}(X \times \mathbb{A}^1).$$

**Example 0.0.6.** We can also show  $\mathbb{A}^1$ -invariance for the Picard group over a PID. Let  $R$  be a PID, then it is a UFD, and we can show that  $\mathrm{Pic}(R) = 0$ , and therefore  $\mathrm{Pic}(R[t_1, \dots, t_n]) = 0$ .

**Example 0.0.7.** [Aso21, 3.7.1.5] Check this doesn't hold for all rings, for example  $R = k[x, y]/(y^2 - x^3)$ .<sup>6</sup>

**Definition 0.0.8.** An *inner product space* over a ring  $R$  is a finitely generated projective  $R$ -module  $M$  and a symmetric bilinear form  $\beta: M \times M \rightarrow R$  for which  $m \mapsto \beta(-, m)$  defines an isomorphism  $M \cong M^*$ .

**Theorem 0.0.9.** (Harder's Theorem, VII.3.13 in Lam's book on Serre's problem) Let  $k$  be a field. Then every inner product space over  $k[t]$  is extended from an inner product space over  $k$ .

**Remark 0.0.10.** The stable analogue of this has to do with  $\mathbb{A}^1$ -invariance of Hermitian  $K$ -theory [reference needed].

Algebraic vector bundles are  $\mathrm{GL}_n$ -torsors (we will talk about torsors in more detail next week), so the Bass–Quillen conjecture is really asking about  $\mathbb{A}^1$ -invariance of torsors over affine schemes. We could ask an analogous question about  $G$ -torsors for any  $G$ . Here's an example result in this direction that we'll see later in the semester:

**Theorem 0.0.11.** [AHW20, 1.3] If  $k$  is a field, and  $G$  is an isotropic reductive group scheme, then  $G$ -torsors in the Nisnevich site are  $\mathbb{A}^1$ -invariant over any smooth affine  $k$ -scheme.

<sup>3</sup>So Bass–Quillen is really a question about *unstable* modules.

<sup>4</sup>The statement for  $K_0$  is originally due to Grothendieck [Aso21, 5.6.1.3]. The statement for  $K_1$  is due to Bass–Heller–Swan [Aso21, 5.8.2.1].

<sup>5</sup>We can get away with weaker assumptions on this, for example in [Aso21, 3.7.13] it is only assumed that  $R$  is a locally factorial Noetherian normal domain.

<sup>6</sup>More generally if  $A$  is reduced, Noetherian, and has a finite normalization, then its Picard group will be  $\mathbb{A}^1$ -invariant if and only if  $A$  is seminormal [Tra70, 3.6].

## 0.0.2 $\mathbb{A}^1$ -homotopy theory

Recall from algebraic topology that  $X \times [0, 1] \rightarrow X$  is a weak homotopy equivalence, which implies that any cohomology theory is insensitive to taking a product with an interval, e.g. for  $H^*(-, \mathbb{Z})$  integral cohomology we get

$$H^*(X, \mathbb{Z}) \xrightarrow{\sim} H^*(X \times [0, 1], \mathbb{Z}).$$

In fact this type of homotopy invariance is an axiom of generalized Eilenberg–Steenrod cohomology theories.

**Example 0.0.12.** Let  $k \subseteq \mathbb{C}$  be a subfield of the complex numbers. Then there is a *Betti realization* functor

$$\begin{aligned} \mathrm{Var}_k &\rightarrow \mathrm{Top} \\ X &\mapsto X(\mathbb{C}) \end{aligned}$$

sending a variety to its underlying analytic space. Note that

$$X \times \mathbb{A}_k^1 \mapsto (X \times \mathbb{A}_k^1)(\mathbb{C}) = X(\mathbb{C}) \times \mathbb{C}.$$

Therefore any homotopy invariant functor out of spaces provides another example of an  $\mathbb{A}^1$ -invariant functor out of  $k$ -varieties, for example

$$\begin{aligned} X &\mapsto H^*(X(\mathbb{C}); \mathbb{Z}) \\ X &\mapsto \pi_*(X(\mathbb{C})). \end{aligned}$$

**Q:** Can we build a homotopy theory of algebraic varieties in which the affine line  $\mathbb{A}^1$  plays the role that the interval  $[0, 1]$  plays in classical topology?

**A:** Yes! This is what’s known as  $\mathbb{A}^1$ -homotopy theory or *motivic homotopy theory*. This dates back to Morel and Voevodsky’s seminal work in 1999, but many ideas date back to work of Karoubi–Villamayor, Jardine, Weibel in the 1980’s, work of Brown, Gersten, Illusie and Joyal in the 1970’s, and of Quillen and Grothendieck in the 1960’s.

**Q:** What can we do with motivic homotopy theory?

**A:** Motivic homotopy theory blends algebraic geometry and topology in a beautiful way, allowing us to do many things. A short list is:

- ▷ we obtain a natural home for these kinds of  $\mathbb{A}^1$ -invariant cohomology theories of varieties, and can study them in a method analogous to algebraic topology (e.g. we can classify cohomology operations, use fiber sequences to carry out computations, unify various spectral sequences, etc.)
- ▷ we have access to obstruction theory for classifying torsors over affine varieties — recent success in this direction is Asok–Bachmann–Hopkins’ resolution of Murthy’s conjecture
- ▷ we can eliminate differentials in the Adams spectral sequence via looking at their motivic depth (citation needed)

**Q:** What techniques will we learn in this class?

**A:** We should be able to do the following things:

- ▷ learn descent and torsors more thoroughly, and unify perspectives on sheaves and stacks using higher categorical language
- ▷ work with  $\infty$ -categories at a very basic level
- ▷ learn how to work with motivic spaces and do computations
- ▷ learn motivic obstruction theory and its applications

### 0.0.3 A bit more about these notes

**Remark 0.0.13.** (*Transparency about pedagogy*) We're going to place affine representability on the horizon and keep our eyes fixed on it in order to motivate the category of motivic spaces. This is of course ahistorical—although affine representability was one of the key early results explored by Morel, motivic spaces and spectra were developed in order to house theories such as algebraic  $K$ -theory, Bloch's higher Chow groups, and to grow new theories such as algebraic cobordism. We chose this route for a few reasons:

1. The Fall 2024 Thursday seminar is on the motivic Steenrod algebra, so we're hoping this class will contrast nicely and provide some foundational background in techniques in motivic obstruction theory.
2. In order to tailor the course to a broader audience, we'd like to unify the class around a key question which has general appeal, so we've picked the classification of torsors over affine varieties using motivic methods as such a question. This has certain advantages, for instance we can pause in the sheaf topos and discuss classifying torsors there before building motivic spaces — this vista is useful to people across many fields.
3. This forces us to spend a bit longer on descent and torsors, two words which strike fear in the heart of many, and are worth investing some more time in.

**Remark 0.0.14.** (*On background*) We're assuming a strong handle of algebraic geometry and category theory, and a fair bit of familiarity with commutative algebra and homotopy theory. We'll see very quickly that  $\infty$ -categories (and/or model categories) are needed in order to develop the setting in which we wish to work. We've elected to take the  $\infty$ -categorical approach, since it streamlines many of the constructions and key ideas, at the cost of being a high technical investment; for this reason we've done our best to make the  $\infty$ -categorical machinery easy to black box. The reader should be aware that while the categorical language will be heavy in the first half of these notes, it will quickly fade into the background as we become familiar with the ambient setting we're working in and can set our focus towards computations.

# Chapter 1

## Torsors

### Contents

---

<b>1.1 Torsors</b>	<b>7</b>
1.1.1 Sheaves and stacks	9
1.1.2 Interlude: representable $G$ -torsors	11
1.1.3 Interlude: representability of torsors in topology	13
1.1.4 Why we like representability of torsors	14
1.1.5 Motivation of what's to come	14
<b>1.2 Torsors II: Simplicial methods</b>	<b>15</b>
1.2.1 More on simplicial sets	17
1.2.2 Torsors via simplicial maps	19
<b>1.3 Torsors III: Sites, sheaves, and Hilbert 90</b>	<b>22</b>
1.3.1 Torsors in finer topologies	23
1.3.2 Torsors in coarser topologies	23
1.3.3 Special algebraic groups	24

---

### 1.1 Torsors

**Assumption 1.1.1.** For this lecture, every time we say “cover” you can assume we are working with Zariski covers, or even open covers of topological spaces, since the intuition will be the same and the results here will be mostly identical. If you know about other topologies, the statements here work for any site. We will go into sheaves and sites more next week, when we will remark that everything here works for other nice sites (étale, Nisnevich, flat, etc.).

**Definition 1.1.2.** Let  $G$  be a group. Then a  $G$ -set  $X$  is called a *torsor* if its  $G$ -action is simple and transitive. Equivalently, the map

$$\begin{aligned} G \times X &\rightarrow X \times X \\ (g, x) &\mapsto (x, gx) \end{aligned} \tag{1.3}$$

is a bijection.

Note there are two types of  $G$ -torsors — sets of the form  $G/e$ , and the empty set. Depending on the convention, we might want to exclude the case of the empty set by including  $X \neq \emptyset$  in the definition.

**Intuition 1.1.4.** A  $G$ -torsor is like a group  $G$  which has remembered its multiplication but forgotten its identity. Any choice of basepoint  $x \in X$  yields a canonical bijection  $G \xrightarrow{\sim} X$  by sending  $g \mapsto g \cdot x$ .

**Example 1.1.5.** In a locally small category  $\mathcal{C}$ , given two objects  $x, y \in \mathcal{C}$ , the set of isomorphisms  $\text{Isom}_{\mathcal{C}}(x, y)$  is a left  $\text{Aut}_{\mathcal{C}}(x)$ -torsor and a right  $\text{Aut}_{\mathcal{C}}(y)$ -torsor.

Let's try to extend this definition to the setting where  $G$  isn't a single group, but a *sheaf of groups*  $\mathcal{G}$  on a site. What is the appropriate analogue of a torsor in this setting? By abuse of notation we will also call this a *torsor*.

**Definition 1.1.6.** [Stacks, 03AH] Let  $\mathcal{G}$  be a sheaf of groups on  $X$ , and let  $\text{Shv}_{\mathcal{G}}(X)$  denote the category of  $\mathcal{G}$ -sheaves, meaning sheaves of sets equipped with a  $\mathcal{G}$ -action, and equivariant morphisms between them. We define the category of  $\mathcal{G}$ -torsors  $\text{Tors}_{\mathcal{G}}(X) \subseteq \text{Shv}_{\mathcal{G}}(X)$  to be the full subcategory on those  $\mathcal{F}$  so that

1. if  $\mathcal{F}(U)$  is non-empty then the action

$$\mathcal{G}(U) \times \mathcal{F}(U) \rightarrow \mathcal{F}(U)$$

is simply transitive.<sup>1</sup>

2. there exists a covering  $\{U_i \rightarrow X\}$  over which  $\mathcal{F}(U_i) \neq \emptyset$ .<sup>2</sup>

**Terminology 1.1.7.** The choice of topology comes into play in that second point. If  $\mathcal{F} \in \text{Shv}_{\mathcal{G}}(X)$ , we say it is  $\tau$ -locally trivial if  $\mathcal{G}$  is a  $\tau$ -sheaf of groups,  $\mathcal{F}$  is a  $\tau$ -sheaf of sets, and point (2) holds for any  $\tau$ -cover.

**Example 1.1.8.** The sheaf  $\mathcal{G}$ , acting on itself by scaling, is called the *trivial*  $\mathcal{G}$ -torsor.

**Example 1.1.9.** For any group scheme  $G$ , we will refer to  $G$ -torsors, meaning torsors the representable functor  $\text{Hom}(-, G)$ .

**Proposition 1.1.10.** A  $\mathcal{G}$ -torsor  $\mathcal{F}$  on  $X$  is trivial if and only if  $\mathcal{F}(X) \neq \emptyset$ , i.e. if it admits a global section.

**Theorem 1.1.11.** Every morphism in  $\text{Tors}_{\mathcal{G}}(X)$  is an isomorphism.

So what are some examples of torsors, and why might we care to classify them?

**Example 1.1.12.** If  $L/k$  is a Galois field extension then  $\text{Spec}(L) \rightarrow \text{Spec}(k)$  is a  $\text{Gal}(L/k)$ -torsor in the étale topology.

▷ *the inverse Galois problem*: which groups  $G$  occur as Galois groups of number fields? This is asking to scratch the surface of understanding  $G$ -torsors over  $\text{Spec}(\mathbb{Q})$  for all finite groups  $G$ .

**Example 1.1.13.** The associated frame bundle of an algebraic vector bundle is a  $\text{GL}_n$ -torsor (say, in the Zariski topology).

*Proof.* Since  $\text{GL}_n$  is affine, every torsor is representable, hence  $\text{GL}_n$ -torsors are just principal  $\text{GL}_n$ -bundles, which are precisely algebraic vector bundles.  $\square$

▷ *Bass–Quillen conjecture*: this can be reframed as asking whether each  $\text{GL}_n$ -torsor over a regular Noetherian ring is trivial

<sup>1</sup>This means that in the category of  $\mathcal{G}(U)$ -sets, we have that  $\mathcal{F}(U) \cong \mathcal{G}(U)$ , but  $\mathcal{F}(U)$  doesn't have a group structure — we might imagine that it has forgotten its identity element. Picking a basepoint  $e \in \mathcal{F}(U)$  defines a group structure on  $\mathcal{F}(U)$ .

<sup>2</sup>In other words, we can find a cover in which to visualize  $\mathcal{F}(U_i)$  as a group for each  $i$ .



▷ *Hartshorne’s conjecture* concerns  $\mathrm{GL}_2$ -torsors over  $\mathbb{P}^n$  for  $n \geq 7$

**Example 1.1.14.** A  $\mathrm{PGL}_n$ -torsor is a Brauer–Severi variety (or a central simple algebra).

▷ the *period-index conjecture* then concerns the complexity of  $\mathrm{PGL}_n$ -torsors.

**Goal 1.1.15.** Develop methods to classify torsors.

Let’s do this, by first considering an alternative perspective on what a torsor is. We learned this from Alex Youcis’ excellent note on torsors [You].

### 1.1.1 Sheaves and stacks

Recall if  $\mathcal{F}$  is a sheaf of sets, this means for every cover  $\{U_i \rightarrow U\}$ , we have that the diagram is an equalizer

$$\mathcal{F}(U) \rightarrow \prod \mathcal{F}(U_i) \rightrightarrows \prod_{i,j} \mathcal{F}(U_i \times_U U_j).$$

This first map is a monomorphism (injection) because it is an equalizer. This means that if  $x, y \in \mathcal{F}(U)$  are equal in  $\mathcal{F}(U_i)$  for each  $i$ , then they are equal in  $\mathcal{F}(U)$ . In other words, the map “reflects equality” — this is literally just what it means for something to be an injection.

Let’s suppose now that  $\mathcal{F}(U)$  is a *category* for every  $U$ . We’ll define this concretely soon once we have more machinery, but for now let’s just pretend that we know what this means — it means we can glue objects and morphisms along covers. Consider the analogous restriction functor:

$$\begin{aligned} \mathcal{F}(U) &\rightarrow \prod \mathcal{F}(U_i) \\ x &\mapsto (x|_{U_i})_i. \end{aligned}$$

**Q:** Does this map need to reflect isomorphisms?<sup>3</sup>

**Example 1.1.16.** Let  $\mathcal{F}$  be the functor sending  $U$  to the category of line bundles over  $U$ . If the cover is picked appropriately small, then all line bundles are isomorphic to the trivial line bundle over  $U_i$ , but they need not be isomorphic globally as line bundles over  $U$ .

**A:** No, by the example above. This means that we can have  $x, y \in \mathcal{F}(U)$  so that  $x|_{U_i} \cong y|_{U_i}$  for each  $i$ , but we *do not have* that  $x$  and  $y$  are isomorphic in the category  $\mathcal{F}(U)$ . In other words, the following two notions are *different*.

- ▷  $x$  and  $y$  agree *globally*, meaning  $x$  and  $y$  are isomorphic in  $\mathcal{F}(U)$ .
- ▷  $x$  and  $y$  agree *locally*, meaning there exists an open cover  $\{U_i \rightarrow U\}$  for which we have isomorphisms  $x|_{U_i} \xrightarrow{\sim} y|_{U_i}$  in  $\mathcal{F}(U_i)$ .<sup>4</sup>

**Remark 1.1.17.** This is a big difference between sheaves of sets (or 1-categories in general) and sheaves of categories (also called *stacks*). Equality is reflected along a cover for sheaves of sets, but isomorphism is not necessarily reflected along a cover.

**Question 1.1.18.** How many isomorphism classes of objects  $y \in \mathcal{F}(U)$  are *locally isomorphic* to  $x$  along a cover, but not globally isomorphic?

We’re going to build a sheaf that measures this! We’ll call this sheaf  $\mathrm{Aut}_{\mathcal{F}}(x)$ , and it is defined by

$$U_i \mapsto \mathrm{Aut}_{\mathcal{F}(U_i)}(x). \quad (1.19)$$

<sup>3</sup>A functor which reflects isomorphisms is called *conservative*.

<sup>4</sup>The notation  $x|_{U_i}$  is shorthand for the image of  $x$  under the restriction functor  $\mathcal{F}(U) \rightarrow \mathcal{F}(U_i)$

**Exercise 1.1.20.** This is a priori just a presheaf of groups. Check it is actually a sheaf of groups.

Recall the following definition.

**Definition 1.1.21.** Let  $\mathcal{G}$  denote a sheaf of groups over  $U$  and  $\mathcal{U} = \{U_i \rightarrow U\}$  a cover.

1. We define a *Čech 1-cocycle* to be a collection of elements  $g_{ij} \in \mathcal{G}(U_i \times_U U_j)$  for each  $i, j$  so that

$$g_{ij}g_{jk} = g_{ik}$$

on triple overlaps.<sup>5</sup>

2. We say two 1-cocycles  $(g_{ij})$  and  $(\gamma_{ij})$  are cohomologous if there are  $\alpha_i \in \mathcal{G}(U_i)$  for each  $i$  so that

$$\alpha_i g_{ij} = \gamma_{ij} \alpha_j.$$

3. We define the *Čech cohomology*  $\check{H}^1(U, \mathcal{G})$  to be the colimit of the Čech cohomology over covers, filtered with respect to refinement.

**Theorem 1.1.22.** There is a bijection between  $H^1(U, \text{Aut}_{\mathcal{F}}(x))$  and isomorphism classes of objects  $y \in \mathcal{F}(U)$  which are locally isomorphic to  $x$ .

*Proof.* Let's first define a map. If  $y$  is locally isomorphic to  $x$ , then there is a cover  $\{U_i\}$  of  $U$  and isomorphisms  $\phi_i: x|_{U_i} \xrightarrow{\sim} y|_{U_i}$  for each  $i$ . If the  $\phi_i$ 's agreed on overlaps then they would glue to a global isomorphism  $\phi: x \xrightarrow{\sim} y$  because  $\text{Aut}(x)$  is a sheaf of groups, so it makes sense to look on overlaps to see what happens. Note that  $\phi_i|_{U_{ij}}$  and  $\phi_j|_{U_{ij}}$  will differ by an automorphism of  $x|_{U_{ij}}$ , call this  $g_{ij}$ :

$$g_{ij} := \left( \phi_i|_{U_{ij}} \right)^{-1} \left( \phi_j|_{U_{ij}} \right) : x|_{U_{ij}} \xrightarrow{\sim} x|_{U_{ij}}.$$

On triple overlaps, it is straightforward to verify that

$$g_{ij}g_{jk} = g_{ik}.$$

In other words we get a 1-cocycle! There was ambiguity here, since we *picked* isomorphisms  $x|_{U_i} \xrightarrow{\sim} y|_{U_i}$  as our starting data. The remaining thing to prove is that any other choice of local isomorphisms gives rise to a cohomologous 1-cocycle.

Suppose we instead picked some  $\psi_i: x|_{U_i} \xrightarrow{\sim} y|_{U_i}$  for each  $i$ , yielding  $\gamma_{ij} = \psi_j^{-1}\psi_i$ . Then  $\psi_i$  and  $\phi_i$  differ by an automorphism of  $y$  which we call  $\alpha_i$ :

$$\begin{array}{ccc} x|_{U_i} & \xrightarrow{\phi_i} & y|_{U_i} \\ & \searrow \alpha_i & \nearrow \psi_i \\ & x|_{U_i} & \end{array}$$

Then on  $U_{ij}$  we have

$$g_{ij} = \phi_i^{-1}\phi_j = (\psi_i\alpha_i)^{-1}(\psi_j\alpha_j) = \alpha_i^{-1}\psi_i^{-1}\psi_j\alpha_j.$$

Hence

$$\alpha_i g_{ij} = \gamma_{ij} \alpha_j$$

And we get cohomologous 1-cocycles. □

**Theorem 1.1.23.** There is a bijection between isomorphism classes of  $\mathcal{G}$ -torsors and  $\check{H}^1(U, \mathcal{G})$ .

---

<sup>5</sup>See e.g. [Mil13, §11].

*Sketch.* Let  $\mathcal{F}$  be a  $\mathcal{G}$ -torsor, and pick  $s_i \in \mathcal{F}(U_i)$  for each  $i$ . Then on the overlap  $U_i \times_U U_j$ , we have that  $s_i$  and  $s_j$  differ by a unique element  $g_{ij} \in \mathcal{G}(U_i \times_U U_j)$ . We run basically an identical argument.  $\square$

So if  $\mathcal{G} = \text{Aut}_{\mathcal{F}}(x)$  then we have a bijection

$$\left\{ \begin{array}{l} \text{iso classes of } y \in \mathcal{F}(U) \\ \text{locally isomorphic to } x \end{array} \right\} \leftrightarrow \check{H}^1(U, \text{Aut}_{\mathcal{F}}(x)) \leftrightarrow \{\text{Aut}_{\mathcal{F}}(x)\text{-torsors}\}.$$

These sorts of arguments are compatible with refinement of the cover, and since Čech and sheaf cohomology agree we see that  $G$ -torsors are in bijection with the first sheaf cohomology  $H^1(U, G)$ .

**Exercise 1.1.24.** Show that every sheaf of groups  $\mathcal{G}$  is of the form  $\text{Aut}_{\mathcal{F}}(x)$  for some sheaf of categories  $\mathcal{F}$ .

**Intuition 1.1.25.** A  $\mathcal{G}$ -torsor is an object whose automorphisms locally look like  $\mathcal{G}(U_i)$ .

### 1.1.2 Interlude: representable $G$ -torsors

When both the presheaf  $\mathcal{G}$  and the sheaf of sets  $\mathcal{F}$  are representable, we get a slightly different characterization.

**Setup 1.1.26.** Suppose  $\mathcal{C} = \text{Sch}_X$  is a site of schemes over  $X$ , and let  $G \in \text{Grp}(\text{Sch}_X)$  be a group scheme over  $X$ . Suppose  $\mathcal{G} = \text{Hom}_X(-, G)$  is a representable sheaf of groups, and let  $\mathcal{F} = \text{Hom}_X(-, Y)$  for some  $Y \in \text{Sch}_X$ , where  $Y$  comes equipped with a  $G$ -action.

In this setup, what does it mean in this case for  $\mathcal{F}$  to be a  $\mathcal{G}$ -torsor?

Condition (1) from [Definition 1.1.6](#) asks that

$$\text{Hom}_X(U, G) \times \text{Hom}_X(U, Y) \rightarrow \text{Hom}_X(U, Y)$$

to be simply transitive for every  $U \in \text{Sch}_X$ . This seems a bit tedious to check, but the following result gives us a cleaner characterization of it, which is a sheafy version of [Equation 1.3](#).

**Proposition 1.1.27.** [[Stacks](#), 0499] In [Setup 1.1.26](#) the following two conditions are equivalent:

1. The map

$$\begin{aligned} G \times_X Y &\rightarrow Y \times_X Y \\ (g, y) &\mapsto (y, gy) \end{aligned}$$

is an isomorphism of  $X$ -schemes.

2. For every  $U \in \text{Sch}_X$ , the induced action

$$\text{Hom}_X(U, G) \times \text{Hom}_X(U, Y) \rightarrow \text{Hom}_X(U, Y)$$

is simply transitive.

What about condition (2) from [Definition 1.1.6](#)? This condition translates, in [Setup 1.1.26](#), to asking for a cover  $\{U_i \rightarrow X\}$  for which  $\text{Hom}_X(U_i, Y) \neq \emptyset$  for each  $i$ . Pick any  $s_i \in \text{Hom}_X(U_i, Y)$ , then it makes the diagram commute:

$$\begin{array}{ccc} & & Y \\ & \nearrow s & \downarrow \\ U_i & \longrightarrow & X. \end{array}$$

Then the pullback torsor  $Y \times_X U_i \rightarrow U_i$  admits a section, which implies it is trivial by [Proposition 1.1.10](#):

$$\begin{array}{ccc} Y \times_X U_i & \longrightarrow & X \\ \downarrow \lrcorner & & \downarrow \\ U_i & \longrightarrow & X. \end{array}$$

We summarize these observations in the following proposition.

**Proposition 1.1.28.** Let  $X$  be a scheme, let  $G$  be a group scheme over  $X$ , and let

$$f: Y \rightarrow X$$

be an  $X$ -scheme equipped with a  $G$ -action. Then  $Y$  is a  $G$ -torsor if and only if

1. The map  $G \times_X Y \rightarrow Y \times_X Y$  is an isomorphism of  $X$ -schemes.
2. There exists a cover  $\{U_i \rightarrow X\}$  for which  $Y \times_X U_i \rightarrow U_i$  is isomorphic to the trivial  $G$ -torsor over  $U_i$ .

In this setting, we call  $Y \rightarrow X$  a *principal  $G$ -bundle*.

**Example 1.1.29.** Let's double back to [Example 1.1.12](#) and actually prove that  $\mathrm{Spec} L \rightarrow \mathrm{Spec} k$  is a principal  $\mathrm{Gal}(L/k)$ -bundle in the étale site when  $L/k$  is a Galois field extension.

*Proof.* We check the conditions from [Proposition 1.1.28](#). The first condition asks that

$$\left( \coprod_{g \in G} \mathrm{Spec}(k) \right) \times_{\mathrm{Spec}(k)} \mathrm{Spec}(L) \rightarrow \mathrm{Spec}(L) \times_{\mathrm{Spec} k} \mathrm{Spec}(L)$$

is an equivalence. Since everything in sight is affine, we can rephrase this as asking for the map

$$\begin{aligned} \coprod_{g \in G} L &\leftarrow L \otimes_k L \\ (x \cdot g(y))_{g \in G} &\leftarrow x \otimes y \end{aligned}$$

to be an equivalence, which we recall is the *normal basis theorem* from Galois theory. The second condition from [Proposition 1.1.28](#) asks us to find a cover of  $\mathrm{Spec}(k)$  trivializing  $\mathrm{Spec}(L) \rightarrow \mathrm{Spec}(k)$ , but if we work in the étale site, then  $\mathrm{Spec}(L) \rightarrow \mathrm{Spec}(k)$  is itself a cover, and it is clear that the product induces a section

$$\begin{array}{ccc} \mathrm{Spec}(L \otimes_k L) & \longrightarrow & \mathrm{Spec}(L) \\ \downarrow \lrcorner & & \downarrow \\ \mathrm{Spec}(L) & \longrightarrow & \mathrm{Spec}(k). \end{array}$$

Hence we have a trivialization, and hence a principal  $\mathrm{Gal}(L/k)$ -bundle (a  $\mathrm{Gal}(L/k)$ -torsor).  $\square$

Often it's just enough to assume the sheaf of groups is representable to get all torsors are representable.

**Proposition 1.1.30.** Let  $G$  be an affine group scheme over  $X$ , and let  $\tau \leq \mathrm{fppf}$ . Then every  $G$ -torsor is representable. (see [\[Stacks, 0497\]](#), [\[You, 3.25\]](#))

*Sketch.* Every  $G$ -torsor is an algebraic space, and algebraic spaces which are locally affine are schemes. Since a  $G$ -torsor is locally isomorphic to  $G$ , which was assumed to be affine, then we conclude every torsor is actually a scheme.  $\square$

**Terminology 1.1.31.** A representable fpqc-torsor for  $G$  is called a *principal homogeneous space*.

**Remark 1.1.32.** If  $t \leq \tau$ , then every  $t$ -cover is a  $\tau$ -cover, hence if we can find a  $t$ -cover trivializing a  $G$ -torsor, then it also trivializes it in the  $\tau$ -topology. hence

$$t \leq \tau \Rightarrow \{t\text{-torsors}\} \subseteq \{\tau\text{-torsors}\}.$$

So a very natural question is *how do we tell when a  $\tau$ -torsor is also a  $t$ -torsor?* We'll discuss this next week.

### 1.1.3 Interlude: representability of torsors in topology

So it's a very reasonable goal to ask for any tools that could help us try to classify torsors. A natural idea, by analogy, is to look to homotopy theory, where we have a suite of tools for studying torsors.

**Definition 1.1.33.** Let  $X$  be a compact Hausdorff topological space and  $G$  a group. Then a *principal  $G$ -bundle* (or we might just say a  $G$ -torsor) is a fiber bundle  $\pi: Y \rightarrow X$  so that  $G$  acts freely and transitively on the fibers.

In topology there is a *universal  $G$ -torsor*, which is denoted  $EG \rightarrow BG$ . This is universal in the sense that, given any map  $f: X \rightarrow BG$ , we can consider the fiber product

$$\begin{array}{ccc} f^*EG & \longrightarrow & EG \\ \downarrow & \lrcorner & \downarrow \\ X & \xrightarrow{f} & BG. \end{array}$$

Then  $f^*EG \rightarrow X$  is a principal  $G$ -bundle, and all principal  $G$ -bundles are obtained in this way. Not only that, but isomorphic principal  $G$ -bundles are given by homotopic classifying maps. In other words we have a bijection

$$\text{Prin}_G(X) \leftrightarrow [X, BG].$$

So the data of a principal  $G$ -bundle is the data of a map  $X \rightarrow BG$ , and an isomorphism of principal  $G$ -bundles is equivalent to a homotopy between two maps  $f, g: X \rightarrow BG$ .

**Example 1.1.34.** We have that  $B(\mathbb{Z}/2) = \mathbb{RP}^\infty$  and  $E\mathbb{Z}/2 = S^\infty$ , so that  $\mathbb{Z}/2$ -torsors are real line bundles. Similarly  $BC^\times = \mathbb{CP}^\infty$ .

A big example comes from quotienting out by a compact subgroup:

**Theorem 1.1.35.** (Samelson, 1941) If  $H \leq G$  is a compact subgroup of a Lie group, then  $G \rightarrow G/H$  is a Serre fibration and principal  $H$ -bundle.

**Corollary 1.1.36.** We have fiber sequences

$$\begin{array}{c} H \rightarrow G \rightarrow G/H \\ G/H \rightarrow BH \rightarrow BG. \end{array}$$

**Example 1.1.37.** For the inclusions  $O(n) \subseteq O(n+1)$  and  $U(n) \subseteq U(n+1)$  we get fiber sequences

$$\begin{array}{c} S^{2n+1} \rightarrow BU(n) \rightarrow BU(n+1) \\ S^n \rightarrow BO(n) \rightarrow BO(n+1). \end{array}$$

This is how Bott periodicity is proved.

### 1.1.4 Why we like representability of torsors

This has a number of huge applications:

- ▷ Given any cohomology theory  $E^*$  and any class  $c \in E^*(BG)$ , if  $f^*(c) \neq g^*(c)$  in  $E^*(X)$ , this means that  $f$  and  $g$  correspond to non-isomorphic torsors. This is the basic idea of characteristic classes.

**Example 1.1.38.** If  $G = \mathrm{GL}_n(\mathbb{C})$ , then  $\mathrm{BGL}_n(\mathbb{C}) = \mathrm{BU}(n) = \mathrm{Gr}_{\mathbb{C}}(n, \infty)$  is a Grassmannian of  $n$ -planes in  $\mathbb{C}^\infty$ . A map  $f: X \rightarrow \mathrm{Gr}_{\mathbb{C}}(n, \infty)$  gives a complex  $n$ -dimensional vector space by pullback.

**Theorem 1.1.39.** (Pontryagin–Steenrod) There is a bijection

$$\mathrm{Vect}_{\mathbb{C}}^n(X) \cong [X, \mathrm{BU}(n)].$$

Since  $H^*(\mathrm{BU}(n); \mathbb{Z}) = \mathbb{Z}[c_1, \dots, c_n]$ , if  $E \rightarrow X$  is any rank  $n$  vector bundle classified by a map  $f: X \rightarrow \mathrm{BU}(n)$ , then its Chern classes are by definition  $c_i(E) = f^*c_i$ .

- ▷ We have access to *obstruction theory* – this lets us break down lifting problems into smaller manageable stages.

**Example 1.1.40.** If  $X$  is a complex  $n$ -dimensional manifold, and  $E \rightarrow X$  is a rank  $n$  complex vector bundle, then it splits off a free summand if and only if  $c_n(E) = 0$ .

**Example 1.1.41.** If we fix  $c_1, \dots, c_n \in H^*(X; \mathbb{Z})$ , we can ask how many isomorphism classes of complex rank  $n$  vector bundles on  $X$  have these given Chern classes. Since  $c_i \in H^{2i}(X; \mathbb{Z}) = [X, K(\mathbb{Z}, 2i)]$  this is equivalent to asking how many lifts there are for

$$\begin{array}{ccc} & & \mathrm{BU}(n) \\ & \nearrow & \downarrow \\ X & \xrightarrow{c_1, \dots, c_n} & \prod_i K(\mathbb{Z}, 2i). \end{array}$$

If  $X$  is a finite CW complex, there are only finitely many such lifts by basic obstruction theory.

- ▷ Suppose we have two groups  $G$  and  $K$ , and we want to study natural ways to create  $K$ -torsors out of  $G$ -torsors over any space. Then representability, combined with the Yoneda lemma, allows us to completely classify all the ways to do this.

**Example 1.1.42.** There is one and only one natural non-trivial function

$$\{\mathrm{GL}_n(\mathbb{R})\text{-torsors}\} \rightarrow \{(\mathbb{Z}/2)\text{-torsors}\},$$

given by the nonzero class in  $[\mathrm{BGL}_n(\mathbb{R}), B\mathbb{Z}/2] = \mathbb{Z}/2$ . This is called the *determinant bundle* or the *first Stiefel–Whitney class*.

### 1.1.5 Motivation of what’s to come

**Question 1.1.43.** By analogy to homotopy theory, we might ask, for a group scheme  $G$ , the following questions:

1. Is there an analogous universal space  $BG$  in algebraic geometry which classifies  $G$ -torsors?
2. If so, can we classify  $G$ -torsors over  $X$  via some “homotopy classes” of maps from  $X$  to  $BG$ ?

The answer to both will be yes, but *not in the category of varieties*. We need more machinery than is available to us there.

## 1.2 Torsors II: Simplicial methods

**Definition 1.2.1.** We denote by  $\Delta$  the category whose objects are finite ordered sets of the form  $[n] = \{0 < 1 < \cdots < n\}$ , and whose morphisms  $[n] \rightarrow [m]$  are order-preserving maps.

**Notation 1.2.2.** Let  $\mathcal{C}$  be a category. Then we refer to a functor  $\Delta \rightarrow \mathcal{C}$  as a *cosimplicial object* of  $\mathcal{C}$ , and a functor  $\Delta^{\text{op}} \rightarrow \mathcal{C}$  as a *simplicial object* of  $\mathcal{C}$ .


**Intuition 1.2.3.** We should think about a (co)simplicial object as a *data type* — it is a combinatorial gadget that is surprisingly convenient for bookkeeping and appears frequently in nature. We'll see quite a few examples, but let's first see how to compress this data.

There are a priori a lot of order-preserving functions  $[n] \rightarrow [m]$ , so we'd like a nice class of easy-to-manage morphisms in  $\Delta$  so that any morphism factors into nice morphisms. These fall in two classes:


### Terminology 1.2.4.

1. We have *coface* maps  $d^i: [n] \rightarrow [n+1]$  for  $0 \leq i \leq n+1$  which are defined by the property that they miss the element  $i \in [n+1]$ .
2. We have *codegeneracy* maps  $s^j: [n] \rightarrow [n-1]$  for  $0 \leq j \leq n-1$  defined by the property that they are surjective and that  $s^j(j) = s_j(j+1)$ .

**Example 1.2.5.** The coface maps  $d^i: [2] \rightarrow [3]$  look like



**Example 1.2.6.** The codegeneracy maps  $[3] \rightarrow [2]$  look like



**Terminology 1.2.7.** Let  $X_\bullet: \Delta^{\text{op}} \rightarrow \mathcal{C}$  denote a simplicial object.

1. We use the notation  $X_n$  to denote the object  $X_\bullet([n])$  in  $\mathcal{C}$ .
2. We denote by  $d_i: X_n \rightarrow X_{n-1}$  for  $0 \leq i \leq n$  the image of the coface map  $d^i$ . We call  $d_i$  a *face map*.
3. Similarly we denote by  $s_i: X_n \rightarrow X_{n+1}$  for  $0 \leq i \leq n$  the image of the codegeneracy map  $s^i$ , and call  $s_i$  a *degeneracy map*.

**Exercise 1.2.8.** Convince yourself that every map in  $\Delta$  factors into face and degeneracy maps, and therefore any simplicial object  $X_\bullet$  in  $\mathcal{C}$  can be described by the data of  $X_n$  for each  $n$ , and its face and degeneracy maps.<sup>6</sup>

**Definition 1.2.9.** If  $\mathcal{C}$  is a locally small category with all finite products (binary products and a terminal object), then a *group object* in  $\mathcal{C}$  is the data of an object  $G \in \mathcal{C}$  together with morphisms

$$\begin{aligned} m: G \times G &\rightarrow G \\ e: 1 &\rightarrow G \\ i: G &\rightarrow G \end{aligned}$$

called multiplication, identity, and inverse, such that the following diagrams commute.

$$\begin{array}{ccccc} G \times G \times G & \xrightarrow{\text{id} \times m} & G \times G & & G & \xrightarrow{e \times \text{id}} & G \times G & & G & \xrightarrow{i \times \text{id}} & G \times G \\ m \times \text{id} \downarrow & & \downarrow m & & \text{id} \times e \downarrow & \searrow \text{id} & \downarrow m & & \text{id} \times i \downarrow & \searrow e & \downarrow m \\ G \times G & \xrightarrow{m} & G & & G \times G & \xrightarrow{m} & G & & G \times G & \xrightarrow{m} & G \end{array}$$

**Example 1.2.10.** Let  $G$  be a group object in a category  $\mathcal{C}$  as above. Then we can define a simplicial object

$$\begin{aligned} B_\bullet G: \Delta^{\text{op}} &\rightarrow \mathcal{C} \\ [n] &\mapsto G^{\times n}, \end{aligned}$$

as:

$$* \begin{array}{c} \xleftarrow{\quad} \\ \xrightarrow{\quad} \end{array} G \begin{array}{c} \xleftarrow{\quad} \\ \xrightarrow{\quad} \end{array} G \times G$$

Explicitly,  $B_n G = G^{\times n}$ , with face maps  $d_i: G^{\times n} \rightarrow G^{\times(n-1)}$  given by

$$d_i(g_1, \dots, g_n) = \begin{cases} (g_2, \dots, g_n) & i = 0 \\ (g_1, \dots, g_i g_{i+1}, g_{i+2}, \dots, g_n) & 1 \leq i \leq n-1 \\ (g_1, \dots, g_{n-1}) & i = n, \end{cases}$$

and degeneracies  $s_i: G^{\times n} \rightarrow G^{\times(n+1)}$  given by

$$s_i(g_1, \dots, g_n) = \begin{cases} (e, g_1, \dots, g_n) & i = 0 \\ (g_1, \dots, g_i, e, g_{i+1}, \dots, g_n) & 1 \leq i \leq n. \end{cases}$$

Note that  $d_1: G^{\times 2} \rightarrow G$  is precisely the group multiplication, while  $s_0: * \rightarrow G$  is the identity on the group.

**Exercise 1.2.11.** Check the simplicial identities hold *precisely* because of the group axioms. In particular, observe that we didn't use anything special about spaces — if  $G$  is a group object in any category  $\mathcal{C}$  we obtain an associated bar construction which is a simplicial object in  $\mathcal{C}$  that we call  $B_\bullet G$ .

**Example 1.2.12.** (Important) For the formal categorical reason in [Exercise 1.2.11](#), any group scheme  $G$  gives rise to a simplicial scheme  $B_\bullet G \in \text{Fun}(\Delta^{\text{op}}, \text{Sch})$ . This dates back at least to work of Friedlander [[Fri82](#), Example 1.2]. We'll use this object frequently.

**Example 1.2.13.** Simplicial objects appeared crucially in Deligne's work on resolution of singularities and mixed Hodge structures. Let  $X$  be a complete<sup>7</sup> singular variety. Then we can “replace”  $X$

<sup>6</sup>Any time we define a simplicial object via this compressed data, we should check that the simplicial identities hold, although we'll mostly omit these arguments here.

<sup>7</sup> $X$  is *complete* if it is proper over the base [[Har77](#), p. 105]. This is a slightly more general notion than projective.



with a simplicial variety

$$\cdots \quad X_2 \begin{matrix} \xrightarrow{\quad} \\ \xleftarrow{\quad} \\ \xrightarrow{\quad} \end{matrix} X_1 \begin{matrix} \xrightarrow{\quad} \\ \xleftarrow{\quad} \\ \xrightarrow{\quad} \end{matrix} X_0,$$

where each  $X_i$  is smooth and projective. There is a cohomological descent spectral sequence computing the cohomology of  $X$  in terms of the cohomology of the  $X_n$ 's. This dates back to SGA 4. Additionally, this replacement is the core idea of *resolution of singularities*. Deligne uses these simplicial methods to endow singular varieties with Hodge structures, so-called *mixed Hodge structures* [Del74].

**Example 1.2.14.** Let  $A_\bullet : \Delta^{\text{op}} \rightarrow \text{Ab}$  be a simplicial abelian group. Then by taking the alternating sum of the face maps, we obtain a chain complex, purely by the simplicial identities. This assignment is functorial:

$$\begin{aligned} \text{sAb} &\rightarrow \text{Ch}_{\geq 0}(\text{Ab}) \\ A_\bullet &\mapsto \left( A_n, \partial = \sum_{i=0}^n (-1)^i d_i \right). \end{aligned}$$

The associated chain complex is called the *Moore complex* of the simplicial abelian group. Two quick remarks about this:

1. This works if we replace  $\text{Ab}$  by any abelian category.
2. This process is invertible — this implies that simplicial objects and connective chain complexes in any abelian category are equivalent.<sup>8</sup> We will use this later, as we will want to construct certain simplicial objects in sheaves of abelian groups, and it will be more direct to construct them first as chain complexes, then pass through this equivalence.

**Example 1.2.15** ( $\check{C}$ ech nerve  $N_\bullet(U)$ ). A simplicial object can conveniently encode the data of an open cover, its intersections, triple intersections, etc.<sup>9</sup> For this, given an open cover  $\{U_i \rightarrow X\}$  of a variety  $X$ , denote by  $U_{ij} := U_i \times_X U_j$  the double overlaps, by  $U_{ijk} := U_i \times_X U_j \times_X U_k$  the triple overlaps, and so on. We define a simplicial object

$$\coprod_i U_i \begin{matrix} \xleftarrow{\quad} \\ \xrightarrow{\quad} \\ \xleftarrow{\quad} \end{matrix} \coprod_{i,j} U_{ij} \begin{matrix} \xleftarrow{\quad} \\ \xrightarrow{\quad} \\ \xleftarrow{\quad} \end{matrix} \coprod_{i,j,k} U_{ijk} \quad \cdots$$

The face maps are defined by omitting the  $j$ th index

$$d_j : U_{i_1 \dots i_n} \rightarrow U_{i_1 \dots \widehat{i_j} \dots i_n},$$

and the degeneracy maps repeat the  $j$ th index

$$s_j : U_{i_1 \dots i_n} \rightarrow U_{i_1 \dots i_{j-1} i_j i_j i_{j+1} \dots i_n}.$$

We call this the  *$\check{C}$ ech nerve* associated to the cover, and we denote it by  $N(\mathcal{U})$ .

### 1.2.1 More on simplicial sets

We denote by

$$\text{sSet} := \text{Fun}(\Delta^{\text{op}}, \text{Set})$$

<sup>8</sup>There's a small lie here, the actual functor exhibiting the equivalence is not the Moore complex but the *normalized chains complex*, although the normalized chain complex maps to the Moore complex in a natural way, and this is a chain homotopy equivalence [GJ99, III.2.4], so they are essentially the same. This equivalence is the *Dold–Kan theorem*. For more detail see [GJ99, §III.2].

<sup>9</sup>Technically it should be an *augmented simplicial object* since we'd also like to remember the data of the maps  $U_i \rightarrow X$ .

the category of *simplicial sets*. Given a simplicial set

$$\begin{aligned} X_\bullet: \Delta^{\text{op}} &\rightarrow \text{Set} \\ [n] &\mapsto X_n, \end{aligned}$$

we call  $X_n$  the set of  $n$ -simplices.

**Example 1.2.16.** Any set  $Y$  gives rise to a constant simplicial set  $\underline{Y}$ , given by sending  $[n] \mapsto Y$ , and every morphism in  $\Delta$  to the identity on  $Y$ .

**Example 1.2.17.** We denote by  $\Delta^n$  the simplicial set

$$\Delta^n := \text{Hom}_\Delta(-, [n]): \Delta^{\text{op}} \rightarrow \text{Set}.$$

By the Yoneda lemma, we have a natural bijection

$$\text{Hom}_{\text{sSet}}(\Delta^n, X_\bullet) \cong X_n$$

for any  $X_\bullet \in \text{sSet}$ .

There is a functor called *geometric realization*, which “assembles” a simplicial set into a topological space: Let  $\Delta_{\text{top}}^n$  denote the *topological  $n$ -simplex*<sup>10</sup>

$$\begin{aligned} | - |: \text{sSet} &\rightarrow \text{Top} \\ X_\bullet &\mapsto \coprod_{n \geq 0} X_n \times \Delta_{\text{top}}^n / \sim, \end{aligned}$$

where  $\sim$  is

$$\begin{aligned} (x, d_i u) &\sim (d_i x, u) & x \in X_n, u \in \Delta_{\text{top}}^{n-1} \\ (y, s_i v) &\sim (s_i y, v) & y \in X_{n-1}, v \in \Delta_{\text{top}}^n. \end{aligned}$$

**Remark 1.2.18.** We discussed this universal space  $BG$  classifying principal  $G$ -bundles last week. We can define this as  $BG := |B_\bullet G|$ , i.e. it is the geometric realization of the bar construction for the group.<sup>11</sup>

**Definition 1.2.19.** We say two maps  $f, g: X_\bullet \rightarrow Y_\bullet$  are *simplicially homotopic* if there is a map

$$H: \Delta^1 \times X_\bullet \rightarrow Y_\bullet$$

so that  $H|_{\{0\} \times X_\bullet} = f$  and  $H|_{\{1\} \times X_\bullet} = g$ .

**Remark 1.2.20.** Unpacking this data, we can verify it is the same as asking for maps for every  $n$ :

$$H_i^n: X_n \rightarrow Y_{n+1} \quad 0 \leq i \leq n,$$

so that  $d_0 H_0^n = f_n$ , and  $d_{n+1} H_n^n = g_n$ , and so that they satisfy the following relations with face and degeneracy maps:<sup>12</sup>

$$\begin{aligned} d_i H_j^n &= \begin{cases} H_{j-1}^{n-1} d_i & i < j \\ d_i H_{j-1}^n & i = j \neq 0 \\ d_{j+1} H_{j+1}^n & i = j + 1, j \neq n \\ H_j^{n-1} d_{i-1} & i > j + 1. \end{cases} \\ s_i H_j^n &= \begin{cases} H_{j+1}^{n+1} s_i & i \leq j \\ H_j^{n+1} s_{i-1} & i > j. \end{cases} \end{aligned}$$

This notion makes sense for maps between simplicial objects in any category, and it is called a *simplicial homotopy*.

<sup>10</sup>This is the locus  $\Delta_{\text{top}}^n := \{(x_0, \dots, x_{n+1}) \in \mathbb{R}^{n+1}: \sum x_i = 1, x_i \geq 0\}$ .

<sup>11</sup>If  $G$  is a discrete group this is literally correct. If  $G$  is a topological group we have to modify the domain of geometric realization to be  $| - |: \text{Fun}(\Delta^{\text{op}}, \text{Top}) \rightarrow \text{Top}$  in order to get the correct definition.

<sup>12</sup>This is in [Wei94, 8.3.11].

1. A priori we should be careful calling it a homotopy, since it doesn't require us to have any notion of a model structure or higher-categorical structure in order to state, however we will see this won't be a problem.
2. In the context of simplicial sets, this coincides with [Definition 1.2.19](#), as we can check (see e.g. [\[Wei94, 8.3.12\]](#)).
3. It is not true (even in simplicial sets) that this is an equivalence relation, and indeed it isn't. Nevertheless it generates one, so when we say “up to simplicial homotopy” we often mean with respect to the transitive closure of this relation.

**Theorem 1.2.21.** Geometric realization preserves products.

**Corollary 1.2.22.** Simplicial homotopies become honest homotopies of spaces after geometric realization.

**Remark 1.2.23.** We can study simplicial sets up to homotopy or spaces up to homotopy, and in a sense that can be made precise, these are essentially the same theory.

**Remark 1.2.24.** Historically this connection between simplicial sets and spaces was part of a research program which used to be called “combinatorial homotopy theory,” led by Kan in the 1950's.

## 1.2.2 Torsors via simplicial maps

The big takeaway of this entire section is the following:

**Theorem 1.2.25.** Let  $\mathcal{U} = \{U_i \rightarrow X\}$  be a cover, and let  $G$  be a group scheme over  $X$ . To any map from the Čech nerve to the bar construction, we can assign a Čech 1-cocycle

$$\mathrm{Hom}_{\mathrm{Fun}(\Delta^{\mathrm{op}}, \mathrm{Var}_k)}(N(\mathcal{U}), B_\bullet G) \rightarrow H^1(\mathcal{U}, G),$$

and this assignment is surjective.

*Proof.* We'll figure out some what's happening in lower degrees first and use this to see how to define maps in general.

**Low-degree intuition:** To specify a map, we need to know what happens on each level, so let's consider the diagram:

$$\begin{array}{ccc} N(\mathcal{U}) = & \coprod_i U_i \rightrightarrows \coprod_{i,j} U_{ij} & \cdots \\ & \downarrow & \downarrow \\ BG = & * \rightrightarrows G & \cdots \end{array}$$

On 0-simplices, there is no data, and on 1-simplices, we are obtaining elements we will call  $(g_{ij} \in \mathrm{Hom}(U_{ij}, G))$ . For each  $i$ , the degeneracy maps require the following diagram to commute.

$$\begin{array}{ccc} U_i & \xrightarrow{s_0} & U_{ii} \\ \downarrow & & \downarrow g_{ii} \\ * & \xrightarrow{e} & G, \end{array}$$

which tells us that  $g_{ii} = e$ . That's about all we can learn from degeneracies. What happens on 2-cells? Let's consider the inner and outermost face maps  $N_2(U) \rightarrow N_1(U)$ :

$$\begin{array}{ccc} U_{jk} & \xleftarrow{d_0} & U_{ijk} \\ g_{jk} \downarrow & & \downarrow \\ G & \xleftarrow{d_0 = \mathrm{proj}_2} & G \times G \end{array} \qquad \begin{array}{ccc} U_{ij} & \xleftarrow{d_2} & U_{ijk} \\ g_{ij} \downarrow & & \downarrow \\ G & \xleftarrow{d_2 = \mathrm{proj}_1} & G \times G \end{array}$$

We've decorated the face maps on the bar construction to recall whether they are projection or multiplication. By the universal property of the product,  $U_{ijk} \rightarrow G \times G$  is determined by its post-composition with the projections, hence it makes sense to call this map  $(g_{ij}, g_{jk})$ . The  $d_1$  relation gives us

$$\begin{array}{ccc} U_{ik} & \xleftarrow{d_1} & U_{ijk} \\ \downarrow g_{ik} & & \downarrow (g_{ij}, g_{jk}) \\ G & \xleftarrow{d_1 = \text{mult}} & G \times G \end{array}$$

This implies that

$$g_{ik} = g_{ij}g_{jk}.$$

This is *exactly the 1-cocycle condition* from [Definition 1.1.21](#). We claim that the remaining data of the map  $N(\mathcal{U})_\bullet \rightarrow B_\bullet G$  is completely determined by this relation. In particular, every Čech 1-cocycle can be extracted from a map  $N_\bullet(\mathcal{U}) \rightarrow B_\bullet G$  by the above process.

**In general:** It is now clear how to define the map  $N_n \mathcal{U} \rightarrow B_n G$  — its component on  $U_{i_0 \dots i_n}$  is given by the tuples of maps

$$\prod_{j=0}^{n-1} g_{ij_{j+1}} : U_{i_0 \dots i_n} \rightarrow G^{\times n}.$$

This clearly commutes with degeneracies, and commutes with face maps by universal property of the product and by the 1-cocycle relation. We see in fact that this map is well-defined if and only if the 1-cocycle condition holds.  $\square$

**Theorem 1.2.26.** Two 1-cocycles for the cover  $\mathcal{U}$  are cohomologous if and only if they are simplicially homotopic.

*Proof sketch.* Let  $f, g : N(\mathcal{U})_\bullet \rightarrow B_\bullet G$  be two maps corresponding to 1-cocycles  $(f_{ij})$  and  $(g_{ij})$ , respectively. Recall simplicial homotopy requires us to define maps  $H_i^n : N(\mathcal{U})_n \rightarrow B_{n+1} G$  for  $0 \leq i \leq n$  so that  $d_0 H_0 = f_n$ ,  $d_{n+1} H_n = g_n$ , and various other relations hold. Again let's start in low degrees to gain some intuition and then prove the theorem directly.

**Low-degree intuition:** Let's see what  $H_0^0$  looks like:

$$\begin{array}{ccccccc} \coprod_i U_i & \xrightleftharpoons{H_0^0} & \coprod_{i,j} U_{ij} & \xrightleftharpoons{H_0^1, H_1^1} & \coprod_{i,j,k} U_{ijk} & \cdots \\ \downarrow & \searrow & \downarrow & \searrow & \downarrow & \searrow \\ * & \xrightleftharpoons{H_0^0} & G & \xrightleftharpoons{H_0^1, H_1^1} & G \times G & \cdots \end{array}$$

At level zero, the map  $H_0^0$  specifies an element  $\alpha_i \in G(U_i)$ . Let's see what the degeneracy relations tell us at level one:

$$\begin{array}{ccc} \coprod_i U_i & \xleftarrow{d^0} \coprod_{i,j} U_{ij} & \\ \downarrow H_0^0 & \searrow & \downarrow H_0^1, H_1^1 \\ G & \xrightleftharpoons[d_2]{d_0} G \times G & \cdots \end{array}$$

Since  $H_i^1$  is mapping into a product, it is determined by its projections, which are post-composing with  $d_2$  and  $d_0$ , respectively. From the relations and the simplicial homotopy condition, we get

$$\begin{aligned} d_0 H_0^1 &= f_{ij} \\ d_2 H_0^1 &= H_0^0 d_1 = \alpha_i, \end{aligned}$$

so  $H_0^1$  is the tuple  $(\alpha_i, f_{ij}): U_{ij} \rightarrow G \times G$ .<sup>13</sup> Similarly we compute

$$\begin{aligned} d_0 H_1^1 &= H_0^0 d_0 = \alpha_j \\ d_2 H_1^1 &= g_{ij}, \end{aligned}$$

so  $H_1^1 = (g_{ij}, \alpha_j): U_{ij} \rightarrow G \times G$ . The remaining relation states that

$$d_1 H_1^1 = d_1 H_0^1,$$

and since  $d_1$  is the multiplication  $G \times G \rightarrow G$ , this tells us that  $\alpha_i f_{ij} = g_{ij} \gamma_j$ , which is the coboundary condition.

**Exercise:** Show that, as maps  $U_{i_0 i_1 i_2} \rightarrow G \times G \times G$ , we have

$$\begin{aligned} H_0^2 &= (\alpha_{i_0}, f_{i_0 i_1}, f_{i_1 i_2}) \\ H_1^2 &= (g_{i_0 i_1}, \alpha_{i_1}, f_{i_1 i_2}) \\ H_2^2 &= (g_{i_0 i_1}, g_{i_1 i_2}, \alpha_{i_2}). \end{aligned}$$

**In general:** We define

$$H_j^n: U_{i_0 i_1 \dots i_n} \rightarrow G^{\times(n+1)}$$

by

$$(g_{i_0 i_1}, \dots, g_{i_{j-1} i_j}, \alpha_{i_j}, f_{i_j i_{j+1}}, \dots, f_{i_{n-1} i_n}).$$

We verify that the relevant relations hold if and only if the coboundary condition  $\alpha_i f_{ij} = g_{ij} \alpha_j$  holds for every  $i, j$ .  $\square$

**Motivation 1.2.27.** Last week we asked for a universal space  $BG$  so that homotopy classes of maps  $X \rightarrow BG$  classifies  $G$ -torsors, and we're getting close! We now have a fantastic candidate for  $BG$ , namely the bar construction  $B_\bullet G$  above. However we (1) didn't have a notion of homotopy of maps to witness two torsors being equivalent, and (2) we were mapping from  $N(\mathcal{U})$ , *not* from  $X$ . To that end, let's write down what we're looking for.

**Wishlist:** We want some nice category  $\mathcal{C}$  where both  $X$  and  $B_\bullet G$  live (so our category should contain simplicial varieties), and we want our category to have a notion of *equivalence* with the following properties:

1. Two maps  $N(\mathcal{U}) \rightarrow B_\bullet G$  are homotopic in  $\mathcal{C}$  if and only if they classify cohomologous  $G$ -torsors. We want to make this notion of homotopy precise.
2. In  $\mathcal{C}$ , we have that  $X \simeq N(\mathcal{U})$ , i.e. a variety is equivalent to the Čech nerve of any cover over it, so that up to homotopy, we can classify torsors via maps  $X \rightarrow B_\bullet G$ .

Let's preview how this is going to go — the category  $\text{Var}_k$  of  $k$ -varieties doesn't support a nice homotopy theory, nor does simplicial varieties  $\text{Fun}(\Delta^{\text{op}}, \text{Var}_k)$ . So what we could do instead is replace each variety by its representable presheaf (functor of points, for the algebraic geometers). The Yoneda embedding  $\text{Var}_k \rightarrow \text{Fun}(\text{Var}_k^{\text{op}}, \text{Set})$  then induces a functor

$$\text{Fun}(\Delta^{\text{op}}, \text{Var}_k) \rightarrow \text{Fun}(\Delta^{\text{op}}, \text{Fun}(\text{Var}_k^{\text{op}}, \text{Set})).$$

Doing some adjoint business, we get that this latter category is equivalent to

$$\text{Fun}(\text{Var}_k^{\text{op}}, \text{Fun}(\Delta^{\text{op}}, \text{Set})) = \text{Fun}(\text{Var}_k^{\text{op}}, \text{sSet}).$$

So altogether we get a Yoneda embedding

$$\text{Var}_k \hookrightarrow \text{Fun}(\text{Var}_k^{\text{op}}, \text{sSet}).$$

This latter category is called the category of *simplicial presheaves*, or in higher category language it is just denoted  $\text{PSh}(\text{Var}_k)$ , and called the  $\infty$ -category of  $(\infty)$ -presheaves. It comes with a notion of homotopy coming from simplicial homotopy theory. We'll see that this was not only a nice

<sup>13</sup>There's abuse of notation here with  $\alpha_i$ .

well-behaved way to access homotopy theory starting from  $\text{Var}_k$ , it was actually the *universal* way to do this (see e.g. [Dug01]).

This category  $\text{PSh}(\text{Var}_k)$  will take care of item (1) on our checklist! After we define what homotopy means there, homotopic maps in this category will give isomorphic torsors.

This doesn't take care of point (2), and we shouldn't really expect it to. The equivalence between a variety and the Čech nerve of the cover should depend on *what covers are permissible*. In other words, it should bake in the Grothendieck topology somehow. We'll see that we can get point (2) (while retaining a notion of homotopy compatible with that of presheaves) by passing to the category of *sheaves* in our site. We'll make all this precise soon.

### 1.3 Torsors III: Sites, sheaves, and Hilbert 90

We have seen that  $H^1(X, G)$  classifies  $G$ -torsors for  $X$ , but we've been a little vague about Grothendieck topologies (i.e. what kinds of covers are we considering for our varieties). Today we'll make things more precise.

**Definition 1.3.1.** Let  $\mathcal{C}$  be a category with pullbacks. Then a *Grothendieck topology*  $\tau$  consists of collections of morphisms  $\{U_i \rightarrow X\}_{i \in I}$  in  $\mathcal{C}$  called *covers* or *coverings*, satisfying the following properties:

1. *Closure under pullbacks:* If  $\{U_i \rightarrow X\}_{i \in I}$  is a covering and  $f: Y \rightarrow X$  is any morphism, then the collection of base change morphisms  $\{U_i \times_X Y \rightarrow Y\}_{i \in I}$  is a covering.
2. *Closure under refinement:* If  $\{U_i \in X\}$  is a covering and  $\{V_{ij} \rightarrow U_i\}_j$  is a covering for each  $i$ , then the composite  $\{V_{ij} \rightarrow U_i \rightarrow X\}_{i,j}$  is a covering.
3. *Isomorphisms:* Any isomorphism  $f: Y \xrightarrow{\sim} X$  gives a one-element cover  $\{Y \rightarrow X\}$ .

A pair of a category and a topology  $(\mathcal{C}, \tau)$  is called a *site*.

**Example 1.3.2.** A *Zariski cover* is a collection of open immersions  $\{U_i \rightarrow U\}$  which are jointly surjective. This generates the *Zariski site* on  $\text{Sch}_S$ .

**Example 1.3.3.** An *étale cover* is a collection of étale morphisms, jointly surjective.

**Example 1.3.4.** A *Nisnevich cover* is a collection of étale morphisms  $\{U_i \rightarrow U\}$  so that for each  $x \in X$  there exists an  $i$  and a  $y \in U_i$  so that  $y \mapsto x$  induces an isomorphism on residue fields.

**Note 1.3.5.** If  $R \rightarrow S$  is an étale ring extension, then  $\text{Spec}(S) \rightarrow \text{Spec}(R)$  is an étale cover but not necessarily a Nisnevich cover. For example  $\text{Spec}(L) \rightarrow \text{Spec}(k)$  is étale if  $L/k$  is a finite separable extension, but not Nisnevich unless  $k = L$ .

**Example 1.3.6.** [Stacks, 021M] An *fppf cover* is a jointly surjective collection of morphisms  $\{U_i \xrightarrow{f_i} U\}$  so that each  $f_i$  is flat and locally of finite presentation.

**Example 1.3.7.** [Stacks, 03NW] An *fpqc cover* is a jointly surjective collection of morphisms  $\{U_i \xrightarrow{f_i} U\}$  so that  $\coprod_i U_i \rightarrow U$  is faithfully flat and quasi-compact.

**Example 1.3.8.** A map  $\text{Spec}(S) \rightarrow \text{Spec}(R)$  is an fpqc cover if and only if the ring map  $R \rightarrow S$  is faithfully flat.

**Terminology 1.3.9.** If  $t$  and  $\tau$  are two Grothendieck topologies, we write  $t \leq \tau$  if every  $t$ -cover is a  $\tau$ -cover. We say  $t$  is *coarser* than  $\tau$  or that  $\tau$  *refines*  $t$  in this case.

**Proposition 1.3.10.** We have that

$$\text{Zar} \leq \text{Nis} \leq \text{et} \leq \text{fppf} \leq \text{fpqc}.$$

For more topologies see [Belmans: topologies comparison](#).

**Question 1.3.11.** If  $t \leq \tau$  then what is the difference between  $t$ -torsors and  $\tau$ -torsors?

Let's break this into two parts

### 1.3.1 Torsors in finer topologies

**Proposition 1.3.12.** Let  $\mathcal{C}$  be some fixed category of schemes, with topologies  $t$  and  $\tau$ . Let  $\mathcal{G}$  be a sheaf of groups in the  $\tau$ -topology. Let  $\mathcal{F}$  be a  $\mathcal{G}$ -torsor in the  $t$ -topology. Then  $\mathcal{F}$  is also a  $\tau$ -torsor if  $\mathcal{F}$  is a  $\tau$ -sheaf.

*Proof.* The only thing that could fail is the sheaf condition in the definition. If  $\mathcal{F}$  is a  $t$ -torsor, then there exists a  $t$ -cover  $\{U_i \rightarrow U\}$  trivializing  $\mathcal{F}$ , and this is also a  $\tau$ -cover since  $\tau$  refines  $t$ .  $\square$

We want to make some guarantee that  $\mathcal{G}$  will still be a sheaf in a finer topology. In our cases, we care about the case where  $\mathcal{G}$  is representable, and a classical result guarantees this for us.

**Theorem 1.3.13.** (Grothendieck, [Stacks, 023Q]) Every representable presheaf is a sheaf in the fpqc topology (and hence in any coarser topology, e.g. Zariski, Nisnevich, étale, syntomic, fppf).

**Corollary 1.3.14.** Let  $G$  be an affine group scheme, and let  $t \leq \tau \leq \text{fppf}$ . Then every  $t$ -torsor is a  $\tau$ -torsor.

*Proof.* Let  $\mathcal{F}$  be a  $G$ -torsor. By [Proposition 1.1.30](#) it is representable, hence it is a sheaf in the  $\tau$  topology as well by [Theorem 1.3.13](#). Therefore by [Proposition 1.3.12](#) it is also a  $\tau$ -torsor.  $\square$

### 1.3.2 Torsors in coarser topologies

Now we're interested in the reverse question — when  $t \leq \tau$ , when is a  $\tau$ -torsor a  $t$ -torsor? The only thing that could fail is condition (2) of [Definition 1.1.6](#), so we get the following answer/definition.

**Definition 1.3.15.** Let  $t \leq \tau$  be topologies on a fixed category of schemes  $\mathcal{C}$ , let  $\mathcal{G}$  be a  $\tau$ -sheaf of groups and let  $\mathcal{F}$  be a  $\mathcal{G}$ -torsor. Then  $\mathcal{F}$  is a  $t$ -torsor if and only if there exists a  $t$ -cover over which  $\mathcal{F}$  is trivialized. We say  $\mathcal{F}$  is *locally trivial in the  $t$ -topology*.

We can now leverage some tools from algebraic geometry to prove this.

**Proposition 1.3.16.** Every smooth morphism of schemes admits a section étale-locally.

**Corollary 1.3.17.** If  $G$  is a smooth group scheme, then there is an equivalence of categories between étale  $G$ -torsors and fppf-torsors.

*Proof.* See [Hal, Proposition 4] for a proof.  $\square$

We can summarize what we've learned in the following cheatsheet:

**Cheatsheet: topologies**

Let  $t \leq \tau$  be two covers, we say that  $\tau$  *refines*  $t$ , or that  $t$  is *coarser* than  $\tau$ . What this means is that

1. Every  $t$ -cover is a  $\tau$ -cover
2. Every  $\tau$ -sheaf is a  $t$ -sheaf
3. If  $\mathcal{F}$  is a  $t$ -torsor then it is a  $\tau$ -torsor if it is a  $\tau$ -sheaf
4. If  $\mathcal{F}$  is a  $\tau$ -torsor it is a  $t$ -torsor if and only if it is  $t$ -locally trivial.

### 1.3.3 Special algebraic groups

**Definition 1.3.18.** We say an algebraic group (a group object in varieties) over a field  $k$  is *linear* if it admits a faithful finite-dimensional representation (c.f. [Mil15, p. 72]).

**Note 1.3.19.** If an algebraic group is linear, it is automatically affine ([Mil15, 1.29]) and finite type. Also, all linear algebraic groups are closed subgroups of  $\mathrm{GL}_n$ .

**Proposition 1.3.20.** Every affine group scheme of finite type over a field  $k$  is linear ([Mil15, 4.8]).

**Definition 1.3.21.** [Gro, p. 5-11] A group scheme is called *special* if it is a linear algebraic group  $G$  with the property that every  $G$ -torsor<sup>14</sup> is locally trivial in the Zariski topology.

**Theorem 1.3.22.** [Gro, Théorème 3] The group  $\mathrm{GL}_n$  is special.

**Corollary 1.3.23.** [Mil13, 11.4] There is a natural bijection

$$H_{\mathrm{Zar}}^1(X, \mathrm{GL}_n) \leftrightarrow H_{\mathrm{Nis}}^1(X, \mathrm{GL}_n) \leftrightarrow H_{\mathrm{et}}^1(X, \mathrm{GL}_n) \leftrightarrow H_{\mathrm{fpqc}}^1(X, \mathrm{GL}_n).$$

Hence any of the groups above can be thought of as parametrizing isomorphism classes of *algebraic vector bundles* on  $X$ .

---

<sup>14</sup>Since  $G$  is affine, every  $G$ -torsor in  $\tau \leq \mathrm{fpf}$  is automatically a principal  $G$ -bundle.



# Chapter 2

## Higher categories

### Contents

---

<b>2.1 Infinity categories</b>	<b>25</b>
2.1.1 Homotopy in an $\infty$ -category	28
2.1.2 Mapping spaces	28
2.1.3 Presentable $\infty$ -categories	30
2.1.4 Localization	31
<b>2.2 Descent</b>	<b>32</b>
2.2.1 Descent, higher categorically	32
2.2.2 cd-structures	33
2.2.3 Sheafification and accessible localizations	34
2.2.4 About the sheaf topos	35
2.2.5 Slice categories	35
2.2.6 Connectivity	36
2.2.7 Whitehead's theorem	37
2.2.8 Equivalences of sheaves and presheaves	38
2.2.9 Torsors revisited	39
<b>2.3 Bonus: more category theory</b>	<b>40</b>
2.3.1 Cofinality	40
2.3.2 Essential smallness	43

---

### 2.1 Infinity categories

**Example 2.1.1.** If  $\mathcal{C}$  is a small 1-category, it gives rise to a simplicial set  $N_{\bullet}\mathcal{C}$ , called the *nerve* of  $\mathcal{C}$ , with the following data:

- ▷ 0-simplices = objects of  $\mathcal{C}$
- ▷ 1-simplices = morphisms in  $\mathcal{C}$
- ▷ 2-simplices = pairs of composable morphisms  $x \xrightarrow{f} y \xrightarrow{g} z$  in  $\mathcal{C}$
- ▷  $\vdots$
- ▷  $n$ -simplices = strings of  $n$ -composable morphisms

Here the degeneracy maps  $(N\mathcal{C})_n \rightarrow (N\mathcal{C})_{n+1}$  insert an identity, while the face maps  $(N\mathcal{C})_n \rightarrow (N\mathcal{C})_{n-1}$  compose maps. Observe that in  $\mathcal{C}$ , composition happens *strictly*, by which we mean there is no notion of homotopy between maps — if  $x \xrightarrow{f} y \xrightarrow{g} z$  is a composite of maps, and  $h: x \rightarrow z$ , then either  $h = g \circ f$ , or it is not equal, and this is encoded by the data of a *unique* 2-cell:

$$\begin{array}{ccc} & y & \\ f \nearrow & & \searrow g \\ x & \xrightarrow{h} & z. \end{array}$$

We think about this 2-cell as a *witness* for the composition. Note that if no 2-cell exists filling the diagram above, this means that  $h$  is not equal to  $g \circ f$ . It is just some other map from  $x$  to  $z$ .

**Example 2.1.2.** If  $\mathcal{C}$  is a one-object groupoid (e.g. a group  $G$ ), then  $N_\bullet \mathcal{C}$  is the bar construction of  $G$ , and its realization  $B\mathcal{C} := |N_\bullet \mathcal{C}|$  is called the *classifying space* of the category. Some examples:

1.  $BC_2 = \mathbb{RP}^\infty$
2.  $B\mathbb{N} = S^1$
3.  $B\mathbb{Z} = S^1$
4.  $B(\bullet \rightarrow \bullet) = \Delta_{\text{top}}^1$
5.  $B(\bullet \xrightarrow{\sim} \bullet) = S^\infty$
6.  $B\text{PBr}_n = \text{Conf}_n(\mathbb{R}^2)$

**Q:** Given a simplicial set, when can you tell whether it arose as the nerve of a 1-category?

**A:** Given any diagram of the form  $\bullet \rightarrow \bullet \rightarrow \bullet$ , it has to fill in uniquely to a 2-cell. But we also need to fill in composites of three morphisms uniquely (to get a tetrahedron), and composites of four morphisms, and so on. To that end, let  $\Lambda_n^k$  be the simplicial set obtained from  $\Delta^n$  by deleting the  $k$ th face. This is called a *horn*.

This isn't a definition of a horn — we might instead characterize the horns by their representable functors, i.e.  $\text{Hom}_{\text{sSet}}(\Lambda_2^1, -)$  represents the set of “composable” edges  $x \xrightarrow{f} y \xrightarrow{g} z$  in any simplicial set.

**Proposition 2.1.3.** A simplicial set  $X_\bullet$  is the nerve of a 1-category if and only if it admits *unique inner horn filling*, meaning for every  $n$  and every  $0 < k < n$ , given any map  $\Lambda_n^k \rightarrow X_\bullet$  it admits a unique lift:

$$\begin{array}{ccc} \Lambda_n^k & \longrightarrow & X_\bullet \\ \downarrow & \nearrow \text{dashed} & \\ \Delta^n & & \end{array}$$

**Example 2.1.4.** Let  $X$  be a topological space. Then it gives rise to a simplicial set called its *fundamental  $\infty$ -groupoid*  $\Pi_\infty X$ , with the data

- ▷ 0-simplices = points  $x \in X$
- ▷ 1-simplices = paths  $x$  to  $y$  in  $X$
- ▷ 2-simplices = homotopies between paths
- ▷ 3-simplices = homotopies between homotopies between paths
- ⋮

Note that a 2-cell is no longer unique! There can be many homotopies between paths. In particular composition of paths isn't well-defined, in the sense that many paths can function naturally as a composite. We might define  $g \circ f$  to be any path together with a 2-cell making the diagram

commute:

$$\begin{array}{ccc} & y & \\ f \nearrow & & \searrow g \\ x & \dashrightarrow & z. \end{array}$$

In order to specify a composite now, we need to give the data not only of the 1-cell but also of the 2-cell! This is the vibe of higher-categorical composition. Note that horns don't fill uniquely here.

**Exercise 2.1.5.** If you're familiar with the singular chains construction

$$|-| : \mathbf{sSet} \hookrightarrow \mathbf{Top} : \mathbf{Sing}(-),$$

convince yourself that  $\mathbf{Sing}(-)$  is the same as  $\Pi_\infty(-)$ .

**Definition 2.1.6.** A *quasicategory* is any simplicial set with (not necessarily unique) inner horn filling. We denote by  $\mathbf{qCat} \subseteq \mathbf{sSet}$  the full subcategory on the quasi-categories.

A natural question to ask is to what extent there is ambiguity in composition – how many choices do we have for horn filling? Do different choices *mean* different things? The following proposition answers this to some extent.

**Proposition 2.1.7.** (Joyal) If  $\mathcal{C}$  is a quasi-category, then the map of simplicial sets

$$\mathbf{Fun}(\Delta^2, \mathcal{C}) \rightarrow \mathbf{Fun}(\Delta_2^1, \mathcal{C})$$

has contractible fibers (i.e., the geometric realization of the fibers under this map are contractible spaces).<sup>1</sup>

**Definition 2.1.8.** A *Kan complex* is a quasi-category which also has *outer horn filling*, meaning we have a lift

$$\begin{array}{ccc} \Lambda_n^k & \longrightarrow & X_\bullet \\ \downarrow & \nearrow \text{dashed} & \\ \Delta^n & & \end{array}$$

not only for  $0 < k < n$ , but also for  $k = 0, n$ . For  $n = 2$ , this means we are also allowed to fill the horns:



**Exercise 2.1.9.** Show a Kan complex is the nerve of a 1-groupoid if and only if its inner horn filling is unique.

**The homotopy hypothesis:** The functor

$$\begin{aligned} (\mathbf{Top}, \text{weak equiv}) &\rightarrow (\mathbf{Kan}, \text{weak equiv}) \\ X &\mapsto \Pi_\infty(X) \end{aligned}$$

yields an equivalence of  $\infty$ -categories.<sup>2</sup> Hence we can think about spaces as Kan complexes without much loss of generality. We use  $\mathcal{S}$  to denote the  $\infty$ -category of spaces.

<sup>1</sup>In fact  $\mathcal{C}$  is a quasi-category if and only if this holds.

<sup>2</sup>We haven't defined what we mean by  $\mathbf{Top}$  or  $\mathbf{Kan}$  as an  $\infty$ -category, and it's a bit subtle. We want to incorporate the weak equivalences, so really we should take the hammock localization of  $\mathbf{Kan}$  at the simplicial weak equivalences, then take its homotopy coherent nerve, but fibrantly replace  $L^W \mathbf{Kan}$  first before taking  $N_\Delta$  so that the resulting simplicial set is an honest quasi-category. An analogous procedure should be carried out with topological spaces, assuming we work with all spaces and not just CW complexes.

**Remark 2.1.10.** (On other models of  $\infty$ -categories) There are a ton of different models of  $\infty$ -categories, we stick with quasi-categories since they have become the standard to some extent. The main advantage is that functors are much easier to describe – they are just maps of the underlying simplicial sets. Functor  $\infty$ -categories  $\mathrm{Fun}(\mathcal{C}, \mathcal{D})$  between two quasi-categories are simply given by the internal hom in simplicial sets.

**Proposition 2.1.11.** If  $\mathcal{C}$  is a quasi-category and  $S$  is any simplicial set, then the internal hom  $\mathrm{Fun}(S, \mathcal{C})$  is also a quasi-category. We think about this as “ $S$ -shaped diagrams in  $\mathcal{C}$ .”

### 2.1.1 Homotopy in an $\infty$ -category

**Definition 2.1.12.** Let  $\mathcal{C}$  be a quasi-category. We define its *homotopy category*, denoted  $h\mathcal{C}$  to be the category freely generated by the 1-truncation  $\tau_{\leq 1}\mathcal{C}$  (i.e. objects and edges), modulo the relations coming from 2-simplices.

**Definition 2.1.13.** We say that a morphism  $f: x \rightarrow y$  in a quasi-category  $\mathcal{C}$  is an *isomorphism/equivalence* if there exists some  $g: y \rightarrow x$  so that  $[gf] = \mathrm{id}_x$  and  $[fg] = \mathrm{id}_y$  in  $h\mathcal{C}$ . Note that  $g$  is *not uniquely defined*, unlike in ordinary 1-category theory.

**Remark 2.1.14.** We should think about this less like isomorphisms in 1-categories, and more like homotopy equivalences in topology.

**Example 2.1.15.** A morphism  $f: X \rightarrow Y$  between CW complexes is a (weak) homotopy equivalence if and only if  $[f]$  is an isomorphism in  $h\mathrm{Top}$ . Hence the “isomorphisms” in the  $\infty$ -category of spaces are not homeomorphisms but rather homotopy equivalences.

**Proposition 2.1.16.** There is an adjunction<sup>3</sup>

$$h: \mathrm{qCat} \rightleftarrows \mathrm{Cat} : N.$$

**Remark 2.1.17.** We recall that the nerve is fully faithful. This is equivalent to the counit of the adjunction being a natural isomorphism:

$$hN\mathcal{C} \xrightarrow{\sim} \mathcal{C},$$

in other words any 1-category can be recovered as the homotopy category of its nerve.

**Example 2.1.18.** Let  $R$  be any ring. Then its category of chain complexes  $\mathrm{Ch}(R)$  is naturally an  $\infty$ -category, and the notion of homotopy recovers the idea of chain homotopy equivalence. We have to be careful constructing this explicitly, refer to §13 of DAG for more info.

### 2.1.2 Mapping spaces

We want to make precise the model of quasi-categories as  $(\infty, 1)$ -categories. The vibe of higher categories is that homs in 1-categories are 0-categories (sets). Homs in 2-categories are 1-categories, homs in 3-categories are 2-categories, etc. Hence homs in  $(\infty, 1)$ -categories should be  $(\infty, 0)$ -categories. From the models we’re working in:

<sup>3</sup>If we consider the codomain of the nerve construction to be all simplicial sets, it still admits a left adjoint called the homotopy category, however it is not given by the formula in [Definition 2.1.12](#). Technically there is a more general construction  $h: \mathrm{sSet} \rightarrow \mathrm{Cat}$ , which is left adjoint to  $N$ , and which agrees with [Definition 2.1.12](#), which we should call  $\tau_{\leq 1}$ , when the simplicial set is a quasi-category. We’ll only apply the homotopy category construction to quasi-categories here so this distinction won’t matter.

$$\begin{aligned} (\infty, 1)\text{-categories} &= \text{quasi-categories} \\ (\infty, 0)\text{-categories} &= \text{Kan complexes,} \end{aligned}$$

hence we want to argue that, for any quasicategory  $\mathcal{C}$ , and any pair of objects (0-simplices)  $x, y \in \mathcal{C}$ , there is a mapping space  $\text{Map}_{\mathcal{C}}(x, y)$  which is a Kan complex.

**Definition 2.1.19.** For  $x, y \in \mathcal{C}$ , where  $\mathcal{C}$  is a quasicategory, we denote by  $\text{Map}_{\mathcal{C}}(x, y)$  the pullback in simplicial sets:

$$\begin{array}{ccc} \text{Map}_{\mathcal{C}}(x, y) & \longrightarrow & \text{Fun}(\Delta^1, \mathcal{C}) \\ \downarrow & \lrcorner & \downarrow (\text{ev}_0, \text{ev}_1) \\ \{x, y\} & \longrightarrow & \mathcal{C} \times \mathcal{C}. \end{array}$$

Here  $\text{Fun}(\Delta^1, \mathcal{C})$  denotes an internal hom from the interval  $\Delta^1$  to  $\mathcal{C}$ . The rightmost vertical map is what's called a *bifibration* (the proof that this map is a bifibration is [Lur09, 2.4.7.11]), which in particular means that  $\text{Map}_{\mathcal{C}}(x, y)$  is a Kan complex.

**Intuition 2.1.20.** The mapping space is intended to generalize the idea of the homotopy category, in the sense that

$$\pi_0 \text{Map}_{\mathcal{C}}(x, y) = \text{Hom}_{h\mathcal{C}}(x, y).$$

In particular its connected components correspond to homotopy classes of maps between  $x$  and  $y$ , but it remembers more information about *how* the homotopies were witnessed, encoded in the higher homotopy type of  $\text{Map}_{\mathcal{C}}(x, y)$ .

**Remark 2.1.21.** Alternatively we may define  $\text{Map}_{\mathcal{C}}(x, y)$  as the simplicial set whose  $n$ -simplices are given by the set of all

$$z: \Delta^{n+1} \rightarrow \mathcal{C},$$

with the  $\{n+1\}$ -vertex mapping to  $y$ , and the vertices  $\{0, \dots, n\}$  mapping to  $x$ . Technically speaking this is the space of *right morphisms* but when  $\mathcal{C}$  is an  $\infty$ -category this models the mapping space (it is canonically isomorphic in the homotopy category). As an exercise, verify that  $\text{Map}_{\mathcal{C}}(x, y)$  is indeed a Kan complex from the definition.

What is  $\text{Map}_{\mathcal{C}}(x, y)$  intended to capture? Its 0-simplices are maps from  $x$  to  $y$  *in the homotopy category*  $h\mathcal{C}$ . In other words, they are equivalence classes of zig-zags of morphisms in  $\mathcal{C}$  from  $x$  to  $y$ , where maps going the wrong way are all invertible.

**Notation 2.1.22.** If  $\mathcal{C}$  is an  $\infty$ -category and  $x, y \in \mathcal{C}$ , we denote by

$$[x, y] := \pi_0 \text{Map}_{\mathcal{C}}(x, y).$$

We call this *homotopy classes of maps* from  $x$  to  $y$ .

**Warning 2.1.23.** It is not true that  $[x, y]_{\mathcal{C}}$  is simply the edges from  $x$  to  $y$  in the quasi-category  $\mathcal{C}$  modulo an equivalence relation, it is more subtle. In the presence of a model structure, we can replace  $x$  and  $y$  by equivalent objects  $Qx$  and  $Ry$  respectively, so that  $[x, y] \cong [Qx, Ry]$ , and this latter set can be literally identified with the 1-cells  $Qx \rightarrow Ry$  in  $\mathcal{C}$  modulo an explicit equivalence relation. We'll come back to this when we talk about sheaves.

**Example 2.1.24.** If  $\mathcal{C}$  is a 1-category, we can view it trivially as an  $\infty$ -category via the nerve construction. In this case  $\text{Map}_{\mathcal{C}}(x, y) = \text{Hom}_{\mathcal{C}}(x, y)$  is just a set (a discrete simplicial set). The homotopy category of  $\mathcal{C}$  is just  $\mathcal{C}$ , because there are no equivalences which aren't isomorphisms.

### 2.1.3 Presentable $\infty$ -categories

Modulo some set-theoretic technicalities, we can now be content with the existence of a model for infinity-categories. All notions of functors, colimits, adjunctions, etc. should now be understood in the higher categorical sense, i.e. up to higher coherence.

**Definition 2.1.25.** [Lur09, 5.4.2.1] An  $\infty$ -category is *accessible* if it is generated under  $\kappa$ -filtered colimits by a small category.

**Example 2.1.26.** The category  $\mathcal{S}$  of spaces is accessible, since it admits all colimits and every space is built out of finite CW complexes.

**Remark 2.1.27.** By [Lur09, 5.4.3.6], a small  $\infty$ -cat is accessible if and only if it is idempotent complete.<sup>4</sup> Observe that the category of finitely generated free  $R$ -modules fail to contain retracts (projectives) so they're not idempotent complete and hence not accessible.

**Definition 2.1.28.** Given any  $\infty$ -category  $\mathcal{C}$ , we denote by  $\mathrm{PSh}(\mathcal{C}) := \mathrm{Fun}(\mathcal{C}^{\mathrm{op}}, \mathcal{S})$  the category of  $(\infty)$ -presheaves.

**Example 2.1.29.** We can think about presheaves of simplicial sets  $\mathrm{Fun}(\mathrm{Sch}_{\mathcal{S}}^{\mathrm{op}}, \mathrm{sSet})$  as the presheaf category  $\mathrm{PSh}(\mathrm{Sch}_{\mathcal{S}})$ . Note what lives in here:

1. Schemes all live in here via the Yoneda embedding  $\mathrm{Sch}_{\mathcal{S}} \hookrightarrow \mathrm{Fun}(\mathrm{Sch}_{\mathcal{S}}^{\mathrm{op}}, \mathrm{Set}) \subseteq \mathrm{Fun}(\mathrm{Sch}_{\mathcal{S}}^{\mathrm{op}}, \mathrm{sSet})$ , by viewing  $\mathrm{Set} \subseteq \mathrm{sSet}$  as discrete simplicial sets (no non-degenerate  $n$ -simplices for  $n \geq 1$ ).
2. *Simplicial schemes* also live in here, by moving some adjoint stuff around:  

$$\mathrm{Fun}(\Delta^{\mathrm{op}}, \mathrm{Sch}_{\mathcal{S}}) \xrightarrow{y} \mathrm{Fun}(\Delta^{\mathrm{op}}, \mathrm{Fun}(\mathrm{Sch}_{\mathcal{S}}^{\mathrm{op}}, \mathrm{Set})) \cong \mathrm{Fun}(\mathrm{Sch}_{\mathcal{S}}^{\mathrm{op}}, \mathrm{Fun}(\Delta^{\mathrm{op}}, \mathrm{Set})) = \mathrm{PSh}(\mathrm{Sch}_{\mathcal{S}}).$$
3. Spaces (viewed as simplicial sets by  $\Pi_{\infty}$ ) live in here as constant presheaves  $\mathrm{sSet} \hookrightarrow \mathrm{Fun}(\mathrm{Sch}_{\mathcal{S}}^{\mathrm{op}}, \mathrm{sSet})$ .

Thus we have a natural home for schemes and spaces, as well as these simplicial scheme data types we've been looking at.

**Definition 2.1.30.** We say an  $\infty$ -category  $\mathcal{C}$  is *presentable* if it is accessible and admits all colimits (cocomplete).

**Example 2.1.31.** By the previous two examples,  $\mathrm{PSh}(\mathcal{C})$  is presentable for any  $\mathcal{C}$ . This is the coYoneda lemma — that any presheaf is a colimit of representable ones.

**Theorem 2.1.32.** (*Adjoint functor theorem*) Let  $F: \mathcal{C} \rightarrow \mathcal{D}$  be a functor between presentable categories. Then

- ▷  $F$  admits a right adjoint if and only if  $F$  preserves all colimits
- ▷  $F$  admits a left adjoint if and only if it preserves all limits and  $\kappa$ -filtered colimits

Really hard to write down functors explicitly in quasi-categories, since we are writing down a map of simplicial sets, which is a lot of data. AFT is nice because it lets us get functors without writing them explicitly, but they are still characterized by being adjoints.

**Notation 2.1.33.** We denote by  $\mathrm{Pr}^L$  the category of presentable  $\infty$ -categories and colimit-preserving functors between them. Note every functor in  $\mathrm{Pr}^L$  is a left adjoint.

**Theorem 2.1.34.** Every presentable category is complete (admits all limits).

<sup>4</sup>Idempotent complete has a number of definitions, in particular it implies that idempotent endomorphisms  $f: X \rightarrow X$  (i.e.  $f \circ f = f$ ) correspond bijectively to retracts of  $X$ , i.e. composites  $Y \hookrightarrow X \rightarrow Y$ . If  $\mathcal{C}$  is idempotent complete then it is closed under retracts.

### 2.1.4 Localization

**Definition 2.1.35.** [Lur09, 5.2.7.2] A functor  $f: \mathcal{C} \rightarrow \mathcal{D}$  is a *localization* if it admits a fully faithful right adjoint.

In many cases a localization is given by inverting a class of morphisms in  $\mathcal{C}$ . In particular let  $S \subseteq \text{mor}\mathcal{C}$  be a class of morphisms in  $\mathcal{C}$ , then we can try to *invert*  $S$  by cooking up a new category  $\mathcal{C}[S^{-1}]$ .

**Example 2.1.36.**

1. The *homotopy category* of spaces is obtained from the category of compactly generated weakly Hausdorff spaces by inverting all homotopy equivalences.
2. The *derived category* of a ring is obtained from the category of chain complexes by inverting the chain homotopy equivalences.
3. A group (as a one-object groupoid) is obtained from a monoid by freely inverting each morphism.

**Definition 2.1.37.** [Lur09, 5.5.4.1] Let  $S \subseteq \text{mor}\mathcal{C}$ . We say  $z \in \mathcal{C}$  is *S-local* if for every  $s: x \rightarrow y$  in  $S$ , the induced map

$$\text{Map}_{\mathcal{C}}(y, z) \rightarrow \text{Map}_{\mathcal{C}}(x, z)$$

is an equivalence.

**Remark 2.1.38.** Let  $\mathcal{C}_0 \subseteq \mathcal{C}$  be the full subcategory of  $S$ -local objects. If this admits a left adjoint, it makes sense to call that adjoint  $L_S$ , that is,  $S$ -localization, since it inverts every morphism in  $S$ . *This is where presentable categories give us an advantage.* In general arguing for the existence of a left adjoint isn't easy, however if  $\mathcal{C}$  is presentable, then the adjoint functor theorem tells us that we just have to check the inclusion  $\mathcal{C}_0 \subseteq \mathcal{C}$  preserves limits and filtered colimits.

Let  $\mathcal{C}_0 \subseteq \mathcal{C}$  be the full subcategory of  $S$ -local objects. If this admits a left adjoint, it makes sense to call that adjoint  $L_S$ , that is,  $S$ -localization, since it inverts every morphism in  $S$ .

**Proposition 2.1.39.** [Lur09, 5.5.4.15] If  $\mathcal{C}$  is presentable and  $S \subseteq \text{mor}\mathcal{C}$  is small, then the inclusion of the full subcategory of  $S$ -local objects admits a left adjoint.<sup>5</sup>

**Example 2.1.40.** In the next talk, our primary application of this machinery will be looking at the presheaf category  $\text{PSh}(\mathcal{C})$ , which is presentable by [Example 2.1.31](#). We can look at full subcategories of presheaves which satisfy a certain sheaf condition and argue this is a reflective subcategory hence we will have an adjoint we call *sheafification*.

**Remark 2.1.41.** Given a class of arrows  $S \subseteq \text{mor}\mathcal{C}$ , we can always form  $\mathcal{C}[S^{-1}]$  by adjoining formal inverses to  $S$  and considering all composites of morphisms in  $\mathcal{C}$  and formal inverses (zig-zags). This is called *Dwyer–Kan localization* or *hammock localization*. This satisfies the correct universal property of localization, but we might encounter size issues. Bousfield localization is a particular example of Dwyer–Kan localization, but where we are able to guarantee that we don't encounter any size issues since the localization is a subcategory of the original category.

**Example 2.1.42.** We define  $\mathcal{S}$  to be the Dwyer–Kan localization of the category  $\text{Top}$  of compactly generated weakly Hausdorff spaces at the weak homotopy equivalences. This has the property that  $h\mathcal{S} = \text{Ho}(\text{Top})$ . See [Lur09, §1.2.16] for more information. As a model category we are invited to think about  $\mathcal{S}$  as

---

<sup>5</sup>The terminology for this is that  $\mathcal{C}_0 \subseteq \mathcal{C}$  is a *reflective subcategory*.

1. simplicial sets with the classical model structure
2. Kan complexes with the classical model structure
3. topological spaces with the classical model structure

Hence we think about  $\mathrm{PSh}(\mathcal{C}) = \mathrm{Fun}(\mathcal{C}^{\mathrm{op}}, \mathcal{S})$  as simplicial presheaves, equipped with a *levelwise* notion of weak equivalence, coming from weak homotopy equivalence of simplicial sets.

## 2.2 Descent

**Goal 2.2.1.** Define the  $\infty$ -topos of Nisnevich sheaves  $\mathrm{Sh}_{\mathrm{Nis}}(\mathrm{Sm}_k)$ .

**Assumption 2.2.2.** We will work over a base scheme  $S$  which is qcqs and Noetherian.

### 2.2.1 Descent, higher categorically

Very roughly speaking, a *sheaf* is a presheaf that glues along covers. We're going to give a general definition, then show how it recovers what we know and remember.

**Notation 2.2.3.** Suppose  $\mathcal{U} = \{U_i \rightarrow X\}_i$  is a cover in  $\mathrm{Sch}_S$ , giving rise to a Čech nerve  $\Delta^{\mathrm{op}} \rightarrow \mathrm{Sch}_S$ . Then if  $\mathcal{C}$  is any  $\infty$ -category and  $F: \mathrm{Sch}_S^{\mathrm{op}} \rightarrow \mathcal{C}$  a presheaf, then we denote by  $F(\mathcal{U}) \in \mathrm{Fun}(\Delta, \mathcal{C})$  the cosimplicial object given by applying  $F$  everywhere in the Čech nerve.<sup>6</sup>

**Definition 2.2.4.** Let  $\mathcal{C}$  be an  $\infty$ -category with all limits, let  $(\mathrm{Sch}_S, \tau)$  be a site, and let  $F: \mathrm{Sch}_S^{\mathrm{op}} \rightarrow \mathcal{C}$  be a presheaf valued in  $\mathcal{C}$ . Then we say  $F$  is a  $\tau$ -*sheaf* if for every  $\tau$ -cover  $\mathcal{U} = \{U_i \rightarrow X\}$ , we have that the induced map

$$F(X) \rightarrow \lim_{\Delta} F(\mathcal{U})$$

is an equivalence.

**Example 2.2.5.** If  $\mathcal{C}$  is a 1-category, e.g. sets, then a higher categorical limit just recovers the notion of an ordinary limit, since there is no higher structure. In particular, the limit reduces to seeing that the pair of parallel morphisms  $[0] \rightrightarrows [1]$ , viewed as a subcategory of  $\Delta$ , is *final* (see [Example 2.3.7](#)). In this case the sheaf condition reduces to asking whether the map

$$F(X) \rightarrow \lim \left( \prod_i F(U_i) \rightrightarrows \prod_{i,j} F(U_{ij}) \right)$$

is an equivalence, i.e. it witnesses  $F(X)$  as a 1-categorical limit (in particular, an equalizer).

**Example 2.2.6.** If  $\mathcal{C}$  is an abelian category, then the equalizer of two maps is just the kernel of their difference, so we get the familiar sheaf condition that

$$0 \rightarrow F(X) \rightarrow \prod_i F(U_i) \rightarrow \prod_{i,j} F(U_{ij})$$

is left exact.

**Example 2.2.7.** If  $\mathcal{C}$  is a 2-category, then  $\Delta_{\leq 2}^{\mathrm{inj}} \subseteq \Delta$  is 2-final (reference needed), so we get that the sheaf condition becomes

$$F(X) \rightarrow \lim \left( \prod_i F(U_i) \rightrightarrows \prod_{i,j} F(U_{ij}) \rightrightarrows \prod_{i,j,k} F(U_{ijk}) \right)$$

---

<sup>6</sup>If  $F$  is product-preserving this is immediate, if not we have to apply  $F$  at each level and then take products.



If  $\mathcal{C} = \mathbf{Grpd}$  is the category of groupoids, viewed as an  $\infty$ -subcategory of  $\mathbf{qCat}$  via the nerve construction, then this is precisely the stack condition! So this is what we meant when we said “sheaf of categories” earlier.

**Remark 2.2.8.** These limits are not 1-categorical limits, they are taking place in a higher categorical sense. The following example is worth thinking about as it makes this more concrete.

**Example 2.2.9.** Let  $R$  be a ring, and  $\langle f, g \rangle = R$  two objects generating the unit ideal, giving rise to a two-object cover  $\{\mathrm{Spec}(R_f) \rightarrow \mathrm{Spec}(R), \mathrm{Spec}(R_g) \rightarrow \mathrm{Spec}(R)\}$ .

1. Argue that the stack condition for this particular cover truncates at the double overlaps, since there are no interesting triple overlaps.
2. See that  $\mathcal{F}: \mathbf{Sch}^{\mathrm{op}} \rightarrow \mathbf{Grpd}$  satisfies descent for this cover if and only if

$$\begin{array}{ccc} \mathcal{F}(\mathrm{Spec}(R)) & \longrightarrow & \mathcal{F}(\mathrm{Spec}(R_f)) \\ \downarrow & \lrcorner & \downarrow \\ \mathcal{F}(\mathrm{Spec}(R_g)) & \longrightarrow & \mathcal{F}(\mathrm{Spec}(R_{fg})) \end{array}$$

is a *pullback of groupoids*.

3. As a particular example, show that  $\mathrm{Mod}(-)$  satisfies descent for two-object Zariski covers. That is,  $\mathrm{Mod}(R)$  is equivalent to the 2-categorical pullback, often called the *category of descent data* attached to the cover.

## 2.2.2 cd-structures

We saw in the previous example how a sheaf condition can simplify on covers with fewer objects. A natural question to ask would be whether descent along a small collection of covers implies descent along all covers. A formalism that often lets us deal with this is the idea of a *cd-structure*. We’re also going to zoom in on presheaves of spaces.

**Definition 2.2.10.** A *cd-structure* is a collection of commutative squares in  $\mathcal{C}$  closed under isomorphism:

$$\begin{array}{ccc} A & \longrightarrow & B \\ \downarrow & & \downarrow \\ C & \longrightarrow & D. \end{array}$$

**Terminology 2.2.11.** Given a cd-structure on  $\mathbf{Sch}_S$ , we define its associated topology  $\tau$  to be the coarsest topology for which  $\{B \rightarrow D, C \rightarrow D\}$  is a  $\tau$ -cover for every distinguished square.

**Example 2.2.12.** The *Zariski cd-structure* is defined by commutative squares of the form

$$\begin{array}{ccc} U \cap V & \longrightarrow & U \\ \downarrow & & \downarrow \\ V & \longrightarrow & U \cup V. \end{array}$$

We will call these squares *distinguished Zariski squares*.

**Theorem 2.2.13.** Let  $\mathcal{F} \in \mathbf{PSh}(\mathbf{Sch}_S^{\mathrm{op}})$  be a presheaf. Then  $\mathcal{F}$  is a Zariski sheaf if and only if  $\mathcal{F}(\emptyset) = *$  and  $\mathcal{F}$  sends every distinguished Zariski square to a (homotopy) pullback square.

**Example 2.2.14.** The Nisnevich topology is generated by a cd-structure given by *distinguished Nisnevich squares*, of the form

$$\begin{array}{ccc} U \times_X V & \longrightarrow & V \\ \downarrow & & \downarrow p \\ U & \xrightarrow{i} & X, \end{array}$$

where  $i$  is an open immersion,  $p$  is étale, and  $p$  restricts to an isomorphism  $p^{-1}(X - U) \rightarrow X - U$ .

**Example 2.2.15.** (*Affine distinguished Nisnevich square*). Suppose  $f : R \rightarrow S$  is a finite étale ring homomorphism, and  $h \in R$  is some element for which  $R/x \cong S/f(x)$  is a ring isomorphism. Then we have a distinguished Nisnevich square:

$$\begin{array}{ccc} \mathrm{Spec}(S_{f(x)}) & \longrightarrow & \mathrm{Spec}(S) \\ \downarrow & \lrcorner & \downarrow \\ \mathrm{Spec}(R_f) & \longrightarrow & \mathrm{Spec}(R). \end{array}$$

*Proof.* The right map is étale, the bottom is an open immersion, and the restriction of  $\mathrm{Spec}(S) \rightarrow \mathrm{Spec}(R)$  to the complement of the open distinguished affine  $D(f) \subseteq \mathrm{Spec}(R)$  is the hypothesis that  $R/x \cong S/f(x)$ .  $\square$

**Definition 2.2.16.** We say  $\mathcal{F} \in \mathrm{PSh}(\mathrm{Sms}_S)$  is a *Nisnevich sheaf* if  $\mathcal{F}(\emptyset) = *$  and  $\mathcal{F}$  sends every distinguished Nisnevich square to a pullback square.

**Exercise 2.2.17.** Show that  $\mathrm{Mod}(-)$  is a Nisnevich sheaf on the site of affine schemes.

### 2.2.3 Sheafification and accessible localizations

**Definition 2.2.18.** [Lur09, 5.4.2.5] We say a functor between accessible  $\infty$ -categories is *accessible* if it is  $\kappa$ -continuous (i.e., preserves  $\kappa$ -limits by a regular cardinal  $\kappa$ ).

**Definition 2.2.19.** If  $\mathcal{C} \subseteq \mathcal{D}$ , then we say a localization  $L : \mathcal{D} \rightarrow \mathcal{C}$  is *accessible* if and only if the composite  $\mathcal{D} \xrightarrow{L} \mathcal{C} \hookrightarrow \mathcal{D}$  is accessible. If  $\mathcal{D}$  is an accessible category, this is equivalent to the statement that  $\mathcal{C} \subseteq \mathcal{D}$  is an accessible subcategory [Lur09, 5.5.4.2].

**Definition 2.2.20.** [Lur09, 6.1.0.4] If  $\mathcal{X}$  is an  $\infty$ -category, we say it is an  $\infty$ -topos if there exists a small category  $\mathcal{C}$  and an accessible left exact localization functor  $\mathrm{PSh}(\mathcal{C}) \rightarrow \mathcal{X}$ .

This is some higher categorical analogue of the fact from topos theory that every (Grothendieck) topos is the category of sheaves of sets on a site.

**Proposition 2.2.21.** [Lur09, 6.2.2.7] If  $\mathcal{C}$  is a (small)  $\infty$ -category with a Grothendieck topology, then  $\mathrm{Shv}_\tau(\mathcal{C})$  is an accessible left exact localization of  $\mathrm{P}(\mathcal{C})$ , in particular it is an  $\infty$ -topos.

*Proof.* The precise statement of [Lur09, 6.2.2.7] is that  $\mathrm{Shv}_\tau(\mathcal{C}) \subseteq \mathrm{PSh}(\mathcal{C})$  is a so-called *topological* localization (defined in [Lur09, 6.2.1.4]). In [Lur09, 6.2.1.6] it is proved that every topological localization of a presentable  $\infty$ -category is accessible and left exact.  $\square$

**Terminology 2.2.22.** We refer to the localization functor

$$L_\tau : \mathrm{PSh}(\mathcal{C}) \hookrightarrow \mathrm{Shv}_\tau : i$$

as *sheafification*.

**Proposition 2.2.23.** [Lur09, 6.2.2.17] If  $\mathcal{C}$  is a small  $\infty$ -category there is a bijection between Grothendieck topologies on  $\mathcal{C}$  and (equivalence classes of) topological localizations of  $\mathrm{PSh}(\mathcal{C})$ .

### 2.2.4 About the sheaf topos

**Corollary 2.2.24.** Some consequences:

1. The *sheafification* functor  $L_\tau: \mathrm{PSh}(\mathcal{C}) \rightarrow \mathrm{Shv}_\tau(\mathcal{C})$  preserves all colimits (being a left adjoint) and all small limits (being left exact).
2. The inclusion functor  $i: \mathrm{Shv}_\tau(\mathcal{C}) \hookrightarrow \mathrm{PSh}(\mathcal{C})$  preserves all limits and all filtered colimits, by the adjoint functor theorem.

Explicitly, by this second point, we have that *limits and filtered colimits of sheaves can be computed in the presheaf category*. This is a crucial fact.

**Example 2.2.25.** (Examples of  $\tau$ -sheaves):

1. Any  $\tau$ -sheaf of *sets* is a  $\tau$ -sheaf of discrete spaces. So we have

$$\mathrm{Sh}_\tau(\mathcal{C}; \mathrm{Set}) \subseteq \mathrm{Sh}_\tau(\mathcal{C}).$$

2. If  $\tau$  is subcanonical, then the representable presheaf

$$h_X := \mathrm{Hom}_{\mathcal{C}}(-, X): \mathcal{C}^{\mathrm{op}} \rightarrow \mathrm{Set}$$

is a sheaf of sets, and hence a sheaf of spaces.

3. Consider any presheaf of groupoids

$$\mathcal{F}: \mathcal{C}^{\mathrm{op}} \rightarrow \mathrm{Grpd}.$$

We denote by  $B\mathcal{F}$  the composite  $\mathcal{C}^{\mathrm{op}} \xrightarrow{\mathcal{F}} \mathrm{Grpd} \xrightarrow{N} \mathrm{sSet}$ . Then  $\mathcal{F}$  is a  $\tau$ -stack if and only if  $B\mathcal{F}$  is a  $\tau$ -sheaf [Hol08, 3.9].

4. If  $Y \in \mathcal{S}$  is any space, we may sheafify the constant presheaf valued at  $Y$  in order to obtain a sheaf  $\underline{Y}$ .

**Terminology 2.2.26.** From the canonical equivalence

$$\mathrm{Fun}(\Delta^{\mathrm{op}}, \mathrm{Fun}(\mathcal{C}^{\mathrm{op}}, \mathrm{Set})) \cong \mathrm{Fun}(\mathcal{C}^{\mathrm{op}}, \mathrm{sSet}),$$

simplicial objects in set-valued presheaves naturally give rise to  $\infty$ -categorical presheaves. Consider an object on the left, of the form  $X_\bullet: \Delta^{\mathrm{op}} \rightarrow \mathrm{PSh}(\mathcal{C}; \mathrm{Set})$ . If this functor factors through  $\mathrm{Sh}_\tau(\mathcal{C}; \mathrm{Set}) \subseteq \mathrm{PSh}(\mathcal{C}; \mathrm{Set})$ , that is if  $X_n$  is a sheaf of sets for each  $n$ , we call this a *simplicial object in sheaves*.

**Warning 2.2.27.** (c.f. [Lur09, 7.1.3.1]) In the literature, we can find objects  $X_\bullet$  of the form in **Terminology 2.2.26** referred to as “simplicial sheaves.” This is overloaded terminology, and suggests that  $\infty$ -categorical sheaves are just those presheaves of simplicial sets which are levelwise sheaves of sets, an erroneous claim that can be found throughout the literature.

### 2.2.5 Slice categories

Recall if  $\mathcal{C}$  is any  $(\infty)$ -category and  $x \in \mathcal{C}$ , we have the under and over categories  $\mathcal{C}_{x/}$  and  $\mathcal{C}_{/x}$ , respectively. If  $*$   $\in \mathcal{C}$  is a terminal object, then it becomes both initial and terminal in  $\mathcal{C}_{*/}$ , that is, it is a zero object. We call a category with a zero object *pointed*.

**Proposition 2.2.28.** We have that  $\Delta^0 \in \mathcal{S}$ , viewed as a constant sheaf, is terminal in the sheaf topos  $\mathrm{Shv}_\tau(\mathcal{C})$  for any site.

*Proof.* The inclusion  $\mathrm{Sh}_\tau(\mathcal{C}) \subseteq \mathrm{PSh}(\mathcal{C})$  will preserve terminal objects, being a limit over an empty diagram, so it suffices to observe that the constant presheaf  $\Delta^0$  is already a sheaf.  $\square$

**Proposition 2.2.29.** Let  $h_S \in \text{PSh}(\text{Sm}_S)$  denote the representable sheaf attached to the base. Then the map  $h_S \xrightarrow{!} \Delta^0$  of presheaves is a *local equivalence*, i.e. it sheafifies to an equivalence.

*Proof.* It suffices to observe that the presheaves are identical, which is true because for any  $U \in \text{Sm}_S$ , we have that  $h_S(U) = \text{Hom}_S(U, S)$ , which is a one-object set consisting of the structure map (since  $S$  is terminal in  $\text{Sm}_S$ ).  $\square$

**Notation 2.2.30.** We denote by  $\text{Shv}_\tau(\mathcal{C})_* := \text{Shv}_\tau(\mathcal{C})_{\Delta^0/}$  the pointed slice category. An object here is a map of sheaves  $\Delta^0 \rightarrow F$ , which we observe is equivalent to picking a basepoint in  $F(U)$  for every  $U \in \mathcal{C}$ . Hence we can think of these as sheaves of *pointed* spaces.

### 2.2.6 Connectivity

**Definition 2.2.31.** Let  $(X, x) \in \text{Shv}_\tau(\mathcal{C})_*$  and  $n \geq 0$ . Then we denote by  $\pi_n(X, x)$  the  $n$ th *homotopy sheaf*, defined to be the  $\tau$ -sheafification of the presheaf of sets

$$\begin{aligned} \mathcal{C}^{\text{op}} &\rightarrow \text{Set} \\ U &\mapsto \pi_n(X(U), x). \end{aligned}$$

**Remark 2.2.32.** There is a more intrinsic definition of  $\pi_n$ , leveraging that an  $\infty$ -topos is cotensored over spaces,<sup>7</sup> and considering the map  $X^{S^n} \rightarrow X$  in the slice topos  $\text{Shv}_{/X}$ , and defining  $\pi_n$  to be its 0-truncation.

**Proposition 2.2.33.** We have that

1.  $\pi_0$  is a sheaf of sets
2.  $\pi_1$  is a sheaf of groups
3.  $\pi_n$  is a sheaf of abelian groups for  $n \geq 2$ .

**Notation 2.2.34.** If  $\mathcal{C}$  is any pointed  $\infty$ -category admitting limits, and  $X \in \mathcal{C}$ , we denote by  $\Omega X$  its *loop space*, defined as the pullback

$$\begin{array}{ccc} \Omega X & \longrightarrow & * \\ \downarrow & \lrcorner & \downarrow \\ * & \longrightarrow & X. \end{array}$$

The loop space interacts with the homotopy groups in a topos in the way we might expect from topology:

**Proposition 2.2.35.** If  $\mathcal{C}$  is an  $\infty$ -topos and  $(X, x) \in \mathcal{C}_*$ , then

$$\pi_n(X, x) = \pi_0(\Omega^n(X, x)).$$

*Proof.* (reference needed)  $\square$

**Example 2.2.36.** Let  $\mathcal{F}: \text{Sm}_S^{\text{op}} \rightarrow \text{Set}$  be any sheaf of sets, groups, abelian groups, etc. Then

$$\pi_n(\mathcal{F}) = \begin{cases} \mathcal{F} & n = 0 \\ * & n > 0. \end{cases}$$

<sup>7</sup>This means that for any  $F \in \text{Shv}_\tau(\text{Sch}_S)$  and any  $X \in \text{sSet}$ , we have a natural object  $F^X \in \text{Shv}_\tau(\text{Sch}_S)$  with natural equivalences of mapping spaces

$$\text{Map}_{\text{Shv}}(G, F^X) \cong \text{Map}_{\text{sSet}}(X, \text{Map}_{\text{Shv}}(G, F)).$$

*Proof.* It suffices to observe that the presheaf  $\pi_n \mathcal{F}$  is identical to the constant presheaf  $\Delta^0$  since  $\mathcal{F}(U)$  has no higher homotopy for each  $U \in \mathbf{Sm}_S$ .  $\square$

**Definition 2.2.37.** [Lur09, 6.5.1.10] Let  $\mathcal{C}$  be an  $\infty$ -topos, and take  $X \in \mathcal{C}$ .

1. We say  $X$  is *n-connective* if  $\pi_k(X, x) = *$  for every  $k < n$  and for every basepoint  $x$ .
2. We say that  $X$  is *connected* if it is 1-connective, meaning  $\tau_{\leq 0}X = *$ .
3. We say  $X$  is *n-truncated* if  $\pi_k(X, x) = *$  for all  $k > n$  and for every basepoint  $x$ .

**Proposition 2.2.38.** [Lur09, 5.5.6.18] Let  $\mathcal{C}$  denote a presentable category and  $\tau_{\leq k}\mathcal{C} \subseteq \mathcal{C}$  the full subcategory spanned by  $k$ -truncated objects. Then the inclusion admits an accessible left adjoint, in other words there is a truncation functor  $\tau_{\leq k} : \mathcal{C} \rightarrow \tau_{\leq k}\mathcal{C}$  which is a localization.

In particular, the collection of  $n$ -truncated spaces is closed under limits.

**Definition 2.2.39.** If  $(\mathcal{C}, \tau)$  is any site, then the category  $\tau_{\leq 0}\mathrm{Sh}_\tau(\mathcal{C}) =: \mathrm{Sh}_\tau(\mathcal{C})_{\leq 0}$  is the category of sheaves of sets  $\mathrm{Sh}_\tau(\mathcal{C}; \mathbf{Set})$ .

**Notation 2.2.40.** We write  $\mathrm{Ab}(\mathrm{Sh}_\tau(\mathcal{C})_{\leq 0})$  for the sheaves of abelian groups.

## 2.2.7 Whitehead's theorem

**Theorem 2.2.41.** (Whitehead's theorem) Let  $f : X \rightarrow Y$  be a map of CW complexes. Then  $f$  is a homotopy equivalence if and only if  $f : \pi_i(X) \rightarrow \pi_i(Y)$  is an isomorphism for each  $i \geq 0$ .

A way to say this is that “ $\infty$ -connective morphisms are homotopy equivalences.” The  $\infty$ -categorical analogue of this is the notion of *hypercompleteness*.

**Definition 2.2.42.** An  $\infty$ -topos is *hypercomplete* if and only if every object is  $\infty$ -connective.

**Definition 2.2.43.** [Lur09, 7.2.11] An  $\infty$ -topos has *homotopy dimension*  $\leq n$  if every  $(n - 1)$ -connected object  $X$  receives a map from the terminal object  $* \rightarrow X$ .

**Theorem 2.2.44.** If an  $\infty$ -topos has finite homotopy dimension then it is hypercomplete.

**Warning 2.2.45.** The étale topos  $\mathrm{Shv}_{\mathrm{et}}(\mathbf{Sm}_S)$  need not be hypercomplete, even for nice  $S$  (reference needed).

**Theorem 2.2.46.** (Voevodsky) If  $S$  is qcqs and Noetherian, then the homotopy dimension of  $\mathrm{Shv}_{\mathrm{Nis}}(\mathbf{Sm}_S)$  is bounded above by  $\dim(S)$ . In particular this implies it is hypercomplete.

Truncation and connectivity fit into fiber sequences

$$\tau_{>n} \rightarrow \mathrm{id} \rightarrow \tau_{\leq n},$$

which allow us to form Postnikov towers out of objects of  $\infty$ -topoi:

$$\begin{array}{ccc} & \tau_{\leq n}X & \\ & \downarrow & \\ & \vdots & \\ X & \longrightarrow & \tau_{\leq 0}X \end{array}$$

Hypercompleteness implies *Postnikov completeness* (reference needed) meaning that the induced map

$$X \rightarrow \lim_n \tau_{\leq n} X$$

is an equivalence. That is, the Postnikov towers converge and we can make sense of obstruction theory.

We'll see soon that this is a valuable perspective, e.g. for classifying torsors via homotopy theory. However the obstruction theory won't be very useful in the sheaf topos setting. This is one of the advantages we gain by passing to motivic spaces.

**Upshot 2.2.47.** In a hypercomplete topos, equivalences can be checked on homotopy sheaves.

**Corollary 2.2.48.** Let  $F \rightarrow G$  be a map in the sheaf topos  $\mathrm{Shv}_{\mathrm{Nis}}(\mathrm{Sch}_S)$  (or any hypercomplete topos) where  $S$  is qcqs and Noetherian. Then the following are equivalent:

1. The map  $F \rightarrow G$  is an equivalence in  $\mathrm{Shv}_{\mathrm{Nis}}(\mathrm{Sm}_S)$
2. The induced maps  $\pi_n(F) \rightarrow \pi_n(G)$  are isomorphisms for every  $n \geq 0$ .

So this gives us a way to check equivalence on homotopy groups.

**Notation 2.2.49.** For  $F, G \in \mathrm{Shv}_{\tau}(\mathrm{Sch}_S)$ , we denote by

$$[F, G]_{\tau} := \pi_0 \mathrm{Map}_{\mathrm{Shv}_{\tau}(\mathrm{Sch}_S)}(F, G).$$

**Remark 2.2.50.** If  $F, G$  are presheaves, then their mapping space in presheaves and the mapping space of their associated sheafifications are quite different. We notice that if  $G$  is already a sheaf, then the unit  $G \rightarrow iL_{\mathrm{Nis}}G$  is an equivalence, inducing an equivalence of mapping spaces

$$\mathrm{Map}_{\mathrm{PSh}(\mathrm{Sch}_S)}(F, G) \cong \mathrm{Map}_{\mathrm{PSh}(\mathrm{Sch}_S)}(F, iL_{\mathrm{Nis}}G) \cong \mathrm{Map}_{\mathrm{Shv}_{\tau}(\mathrm{Sch}_S)}(L_{\mathrm{Nis}}F, L_{\mathrm{Nis}}G).$$

We think of this as a higher categorical analogue of the universal property of sheafification: that a map from a presheaf to a sheaf factors uniquely through its sheafification.

## 2.2.8 Equivalences of sheaves and presheaves

Recall that if  $f: F \rightarrow G$  is a map in  $\mathrm{PSh}(\mathrm{Sch}_S)$ , then it is an equivalence if and only if  $F(U) \rightarrow G(U)$  is an equivalence of spaces for each  $U \in \mathrm{Sch}_S$ , also called a *sectionwise equivalence*. If  $f: F \rightarrow G$  is an equivalence, then so is the associated map after sheafification. The converse doesn't hold though, since different presheaves can admit the same sheafification. To that end, we introduce a new notion.

**Definition 2.2.51.** Let  $F, G \in \mathrm{PSh}(\mathrm{Sch}_S)$  be presheaves, and let  $f: F \rightarrow G$  be a map of presheaves. We say that  $f$  is a *local equivalence* if

$$L_{\mathrm{Nis}}(f): L_{\mathrm{Nis}}F \rightarrow L_{\mathrm{Nis}}G$$

is an equivalence of sheaves.

**Example 2.2.52.** The unit of the sheafification adjunction  $L_{\mathrm{Nis}} \dashv i$  has components

$$F \rightarrow iL_{\mathrm{Nis}}F,$$

which are all local equivalences.

*Proof.* Since  $i$  is fully faithful, we have that the counit  $L_{\mathrm{Nis}}i \rightarrow \mathrm{id}$  is a natural equivalence. We'd like to argue that

$$L_{\mathrm{Nis}}F \rightarrow L_{\mathrm{Nis}}iL_{\mathrm{Nis}}F$$

is an equivalence. Since  $i$  is fully faithful, it reflects equivalences, so it suffices to check that

$$iL_{\text{Nis}}F \rightarrow iL_{\text{Nis}}iL_{\text{Nis}}F$$

is a natural equivalence, which follows from the counit being a natural equivalence.  $\square$

**Notation 2.2.53.** Let  $X$  be a scheme and  $x \in X$ . We denote by  $\text{Hen}_{X,x}$  the category of maps  $f: (Y, y) \rightarrow (X, x)$  where  $f: Y \rightarrow X$  is étale,  $f(y) = x$ , and  $f$  induces an isomorphism  $k(x) \xrightarrow{\sim} k(y)$ .

**Definition 2.2.54.** Let  $X$  be a scheme and  $x \in X$ . We denote by

$$X_x^h := \lim_{(Y,y) \in \text{Hen}_{X,x}} Y = \lim_{(Y,y) \in \text{Hen}_{X,x}} Y_y$$

See [Stacks, 04GV], or [Bac, 2.22].

**Notation 2.2.55.** Denote by  $\text{Sm}_S \subseteq \text{Sch}_S$  the full subcategory of smooth  $S$ -schemes. Here  $S$  is still qcqs and Noetherian as assumed. We will restrict our attention here as it will be needed for upcoming results.

**Definition 2.2.56.** Let  $f: F \rightarrow G$  be a map of presheaves in  $\text{PSh}(\text{Sm}_S)$ . Then we say  $f$  is a *stalkwise (Nisnevich) weak equivalence* if for every  $X \in \text{Sm}_S$  and every  $x \in X$ , the induced map

$$\text{colim}_{\text{Hen}_{X,x}} F(Y) \rightarrow \text{colim}_{\text{Hen}_{X,x}} G(Y)$$

is an equivalence of spaces.

**Warning 2.2.57.** This is *not* the same as saying that  $F(X_x^h) \rightarrow G(X_x^h)$  is an equivalence.

**Theorem 2.2.58.** (Voevodsky) Let  $S$  be qcqs and Noetherian, and let  $f: F \rightarrow G$  in  $\text{PSh}(\text{Sm}_S)$ . Then  $f$  is a local equivalence if and only if it is a stalkwise equivalence.

*Proof.* (todo — hard)  $\square$

As a particular case of [Theorem 2.2.58](#) applied to [Example 2.2.52](#), we recover the following familiar result.

**Corollary 2.2.59.** A presheaf and its Nisnevich sheafification admit the same stalks in the Nisnevich site.

## 2.2.9 Torsors revisited

**Proposition 2.2.60.** Let  $S$  be qcqs and Noetherian, and let  $\tau$  be a topology on  $\text{Sch}_S$ . Then given any  $\tau$ -cover  $\mathcal{U} := \{U_i \rightarrow X\}$ , we can look at the induced map from the Čech nerve (viewed as a simplicial object of representable presheaves) to  $X$  viewed as a discrete representable simplicial presheaf. This map

$$N_\bullet(\mathcal{U}) \rightarrow X$$

is a local equivalence of presheaves.

**Remark 2.2.61.** (How to prove this)

1. One way to prove this is by identifying the sheaf topos, as a simplicial model category, with the Bousfield localization of the category of simplicial presheaves, endowed with the projective model structure, at the class of hypercovers in the topology. We insist on  $S$  being qcqs and Noetherian so we don't have to stress about the difference between covers and hypercovers here, although there is a more general statement over any base. This follows a body of work by Jardine, Bousfield and Kan, Dugger, Hollander and Isaksen.

2. Another direction is by formal nonsense of  $\infty$ -topos theory. For instance if  $X \in \mathrm{Sm}_S$ , we can apply [Lur, A.5.3.1] to the Čech nerve of a cover after pulling it back to the slice topos  $\mathrm{Shv}_\tau(\mathrm{Sm}_S)_{/X}$ .

What about classifying spaces  $B_\bullet \mathcal{G} \in \mathrm{PSh}(\mathrm{Sch}_S)$ ? We would like to understand them in the sheaf topos.

**Notation 2.2.62.** If  $\tau$  is a topology, we denote by  $B_\tau \mathcal{G} := L_\tau B_\bullet \mathcal{G}$  the sheafification of the bar construction in the  $\tau$  topology.

**Example 2.2.63.** For any  $\mathcal{G}$ , we have a stack of  $\mathcal{G}$ -torsors, which we denote by

$$\mathrm{Tors}_\tau(\mathcal{G}): \mathcal{C}^{\mathrm{op}} \rightarrow \mathrm{Grpd}.$$

This gives rise to a sheaf  $B\mathrm{Tors}_\tau(\mathcal{G})$  by post-composing with the nerve (Example 2.2.25).

**Theorem 2.2.64.** [AHW18, 2.3.2] There is a morphism of simplicial presheaves  $B_\bullet \mathcal{G} \rightarrow B\mathrm{Tors}_\tau(\mathcal{G})$ , defined on sections by sending the unique vertex of  $B_\bullet \mathcal{G}(U)$  to the trivial  $\mathcal{G}$ -torsor over  $U$ . This map is a local equivalence.

**Corollary 2.2.65.** For any  $\tau$ -sheaf of groups  $\mathcal{G}$ , there is a natural isomorphism of sheaves of sets

$$[-, B_\tau \mathcal{G}]_\tau = \pi_0 \mathrm{Map}_{\mathrm{Shv}_\tau(\mathrm{Sch}_S)}(-, B_\tau \mathcal{G}) \cong H_\tau^1(-, \mathcal{G}).$$

This is the result we've been hoping for. It tells us we can classify torsors in the sheaf topos  $\mathrm{Sh}_\tau(\mathrm{Sch}_S)$ , and it is a completely topos-theoretic fact. In particular we gain access to all the homotopically flavored tools available to us in an  $\infty$ -topos. For instance, we could attempt to leverage Postnikov towers to deal with obstruction theory for the classifying sheaf  $B_\tau \mathcal{G}$ . The following result indicates that this tower won't contain any interesting information.

**Example 2.2.66.** If  $\mathcal{G}$  is a sheaf of discrete groups, we have that

$$\pi_n B_\tau \mathcal{G} = \begin{cases} \mathcal{G} & i = 1 \\ 0 & \text{otherwise} \end{cases}$$

*Proof.* By Theorem 2.2.64 we have that  $B_\tau \mathcal{G} \simeq B\mathrm{Tors}_\tau \mathcal{G}$ , and the latter is a sheaf whose sections are exactly the nerves of discrete groupoids, and hence  $K(\pi, 1)$ 's.  $\square$

## 2.3 Bonus: more category theory

### 2.3.1 Cofinality

**Definition 2.3.1.** We say a subcategory  $I \subseteq J$  is *cofinal* if for any functor  $F: J \rightarrow \mathcal{C}$ , the induced map on colimits

$$\mathrm{colim}_I F \rightarrow \mathrm{colim}_J F \tag{2.2}$$

is an isomorphism.<sup>8</sup> In other words, in order to compute a  $J$ -shaped colimit, it suffices to restrict to the subdiagram  $I \subseteq J$ .

**Remark 2.3.3.** This definition makes sense in 1-category theory as well as it does in  $\infty$ -category theory, however we remark that the two notions are different, so let's differentiate between the two, continuing the story of Definition 2.3.1:

---

<sup>8</sup>We dually say  $I \subseteq J$  is *final* if the natural map  $\lim_I F \rightarrow \lim_J F$  is an isomorphism for any  $F$ .



- ▷  $I \subseteq J$  is 1-cofinal if for any 1-category  $\mathcal{C}$  and 1-functor  $F: J \rightarrow \mathcal{C}$ , the induced map Equation 2.2 is an isomorphism.
- ▷  $I \subseteq J$  is cofinal ( $\infty$ -cofinal if we want to be really pedantic) if for any  $\infty$ -category and  $\infty$ -functor  $F: J \rightarrow \mathcal{C}$ , the induced map Equation 2.2 is an equivalence.

Note this latter definition extends to the case where  $I$  and  $J$  are themselves  $\infty$ -categories, or even just simplicial sets.

**Example 2.3.4.** The subcategories  $2\mathbb{N} \subseteq \mathbb{N}$  and  $2\mathbb{N} + 1 \subseteq \mathbb{N}$  are both 1-cofinal and  $\infty$ -cofinal.

It is straightforward to check when a subdiagram is 1-cofinal:

**Proposition 2.3.5.** [Stacks, 04E6]  $I \subseteq J$  is 1-cofinal if

- ▷ every  $j \in J$  has some  $i \in I$  with a morphism  $j \rightarrow i$
- ▷ for every  $j \in J$  and pair of objects  $i, i' \in I$ , there is a zig-zag of morphisms between  $i, i' \in I$  and maps from  $j$  into the zig-zag making the diagram commute:

$$\begin{array}{ccccccc} & & & j & & & \\ & & \swarrow & \downarrow & \searrow & & \\ \cdots & \longrightarrow & i_n & \longleftarrow & i_{n+1} & \longrightarrow & i_{n+2} \longleftarrow \cdots \end{array}$$

**Notation 2.3.6.** Let  $\Delta^{\text{inj}} \subseteq \Delta$  be the subcategory of injective maps, and let  $\Delta_{\leq n} \subseteq \Delta$  denote the full subcategory of objects  $[k]$  for  $k \leq n$ . For example  $\Delta_{\leq 1}^{\text{inj}}$  is just a the category with two parallel arrows, i.e. the “(co)equalizer” category:

$$\Delta_{\leq 1}^{\text{inj}} := \bullet \rightrightarrows \bullet$$

**Example 2.3.7.** We have that  $\Delta_{\leq 1}^{\text{inj}, \text{op}} \subseteq \Delta^{\text{op}}$  is 1-cofinal (c.f. [Rie14, 8.3.8]).

**Proposition 2.3.8.** The inclusion  $\Delta_{\leq n}^{\text{inj}, \text{op}} \subseteq \Delta^{\text{op}}$  is  $n$ -cofinal for any  $1 \leq n \leq \infty$ .

It turns out by a souped-up extension of Quillen Theorem A, originally due to Joyal, we have a necessary condition for cofinality in the  $\infty$ -categorical setting.

**Theorem 2.3.9.** (Joyal, Quillen) Let  $f: I \rightarrow J$  be a functor of 1-categories which is  $(\infty)$ -cofinal. Then the induced map on classifying spaces

$$BI \rightarrow BJ$$

is a weak homotopy equivalence [Lur09, 4.1.3.1, 4.1.3.3].

**Example 2.3.10.** The same subcategory  $\Delta_{\leq 1}^{\text{inj}, \text{op}} \subseteq \Delta^{\text{op}}$  is *not*  $\infty$ -cofinal.

*Proof.* The classifying space of the coequalizer diagram is  $S^1$ , however since  $[0] \in \Delta$  is terminal, it is initial in  $\Delta^{\text{op}}$ , hence  $B\Delta^{\text{op}} \simeq *$  is contractible.  $\square$

**Remark 2.3.11.** Removing this injectivity hypothesis is also interesting, since we include the opposite of the map  $[1] \rightarrow [0]$  — the category  $\Delta_{\leq 1}^{\text{op}}$  is the *split coequalizer* category:

$$\bullet \begin{array}{c} \rightrightarrows \\ \lhookrightarrow \end{array} \bullet$$

**Proposition 2.3.12.** Each of the composites

$$\Delta_{\leq 1}^{\text{inj}, \text{op}} \subseteq \Delta_{\leq 1}^{\text{op}} \subseteq \Delta^{\text{op}}$$

is 1-cofinal, however none of these inclusions are  $\infty$ -cofinal.

*Proof.* The universal property of the 1-categorical colimit for both  $\Delta_{\leq 1}^{\text{inj}, \text{op}}$  and  $\Delta_{\leq 1}^{\text{op}}$  agree, so this is direct. The other inclusion now follows by [Example 2.3.7](#).  $\square$

In general we cannot truncate in order to obtain an  $\infty$ -cofinal diagram. We can, however, restrict only to face maps and throw out degeneracies:

**Lemma 2.3.13.** [[Lur09](#), 6.5.3.7] The inclusion  $\Delta^{\text{inj}, \text{op}} \subseteq \Delta$  is cofinal.

**Definition 2.3.14.** An  $\infty$ -category  $\mathcal{C}$  is *sifted* if the diagonal map  $\mathcal{C} \rightarrow \mathcal{C} \times \mathcal{C}$  is cofinal [[Lur09](#), 5.5.8.1].

**Proposition 2.3.15.** [[Lur09](#), 5.5.8.11] Sifted colimits valued in  $\mathbf{Set}$  commute with finite products.

**Remark 2.3.16.** In the 1-categorical setting, the converse of [Proposition 2.3.15](#) holds, meaning we can take this to be the definition of sifted colimits.

**Example 2.3.17.**

1. The category  $\Delta^{\text{op}}$  is sifted
2. Any filtered category is sifted

The examples above are the only interesting examples, in the following more precise sense.

**Proposition 2.3.18.** A category  $\mathcal{C}$  admits all sifted colimits if and only if it admits all filtered colimits and it admits geometric realizations (meaning  $\Delta^{\text{op}}$ -indexed colimits).

**Corollary 2.3.19.** A 1-category admits all sifted colimits if and only if it admits all filtered colimits and it has coequalizers.

*Proof.*  $\Delta^{\text{op}}$ -indexed colimits are just coequalizers in 1-categories by [Example 2.3.7](#).  $\square$

Moreover, we may add in *finite coproducts* to capture all colimits.

**Proposition 2.3.20.** Let  $\mathcal{C}$  be a category,  $\mathcal{D} \subset \mathcal{C}$ , a full subcategory, and  $X$  an object. Then, the following are equivalent.

- (a)  $X$  is a geometric realization of coproducts of elements in  $\mathcal{D}$ .
- (b)  $X$  is a sifted colimit of finite coproducts of elements in  $\mathcal{D}$ .
- (c)  $X$  is a colimit of object in  $\mathcal{D}$ .

*Proof.* The implication (a)  $\implies$  (b) is essentially [Proposition 2.3.18](#) (as coproducts are filtered colimits of finite coproducts) and the implication (b)  $\implies$  (c) is the fact that colimits of colimits are colimits. The implication (c)  $\implies$  (a) follows from the Bousfield-Kan formula for a limit of  $p: K \rightarrow \mathcal{C}$ :

$$\text{colim}_K p \xleftarrow{\sim} \text{colim} \left( \coprod_{x \in K_0} p(x) \xleftarrow{\quad} \coprod_{\alpha \in K_1} p(\alpha(0)) \xleftarrow{\quad} \coprod_{\alpha \in K_2} p(\alpha(0)) \xleftarrow{\quad} \cdots \right)$$

see e.g. [[Sha23](#), Cor 12.5].  $\square$

### 2.3.2 Essential smallness

Being small is not a property of categories that is invariant under equivalence, so it is more meaningful to ask whether a category is *essentially small* (whether it is equivalent to a small category). This is equivalent to a category admitting a small skeleton, although the axiom of choice is required in order to pick representatives for each isomorphism class of object.

**Theorem 2.3.21.** The category of finite type  $S$ -schemes is essentially small.

*Proof sketch.* (see [MO251044](#) for details) If  $S = \operatorname{Spec}(A)$  is affine, then the category  $\operatorname{Alg}_A^{\text{f.t.}}$  of finite type  $A$ -algebras is essentially small, since all the objects are isomorphic to algebras of the form  $A[x_1, \dots, x_n]/(f_1, \dots, f_r)$ , of which there are a set. We can glue finite type  $S$ -schemes over affines, and then bootstrap to the more general case of the base  $S$  not being affine by gluing finite type schemes over all affine subschemes of  $S$ .  $\square$

**Remark 2.3.22.** The category of all  $S$ -schemes is not essentially small.

# Chapter 3

## Motivic spaces

### Contents

---

<b>3.1 Motivic spaces</b>	<b>45</b>
3.1.1 Singular chains	46
3.1.2 The category of motivic spaces	48
3.1.3 Properties of motivic localization	49
3.1.4 Motivic equivalences	49
<b>3.2 Jouanolou devices and motivic equivalences</b>	<b>50</b>
3.2.1 Checking motivic equivalences on affines	51
<b>3.3 Pointed motivic spaces</b>	<b>52</b>
3.3.1 Motivic spheres	53
<b>3.4 <math>\mathbb{A}^1</math>-homotopy classes of maps</b>	<b>54</b>
3.4.1 Connected components	54
<b>3.5 Strong and strict <math>\mathbb{A}^1</math>-invariance</b>	<b>55</b>
3.5.1 Unramified sheaves	56
3.5.2 Unramified sheaves from theories on fields	57
<b>3.6 Milnor <math>K</math>-theory</b>	<b>59</b>
3.6.1 Basic symbol algebra in $K_*^M$	60
3.6.2 Computations	61
<b>3.7 Milnor–Witt <math>K</math>-theory</b>	<b>61</b>
3.7.1 Symbol algebra in Milnor–Witt $K$ -theory	62
3.7.2 Comparison to Milnor $K$ -theory and Grothendieck–Witt	64
<b>3.8 The Milnor conjecture</b>	<b>65</b>
<b>3.9 Eilenberg–MacLane spaces</b>	<b>67</b>
3.9.1 Strong and strict invariance revisited	68
3.9.2 Cofiber sequences	69
<b>3.10 Contraction</b>	<b>70</b>
3.10.1 Examples of contraction	72
3.10.2 Contraction is exact	73
<b>3.11 Homotopy modules</b>	<b>73</b>
3.11.1 Monoidal structure on $\mathrm{HI}(k)$	73

3.11.2 The definition of homotopy modules . . . . .	74
3.11.3 Monoidal structure on homotopy modules . . . . .	75
<b>3.12 Gersten complexes</b> . . . . .	<b>76</b>
3.12.1 Vanishing of $I^j$ -cohomology . . . . .	78
3.12.2 Contraction yoga . . . . .	78
<b>3.13 The fundamental group of <math>\mathbb{P}^1</math></b> . . . . .	<b>79</b>
3.13.1 The fundamental group of $\mathbb{P}^1$ . . . . .	81
<b>3.14 The Brouwer degree</b> . . . . .	<b>82</b>
3.14.1 Explicit computations . . . . .	84
<b>3.15 Naive homotopies</b> . . . . .	<b>85</b>
3.15.1 Cazanave's theorem . . . . .	87
<b>3.16 Unstable connectivity</b> . . . . .	<b>87</b>
<b>3.17 Simplicial Freudenthal</b> . . . . .	<b>89</b>
<b>3.18 Motivic homotopy groups of spheres</b> . . . . .	<b>90</b>
3.18.1 The Steinberg relation . . . . .	91
3.18.2 Digression: the Hopf construction . . . . .	92
3.18.3 Splitting cofiber sequences . . . . .	92

## 3.1 Motivic spaces

We are interested in studying  $\mathbb{A}^1$ -invariance of presheaves. There are a few ways we might impose this, the first being the most naive — we can study presheaves  $F$  for which

1. any projection  $X \times \mathbb{A}^1 \rightarrow X$  induces an equivalence  $F(X) \xrightarrow{\sim} F(X \times \mathbb{A}^1)$
2. any vector bundle torsor  $E \rightarrow X$  induces an equivalence  $F(X) \xrightarrow{\sim} F(E)$
3. any algebraic vector bundle *torsor*  $Y \rightarrow X$  induces an equivalence  $F(X) \rightarrow F(Y)$ .

We have a strengthening of conditions here, but we notice that in the context of sheaves they all become equivalent, so the definition is only really important for presheaves. We will work with the first one as it is classically what people work with, and the last one since it will simplify some proofs. Let's first define vector bundle torsors explicitly.

**Definition 3.1.1.** [Wei89, 4.2] A *vector bundle torsor* is an affine map  $Y \rightarrow X$  which is a torsor for an algebraic vector bundle  $E \rightarrow X$ . Explicitly,  $Y \rightarrow X$  is a Zariski locally trivial affine morphism with fibers isomorphic to affine space.

**Example 3.1.2.** If  $X = \text{Spec}(A)$  is an affine scheme, then algebraic vector bundle torsors over  $X$  are the same as vector bundles over  $X$ .

todo - find an example of a non-affine  $X$  for which vector bundle torsors and vector bundles don't agree

**Notation 3.1.3.** Let  $\text{Sm}_S$  denote the category of smooth  $S$ -schemes of finite type over  $S$ . The finite type assumption is needed in order to guarantee that  $\text{Sm}_S$  is essentially small (see [Theorem 2.3.21](#)).

**Definition 3.1.4.** Let  $F \in \text{PSh}(\text{Sm}_S)$  be a presheaf.

1. We say  $F$  is  $\mathbb{A}^1$ -invariant if for every  $X \in \text{Sm}_S$ , the projection map  $X \times \mathbb{A}^1 \rightarrow X$  induces an equivalence

$$F(X) \xrightarrow{\sim} F(X \times \mathbb{A}^1).$$

2. We say  $F$  is *strongly homotopy invariant* if for every vector bundle torsor  $Y \rightarrow X$ , the restriction map

$$F(X) \xrightarrow{\sim} F(Y)$$

is an equivalence.

We denote by  $\mathrm{PSh}_{\mathbb{A}^1}(\mathrm{Sm}_S)$  and  $\mathrm{PSh}_{\mathrm{htp}}(\mathrm{Sm}_S)$  the full subcategories of  $\mathrm{PSh}(\mathrm{Sm}_S)$  spanned by the  $\mathbb{A}^1$ -invariant and strongly homotopy invariant presheaves, respectively.

**Remark 3.1.5.** It is clear that we have containments

$$\mathrm{PSh}_{\mathrm{htp}}(\mathrm{Sm}_S) \subseteq \mathrm{PSh}_{\mathbb{A}^1}(\mathrm{Sm}_S) \subseteq \mathrm{PSh}(\mathrm{Sm}_S),$$

but neither of the reverse inclusions hold.

**Proposition 3.1.6.** Both of  $\mathrm{PSh}_{\mathbb{A}^1}(\mathrm{Sm}_S)$  and  $\mathrm{PSh}_{\mathrm{htp}}(\mathrm{Sm}_S)$  are accessible subcategories of  $\mathrm{PSh}(\mathrm{Sm}_S)$ , and therefore their inclusions admit left adjoints.

*Proof.* By [Proposition 2.1.39](#) it suffices to check they are defined by being local with respect to a set of maps. Note that  $\mathrm{Sm}_S$  has a small skeleton, hence we may pick a form a set  $S$  containing a projection  $X \times \mathbb{A}^1 \rightarrow X$  for each isomorphism class of smooth schemes  $X$ . Now note that a presheaf  $F$  is  $\mathbb{A}^1$ -invariant if and only if it is  $S$ -local, meaning that

$$\mathrm{Map}_{\mathrm{PSh}(\mathrm{Sm}_S)}(h_X, F) \xrightarrow{\sim} \mathrm{Map}_{\mathrm{PSh}(\mathrm{Sm}_S)}(h_{X \times \mathbb{A}^1}, F)$$

is an equivalence. We have used here the Yoneda lemma, and we conclude by noting that the Yoneda embedding preserves finite products. This proves the statement for  $\mathbb{A}^1$ -invariant presheaves, and an analogous argument works for  $\mathrm{PSh}_{\mathrm{htp}}(\mathrm{Sm}_S)$ .  $\square$

**Terminology 3.1.7.** In light of the previous remark, we might also call presheaves or sheaves  $\mathbb{A}^1$ -local instead of  $\mathbb{A}^1$ -invariant. This is the terminology used in [\[MV99\]](#).

**Example 3.1.8.** (*Not every representable is  $\mathbb{A}^1$ -invariant*) Representable presheaves need not be  $\mathbb{A}^1$ -invariant. For example:

- ▷  $\mathbb{G}_m$  is  $\mathbb{A}^1$  invariant assuming the base is reduced. This is because it represents units, which are  $\mathbb{A}^1$ -invariant.
- ▷  $\mathbb{A}^1$  is not  $\mathbb{A}^1$ -invariant, since it represents global sections.

**Notation 3.1.9.** We denote by

$$\begin{aligned} L_{\mathbb{A}^1} &: \mathrm{PSh}(\mathrm{Sm}_S) \rightarrow \mathrm{PSh}_{\mathbb{A}^1}(\mathrm{Sm}_S), \\ L_{\mathrm{htp}} &: \mathrm{PSh}(\mathrm{Sm}_S) \rightarrow \mathrm{PSh}_{\mathrm{htp}}(\mathrm{Sm}_S) \end{aligned}$$

the associated localizations which are left adjoint to the inclusions.

### 3.1.1 Singular chains

Here we develop an explicit formula for  $L_{\mathbb{A}^1}$  which will help us do computations.

**Notation 3.1.10.** We denote by  $\Delta^n$  the *algebraic  $n$ -simplex*

$$\Delta^n := \mathrm{Spec}(\mathbb{Z}[t_0, \dots, t_n] / (\sum t_i - 1)).$$

These give a cosimplicial scheme  $\Delta^\bullet \in \mathrm{Fun}(\Delta, \mathrm{Sch})$ .

**Definition 3.1.11.** We define the *singular chains* construction

$$\mathrm{Sing}: \mathrm{PSh}(\mathrm{Sm}_S) \rightarrow \mathrm{PSh}(\mathrm{Sm}_S)$$

by the formula

$$\mathrm{Sing}(F)(X) = \mathrm{colim}_{\Delta^{\mathrm{op}}} F(X \times \Delta^n)$$

**Proposition 3.1.12.** We have that  $\text{Sing}(F)$  is  $\mathbb{A}^1$ -invariant for any  $F$ .

*Proof sketch.* We want to prove for any  $X \in \text{Sch}_S$  that the projection map  $\pi: X \times \mathbb{A}^1 \rightarrow X$  induces an equivalence

$$\pi^*: (\text{Sing}F)(X \times \mathbb{A}^1) \rightarrow (\text{Sing}F)(X).$$

Let  $z: X \rightarrow X \times \mathbb{A}^1$  denote the zero section. Then we claim  $z^*$  exhibits a simplicial homotopy equivalence with  $\pi^*$ . Since  $\pi z = \text{id}$ , it is clear that  $z^* \pi^* = \text{id}$ , so it suffices to exhibit a simplicial homotopy  $\pi^* z^* \simeq \text{id}$ . Since any functor will preserve simplicial homotopies, and geometric realizations will send simplicial homotopies to honest homotopies, it suffices to exhibit a homotopy of cosimplicial simplicial varieties

$$\text{id}, z \circ \pi: X \times \mathbb{A}^1 \times \Delta^\bullet \rightarrow X \times \mathbb{A}^1 \times \Delta^\bullet,$$

meaning maps whose opposites satisfy the simplicial homotopy identities. (todo: add explicit simplicial homotopy)  $\square$

**Proposition 3.1.13.** If  $F$  is already  $\mathbb{A}^1$ -invariant, then the natural map

$$F \rightarrow \text{Sing}(F)$$

is an equivalence of presheaves.

*Proof.* Immediate since  $F$  is  $\mathbb{A}^1$ -invariant, so we can turn  $F(X \times \Delta^\bullet)$  into a constant diagram.  $\square$

**Proposition 3.1.14.** There is a natural equivalence of functors  $L_{\mathbb{A}^1} \simeq \text{Sing}$ .

*Proof.* We note that the essential image of  $\text{Sing}$  is precisely  $\text{PSh}_{\mathbb{A}^1}(\text{Sm}_k)$  by [Proposition 3.1.13](#). By that same result, the natural transformation

$$\eta: \text{id} \rightarrow \text{Sing}$$

induces an equivalence between the maps  $\text{Sing}(\eta_{(-)})$  and  $\eta_{\text{Sing}(-)}$  on its components. By [\[Lur09, 5.2.7.4\]](#), this implies that  $\text{Sing}(-)$  is a left adjoint, with right adjoint given by the fully faithful inclusion of its essential image  $\text{PSh}_{\mathbb{A}^1}(\text{Sm}_k) \subseteq \text{PSh}(\text{Sm}_k)$ . By uniqueness of adjoints, this implies  $\text{Sing} \simeq L_{\mathbb{A}^1}$ .  $\square$

**Corollary 3.1.15.**  $L_{\mathbb{A}^1}$  preserves finite products.

*Proof.* This is not immediate from the definition, but it follows after identifying  $L_{\mathbb{A}^1}$  with  $\text{Sing}(-)$ . Since  $\Delta^{\text{op}}$  is sifted (c.f. [\[Lur09, 5.5.4.8\]](#)), colimits indexed over  $\Delta^{\text{op}}$  commute with products, hence the result follows.  $\square$

**Proposition 3.1.16.** For any  $X \in \text{Sm}_S$ , we have that  $X \times \mathbb{A}_S^n \rightarrow X$  is an equivalence after  $\mathbb{A}^n$ -localization.

*Proof.* For  $n = 1$  this is by definition of the localization, and for higher  $n$  this follows from  $L_{\mathbb{A}^1}$  preserving finite products.  $\square$

**Remark 3.1.17.** An analogous formula holds for  $L_{\text{htp}}$  — if we let  $\text{VBT}_{/X}$  denote the category of vector bundle torsors over  $X$ , then  $\text{VBT}_{/X}$  is cosifted, and we obtain an identification

$$(L_{\text{htp}}F)(X) = \text{colim}_{Y \in \text{VBT}_{/X}} F(Y), \tag{3.18}$$

with the same properties that it preserves finite products and is locally cartesian [\[Hoy17, 3.5\]](#)

### 3.1.2 The category of motivic spaces

**Definition 3.1.19.** We define the category of *motivic spaces*  $\mathrm{Spc}(k)$  as

$$\mathrm{Spc}(k) = \mathrm{Shv}_{\mathrm{Nis}}(\mathrm{Sm}_k) \cap \mathrm{PSh}_{\mathbb{A}^1}(\mathrm{Sm}_k) \subseteq \mathrm{PSh}(\mathrm{Sm}_k),$$

that is, the full subcategory of presheaves which are both Nisnevich sheaves and are  $\mathbb{A}^1$ -invariant.

**Remark 3.1.20.** Since every affine bundle torsor is locally trivialized, once we impose the sheaf condition, the properties of being  $\mathbb{A}^1$ -invariant and strongly homotopy invariant can be checked locally, and hence agree:

$$\mathrm{Shv}_{\mathrm{Nis}}(\mathrm{Sm}_S) \cap \mathrm{PSh}_{\mathbb{A}^1}(\mathrm{Sm}_S) = \mathrm{Shv}_{\mathrm{Nis}}(\mathrm{Sm}_S) \cap \mathrm{PSh}_{\mathrm{htp}}(\mathrm{Sm}_S).$$

In other words we could equivalently define motivic spaces via  $\mathrm{PSh}_{\mathbb{A}^1}$ , or via  $\mathrm{PSh}_{\mathrm{htp}}$ . The former is more common, although the latter has some nice advantages (see [Hoy17, p. 204] for a discussion, and [Hoy17, 3.13] for the equality above).

**Problem:** Nisnevich sheafifying an  $\mathbb{A}^1$ -invariant presheaf needs not preserve  $\mathbb{A}^1$ -invariance, and  $\mathbb{A}^1$ -localizing a sheaf may break the sheaf condition.

**Example 3.1.21.** [MV99, 3.2.7] Let  $U_0 = \mathbb{A}^1 - 0$  and  $U_1 = \mathbb{A}^1 - 1$ , and let  $U_{01} = U_0 \cap U_1 \subseteq \mathbb{A}^1$ . Since both  $U_0$  and  $U_1$  are  $\mathbb{A}^1$ -invariant, we claim that  $U_{01}$  is as well. Pick a closed embedding of  $U_{01}$  in  $\mathbb{A}^n$  for some  $n$ , and consider the non-smooth scheme

$$Y := (U_0 \times \mathbb{A}^n) \amalg_{U_{01}} (U_1 \times \mathbb{A}^n) \in \mathrm{Sch}_S.$$

Then for any connected  $X \in \mathrm{Sm}_S$ , we have that

$$\mathrm{Hom}_{\mathrm{Sch}_S}(X, Y) = \mathrm{Hom}_{\mathrm{Sm}_S}(X, U_0 \times \mathbb{A}^n) \coprod_{\mathrm{Hom}_{\mathrm{Sm}_S}(X, U_{01})} \mathrm{Hom}_{\mathrm{Sm}_S}(X, U_1 \times \mathbb{A}^n).$$

Since  $L_{\mathbb{A}^1}$  preserves pushouts, we have that

$$L_{\mathbb{A}^1} h_Y = L_{\mathbb{A}^1}(U_0 \times \mathbb{A}^n) \amalg_{L_{\mathbb{A}^1}(U_{01})} L_{\mathbb{A}^1}(U_1 \times \mathbb{A}^n).$$

By [Proposition 3.1.16](#) we can contract away the  $\mathbb{A}^n$ 's, and we invoke that  $U_0$ ,  $U_1$ , and  $U_{01}$  were already  $\mathbb{A}^1$ -local to get that the above is equivalent to

$$\simeq L_{\mathbb{A}^1}(U_0) \coprod_{L_{\mathbb{A}^1}(U_{01})} L_{\mathbb{A}^1}(U_1) = U_0 \coprod_{U_{01}} \mathbb{A}^1.$$

If this were a sheaf, it would agree with its sheafification, and since sheafification preserves pushouts, we would have that it is equal to the *pushout of sheaves*

$$U_0 \amalg_{U_{01}} U_1 = \mathbb{A}^1,$$

which is the representable sheaf given by  $\mathbb{A}^1$  by Zariski descent. But note that  $\mathbb{A}^1$  is not  $\mathbb{A}^1$ -invariant, so we get that  $L_{\mathbb{A}^1} h_Y$  cannot be a sheaf.

**Proposition 3.1.22.** The category  $\mathrm{Spc}(S) \subseteq \mathrm{PSh}(\mathrm{Sm}_S)$  is an accessible localization, hence the inclusion admits a left adjoint.

We call this adjoint *motivic localization*, and we can describe it explicitly as the infinite composition of both functors.

**Definition 3.1.23.** We define  $L_{\mathrm{mot}} : \mathrm{PSh}(\mathrm{Sm}_S) \rightarrow \mathrm{Spc}(S)$  by the formula

$$L_{\mathrm{mot}} := \mathrm{colim} (L_{\mathrm{Nis}} \rightarrow L_{\mathbb{A}^1} L_{\mathrm{Nis}} \rightarrow L_{\mathrm{Nis}} L_{\mathbb{A}^1} L_{\mathrm{Nis}} \rightarrow \cdots)$$

where this colimit is computed in the presheaf category.



**Remark 3.1.24.** By cofinality (Example 2.3.4), this is canonically equivalent to the following two colimits:

$$\begin{aligned} \operatorname{colim} (L_{\mathrm{Nis}} \circ L_{\mathbb{A}^1} \rightarrow (L_{\mathrm{Nis}} \circ L_{\mathbb{A}^1})^{\circ 2} \rightarrow \cdots) \\ \operatorname{colim} (L_{\mathbb{A}^1} \circ L_{\mathrm{Nis}} \rightarrow (L_{\mathbb{A}^1} \circ L_{\mathrm{Nis}})^{\circ 2} \rightarrow \cdots). \end{aligned}$$

The former is a colimit computed in  $\operatorname{Shv}_{\mathrm{Nis}}(\operatorname{Sm}_k)$ , which is closed under filtered colimits by Proposition 3.1.6, hence the resulting object is a sheaf. The latter is a colimit computed in  $\operatorname{PSh}_{\mathbb{A}^1}(\operatorname{Sm}_k)$ , which is also closed under filtered colimits (again by Proposition 3.1.6), hence it is  $\mathbb{A}^1$ -invariant.

### 3.1.3 Properties of motivic localization

We know that  $L_{\mathrm{mot}}$  preserves colimits, being a left adjoint. What other properties does it admit?

**Proposition 3.1.25.** [Hoy14, C.6] We have that  $L_{\mathrm{mot}}$  preserves finite products.

*Proof.* This follows by both  $L_{\mathrm{Nis}}$  and  $L_{\mathbb{A}^1}$  preserving finite products, together with transfinite composition preserving products.  $\square$

**Proposition 3.1.26.** [Hoy17, 3.15]  $L_{\mathrm{mot}}$  is locally cartesian, meaning that for a cospan  $X \rightarrow Y \leftarrow Z$  in  $\operatorname{PSh}(\operatorname{Sm}_S)$ , if both  $X$  and  $Y$  are motivic spaces, then the natural map

$$L_{\mathrm{mot}}(X \times_Y Z) \rightarrow X \times_Y L_{\mathrm{mot}}(Z)$$

is an equivalence. Note the pullback on the domain is in the category of presheaves, while the pullback on the right is in the category of motivic spaces.

**Corollary 3.1.27.** Colimits in  $\operatorname{Spc}(S)$  are universal (pullback-stable).

*Proof.* Let  $X$  and  $Y$  be some motivic spaces, and let  $j: I \rightarrow \operatorname{Spc}(S)$  be a diagram, whose colimit maps to  $Y$ . We claim that

$$\operatorname{colim}_{i \in I} (X \times_Y j(i)) \rightarrow X \times_Y \operatorname{colim}_{i \in I} j(i)$$

is an equivalence in  $L_{\mathrm{mot}}$ . This is true in the ambient presheaf category since it is a topos, but this doesn't immediately imply the result for motivic spaces, since pullbacks of presheaves do not in general yield pullbacks of motivic spaces. However by applying  $L_{\mathrm{mot}}$  to both sides, and using the fact that it is locally cartesian, the result follows.  $\square$

**Proposition 3.1.28.** We have that  $L_{\mathrm{mot}}$  is *not left exact*. In particular  $\operatorname{Spc}(k)$  is not an  $\infty$ -topos.

We will come back to prove Proposition 3.1.28 later when we have a little more machinery.

### 3.1.4 Motivic equivalences

**Definition 3.1.29.** We say that  $f: F \rightarrow G$  in  $\operatorname{PSh}(\operatorname{Sm}_S)$  is a *motivic equivalence* if  $L_{\mathrm{mot}}f$  is an equivalence in  $\operatorname{Spc}(S)$ .

**Proposition 3.1.30.** We have that  $\Delta^0 \cong S$  in  $\operatorname{Spc}(S)$ , and moreover these are terminal, hence we will often denote them by  $*$ .

*Proof.* This follows from  $\operatorname{Spc}(S) \subseteq \operatorname{Shv}_{\mathrm{Nis}}(\operatorname{Sm}_S)$  being an accessible subcategory (hence preserving terminal objects) together with Proposition 2.2.29.  $\square$

**Proposition 3.1.31.** The map to the terminal object in schemes  $\mathbb{A}_S^n \rightarrow S$  is a motivic equivalence.

*Proof.* It is clear that  $\mathbb{A}_S^1 \rightarrow S$  is a motivic equivalence essentially by definition. The more general statement follows from both the Yoneda embedding and motivic localization preserving finite products.  $\square$

The following proposition is also essentially by definition.

**Proposition 3.1.32.** For any  $X \in \mathrm{Sm}_S$ , the projection map  $X \times \mathbb{A}_S^n \rightarrow X$  is a motivic equivalence.

**Proposition 3.1.33.** For any  $F \in \mathrm{PSh}(\mathrm{Sm}_S)$ , the projection map  $F \times \mathbb{A}^n \rightarrow F$  is a motivic equivalence.

*Proof.* By the co-Yoneda lemma, any presheaf is a colimit of representable presheaves. Since  $L_{\mathrm{mot}}$  is locally cartesian, colimits are pullback stable, and hence product-stable, meaning that

$$F \times \mathbb{A}^n \cong (\mathrm{colim}_U h_U) \times \mathbb{A}^n.$$

Since colimits are universal in any  $\infty$ -topos (in particular in the presheaf category), we have that we can distribute the colimit over the product. Then the result follows by [Proposition 3.1.32](#).  $\square$

## 3.2 Jouanolou devices and motivic equivalences

**Theorem 3.2.1.** (Jouanolou–Thomason trick) Suppose that  $S$  is qcqs and further suppose either  $S$  is affine or it is noetherian, regular, and separated. Then for any quasi-projective  $X \in \mathrm{Sch}_S$ , there exists an affine bundle  $Y \rightarrow X$ , where  $Y$  is affine.

**Remark 3.2.2.** The above result probably holds under weaker assumptions – we’re citing [\[Hoy17, 2.20\]](#), pulling our assumptions on  $S$  from [\[Hoy17, 2.8\]](#) and letting  $G$  be trivial. In most cases we apply this,  $S$  will be a field, and  $X$  will be smooth, and we can apply this trick by e.g. [\[AF14b, 3.3.3\]](#).

**Example 3.2.3.** The easiest example is  $\mathbb{P}_k^1$  over a field  $k$ . There is a map

$$\mathrm{Spec} \left( \frac{k[x, y, z, w]}{x + w - 1, xw - yz} \right) \rightarrow \mathbb{P}^1$$

$$(x, y, z, w) \mapsto \begin{cases} [x : y] & (x, y) \neq (0, 0) \\ [z : w] & (z, w) \neq (0, 0) \end{cases}$$

We claim this is well-defined, exhibiting an affine torsor over  $\mathbb{P}^1$ .

**Example 3.2.4.** In a similar vein, the map

$$\mathrm{SL}_2 \rightarrow \mathbb{A}^2 \setminus 0$$

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \mapsto (b, d)$$

is an affine vector bundle torsor over  $\mathbb{A}^2 \setminus 0$ . More generally

$$\mathrm{SL}_n \rightarrow \mathbb{A}^n \setminus \{0\}$$

is an affine vector bundle torsor, by sending an invertible determinant one matrix to its last column.

**Proposition 3.2.5.** Any affine vector bundle torsor  $E \rightarrow B$  is a motivic equivalence.

*Proof.* It suffices to verify on a local cover where the bundle is trivialized, at which point it follows by [Proposition 3.1.32](#).  $\square$

### 3.2.1 Checking motivic equivalences on affines

**Proposition 3.2.6.** The category  $\mathrm{Spc}(S)$  is generated under sifted colimits by (the motivic localization of the representable presheaves attached to) affine  $S$ -schemes in  $\mathrm{Sm}_S$  [Hoy17, 3.16].

*Proof.* We do this in two steps:

1. First we argue that  $\mathrm{Spc}(S)$  is generated under sifted colimits by  $\mathrm{Sm}_S$ . Let  $F \in \mathrm{Spc}(S)$  be any motivic space, then, considered as a presheaf, it is a colimit of representable presheaves. What is furthermore true is that it is a sifted colimit of finite coproducts of representable presheaves (Proposition 2.3.20). While it is not true that  $h_U \amalg h_V$  is  $h_{U \amalg V}$  in the category of presheaves, it *is true* in the category of sheaves (and hence in the category of motivic spaces), by descent. Hence any motivic localization is in fact a sifted colimit of the motivic localization of representable sheaves.
2. Second we argue that  $\mathrm{Sm}_S \subseteq \mathrm{Spc}(S)$  is generated under sifted colimits by affine  $S$ -schemes in  $\mathrm{Sm}_S$ . Given any  $X \in \mathrm{Sm}_S$ , we can write it as an  $\Delta^{\mathrm{op}}$ -indexed colimit in the category  $\mathrm{Spc}(S)$  over the Čech nerve of a Nisnevich cover. Each fiber product appearing in the cover is equivalent to a smooth affine scheme by Jouanolou's trick, and we can get rid of finite coproducts by the same trick. Hence  $X$  can be written as a  $\Delta^{\mathrm{op}}$ -indexed colimit over smooth affine  $S$ -schemes.

As sifted colimits are combinations of filtered colimits and geometric realizations (Proposition 2.3.18), the result follows.  $\square$

This gives us a powerful way to check motivic equivalences between presheaves.

**Proposition 3.2.7.** [Hoy17, 3.16] Let  $f: F \rightarrow G$  be a morphism in  $\mathrm{PSh}(\mathrm{Sm}_S)$ . If

$$F(U) \rightarrow G(U)$$

is an equivalence for every affine  $U \in \mathrm{Sm}_S$ , then  $f$  is a motivic equivalence.

*Proof.* By Yoneda it suffices to argue that

$$\mathrm{Map}_{\mathrm{Spc}(S)}(-, L_{\mathrm{mot}}F) \rightarrow \mathrm{Map}_{\mathrm{Spc}(S)}(-, L_{\mathrm{mot}}G)$$

is a natural equivalence. Since  $\mathrm{Map}(-, -)$  sends colimits to limits in the first variable, by Proposition 3.2.6 it suffices to argue that

$$\mathrm{Map}_{\mathrm{Spc}(S)}(L_{\mathrm{mot}}h_U, L_{\mathrm{mot}}F) \rightarrow \mathrm{Map}_{\mathrm{Spc}(S)}(L_{\mathrm{mot}}h_U, L_{\mathrm{mot}}G)$$

is an equivalence for each affine  $U \in \mathrm{Sm}_S$ . By adjunction, this is equivalent to asking that  $L_{\mathrm{mot}}F \rightarrow L_{\mathrm{mot}}G$  is a sectionwise equivalence of presheaves when restricted to affines.

We first argue  $L_{\mathrm{htp}}F \rightarrow L_{\mathrm{htp}}G$  is a sectionwise equivalence. Since every  $X \in \mathrm{Sm}_S$  admits an affine bundle torsor  $U \rightarrow X$  which is itself affine by the Jouanolou trick, we have that  $(L_{\mathrm{htp}}F)(X) \xrightarrow{\sim} (L_{\mathrm{htp}}F)(U)$  is an equivalence, so it suffices to check that  $L_{\mathrm{htp}}F \rightarrow L_{\mathrm{htp}}G$  is a sectionwise equivalence on affines. Since every vector bundle torsor over an affine is affine<sup>1</sup>, we can leverage the formula for  $L_{\mathrm{htp}}$  (Equation 3.18) in order to write

$$(L_{\mathrm{htp}}F)(U) = \mathrm{colim}_{V \in \mathrm{VBT}/U} F(V).$$

Since  $F(V) \rightarrow G(V)$  is an equivalence for each  $V$  above, it is clear that  $(L_{\mathrm{htp}}F)(U) \rightarrow (L_{\mathrm{htp}}G)(U)$  is an equivalence for each affine  $U$ , hence  $L_{\mathrm{htp}}F \rightarrow L_{\mathrm{htp}}G$  is a sectionwise equivalence.

<sup>1</sup>Over an affine scheme, vector bundle torsors are just vector bundles. If  $\mathcal{F}$  is a vector bundle (considered as a sheaf of modules) over  $X = \mathrm{Spec}(A)$ , then the underlying scheme is  $\mathrm{Spec}_X(\mathrm{Sym}(\mathcal{F}^*))$  which is affine over  $X$ , and hence is itself affine over the base.

Finally, since a sectionwise equivalence of presheaves is a local equivalence, we have that

$$L_{\text{Nis}}L_{\text{htp}}F \rightarrow L_{\text{Nis}}L_{\text{htp}}G$$

is an equivalence of presheaves. Passing to the colimit it is now clear that  $F \rightarrow G$  is a motivic equivalence.  $\square$

### 3.3 Pointed motivic spaces

**Notation 3.3.1.** By abuse of notation, if  $X \in \text{Sm}_S$ , then we also denote by  $X \in \text{Spc}(S)$  the motivic space  $L_{\text{mot}}h_X$ .

**Notation 3.3.2.** We denote by  $\text{Spc}(S)_*$  the category of pointed motivic spaces. This has a zero object, which we denote by  $*$ . This comes with an adjunction

$$(-)_+ : \text{Spc}(S) \rightleftarrows \text{Spc}(S)_* : U,$$

where  $X_+$  is the coproduct  $X \amalg S$ , pointed at that copy of  $S$ , and the right adjoint forgets the basepoint.

**Notation 3.3.3.** If  $Y \rightarrow X$  is a map of (pointed) motivic spaces, we denote by  $X/Y$  the cofiber, computed as the pushout

$$\begin{array}{ccc} Y & \longrightarrow & X \\ \downarrow & & \downarrow \\ * & \longrightarrow & X/Y. \end{array}$$

We denote by  $X \vee Y$  the coproduct in the category  $\text{Spc}(S)_*$  of pointed motivic spaces, and by

$$X \wedge Y = \frac{X \times Y}{X \vee Y}$$

the *smash product*.

**Notation 3.3.4.** Since  $\text{Spc}(S)_*$  has a zero object, we can denote by  $\Sigma X$  the pushout

$$\begin{array}{ccc} X & \longrightarrow & * \\ \downarrow & \lrcorner & \downarrow \\ * & \longrightarrow & \Sigma X, \end{array}$$

for any motivic space  $X$ .

**Proposition 3.3.5.** If  $X$  is a pointed motivic space, then there is a canonical equivalence

$$\Sigma X \simeq S^1 \wedge X.$$

*Proof.* Since  $L_{\text{mot}}$  preserves coproducts and  $\text{PSh}(\text{Sm}_S)$  is cocomplete, it suffices to compute the pushout at the level of simplicial presheaves. In this setting, the coproduct is computed levelwise, in which case it is clear that  $\Sigma X(U) = S^1 \wedge X(U)$  for any  $U \in \text{Sm}_S$ . Hence as pointed presheaves, we have that  $\Sigma X \simeq S^1 \wedge X$ . Finally,  $L_{\text{mot}}$  preserves the smash product construction since it preserves finite products, as well as coproducts and cofibers.  $\square$

*Proof 2.* We can first show it for  $X = S^0$ , then leverage universality of colimits and [PeterSanath] to argue in general.  $\square$

By descent, Nisnevich covers of schemes give rise to colimits of motivic spaces. One of the most immediate examples is the following:

**Example 3.3.6.** The following diagram of (pointed) motivic spaces is a pushout:

$$\begin{array}{ccc} \mathbb{G}_m & \xrightarrow{z} & \mathbb{A}^1 \\ z^{-1} \downarrow & & \downarrow \\ \mathbb{A}^1 & \longrightarrow & \mathbb{P}^1. \end{array}$$

Since  $\mathbb{A}^1 \xrightarrow{\sim} *$  is trivial, and since all our colimits are  $\infty$ -colimits, we can replace the above diagram with the weakly equivalent span  $* \leftarrow \mathbb{G}_m \rightarrow *$  and compute the same colimit. This implies that the natural map

$$\Sigma \mathbb{G}_m \xrightarrow{\sim} \mathbb{P}^1 \tag{3.7}$$

is a motivic equivalence.

### 3.3.1 Motivic spheres

**Terminology 3.3.8.** We have *two kinds of spheres* in motivic homotopy theory: ones coming from algebraic geometry (the multiplicative group scheme  $\mathbb{G}_m$  or the projective line  $\mathbb{P}^1$ ) and ones coming from algebraic topology (the constant presheaf at  $S^1$  or at  $S^n$  for any  $n$ ). So we get bigradings on the spheres. There are competing conventions in the literature for motivic grading, but the increasingly standardized convention is to write

$$\begin{aligned} S^{1,1} &:= \mathbb{G}_m \\ S^{1,0} &= S^1 \\ S^{2,1} &= \mathbb{P}^1. \end{aligned}$$

**Example 3.3.9.** We have that  $S^1 \simeq \mathbb{A}^1 / \{0, 1\}$ .

*Proof.* We can consider the cofiber diagram

$$\begin{array}{ccc} S^0 & \xrightarrow{(0,1)} & \mathbb{A}^1 \\ \downarrow & \lrcorner & \downarrow \\ * & \longrightarrow & S^1. \end{array}$$

□

**Exercise 3.3.10.** The diagram

$$\begin{array}{ccc} X \times Y & \longrightarrow & X \\ \downarrow & \lrcorner & \downarrow \\ Y & \longrightarrow & \Sigma(X \wedge Y) \end{array}$$

is a pushout.

**Proposition 3.3.11.** There is a motivic equivalence

$$\mathbb{A}^n \setminus \{0\} \simeq (S^1)^{\wedge(n-1)} \wedge (\mathbb{G}_m)^{\wedge n} = S^{2n-1,n}.$$

**Corollary 3.3.12.** We have that

$$\frac{\mathbb{A}^n}{\mathbb{A}^n \setminus \{0\}} \simeq S^{2n,n}.$$

**Proposition 3.3.13.** We have that

$$\frac{\mathbb{P}^n}{\mathbb{P}^n - 0} \simeq S^{n,n}.$$

*Proof.* By covering  $\mathbb{P}^n$  with  $\mathbb{A}^n$  and  $\mathbb{P}^n \setminus \{0\}$ , we get a pushout diagram

$$\begin{array}{ccc} \mathbb{A}^n \setminus \{0\} & \longrightarrow & \mathbb{P}^n \setminus \{0\} \\ \downarrow & \lrcorner & \downarrow \\ \mathbb{A}^n & \longrightarrow & \mathbb{P}^n. \end{array}$$

The cofibers of the vertical maps are equivalent, yielding

$$S^{2n,n} \xrightarrow{\sim} \frac{\mathbb{P}^n}{\mathbb{P}^n - 0}.$$

□

### 3.4 $\mathbb{A}^1$ -homotopy classes of maps

Given two simplicial presheaves  $F, G \in \text{PSh}(\text{Sm}_S)$ , by abuse of notation we denote by  $[F, G]_{\mathbb{A}^1}$  homotopy classes of maps of their associated motivic localizations

$$[F, G]_{\mathbb{A}^1} := \pi_0 \text{Map}_{\text{Spc}(S)}(L_{\text{mot}} F, L_{\text{mot}} G).$$

A goal is to understand how to compute this. There are some easy cases first:

**Example 3.4.1.** Let  $X$  be a representable sheaf and  $\mathcal{Y}$  be a motivic space. Then via the Yoneda lemma, we have

$$\text{Map}_{\text{Spc}(S)}(L_{\text{mot}} h_X, \mathcal{Y}) \cong \text{Map}_{\text{PSh}(\text{Sm}_S)}(h_X, \mathcal{Y}) \cong \mathcal{Y}(X).$$

Hence  $[X, \mathcal{Y}]_{\mathbb{A}^1} = \pi_0 \mathcal{Y}(X)$ .

**Example 3.4.2.** We have a bijection

$$[X, \mathbb{G}_m]_{\mathbb{A}^1} = \mathbb{G}_m(X)$$

for any  $X \in \text{Sm}_S$ .

**Example 3.4.3.** If  $X$  and  $Y$  are both varieties, then

$$[X, Y]_{\mathbb{A}^1} = \pi_0(L_{\text{mot}} h_Y)(X).$$

This is much harder to access in general, since we don't have control over the sections of the presheaf  $L_{\text{mot}} h_Y$ .

#### 3.4.1 Connected components

Given a motivic space  $X$ , we can consider the presheaf of sets

$$\begin{aligned} \text{Sm}_S^{\text{op}} &\rightarrow \text{Set} \\ U &\mapsto [U, X]_{\mathbb{A}^1}. \end{aligned}$$

Its sheafification is called the *sheaf of connected components*, and we denote it by

$$\pi_0^{\mathbb{A}^1}(X) \in \text{Shv}_{\text{Nis}}(\text{Sm}_k).$$

**Note 3.4.4.** We have that  $\pi_0^{\mathbb{A}^1}(X) = \pi_0(X)$  is the same as the homotopy sheaf in the sheaf topos, provided  $X$  is a motivic space. We have to be a bit careful what we mean here — if  $F$  is an arbitrary presheaf, then  $\pi_0(F)$  will mean  $\pi_0(L_{\text{Nis}} F)$  by convention, while  $\pi_0^{\mathbb{A}^1} F$  will mean  $\pi_0(L_{\text{mot}} F)$ .

**Proposition 3.4.5.** We have that  $\pi_0^{\mathbb{A}^1}(\text{SL}_n) \simeq *$ .

*Proof sketch.* It suffices to argue the inclusion of any element is homotopic to the identity. First suppose  $M$  is an elementary matrix. Then there is a map

$$\mathbb{A}^1 \rightarrow \mathrm{SL}_n,$$

sending 0 to the identity and 1 to  $M$ . Since  $\mathrm{SL}_n$  is generated by elementary matrices, the result follows.  $\square$

**Definition 3.4.6.** If  $(X, x)$  is a pointed motivic space, we have that  $\pi_n^{\mathbb{A}^1}(X, x)$  is defined to be the Nisnevich presheaf of

$$U \mapsto [\Sigma^n U_+, X]_{\mathrm{Spc}(S)_*}.$$

**Proposition 3.4.7.** If  $(X, x)$  is a motivic space, then  $\pi_n^{\mathbb{A}^1}(X, x) = \pi_n(X, x)$ .

*Proof.* We show the presheaves are identical before sheafifying. By adjunction we have that

$$[\Sigma^n U_+, X]_{\mathrm{Spc}(S)_*} \cong [U_+, \Omega^n X]_{\mathrm{Spc}(S)_*} = [U, \Omega^n X]_{\mathrm{Spc}(S)}.$$

The latter is the presheaf attached to  $\pi_n(X, x)$  by [Proposition 2.2.35](#).  $\square$

**Proposition 3.4.8.** Let  $f: F \rightarrow G$  be a map of simplicial presheaves. Then it is a motivic equivalence if and only if

$$\pi_n^{\mathbb{A}^1}(f): \pi_n^{\mathbb{A}^1}(F, x) \rightarrow \pi_n^{\mathbb{A}^1}(G, f(x))$$

is an equivalence for all  $n \geq 0$  and all basepoints  $x \in F$ .

*Proof.* The forward direction is clear, since  $f$  being a motivic equivalence would induce a natural isomorphism  $[-, X]_{\mathbb{A}^1} \cong [-, Y]_{\mathbb{A}^1}$ , hence the associated homotopy (pre)sheaves would be identical. For the backwards direction, this condition on homotopy groups unwinds to tell us that  $L_{\mathrm{mot}} F \rightarrow L_{\mathrm{mot}} G$  is an equivalence of sheaves by [Corollary 2.2.48](#), which in particular implies it is an equivalence of motivic spaces.  $\square$

**Terminology 3.4.9.** We say that a motivic space  $X$  is  $\mathbb{A}^1$ - $n$ -connected if  $\pi_i^{\mathbb{A}^1}(X, x)$  is trivial for all  $i \leq n$  and all basepoints. Some special cases:

$$\begin{aligned} \mathbb{A}^1\text{-connected} &= \mathbb{A}^1\text{-0-connected} \\ \mathbb{A}^1\text{-simply connected} &= \mathbb{A}^1\text{-1-connected} \\ \mathbb{A}^1\text{-contractible} &= \mathbb{A}^1\text{-}\infty\text{-connected.} \end{aligned}$$

**Example 3.4.10.** We have that  $S^{i+j,j}$  is  $\mathbb{A}^1$ -( $i-1$ )-connected (see [\[AO19, 2.4.5\]](#)). In particular it depends only on the number of copies of  $S^{1,0}$  and not on the number of copies of  $\mathbb{G}_m$ .

### 3.5 Strong and strict $\mathbb{A}^1$ -invariance

**Goal 3.5.1.** We have seen that in order to check a map is a motivic equivalence, it suffices to look at homotopy sheaves. We'd therefore like to better understand when homotopy sheaves are equivalent. It turns out homotopy sheaves (for  $n \geq 2$ ) are a prototypical example of a particularly nice class of sheaves of abelian groups called *strictly invariant* sheaves. These have nice properties that make them easier to work with.

Note that  $\pi_n^{\mathbb{A}^1}(X)$  is a Nisnevich sheaf of sets, but it is not necessarily a motivic space. We can ask to what extent it is  $\mathbb{A}^1$ -invariant.

It was conjectured by Morel that  $\pi_0^{\mathbb{A}^1}(X)$  was always  $\mathbb{A}^1$ -invariant, but a counterexample was found by Ayoub (todo). Nevertheless we can ask for  $\pi_i^{\mathbb{A}^1}(X)$  for  $i \geq 1$ . A more general question is to understand conditions that tell us a sheaf of groups is invariant.

Let  $\mathcal{G}$  denote a Nisnevich sheaf of groups. Since  $\mathcal{G}(X) = H_{\text{Nis}}^0(X, \mathcal{G})$ , the condition that  $\mathcal{G}$  is  $\mathbb{A}^1$ -invariant is equivalent to asking whether

$$H_{\text{Nis}}^0(X; \mathcal{G}) \rightarrow H_{\text{Nis}}^0(X \times \mathbb{A}^1; \mathcal{G})$$

is  $\mathbb{A}^1$ -invariant.

**Definition 3.5.2.** Let  $\mathcal{G}$  be a Nisnevich sheaf of groups over  $S$ . Then we say

1.  $\mathcal{G}$  is  $\mathbb{A}^1$ -invariant if  $H_{\text{Nis}}^0(X; \mathcal{G}) \rightarrow H_{\text{Nis}}^0(X \times \mathbb{A}^1; \mathcal{G})$  is an equivalence for every  $X \in \text{Sm}_S$
2.  $\mathcal{G}$  is *strongly*  $\mathbb{A}^1$ -invariant if

$$H_{\text{Nis}}^i(X; \mathcal{G}) \rightarrow H_{\text{Nis}}^i(X \times \mathbb{A}^1; \mathcal{G})$$

is an equivalence for  $i = 0, 1$  and for all  $X$

3. if  $\mathcal{G}$  is a sheaf of abelian groups, we say it is *strictly*  $\mathbb{A}^1$ -invariant if

$$H_{\text{Nis}}^i(X; \mathcal{G}) \rightarrow H_{\text{Nis}}^i(X \times \mathbb{A}^1; \mathcal{G})$$

is an equivalence for all  $i$  and for all  $X$ .

**Example 3.5.3.** We have that  $\mathbb{G}_m$  is strongly invariant, since both units and the Picard group are  $\mathbb{A}^1$ -invariant over a base field.

**Theorem 3.5.4.** (Morel) If  $X$  is a motivic space over a field  $k$ , then  $\pi_1^{\mathbb{A}^1}(X)$  is strongly  $\mathbb{A}^1$ -invariant.

**Theorem 3.5.5.** (Morel) If  $k$  is a perfect field, and  $\mathcal{A}$  is a sheaf of abelian groups on  $\text{Sm}_k$ , then it is strong if and only if it is strict.

The proof of these theorems is hard — to see them worked out in detail we refer the reader to an amazing recent survey paper of Bachmann [Bac24].

**Assumption 3.5.6.** From here on out, we'll assume  $k$  is a perfect field, so that we can access these results.

**Corollary 3.5.7.** If  $X \in \text{Spc}(k)_*$ , then  $\pi_n^{\mathbb{A}^1}(X)$  is strictly invariant for  $n \geq 2$  and strongly invariant for  $n = 1$ .

*Proof.* We apply [Theorem 3.5.4](#) to  $\Omega^{n-1}X$  and get that

$$\pi_n(X) = \pi_1(\Omega X)$$

is strongly invariant, which is strictly invariant for  $n > 1$  by [Theorem 3.5.5](#). □

### 3.5.1 Unramified sheaves

Let  $\mathcal{F}$  be a sheaf of sets. We say it *ramifies* if, for some  $X$ , the map

$$\mathcal{F}(X) \rightarrow \bigcap_{x \in X^{(1)}} \mathcal{F}(\text{Spec } \mathcal{O}_{X,x})$$

has nontrivial kernel (here the intersection takes place in  $\mathcal{F}(\kappa(X))$ ). This is very related to the idea of *purity* for torsors.

**Definition 3.5.8.** [Mor12, 2.1] An *unramified presheaf of sets* on  $\text{Sm}_k$  is a presheaf  $\mathcal{F}$  so that

1. If  $X$  has irreducible components  $\{X_\alpha\}$ , then

$$\mathcal{F}(X) \rightarrow \prod_{\alpha} \mathcal{F}(X_\alpha)$$

is a bijection



2. If  $U \subseteq X$  is open and dense, then the restriction map

$$\mathcal{F}(X) \rightarrow \mathcal{F}(U)$$

is injective

3. The map

$$\mathcal{F}(X) \rightarrow \bigcap_{x \in X^{(1)}} \mathcal{F}(\mathrm{Spec} \mathcal{O}_{X,x})$$

is an injection.

**Example 3.5.9.** Any unramified presheaf of sets is automatically a Zariski sheaf.

**Theorem 3.5.10.** (Morel) Any strictly  $\mathbb{A}^1$ -invariant sheaf is unramified.

**Example 3.5.11.** Some other examples of unramified sheaves:

1. Homotopy modules with transfers
2. Rost cycle modules
3. Unramified étale cohomology (ref needed)

Unramified sheaves are determined in a precise sense by their values on fields, which we will now explain (and eventually reach in Theorem 3.5.17).

### 3.5.2 Unramified sheaves from theories on fields

**Definition 3.5.12.** A *homotopy sheaf* is a strongly (=strictly, therefore unramified)  $\mathbb{A}^1$ -invariant sheaf of abelian groups. We denote by  $\mathrm{HI}(k) \subseteq \mathrm{Ab}_{\mathrm{NIS}}(k)$  the full subcategory of homotopy sheaves.

**Theorem 3.5.13.** [Mor12, 6.24] The subcategory  $\mathrm{HI}(k) \subseteq \mathrm{Ab}_{\mathrm{NIS}}(k)$  is abelian, with exact inclusion.

We will see that elements in  $\mathrm{HI}(k)$  are completely determined by their value on certain fields over  $k$ . Let's fix some notation.

**Notation 3.5.14.**

1. We let  $\mathcal{F}_k \subseteq \mathrm{Alg}_k$  be the full subcategory of fields  $F$  so that  $F/k$  has finite transcendence degree [Mor12, p. vi].
2. We let  $\mathcal{E}_k \subseteq \mathcal{F}_k$  be the full subcategory of those fields  $F$  which are further assumed to be finitely generated over  $k$  [Dég07, p. 43]. These are precisely the *algebraic function fields* over  $k$ , equivalently given as finite field extensions of  $k(t_1, \dots, t_n)$  for some  $n$ .

**Remark 3.5.15.** Since every  $F \in \mathcal{F}_k$  is a (filtered) colimit over its finitely generated subfields, we conclude that  $\mathcal{F}_k$  is generated by  $\mathcal{E}_k$  under filtered colimits. We will be mostly concerned with continuous functors out of  $\mathcal{F}_k$ , which by definition are then determined by their values on  $\mathcal{E}_k$ .

We'll work with unramified sheaves of abelian groups here (c.f. [Fel21, §1.5]), but the more general definitions are for sheaves of pointed sets.

**Definition 3.5.16.** [Mor12, 2.6, 2.9] An *unramified  $\mathcal{F}_k$ -datum* is the data of

**D1** A continuous functor  $M: \mathcal{F}_k \rightarrow \mathrm{Ab}$ .

**D2** For every  $F \in \mathcal{F}_k$  and discrete valuation  $v$  on  $F$ , a subgroup

$$M(\mathcal{O}_v) \subseteq M(F),$$

**D3** For every  $F \in \mathcal{F}_k$  and valuation  $v$  on  $F$  a specialization map

$$s_v: M(\mathcal{O}_v) \rightarrow M(\kappa(v)).$$

This data is subject to the axioms

- A1** If  $(E, v) \subseteq (F, w)$  is a separable extension of discretely valued fields (so that  $w|_E = v$ ), and  $v$  has ramification index 1 on  $E$ <sup>2</sup> then the square commutes

$$\begin{array}{ccc} M(\mathcal{O}_w) & \dashrightarrow & M(\mathcal{O}_w) \\ \downarrow & & \downarrow \\ M(E) & \longrightarrow & M(F). \end{array}$$

If the field extension  $\kappa(v) \rightarrow \kappa(w)$  is an isomorphism then this square is cartesian.

- A2** If  $X$  is irreducible with function field  $F$ , and  $m \in M(F)$ , then  $m$  lies in all but a finite number of  $S(\mathcal{O}_{X,x})$ , where  $x \in X^{(1)}$ .

- A3(i)** If  $(E, w) \subseteq (F, v)$  is an extension in  $\mathcal{F}_k$ , then the diagram commutes:

$$\begin{array}{ccc} M(\mathcal{O}_v) & \dashrightarrow & M(\mathcal{O}_w) \\ \downarrow & & \downarrow \\ M(\kappa(v)) & \longrightarrow & M(\kappa(w)). \end{array}$$

- A3(ii)** If  $E \subseteq F$  is an extension in  $\mathcal{F}_k$  and  $v$  a discrete valuation on  $F$  restricting to zero on  $E$ , and  $j: E \hookrightarrow \kappa(v)$  the induced field extension, then the diagram commutes

$$\begin{array}{ccc} & & M(\kappa(v)) \\ & \nearrow^{M(j)} & \uparrow s_v \\ M(E) & \dashrightarrow & M(\mathcal{O}_v) \\ & \searrow & \downarrow \\ & & M(F). \end{array}$$

- A4(i)** If  $X \in \text{Sm}_k$  is local of dimension two with closed point  $z$ , and  $y_0 \in X^{(1)}$  has smooth reduced scheme  $\bar{y}_0$ , then the diagram commutes

$$\begin{array}{ccc} \bigcap_{y \in X^{(1)}} M(\mathcal{O}_y) & \dashrightarrow & M(\mathcal{O}_{\bar{y}_0, z}) \\ \downarrow & & \downarrow \\ M(\mathcal{O}_{y_0}) & \xrightarrow{s_{y_0}} & M(\kappa(y_0)). \end{array}$$

- A4(ii)** The composite

$$\bigcap_{y \in X^{(1)}} M(\mathcal{O}_y) \rightarrow M(\mathcal{O}_{\bar{y}_0, z}) \rightarrow M(\kappa(z))$$

doesn't depend on the choice of  $y_0$ .

We say that  $M$  is *strongly unramified* if it satisfies some further axioms (see [Mor12, §2]).

**Theorem 3.5.17.** [Mor12, 2.27] By restricting  $\mathcal{F} \in \text{Ab}_{\text{Nis}}(k)$  to its values on  $\mathcal{F}_k$ , we obtain the following results.

1. There is an equivalence of categories between unramified sheaves of abelian groups and unramified  $\mathcal{F}_k$ -data.
2. There is an equivalence of categories between  $\text{HI}(k)$  and *strongly unramified*  $\mathcal{F}_k$  data.

This lets us take theories on fields and bootstrap them up to presheaves of abelian groups! Suppose

---

<sup>2</sup>This means that  $\pi_v = u\pi_w$  for some unit  $u$ , c.f. [Stacks, 09E4].

that  $M$  is an unramified sheaf, and  $X \in \mathrm{Sm}_k$  is an irreducible smooth scheme. Then we can define

$$M(X) := \bigcap_{x \in X^{(1)}} M(\mathcal{O}_x) \subseteq M(F).$$

If  $X$  is smooth with irreducible components  $X_\alpha$ , we define

$$M(X) := \prod_{\alpha \in X^{(0)}} M(X_\alpha).$$

This defines  $M$  on objects. Now if  $f: X \rightarrow Y$  is any morphism in  $\mathrm{Sm}_k$  we can factor it as

$$\begin{array}{ccc} X & \hookrightarrow & Z \\ & \searrow f & \downarrow \\ & & Y, \end{array}$$

where the first map is a closed immersion and the latter is a smooth projection. Factoring the closed immersion as a sequence of closed immersions, each of which is codimension one, we can define  $M(X) \rightarrow M(Z)$  [Mor12, 2.13]. To define  $M(Z) \rightarrow M(Y)$ , we leverage that  $Z \rightarrow Y$  is smooth and define it in terms of the induced map on function fields [Mor12, p. 17].

**Motivation 3.5.18.** We'll leverage this equivalence of categories to construct certain unramified sheaves (Milnor  $K$ -theory, Milnor–Witt  $K$ -theory, Witt theory, etc.) by defining them on fields. We'll then be able to have a strong handle on cohomology with coefficients in these elements in  $\mathrm{HI}(k)$ , and we will better understand how to construct Eilenberg–MacLane spaces for them in the category of motivic spaces.

As an immediate application of this equivalence of categories, we obtain the following.

**Theorem 3.5.19.** Let  $f: F \rightarrow G$  be a morphism of strictly invariant sheaves on  $\mathrm{Sm}_k$ , where  $k$  is a perfect field. Then  $f$  is an isomorphism if and only if

$$f(L): F(L) \rightarrow G(L)$$

is an equivalence for every finitely generated (separable) field extension  $L/k$  ([Mor12, 2.3, 2.8], c.f. [Hoy15, 2.7]).

*Proof.* By the equivalence above, we can restrict to  $\mathcal{F}_k$ , and continuity allows us to further restrict to finite separable field extensions.  $\square$

**Corollary 3.5.20.** Let  $f: X \rightarrow Y$  be a map of motivic spaces in  $\mathrm{Spc}(k)$ , for  $k$  perfect. Then it is an equivalence if and only if it induces an isomorphism on  $\pi_0$ , and for  $n \geq 1$  and every finitely generated field extension  $L/k$  we have that

$$\pi_n(X, x)(L) \rightarrow \pi_n(Y, y)(L)$$

is an equivalence.

## 3.6 Milnor $K$ -theory

By Theorem 3.5.17, we can define unramified sheaves of abelian groups via their values on fields. In the following few sections we provide some examples.

**Definition 3.6.1.** *Milnor  $K$ -theory* of a field  $F$  is defined to be the graded algebra  $K_*^M$  generated by symbols  $\{a\} \in K_1^M$  for  $a \in F^\times$  subject to the relations

- (M1)  $\{a\} \cdot \{1 - a\} = 0$
- (M2)  $\{ab\} = \{a\} + \{b\}.$

**Notation 3.6.2.** We denote by

$$\{a_1, \dots, a_n\} := \{a_1\} \cdots \{a_n\} \in K_n^M(F).$$

**Proposition 3.6.3.** For any field  $F$  we have that

$$\begin{aligned} K_{-n}^M(F) &= 0 \\ K_0^M(F) &= \mathbb{Z} \\ K_1^M(F) &= F^\times. \end{aligned}$$

**Notation 3.6.4.** We denote by  $K_*^M(F) = \bigoplus_{n \geq 0} K_n^M(F)$ . This is a graded abelian group, with multiplication coming from the multiplication of symbols.

### 3.6.1 Basic symbol algebra in $K_*^M$

Much of this can be found in [GS17, §7.1].

An immediate corollary of bilinearity of symbols is the following:

**Proposition 3.6.5.** In  $K_1^M(F)$  we have that  $\{1\} = 0$ .

*Proof.* By (M2), we have that  $\{1 \cdot 1\} = \{1\} + \{1\}$ , hence  $0 = \{1\}$  by subtracting a copy of  $\{1\}$  from either side.  $\square$

**Proposition 3.6.6.** In  $K_2^M(F)$  we have that

1.  $-\{x, y\} = \{x^{-1}, y\}$ .
2. More generally for any  $i, j \in \mathbb{Z}$  we have

$$\{x^i, y^j\} = (i + j) \{x, y\}.$$

**Proposition 3.6.7.** [Mil70, 1.1] Multiplication on  $K_*^M(F)$  is graded commutative, meaning if  $\alpha \in K_m^M(F)$  and  $\beta \in K_n^M(F)$ , we have that

$$\alpha\beta = (-1)^{mn}\beta\alpha.$$

**Proposition 3.6.8.** [Mil70, 1.1, 1.2] We have that

$$\begin{aligned} \{x, -x\} &= 0 \\ \{x, x\} &= \{x, -1\}. \end{aligned}$$

In particular  $2\{x, x\} = 0$  for any  $x$ .

*Proof.* Note for  $x \neq 1$ , we have that

$$-x = \frac{1-x}{1-x^{-1}}.$$

Hence

$$\{x, -x\} = \{x, 1-x\} - \{x, 1-x^{-1}\} = -\{x, 1-x^{-1}\} = \{x^{-1}, 1-x^{-1}\} = 0.$$

This last equality uses Proposition 3.6.6. The argument that  $\{x, x\} = \{x, -1\}$  is similar. The last statement follows from observing that

$$2\{x, x\} = 2\{x, -1\} = \{x, 1\} = 0,$$

since  $\{1\} = 0$ .  $\square$

### 3.6.2 Computations

**Example 3.6.9.** [Wei13, III.6.1, III.7.2] We have that  $K_n^M(\mathbb{F}_q) = 0$  for  $n \geq 2$ .

*Proof.* We will show that it vanishes for  $n = 2$ . Pick  $x$  to generate  $\mathbb{F}_q^\times$ , then any element in  $K_2^M(\mathbb{F}_q)$  is of the form  $\{x^i, x^j\}$ . By Proposition 3.6.6, this is equal to  $(i + j)\{x, x\}$  so it will suffice to verify that  $\{x, x\} = \{x, -1\}$  vanishes. This has order dividing two. We want to show it is killed by an odd number as well, and we'll be done. If  $q = 2^m$ , then  $x^{2^m-1} = 1$ , and hence

$$0 = \{1, x\} = \{x^{2^m-1}, x\} = (2^m - 1)\{x, x\},$$

which concludes the proof. If  $q$  has odd exponential characteristic, we can find two non-squares in  $\mathbb{F}_q$  which sum to 1 (c.f. [GS17, 1.3.6]). This gives us  $x^k + x^\ell = 1$  for  $k, \ell$  odd. Hence we get

$$0 = \{x^k, x^\ell\} = (k + \ell)\{x, x\},$$

and we are done.  $\square$

**Example 3.6.10.** [Wei13, III.7.2] If  $F = \bar{F}$  is algebraically closed, then  $K_n^M(F)$  is uniquely divisible for  $n \geq 1$ . In particular  $K_n^M(F)/\ell = 0$  for any  $n \geq 1$  and any  $\ell \neq 0$ .

These come equipped with subrings

$$K_*^M(\mathcal{O}_v) \subseteq K_*^M(F)$$

for every discrete valuation  $v$  on  $F$ , and specialization maps, which we omit (see Milnor's original paper or [Wei13, 7.3]). These satisfy the axioms for being an unramified  $\mathcal{F}_k$ -datum, proving the following.

**Proposition 3.6.11.** Milnor  $K$ -theory gives rise to an unramified sheaf of abelian groups  $\mathbf{K}_n^M$  for every  $n$ .

We will see later that the sheaf cohomology of  $\mathbf{K}_n^M$  computes the Chow groups of a scheme.

**Proposition 3.6.12.** There is a *symbol map* for any field  $F$  valued in Quillen  $K$ -theory

$$K_n^M(F) \rightarrow K_n(F),$$

which is an isomorphism for  $n \leq 2$ . (For  $n = 0, 1$  this is easy, for  $n = 2$  this is *Matsumoto's theorem*).

## 3.7 Milnor–Witt $K$ -theory

We refer the reader to [Dég23; Car23] for more in-depth discussions of what's found here.

**Definition 3.7.1.** *Milnor–Witt  $K$ -theory* of a field  $F$  is defined to be the graded algebra  $K_*^{\text{MW}}(F)$  defined by symbols  $[a] \in K_1^{\text{MW}}(F)$  for  $a \in F^\times$  and  $\eta \in K_{-1}^{\text{MW}}(F)$ , modulo the relations:

- (MW1)  $[a][1 - a] = 0$  for  $a \neq 0, 1$
- (MW2)  $[ab] = [a] + [b] + \eta[a][b]$
- (MW3)  $\eta[a] = [a]\eta$
- (MW4)  $\eta(2 + \eta[-1]) = 0$ .

It will benefit us to have some notation for various special elements in Milnor–Witt  $K$ -theory.

**Notation 3.7.2.** (*Special elements in  $K_*^{\text{MW}}(F)$* )

1. For any  $a \in F$  we denote by

$$\langle a \rangle := 1 + \eta[a] \in K_0^{\text{MW}}(F).$$

2. We denote by  $h := 1 + \langle -1 \rangle = 2 + \eta[-1]$  the *hyperbolic element*.
3. We denote by  $\epsilon$  the element

$$\epsilon = -\langle -1 \rangle = -(1 + \eta[-1]) \in K_0^{\text{MW}}(F).$$

**Remark 3.7.3.** Observe that relation MW4 can take either of the following equivalent forms

$$\text{(MW4)} \quad \eta h = 0$$

$$\text{(MW4)} \quad \epsilon \eta = \eta.$$

### 3.7.1 Symbol algebra in Milnor–Witt $K$ -theory

We will develop some basic properties, starting in lower degrees and going to higher degrees.

**Proposition 3.7.4.** (*Properties in  $K_0^{\text{MW}}$* )

1.  $\langle a \rangle \langle b \rangle = \langle ab \rangle$  for any  $a, b \in F^\times$ .
2.  $1 = \langle 1 \rangle$  is the multiplicative unit.
3.  $\eta[1] = 0$ .
4.  $\epsilon^2 = 1$

*Proof.*

1. We see that

$$\begin{aligned} \langle a \rangle \langle b \rangle &= (1 + \eta[a]) (1 + \eta[b]) = 1 + \eta([a] + [b]) + \eta^2[a][b] \\ &\stackrel{(\text{MW2})}{=} 1 + \eta([ab] - \eta[a][b]) + \eta^2[a][b] = 1 + \eta[ab] = \langle ab \rangle. \end{aligned}$$

2. By definition, we have that  $\eta[1] = \langle 1 \rangle - 1$ . Multiplying [1] into (MW4) we get that

$$\begin{aligned} 0 &= \eta[1](1 + \langle -1 \rangle) = (\langle 1 \rangle - 1)(1 + \langle -1 \rangle) \\ &= \langle 1 \rangle - 1 + \langle -1 \rangle - \langle -1 \rangle \\ &= \langle 1 \rangle - 1. \end{aligned}$$

Hence  $1 = \langle 1 \rangle$ .

3. Since  $\eta[1] = \langle 1 \rangle - 1$ , we get that  $\eta[1] = 0$  by the previous result.
4. By 1, it is clear that  $\epsilon^2 = \langle 1 \rangle$ , which is equal to 1 by 2.

□

In degree one, we have that  $[1] = 0$ , completely analogous to Milnor  $K$ -theory, and various other properties:

**Proposition 3.7.5.** (*Properties in  $K_1^{\text{MW}}$* ) Some basic properties to record are that  $[1]$  vanishes, analogous to Milnor  $K$ -theory, as well as the following commutativity relations:

1.  $[1] = 0$ .
2.  $\eta[a][b] = \eta[b][a]$
3.  $[a] \langle b \rangle = \langle b \rangle [a]$ .
4.  $\epsilon[a] = [a]\epsilon$

Some further relations let us expand various degree one elements, and are useful in further computation:

4.  $[a^2] = (1 + \langle a \rangle)[a]$
5.  $[a] = -\langle a \rangle [a^{-1}]$
6.  $[ab] = [a] + \langle a \rangle [b]$
7.  $[a/b] = [a] - \langle a/b \rangle [b]$

*Proof.*

1. By MW2 we get

$$[1 \cdot 1] = [1] + [1] + \eta[1][1].$$

Since  $\eta[1] = 0$  by [Proposition 3.7.4](#), we conclude that  $[1] = [1] + [1]$ , from which the result follows.

2. Since  $ab = ba$  in  $F$ , we have that  $[ab] = [ba]$ . Expanding each of these using MW2 gives the desired result.
3. By applying [item 2](#), we see that

$$[a] \langle b \rangle = [a] (1 + \eta[b]) = [a] + \eta[a][b] = [a] + \eta[b][a] = \langle b \rangle [a].$$

4. This follows from [Proposition 3.7.5\(3\)](#) with  $b = -1$  and a negative sign.
5. Applying MW2 we get

$$[a^2] = 2[a] + \eta[a][a] = (2 + \eta[a])[a] = (1 + \langle a \rangle)[a].$$

6. Applying MW2 to  $a, a^{-1}$  we get

$$0 = [1] = [a^{-1}] + [a] + \eta[a^{-1}][a] = [a^{-1}] + \langle a^{-1} \rangle [a],$$

Subtracting  $[a^{-1}]$  from both sides then replacing  $a$  with  $a^{-1}$  yields the desired result.

7. We see that

$$[ab] = [a] + [b] + \eta[a][b] = [a] + \langle a \rangle [b].$$

8. This is a previous case of the previous result:

$$[a] = \left[ \frac{a}{b} b \right] = \left[ \frac{a}{b} \right] + \langle a/b \rangle [b].$$

Rearranging, the result follows. □

In degree two, we see the value of  $\epsilon$ , that it measures the failure of commutativity:

**Proposition 3.7.6.** (*Properties in  $K_2^{\text{MW}}$* )

1.  $[a][-a] = 0$
2.  $[a][a] = \epsilon[a][-1] = \epsilon[-1][a]$
3.  $[a][-1] = [-1][a]$ .
4.  $[a][b] = \epsilon[b][a]$ .

*Proof.*

1. We do the same trick as in [Proposition 3.6.8](#), and write  $-a = \frac{1-a}{1-a^{-1}}$ . By [item 7](#), we get

$$[-a] = \left[ \frac{1-a}{1-a^{-1}} \right] = [1-a] - \langle -a \rangle [1-a^{-1}].$$

Scaling through by  $[a]$  we get

$$[a][-a] = -[a] \langle -a \rangle [1-a^{-1}].$$

So this reduces to showing that

$$[a] \langle -a \rangle [1-a^{-1}] = 0.$$

We can commute  $[a]$  and  $\langle a \rangle$  by [Proposition 3.7.5\(3\)](#), then expand  $[a]$  using [Proposition 3.7.5\(5\)](#) to get

$$\langle -a \rangle [a][1-a^{-1}] = -\langle -a \rangle \langle a \rangle [a^{-1}][1-a^{-1}].$$

The latter two terms multiply to zero by MW1.

2. By [Proposition 3.7.5\(6\)](#), we have

$$[-a] = [-1] + \langle -1 \rangle [a]. \tag{3.7}$$

Multiplying Equation 3.7 by  $[a]$  on the left and applying 1 we get

$$0 = [a][-1] + [a] \langle -1 \rangle [a] = [a][-1] + \langle -1 \rangle [a][a].$$

Rearranging, we get

$$\langle -1 \rangle [a][a] = -[a][-1],$$

and multiplying through by  $\langle -1 \rangle$  gives the desired result.

Multiplying Equation 3.7 by  $[a]$  on the right instead, we get

$$0 = [-1][a] + \langle -1 \rangle [a][a],$$

which gives us a similar equality.

3. By Proposition 3.7.6(2) we have  $\epsilon[a][-1] = \epsilon[-1][a]$ . Multiplying both sides by  $\epsilon$  and using that  $\epsilon^2 = 1$  (Proposition 3.7.4(4)) gives the desired result.
4. We can write

$$\begin{aligned} 0 &= [ab][-ab] = ([a] + \langle a \rangle [b])([-a] + \langle -a \rangle [b]) \\ &= \langle -a \rangle [a][b] + \langle a \rangle [b][-a] + \langle -1 \rangle [b][b]. \end{aligned}$$

Scaling through by  $\langle a \rangle$  we get

$$0 = -\epsilon[a][b] + [b][-a] + \langle a \rangle \langle -1 \rangle [b][b]$$

Since  $[-a] = [a] + \langle a \rangle [-1]$  by Proposition 3.7.5(6), and  $\langle -1 \rangle [b][b] - [b][-1]$  by Proposition 3.7.6(2), we can plug these into the above equation to get

$$\begin{aligned} 0 &= -\epsilon[a][b] + [b]([a] + \langle a \rangle [-1]) - \langle a \rangle [b][-1] \\ &= -\epsilon[a][b] + [b][a] + \langle a \rangle [b][-1] - \langle a \rangle [b][-1] \\ &= -\epsilon[a][b] + [b][a]. \end{aligned}$$

□

### 3.7.2 Comparison to Milnor $K$ -theory and Grothendieck–Witt

**Proposition 3.7.8.** There is a surjective homomorphism of graded algebras

$$\begin{aligned} K_*^{\text{MW}}(F) &\rightarrow K_*^M(F) \\ [a] &\mapsto \{a\} \\ \eta &\mapsto 0. \end{aligned}$$

That is, Milnor  $K$ -theory is obtained from Milnor–Witt  $K$ -theory by killing  $\eta$ .

*Proof.* Relations M1 and M2 are just relations MW1 and MW2 after modding out by  $\eta$ , so the map is well-defined, and it is clearly surjective since  $\{u_1, \dots, u_n\}$  is hit by  $[u_1, \dots, u_n]$ . □

**Definition 3.7.9.** Let  $F$  be a field of characteristic  $\neq 2$ . We define the *Grothendieck–Witt ring* of  $F$ , denoted  $\text{GW}(F)$  to be the ring of isomorphism classes of non-degenerate symmetric bilinear forms, group completed.

**Proposition 3.7.10.**  $\text{GW}(F)$  is generated by rank one forms

$$\begin{aligned} \langle a \rangle : k \times k &\rightarrow k \\ (x, y) &\mapsto axy, \end{aligned}$$

modulo the relations

- (GW1)  $\langle ab^2 \rangle = \langle a \rangle$  for any  $a, b \in F^\times$
- (GW2)  $\langle a \rangle + \langle b \rangle = \langle ab(a+b) \rangle + \langle a+b \rangle$
- (GW3)  $\langle a \rangle \langle b \rangle = \langle ab \rangle$ .



**Exercise 3.7.11.** Show that

$$\langle 1 \rangle + \langle -1 \rangle = \langle a \rangle + \langle -a \rangle$$

for any  $a \in F^\times$ . This is sometimes taken as a relation, but it is implied by (GW1) and (GW2).

**Notation 3.7.12.** We often save space and write

$$\begin{aligned} \{a_1, \dots, a_n\} &:= \{a_1\} \cdots \{a_n\} \in K_n^{\text{MW}}(F) \\ [a_1, \dots, a_n] &:= [a_1] \cdots [a_n] \in K_n^M(F) \\ \langle a_1, \dots, a_n \rangle &:= \langle a_1 \rangle + \dots + \langle a_n \rangle \in \text{GW}(F). \end{aligned}$$

**Proposition 3.7.13.** There is a ring isomorphism

$$\begin{aligned} \text{GW}(F) &\rightarrow K_0^{\text{MW}}(F) \\ \langle a \rangle &\mapsto \langle a \rangle. \end{aligned}$$

*Proof.* We first check this is well-defined, in that the map respects the relations for the Grothendieck–Witt ring. Clearly **(GW3)** holds by **Proposition 3.7.4(1)**. Since  $\langle a \rangle$  is multiplicative, checking **(GW1)** reduces to showing that  $\langle b^2 \rangle = 1$  in  $K_0^{\text{MW}}$ , which reduces to checking that  $\eta[b^2] = 0$ . By MW2, and **Proposition 3.7.5(4)** we have that

$$[b^2] = 2[b] + \eta[b][b] = 2[b] + \eta[-1][b] = (2 + \eta[-1])[b].$$

Multiplying by  $\eta$  and applying MW4 gives zero. Finally we want to check **(GW2)**, and by multiplicativity we can assume that  $b = 1 - a$ , from which we get

$$\begin{aligned} \langle a \rangle + \langle 1 - a \rangle &= 1 + \eta[a] + 1 + \eta[1 - a] \\ &= 2 + \eta([a] + [1 - a]) \\ &\stackrel{(\text{MW2})}{=} 2 + \eta([a(1 - a)] - \eta[a][1 - a]) \\ &\stackrel{(\text{MW1})}{=} 2 + \eta[a(1 - a)] \\ &\stackrel{[1]=0}{=} 1 + \eta[a(1 - a)] + 1 + \eta[1] \\ &= \langle a(1 - a) \rangle + \langle 1 \rangle. \end{aligned}$$

It is clear by construction that this map is injective, so it suffices to see it is surjective by verifying that every element in  $K_0^{\text{MW}}(F)$  is a sum of elements in the image of the homomorphism produced above. Since  $1 = \langle 1 \rangle$ , we have that  $\eta[a] = \langle a \rangle - \langle 1 \rangle$  is in the image of the homomorphism above. Iterated application of (MW2) and (MW3) yields the desired result.  $\square$

Via the equivalence of categories between unramified sheaves of groups and unramified data for fields, we get sheaves corresponding to each of the invariants above.

**Proposition 3.7.14.** [Mor12, p. 71]  $K_n^{\text{MW}}$  is unramified and strongly  $\mathbb{A}^1$ -invariant.

**Warning 3.7.15.** This is a bit of a lie. Unlike the case of Milnor  $K$ -theory, the residue homomorphisms for Milnor–Witt  $K$ -theory depend on a choice of uniformizing parameter, so we have to be careful about twists here.

## 3.8 The Milnor conjecture

**Definition 3.8.1.** We define the *Witt ring* to be the quotient  $W(F) := \text{GW}(F)/h$ .

**Definition 3.8.2.** We define the *fundamental ideal*  $I(F)$  to be the kernel of the modulo two rank homomorphism  $W(F) \rightarrow \mathbb{Z}/2$ . We denote by  $I^j$  the powers of the fundamental ideal.

**Proposition 3.8.3.** We obtain unramified sheaves of groups

$$\mathbf{GW}, \mathbf{W}, \mathbf{I}^j,$$

associated to each of these  $\mathcal{F}_k$ -data.

**Proposition 3.8.4.** There is a pullback square of abelian groups for any field  $F$

$$\begin{array}{ccc} \mathbf{GW}(F) & \longrightarrow & \mathbb{Z} \\ \downarrow & \lrcorner & \downarrow \\ \mathbf{W}(F) & \longrightarrow & \mathbb{Z}/2. \end{array}$$

This extends to a pullback of unramified sheaves of groups.

**Proposition 3.8.5.**

1. The fundamental ideal is equivalently the kernel of the rank homomorphism  $\mathbf{GW}(F) \rightarrow \mathbb{Z}(F)$ , which we denote by  $\hat{I}(F)$ . Explicitly, there is an isomorphism

$$\begin{array}{ccccc} \hat{I}(F) & \longrightarrow & \mathbf{GW}(F) & \longrightarrow & \mathbb{Z} \\ \cong \downarrow & & \downarrow & \lrcorner & \downarrow \\ I(F) & \longrightarrow & \mathbf{W}(F) & \longrightarrow & \mathbb{Z}/2. \end{array}$$

2.  $\hat{I}(F)$  is generated by elements of the form  $\langle 1 \rangle - \langle a \rangle$  for  $a \in F^\times$
3. The isomorphism  $\hat{I}(F) \rightarrow I(F)$  is given by sending

$$\begin{aligned} \hat{I}(F) &\rightarrow I(F) \\ \langle 1 \rangle - \langle a \rangle &\mapsto \langle 1, -a \rangle. \end{aligned}$$

**Definition 3.8.6.** For  $a \in F^\times$  we denote by  $\langle\langle a \rangle\rangle := \langle 1 \rangle - \langle a \rangle$  the *Pfister form* attached to  $a$ .

**Remark 3.8.7.** (On notation) Authors in the literature use  $\langle\langle a \rangle\rangle$  to denote  $\langle 1 \rangle - \langle a \rangle \in \mathbf{GW}(F)$  as we have done, or to denote  $\langle 1, -a \rangle \in \mathbf{W}(F)$  (see e.g. [EKM08, p. 24]. We should be careful about the context when using this notation, and we also warn the reader that a different sign convention is used in [Lam05, X.1.1].

**Example 3.8.8.** (*Examples of fundamental ideals*)

1. Every Pfister form is hyperbolic over an algebraically closed field, or even just a quadratically closed field. Hence  $I(F) = 0$  if  $F = \bar{F}$ .
2.  $I(\mathbb{R}) \cong \mathbb{Z}$ , generated by  $\langle\langle -1 \rangle\rangle$ . We can check that

$$\langle\langle -1 \rangle\rangle^n = 2^{n-1} \langle\langle -1 \rangle\rangle,$$

therefore  $I(\mathbb{R}) \cong I^2(\mathbb{R}) \cong \dots \cong I^n(\mathbb{R}) \cong \mathbb{Z}$ , and we have that

$$I^n(\mathbb{R})/I^{n+1}(\mathbb{R}) \cong \mathbb{Z}/2\mathbb{Z}.$$

3. This is a more general fact for any field that

$$I(F)/I^2(F) \cong F^\times / (F^\times)^2.$$

This is a result of Pfister (c.f. [Lam05, II.2.3]).

4. For any  $F$  we have that

$$I^2(F)/I^3(F) \cong \mathrm{Br}(F)[2]$$

is the 2-torsion in the Brauer group.

**Proposition 3.8.9.** Given two quadratic forms  $\alpha, \beta \in \mathbf{GW}(F)$ , they agree if and only if they agree modulo  $I^n(F)$  for every  $n$ , and therefore if and only if they agree in the associated graded  $I^n(F)/I^{n+1}(F)$  for every  $n$ .

*Proof idea.* It suffices to argue that

$$\bigcap_{n=0}^{\infty} I^n(F) = 0,$$

since we can show that if  $\alpha - \beta \in \bigcap_{n=0}^{\infty} I^n(F)$ , then  $\alpha = \beta$ . The question of whether this intersection is zero was first raised by Milnor [Mil70, 4.4], and proven shortly thereafter by Aarason and Pfister as a consequence of their Hauptsatz [Lam05, X.5.1].  $\square$

**Proposition 3.8.10.** (Milnor) There is a homomorphism

$$\begin{aligned} K_n^M(F)/2 &\rightarrow I^n(F)/I^{n+1}(F) \\ \{a_1, \dots, a_n\} &\mapsto \prod_{i=1}^n \langle\langle a_i \rangle\rangle. \end{aligned} \tag{3.11}$$

This homomorphism was constructed and shown to be surjective by Milnor [Mil70, 4.1], who conjectured it was bijective for all  $n$  [Mil70, 4.3].

**Theorem 3.8.12.** (Milnor Conjecture, Orlov–Vishik–Voevodsky) The homomorphism Equation 3.11 is an isomorphism. Moreover, we obtain an isomorphism of unramified sheaves of groups

$$\mathbf{K}_n^M(F)/2 \xrightarrow{\sim} \mathbf{I}^n/\mathbf{I}^{n+1}.$$

**Motivation 3.8.13.** We'll double back and compute cohomology in these theories. We first want to show these cohomology groups are *representable* by Eilenberg–MacLane spaces. Formal consequences of representability will motivate the construction of complexes which resolve these sheaves and let us carry out computations more directly.

**Corollary 3.8.14.** There is a pullback diagram of unramified sheaves

$$\begin{array}{ccc} \mathbf{K}_n^{\text{MW}} & \longrightarrow & \mathbf{K}_n^M \\ \downarrow & \lrcorner & \downarrow \\ \mathbf{I}^n & \longrightarrow & \mathbf{K}_n^M/2. \end{array}$$

**Remark 3.8.15.**

1. Note that when  $n = 0$ , the pullback diagram in Corollary 3.8.14 recovers that of Proposition 3.8.4.
2. The map  $\mathbf{K}_n^{\text{MW}} \rightarrow \mathbf{I}^n$  is given by sending  $[a]$  to  $\langle\langle a \rangle\rangle$ .

## 3.9 Eilenberg–MacLane spaces

We denote by  $B_{\text{mot}}G := L_{\text{mot}}BG$ , and we have seen the following:

**Proposition 3.9.1.** If  $X$  is a scheme, then

$$\pi_0 \text{Map}_{\text{Spc}(k)}(L_{\text{mot}}h_X, B_{\text{mot}}G) \cong H_{\text{Nis}}^1(X, G).$$

We wonder whether higher Nisnevich cohomology is represented by Eilenberg–MacLane spaces? The answer is yes!

**Theorem 3.9.2.** (Dold–Kan) If  $\text{Ab}_{\text{Nis}}(k)$  denotes the category of Nisnevich sheaves of abelian groups on  $X$ , there is an equivalence of categories

$$\text{Ch}_{\geq 0}(\text{Ab}_{\text{Nis}}(k)) \cong \text{Fun}(\Delta^{\text{op}}, \text{Ab}_{\text{Nis}}(k)).$$

Given a chain complex of abelian sheaves, we can view it as an object on the right hand side. Forgetting the levelwise sheaf structure, we can view it as a simplicial presheaf of abelian groups, and therefore a presheaf of simplicial sets:

$$\mathrm{Fun}(\Delta^{\mathrm{op}}, \mathrm{Ab}_{\mathrm{Nis}}(k)) \subseteq \mathrm{Fun}(\Delta^{\mathrm{op}}, \mathrm{Fun}(\mathrm{Sm}_k^{\mathrm{op}}, \mathrm{Ab})) \rightarrow \mathrm{Fun}(\Delta^{\mathrm{op}}, \mathrm{Fun}(\mathrm{Sm}_k^{\mathrm{op}}, \mathrm{Set})) \cong \mathrm{PSh}(\mathrm{Sm}_k).$$

We'll note by DK this composite:

$$\mathrm{DK}: \mathrm{Ch}_{\geq 0}(\mathrm{Ab}_{\mathrm{Nis}}(k)) \rightarrow \mathrm{PSh}(\mathrm{Sm}_k).$$

**Proposition 3.9.3.** If  $\mathcal{A} \in \mathrm{Ch}_{\geq 0}(\mathrm{Ab}_{\mathrm{Nis}}(k))$ , then there is an isomorphism

$$H_n(\mathcal{A}) \cong \pi_n(L_{\mathrm{Nis}}\mathrm{DK}(\mathcal{A})).$$

*Proof.* This follows from the more general statement that the homology of a chain complex agrees with the simplicial homotopy groups of the associated simplicial abelian group produced by the Dold–Kan correspondence (c.f. [GJ99, III.2.5]).  $\square$

**Definition 3.9.4.** For any  $A \in \mathrm{Ab}_{\mathrm{Nis}}(k)$  we denote by

$$K(A, n) \in \mathrm{PSh}(\mathrm{Sm}_k)$$

the space  $K(A, n) := \mathrm{DK}(A[n])$  given by applying the Dold–Kan construction to the chain complex with  $A$  concentrated in degree  $n$ .

**Proposition 3.9.5.** (*Properties of  $K(A, n)$* )

1. We have that  $K(A, n)$  is already an object in the sheaf topos  $\mathrm{Shv}_{\mathrm{Nis}}(\mathrm{Sm}_k)$
2. We have that

$$\pi_i K(A, n) = \begin{cases} A & i = n \\ 0 & \text{else} \end{cases}$$

There is a natural identification

$$\pi_0 \mathrm{Map}_{\mathrm{Shv}_{\mathrm{Nis}}}(-, K(A, n)) \cong H_{\mathrm{Nis}}^n(-, A).$$

*Proof references.* The classical way to approach this is using explicit model structures on simplicial presheaves and chain complexes of sheaves of abelian groups, and arguing that the construction of an Eilenberg–MacLane object preserves fibrancy, which implies that  $K(A, n)$  is a sheaf in the local model structure. See [Mor12, Chapter 6] or [MV99, pp.56–59] for this approach. The high-level perspective on Eilenberg–MacLane objects in a general  $\infty$ -topos is in [Lur09, §7.2.2], from which these properties are formal.  $\square$

### 3.9.1 Strong and strict invariance revisited

With EM spaces in hand in the sheaf topos, we can reframe our definitions of strong and strict invariance.

**Proposition 3.9.6.** Let  $\mathcal{G}$  be a sheaf of groups and  $\mathcal{A}$  a sheaf of abelian groups. Then

1.  $\mathcal{G}$  is strongly invariant if and only if  $B_{\mathrm{Nis}}\mathcal{G}$  is  $\mathbb{A}^1$ -local
2.  $\mathcal{A}$  is strictly invariant if and only if  $K(\mathcal{A}, n)$  is  $\mathbb{A}^1$ -local for every  $n \geq 0$ .

*Proof.* The backwards direction is immediate by representability of cohomology. The forwards direction needs a nontrivial argument, see [Bac24, 1.5]. For the forwards direction, we want to see that

$$(B_{\mathrm{Nis}}G)(X) \rightarrow (B_{\mathrm{Nis}}G)(X \times \mathbb{A}^1)$$

is an equivalence of spaces. Each space is 1-truncated and we have an isomorphism on  $\pi_0$  since  $H^0(-, G)$  is  $\mathbb{A}^1$  invariant. So we just want to show an isomorphism on  $\pi_1$  for each choice of basepoint  $x \in X$ . Since the presheaf

$$(\mathrm{Sm}_k)_{/X} \rightarrow \mathrm{Grp} \\ Y \mapsto \pi_1((B_{\mathrm{Nis}}G)(Y), x)$$

is a Nisnevich sheaf (equivalent to  $\Omega_x(B_{\mathrm{Nis}}G)_{|X}$ ), we can check it is  $\mathbb{A}^1$ -invariant locally. Locally  $x$  is a trivial torsor, in which case the presheaf above is  $H^0(-, G)$  which we assumed  $\mathbb{A}^1$ -invariant.

An analogous argument works for the forward direction of  $K(\mathcal{A}, n)$ .  $\square$

We now have a list of nice properties that homotopy sheaves satisfy. For  $\mathcal{A} \in \mathrm{HI}(k)$  we have that

1.  $\mathcal{A}$  is determined by its underlying  $\mathcal{F}_k$ -datum
2.  $K(\mathcal{A}, n)$  is a motivic space for each  $n \geq 0$
3.  $H_{\mathrm{Nis}}^n(-, \mathcal{A})$  is  $\mathbb{A}^1$ -invariant for any  $n \geq 0$
4. Isomorphisms between  $\mathcal{A} \rightarrow \mathcal{B}$  in  $\mathrm{HI}(k)$  can be checked on separable field extensions of the base
5.  $\mathcal{A}$  lives in an abelian category  $\mathrm{HI}(k)$

**Proposition 3.9.7.** For any sheaf of abelian groups, we have that

$$\Omega K(A, n) \cong K(A, n-1).$$

*Proof.* The statement in the sheaf topos is a formal consequence of the fact that  $\pi_i(\Omega X) = \pi_{i+1}X$ .  $\square$

If  $A$  is strictly invariant, we will prove that this is an equivalence of motivic spaces as well. It's not immediately obvious that this is the case — we first need to know that computing loops in the sheaf topos agrees with computing loops in motivic spaces. The following argument lets us prove this.

**Remark 3.9.8.** If  $B \in \mathrm{Shv}_{\mathrm{Nis}}(\mathrm{Sm}_k)$  is  $\mathbb{A}^1$ -local, and  $\Omega B$  is  $\mathbb{A}^1$ -local, then

$$\Omega B = \lim_{\mathrm{Spc}(k)} (* \rightarrow B \leftarrow *) = \lim_{\mathrm{Shv}_{\mathrm{Nis}}(\mathrm{Sm}_k)} (* \rightarrow B \leftarrow *).$$

**Corollary 3.9.9.** If  $A$  is strictly  $\mathbb{A}^1$ -invariant, we have an identification in  $\mathrm{Spc}(k)$  of the form

$$\Omega^{1,0} K(A, n) \simeq K(A, n-1).$$

### 3.9.2 Cofiber sequences

We say that

$$X \rightarrow Y \rightarrow C$$

is a *cofiber sequence* in  $\mathrm{Spc}(k)_*$  if the following diagram is a pushout

$$\begin{array}{ccc} X & \longrightarrow & Y \\ \downarrow & & \downarrow \\ * & \longrightarrow & C. \end{array}$$

Since adding a disjoint basepoint is a left adjoint, it doesn't matter if  $X$  and  $Y$  are pointed or not, only  $C$ . In this setting, we have the following result.

**Proposition 3.9.10.** For  $G$  any strongly invariant sheaf of groups and any cofiber sequence  $X \rightarrow Y \rightarrow C$ , we obtain a long exact sequence

$$\begin{aligned} 0 \rightarrow H_{\mathrm{Nis}}^0(C; G) \rightarrow H_{\mathrm{Nis}}^0(Y, G) \rightarrow H_{\mathrm{Nis}}^0(X; G) \\ \rightarrow H_{\mathrm{Nis}}^1(C; G) \rightarrow H_{\mathrm{Nis}}^1(Y, G) \rightarrow H_{\mathrm{Nis}}^1(X; G) \end{aligned}$$

if  $G$  is a sheaf of abelian groups (hence strictly invariant), this continues to  $H^2$  and so on.

**Proposition 3.9.11.** If  $X \rightarrow Y \rightarrow C$  is a cofiber sequence in  $\mathrm{Spc}(k)$ , and  $B \in \mathrm{Spc}(k)$  is arbitrary, then

$$X \times B \rightarrow Y \times B \rightarrow C \wedge B_+$$

is a cofiber sequence.

*Proof.* This is a direct consequence of universality of colimits in  $\mathrm{Spc}(k)$ . Alternatively, we can use that  $B \times -$  is a left adjoint, to see that

$$\begin{array}{ccc} X \times B & \longrightarrow & Y \times B \\ \downarrow & \lrcorner & \downarrow \\ B & \longrightarrow & C \times B \end{array}$$

is a pushout. The induced maps on the cofibers of the horizontal arrows will be an equivalence, and the bottom is clearly  $C \wedge B_+$ .  $\square$

**Example 3.9.12.** For any motivic space  $B$ , we have a cofiber sequence

$$\mathbb{G}_m \times B \rightarrow B \rightarrow \mathbb{P}^1 \wedge B_+.$$

This follows from [Example 3.3.6](#) and [Proposition 3.9.11](#).

### 3.10 Contraction

**Definition 3.10.1.** If  $F$  is a sheaf of pointed sets, we define the *contraction* of  $F$  to be the sheafification of the presheaf  $F_{-1}$ , where  $F_{-1}$  is defined to be the kernel in the short exact sequence

$$0 \rightarrow F_{-1}(U) \rightarrow F(U \times \mathbb{G}_m) \rightarrow F(U) \rightarrow 0,$$

and the latter map is  $\mathrm{id} \times 1: U \rightarrow U \times \mathbb{G}_m$ . See [\[ABH23, 2.1.10\]](#).

**Remark 3.10.2.** Since  $F(U \times \mathbb{G}_m) \rightarrow F(U)$  is split by the inclusion of units, contraction is equivalently defined as the cokernel of the projection off of  $\mathbb{G}_m$ :

$$0 \rightarrow F(U) \rightarrow F(U \times \mathbb{G}_m) \rightarrow F_{-1}(U) \rightarrow 0.$$

**Proposition 3.10.3.** For any  $U \in \mathrm{Sm}_k$  and any strongly  $\mathbb{A}^1$ -invariant sheaf of groups  $G$ , we have that  $H^0(\mathbb{P}^1 \wedge U_+, G) = 0$ .

*Proof.* This follows by connectivity – smashing with  $\mathbb{P}^1$  (and hence  $S^1$ ) makes  $\mathbb{P}^1 \wedge U_+$  connected, so every map into  $G = K(G, 0)$  is trivial.  $\square$

**Remark 3.10.4.** If  $G$  is strongly  $\mathbb{A}^1$ -invariant, we have our cofiber sequence

$$\mathbb{G}_m \times U \rightarrow U \rightarrow \mathbb{P}^1 \wedge U_+.$$

Let's look at the long exact sequence with coefficients in  $G$ , together with the vanishing of  $H^0(\mathbb{P}^1 \wedge U_+, G)$  by [Proposition 3.10.3](#). Then there is an induced map out of the contraction

$$\begin{array}{ccccccc} 0 & \longrightarrow & G(U) & \hookrightarrow & G(U \times \mathbb{G}_m) & \longrightarrow & H^1(\mathbb{P}^1 \wedge U_+) \longrightarrow \cdots \\ & & & & \downarrow & \nearrow \text{dashed} & \\ & & & & G_{-1}(U) & & \end{array}$$

**Lemma 3.10.5.** [Mor12, 2.34] Let  $G$  be strongly  $\mathbb{A}^1$ -invariant and  $U \in \text{Sm}_k$ . Then the canonical map:

$$G_{-1}(U) \rightarrow H^1(\mathbb{P}^1 \wedge U_+, G).$$

is a bijection (an isomorphism if  $G$  is abelian).

*Proof.* The map

$$H^1(U, G) \rightarrow H^1(\mathbb{G}_m \times U, G)$$

is split by evaluation at one and hence injective, so the sequence

$$0 \rightarrow G(U) \rightarrow G(U \times \mathbb{G}_m) \rightarrow H^1(\mathbb{P}^1 \wedge U_+) \rightarrow 0$$

is exact, from which the result follows.  $\square$

**Corollary 3.10.6.** For any strongly  $\mathbb{A}^1$ -invariant sheaf of abelian groups, we have that

$$\Omega^{2,1}K(G, 1) = K(G_{-1}, 0) = G_{-1}.$$

*Proof.* We just verified there is a canonical bijection of presheaves

$$G_{-1}(-) \cong H^1(\mathbb{P}^1 \wedge (-)_+, G) = \pi_0 \text{Map}(\Sigma^{2,1}(-)_+, BG) = \pi_0 \text{Map}((-)_+, \Omega^{2,1}BG).$$

$\square$

This suggests that  $\mathbb{A}^1$ -invariance of a sheaf is related to  $\mathbb{A}^1$ -invariance of its contraction.

**Proposition 3.10.7.** (Properties of contraction), c.f. [Bac24, §4.1]

1. If  $F$  is a sheaf of (abelian) groups, so is  $F_{-1}$
2. If  $F$  is  $\mathbb{A}^1$ -invariant, so is  $F_{-1}$ .

*Proof.* The first statement is clear from its definition as a presheaf. The second statement follows from Corollary 3.10.6 — if  $G$  is strongly invariant, then  $K(G_{-1}, 0) = G_{-1}$  is a motivic space, hence  $\mathbb{A}^1$ -invariant.  $\square$

**Lemma 3.10.8.** If  $G$  is a strongly  $\mathbb{A}^1$ -invariant sheaf of groups, then

$$\Omega_{\mathbb{G}_m} K(G, 1) \simeq K(G_{-1}, 1).$$

*Proof sketch.* Easy to check that  $\pi_1 K(G_{-1}, 1) = G_{-1}$ . What's less obvious is  $\Omega_{\mathbb{G}_m} K(G, 1)$  is connected. We've learned we can check connectivity on fields, so this reduces to asking that

$$[\text{Spec}(F)_+, \Omega_{\mathbb{G}_m} K(G, 1)]_{\mathbb{A}^1}$$

is trivial for each  $F/k$  finitely generated. By adjunction and representability of cohomology, this is asking whether  $H^1(\mathbb{G}_m(F), G)$  is trivial. We can pass to the Zariski site by some formal nice properties of strongly invariant sheaves of groups (c.f. [Mor12, 2.24]), and then we can conclude by an argument involving 1-cocycles (see e.g. [Bac24, 4.2]).  $\square$

**Corollary 3.10.9.** 1. If  $G$  is a strongly  $\mathbb{A}^1$ -invariant sheaf of groups, then so is  $G_{-1}$ .

2. If  $A$  is a strictly  $\mathbb{A}^1$ -invariant sheaf of abelian groups. Then there is a canonical equivalence

$$\Omega_{\mathbb{G}_m} K(A, n) \xrightarrow{\sim} K(A_{-1}, n).$$

*Proof.*

1. Since  $G$  was strongly  $\mathbb{A}^1$ -invariant, we have that  $B_{\text{Nis}}G = K(G, 1)$  is  $\mathbb{A}^1$ -local by Proposition 3.9.6. Since  $\Omega_{\mathbb{G}_m}$  preserves  $\mathbb{A}^1$ -invariant objects, we have that  $\Omega_{\mathbb{G}_m} B_{\text{Nis}}G$  is  $\mathbb{A}^1$ -local as well. Hence the result follows.

2. It is clear by adjunction that

$$\pi_n \Omega^{1,1} K(A, n) = A_{-1},$$

and the higher homotopy groups vanish. To check that  $\pi_i \Omega^{1,1} K(A, n) = 0$  for  $i < n$ , it suffices to see that  $H^i(\mathbb{G}_m(F), A) = 0$  for all  $i < n$ . This is clear for  $i \geq 2$  since  $\mathbb{G}_m$  has Krull dimension one, and the case  $i = 1$  follows by the proof of [Lemma 3.10.8](#).  $\square$

**Remark 3.10.10.** It is clear from [Corollary 3.10.9\(1\)](#) that if  $A$  is strictly  $\mathbb{A}^1$ -invariant then so is  $A_{-1}$  by [Theorem 3.5.5](#). However the argument in [Corollary 3.10.9\(2\)](#) also allows us to prove that contraction preserves strict invariance without reference to the “strong=strict” theorem.

**Corollary 3.10.11.** We have that

$$\Omega^{2,1} K(A, n) \cong K(A_{-1}, n - 1).$$

This is a higher-dimensional generalization of [Corollary 3.10.6](#).

### 3.10.1 Examples of contraction

We want to compute contractions of these sheaves we’ve developed, being Milnor/Milnor–Witt  $K$ -theory, etc.. To do this, we introduce the so-called *Milnor exact sequence*.

**Theorem 3.10.12.** [[Mor12](#), 3.24] For any field  $F$ , we have a split short exact sequence of abelian groups (actually of  $K_*^{\text{MW}}(F)$ -modules):

$$0 \rightarrow K_n^{\text{MW}}(F) \rightarrow K_n^{\text{MW}}(F(t)) \xrightarrow{\sum \partial_{(p)}} \oplus_p K_{n-1}^{\text{MW}}(F[t]/p) \rightarrow 0,$$

where  $p$  runs over monic irreducible polynomials  $p(t) \in F[t]$ . Phrased differently, this is a short exact sequence

$$0 \rightarrow K_n^{\text{MW}}(F) \rightarrow K_n^{\text{MW}}(F(t)) \rightarrow \oplus_{x \in (\mathbb{A}^1)^{(1)}} K_{n-1}^{\text{MW}}(\kappa(x)) \rightarrow 0.$$

**Corollary 3.10.13.** We have that

$$(\mathbf{K}_n^{\text{MW}})_{-1} \cong \mathbf{K}_{n-1}^{\text{MW}}.$$

*Proof.* It suffices to check on fields. Take another exact sequence almost identical to the Milnor one, but without the valuation at zero.

$$\begin{array}{ccccc} K_n^{\text{MW}}(F \times \mathbb{G}_m) & \longrightarrow & K_n^{\text{MW}}(F(t)) & \longrightarrow & \oplus_{x \neq 0} K_{n-1}^{\text{MW}}(\kappa(x)) \\ \uparrow & & \parallel & & \\ K_n^{\text{MW}}(F) & \longrightarrow & K_n^{\text{MW}}(F(t)) & \longrightarrow & \oplus_x K_{n-1}^{\text{MW}}(\kappa(x)) \end{array}$$

Since all these sequences are split exact, we get

$$K_n^{\text{MW}}(F \times \mathbb{G}_m) = K_n^{\text{MW}}(F) \oplus K_{n-1}^{\text{MW}}(F),$$

where the latter is the target of the residue map at zero. Since the contraction  $F_{-1}(U)$  is alternatively described as a summand in  $F_{-1}(\mathbb{G}_m \times U)$  complementary to  $F_{-1}(U)$ , the result follows.  $\square$

**Corollary 3.10.14.** We have that

$$(\mathbf{K}_n^{\text{M}})_{-1} \cong \mathbf{K}_{n-1}^{\text{M}} \text{ and } (\mathbf{I}_n)_{-1} \cong \mathbf{I}_{n-1}.$$

*Proof.* A nearly identical argument to [Corollary 3.10.13](#) works. Alternatively, we can leverage some techniques of dévissage and compactly supported cohomology to prove it, as in [[AF14c](#), 2.9].  $\square$



**Note 3.10.15.** There should(?) be a more direct algebraic argument that  $(\mathbf{K}_n^M)_{-1} \cong \mathbf{K}_{n-1}^M$  which doesn't make reference to the Milnor exact sequence, and just leverages the symbol algebra of Milnor  $K$ -theory.

### 3.10.2 Contraction is exact

Contraction yields a functor from homotopy sheaves to itself:

$$(-)_{-1} : \mathrm{HI}(k) \rightarrow \mathrm{HI}(k).$$

**Proposition 3.10.16.** For any  $A \in \mathrm{HI}(k)$  and any function field  $E \in \mathcal{E}_k$  we have that  $H^1(\mathbb{G}_m \times \mathrm{Spec}(E); A) = 0$  (see [Dég07, 4.10]).

**Proposition 3.10.17.** As an endofunctor on  $\mathrm{HI}(k)$ , contraction is exact.

*Proof.* Suppose that

$$A \rightarrow B \rightarrow C$$

is an exact sequence in  $\mathrm{HI}(k)$ . Since  $H^1(\mathbb{G}_m \times -, A) = 0$  by Proposition 3.10.16, the sequence

$$0 \rightarrow A(\mathbb{G}_m \times -) \rightarrow B(\mathbb{G}_m \times -) \rightarrow C(\mathbb{G}_m \times -) \rightarrow 0$$

is exact as well. Finally since  $A_{-1}(U) \oplus A(U) \cong A(\mathbb{G}_m \times U)$  and this splitting is natural in  $A$  and  $U$ , the result follows.  $\square$

**Corollary 3.10.18.** Exactness of contraction implies that  $(M/n)_{-1} = M_{-1}/n$  for any  $M \in \mathrm{HI}(k)$  and any  $n \in \mathbb{Z}$ . It also implies quotients pass through contraction — in particular the isomorphism

$$\mathbf{K}_n^M/2 \cong \mathbf{I}^n/\mathbf{I}^{n+1}$$

are compatible with contraction.

## 3.11 Homotopy modules

We now want to take these properties that  $\mathbf{K}_n^{\mathrm{MW}}$ ,  $\mathbf{K}_n^M$ , and  $\mathbf{I}^n$  all satisfy and axiomatize them. First we establish further properties about homotopy sheaves.

### 3.11.1 Monoidal structure on $\mathrm{HI}(k)$

We claim that  $\mathrm{HI}(k)$  admits a closed symmetric monoidal structure. The origin of this perhaps makes sense with reference to other objects — it is the unique symmetric monoidal structure making the forgetful map from the category of “sheaves with transfers” symmetric monoidal [Dég11, Lemme 1.8], alternatively it is the unique symmetric monoidal structure induced from that on the  $\mathbb{A}^1$ -derived category c.f. [Fel21, 1.5.1.20]. We won't define it explicitly, but just comment that it exists, and explore some key properties:

**Theorem 3.11.1.** The category  $\mathrm{HI}(k)$  admits a closed symmetric monoidal structure, which we denote by  $\otimes_{\mathrm{HI}}$ . The unit for the monoidal structure is given by the constant sheaf  $\mathbb{Z}$ .

We'll come back and add a proof for this later. The slick proofs use a bit more machinery than we've developed so far, see e.g. [Fel21, 4.2.1.20].

Since the monoidal structure is closed, we have an internal hom object  $\underline{\mathrm{Hom}}_{\mathrm{HI}(k)}(-, -)$ . Since contraction was a loop space construction, we might expect it to be a right adjoint, and indeed this is true!

**Proposition 3.11.2.** For any  $M \in \mathrm{HI}(k)$ , we have an isomorphism

$$M_{-1} \cong \underline{\mathrm{Hom}}_{\mathrm{HI}(k)}(\mathbf{K}_1^{\mathrm{MW}}, M),$$

which is natural in  $M$ .

There is a  $\mathbf{K}_0^{\mathrm{MW}}$ -module structure on any  $M_{-1}$  coming from its action on  $\mathbf{K}_1^{\mathrm{MW}}$ :

$$\begin{aligned} \mathbf{K}_0^{\mathrm{MW}} \times \mathbf{K}_1^{\mathrm{MW}} &\rightarrow \mathbf{K}_1^{\mathrm{MW}} \\ (\langle a \rangle, \beta) &\mapsto \langle a \rangle \cdot \beta. \end{aligned}$$

So multiplication by  $\langle a \rangle$  induces an endomorphism of  $\mathbf{K}_1^{\mathrm{MW}}$ , which therefore induces an endomorphism

$$M_{-1} \cong \underline{\mathrm{Hom}}_{\mathrm{HI}}(\mathbf{K}_1^{\mathrm{MW}}, M) \xrightarrow{\langle a \rangle \cdot -} \underline{\mathrm{Hom}}_{\mathrm{HI}}(\mathbf{K}_1^{\mathrm{MW}}, M) \cong M_{-1}.$$

**Corollary 3.11.3.** There is an adjunction

$$\mathbf{K}_1^{\mathrm{MW}} \otimes_{\mathrm{HI}} - : \mathrm{HI}(k) \rightleftarrows \mathrm{HI}(k) : (-)_{-1}.$$

Since contraction is exact, the identification above states that  $\mathrm{Hom}(\mathbf{K}_1^{\mathrm{MW}}, -)$  is exact, implying that  $\mathbf{K}_1^{\mathrm{MW}}$  is projective as an object in the abelian category  $\mathrm{HI}(k)$ . This implies it is flat (since  $\mathrm{HI}(k)$  is a module category), and it is moreover faithfully flat.

**Proposition 3.11.4.** [Dég11, 1.15] We have that  $\mathbf{K}_1^{\mathrm{MW}} \otimes_{\mathrm{HI}} -$  is fully faithful.

A key property of the tensor product is the following

**Proposition 3.11.5.** We have that

$$\mathbf{K}_m^{\mathrm{MW}} \otimes_{\mathrm{HI}(k)} \mathbf{K}_n^{\mathrm{MW}} \cong \mathbf{K}_{m+n}^{\mathrm{MW}}.$$

We had these isomorphisms  $\mathbf{I}^{n-1} \xrightarrow{\sim} \mathbf{I}_{-1}^n$ . So now under adjunction it seems natural to look at their mates

$$\mathbf{K}_1^{\mathrm{MW}} \otimes_{\mathrm{HI}} \mathbf{I}^{n-1} \rightarrow \mathbf{I}^n.$$

In particular we want to take this analogy kind of seriously:

$$\begin{aligned} \mathbf{K}_1^{\mathrm{MW}} &\rightsquigarrow \Sigma \\ (-)_{-1} &\rightsquigarrow \Omega \end{aligned}$$

a sequence of homotopy sheaves  $\{\mathbf{K}_n^{\mathrm{MW}} \otimes_{\mathrm{HI}} M\}_{n \geq 0} \rightsquigarrow$  a suspension spectrum  $\{\Sigma^n X\}_{n \geq 0}$

a sequence where the mates are equivalences  $\rightsquigarrow$  a spectrum

In particular  $\mathbf{I}^*$ ,  $\mathbf{K}_*^{\mathrm{MW}}$  and  $\mathbf{K}_*^{\mathrm{M}}$  would all be *spectra* in the above analogy. This idea of spectra admits a name, called *homotopy modules*. Let's make this definition precise.

### 3.11.2 The definition of homotopy modules

**Definition 3.11.6.** (c.f. [Fel21, 3.4.1.2]) A *homotopy module* is a pair  $(M_*, \omega_*)$  where  $M_*$  is a  $\mathbb{Z}$ -graded strictly invariant sheaf of graded abelian groups (i.e.  $M_n \in \mathrm{HI}(k)$  for each  $n$ ), and

$$\omega_n : M_{n-1} \xrightarrow{\sim} (M_n)_{-1}$$

is an isomorphism called a *desuspension*. A morphism of homotopy sheaves is a levelwise homomorphism compatible with desuspensions. This gives a category  $\mathrm{HM}(k)$ , we call the category of *homotopy modules*.

**Remark 3.11.7.** Alternatively, we can describe  $\mathrm{HM}(k)$  as the localization  $\mathrm{HM}(k) = \mathrm{HI}(k) [\mathbf{K}_1^{\mathrm{MW}} \otimes_{\mathrm{HI}} -]$ . It is also abelian, and we will prove it is symmetric monoidal.

Let's note some natural functors. We have

$$\begin{aligned}\omega^\infty: \mathrm{HM}(k) &\rightarrow \mathrm{HI}(k) \\ M_* &\mapsto M_0,\end{aligned}$$

which just picks out the zero space. Let's try to build a map back the other way — in trying to do so, we bump into a puzzle: given  $M$ , how do we cook up an  $N$  for which  $M \cong N_{-1}$ ? With our new understanding of the monoidal structure on  $\mathrm{HI}(k)$ , this becomes easier to answer.

**Corollary 3.11.8.** For any  $M, N \in \mathrm{HI}(k)$ , if

$$f: \mathbf{K}_1^{\mathrm{MW}} \otimes_{\mathrm{HI}} M \xrightarrow{\sim} N$$

is an equivalence, then the mate

$$M \xrightarrow{\sim} \underline{\mathrm{Hom}}_{\mathrm{HI}}(\mathbf{K}_1^{\mathrm{MW}}, N) = N_{-1}$$

is an isomorphism as well.

*Proof.* It is not true in general that the mate of an isomorphism needs to be an isomorphism. It is however true in this case since  $\mathbf{K}_1^{\mathrm{MW}}$  is fully faithful. The mate is defined by

$$M \xrightarrow{\eta_M} \underline{\mathrm{Hom}}_{\mathrm{HI}}(\mathbf{K}_1^{\mathrm{MW}}, \mathbf{K}_1^{\mathrm{MW}} \otimes_{\mathrm{HI}} M) \xrightarrow{\underline{\mathrm{Hom}}_{\mathrm{HI}}(\mathbf{K}_1^{\mathrm{MW}}, f)} \underline{\mathrm{Hom}}_{\mathrm{HI}}(\mathbf{K}_1^{\mathrm{MW}}, N) = N_{-1}.$$

The latter map is an isomorphism by hypothesis, and the first is because a left adjoint is fully faithful if and only if the unit of the adjunction is a natural isomorphism.  $\square$

**Corollary 3.11.9.** If  $M \in \mathrm{HI}(k)$ , then  $M$  is a contraction of  $M \otimes_{\mathrm{HI}} \mathbf{K}_1^{\mathrm{MW}}$ .

**Proposition 3.11.10.** There is a functor

$$\sigma^\infty: \mathrm{HI}(k) \rightarrow \mathrm{HM}(k),$$

defined by the property that

$$(\sigma^\infty M)_n = \begin{cases} M \otimes_{\mathrm{HI}} \mathbf{K}_n^{\mathrm{MW}} & n \geq 0 \\ M_{-j} & n = -j < 0 \end{cases}$$

*Proof.* The gluing maps

$$(\sigma^\infty M)_n \rightarrow ((\sigma^\infty M)_n)_{-1}$$

are defined to be mates to the natural equivalences

$$\mathbf{K}_1^{\mathrm{MW}} \otimes \mathbf{K}_n^{\mathrm{MW}} \otimes M \xrightarrow{\sim} \mathbf{K}_{n+1}^{\mathrm{MW}} \otimes M.$$

$\square$

### 3.11.3 Monoidal structure on homotopy modules

**Definition 3.11.11.** An object  $X \in \mathcal{C}$  in a symmetric monoidal 1-category (resp.  $\infty$ -category) is called *cyclic* if some permutation of  $X^{\otimes n}$  is equal to (resp. homotopic to) the identity.

We'll phrase this in two settings. The  $\infty$ -categorical one will come in use later.

**Theorem 3.11.12.**

1. If  $\mathcal{C}$  is a closed symmetric monoidal 1-category and  $X \in \mathcal{C}$  is cyclic, then the localization  $\mathcal{C}[X^{-1}]$  exists, defined as the colimit

$$\mathcal{C} \xrightarrow{X \otimes -} \mathcal{C} \xrightarrow{X \otimes -} \dots$$

It is symmetric monoidal, and the localization  $\mathcal{C} \rightarrow \mathcal{C}[X^{-1}]$  is strong monoidal.

2. If  $\mathcal{C}$  is a presentably symmetric monoidal category, then if  $X$  is cyclic, the localization  $\mathcal{C}[X^{-1}]$  is still presentably symmetric monoidal, with monoidal localization functor [Rob15, §2].

**Proposition 3.11.13.** We have that  $\mathbf{K}_1^{\text{MW}} \in \text{HI}(k)$  is cyclic.

*Proof.* We claim the permutation  $(1\ 2\ 3)$  going from

$$\mathbf{K}_3^{\text{MW}} \cong (\mathbf{K}_1^{\text{MW}})^{\otimes 3} \rightarrow (\mathbf{K}_1^{\text{MW}})^{\otimes 3} \cong \mathbf{K}_3^{\text{MW}}$$

is equal to the identity, proving  $\mathbf{K}_1^{\text{MW}}$  is cyclic. Since  $\mathbf{K}_n^{\text{MW}}$  is generated by symbols  $[u_1, \dots, u_n]$  for  $n \geq 1$ , we are reduced to checking that  $[c, a, b] = [a, b, c]$  for any  $a, b, c \in F^\times$ . We use that  $\epsilon$  commutes with  $[u]$  for any  $u$  by Proposition 3.7.5(4), and that  $\epsilon^2 = 1$  by Proposition 3.7.4(4). This is now an immediate check by what we've done:

$$[c, a, b] := [c][a][b] = \epsilon[a][c][b] = \epsilon[a]\epsilon[b][c] = \epsilon^2[a][b][c] = [a, b, c].$$

□

Combining Proposition 3.11.13 and Theorem 3.11.12, we immediately obtain the following.

**Corollary 3.11.14.** The category of homotopy modules  $\text{HM}(k)$  is symmetric monoidal, and

$$\sigma^\infty: \text{HI}(k) \rightarrow \text{HM}(k)$$

is strong symmetric monoidal.

**Proposition 3.11.15.** We have that  $\sigma^\infty$  is fully faithful and  $\omega^\infty$  is exact.

*Proof.* It is clear  $\sigma^\infty$  is fully faithful by construction, since the data of a map  $\sigma^\infty M \rightarrow \sigma^\infty N$  is the data of a map  $M \rightarrow N$  in  $\text{HI}(k)$ . Meanwhile, exactness in  $\text{HM}(k)$  means exactness at each level, which implies exactness at level zero, hence  $\omega^\infty$  is exact. □

Let's also remark something interesting — if  $M_*$  is a homotopy module, then the mates to the structure isomorphisms are all of the form

$$\mathbf{K}_1^{\text{MW}} \otimes_{\text{HI}} M_* \rightarrow M_{*+1},$$

and by iterating this we get

$$\mathbf{K}_n^{\text{MW}} \otimes_{\text{HI}} M_* \rightarrow M_{*+n}.$$

**Remark 3.11.16.** Every homotopy module is a graded  $\mathbf{K}_*^{\text{MW}}$ -module.

## 3.12 Gersten complexes

If  $F$  is a homotopy sheaf and  $X \in \text{Sm}_k$ , we want to *resolve*  $F(X) = H^0(X, F)$  by its values on codimension one, two, three, points. By this we mean its values on the function fields of those points.

The goal is to write down a complex that looks like

$$\bigoplus_{x \in X^{(0)}} F(\kappa(x)) \rightarrow \bigoplus_{x \in X^{(1)}} F_{-1}(\kappa(x)) \rightarrow \bigoplus_{x \in X^{(2)}} F_{-2}(\kappa(x)) \rightarrow \cdots$$

The primary difficulty is defining the differentials and showing that they actually yield a complex. This is one of the main complicating factors in the literature. Following the slogan that homotopy sheaves are easier once they are contractions, we might approach this when a homotopy sheaf is a 1-fold or 2-fold contraction and leverage the nice properties it inherits to try to construct this

complex. This is the focus of a large chunk of Morel's book – building these complexes for  $M_{-1}$  and for  $M_{-2}$  when  $M$  is a homotopy sheaf.

For the purposes of this notes, we will work with a smaller class of homotopy sheaves, namely those which are infinite contractions — i.e. homotopy modules!

**Theorem 3.12.1.** If  $M_*$  is a homotopy module, then for any  $n$ , we can define well-defined differentials so that

$$C^*(X, M_n) := \left( \bigoplus_{x \in X^{(0)}} M_n(\kappa(x)) \rightarrow \bigoplus_{x \in X^{(1)}} M_{n-1}(\kappa(x)) \rightarrow \cdots \right)$$

is a complex, natural in  $X$ , called the *Rost-Schmid complex* [Mor12, 5.31]. This is an acyclic resolution, so the  $j$ th cohomology group of this complex is *precisely* the sheaf cohomology

$$H^j C^*(X, M_n) = H_{\text{Nis}}^j(X, M_n).$$

**Remark 3.12.2.**

1. We are being vague about differentials and twists — these need to be carefully considered although for the sake of time we are omitting them.
2. A Rost-Schmid complex exists for a strongly invariant sheaf of abelian groups, and its construction is a key input in proving the strongly=strictly theorem. See [Bac24, §6] for details.

**Definition 3.12.3.** The full subcategory

$$\text{Mod}_{\mathbf{K}_*^{\text{M}}} \subseteq \text{HM}(k)$$

of modules over Milnor  $K$ -theory (or modules over  $\mathbf{K}_*^{\text{MW}}$  on which  $\eta$  acts trivially) is called the category of *homotopy modules with (Voevodsky) transfers*. There is an equivalence of categories between these and so-called *Rost cycle modules*.

**Remark 3.12.4.**

1. It is easier to define differentials for homotopy modules with transfers since we don't have to stress about twists.
2. Rost cycle modules were used famously in the proof of the Milnor conjecture. Moreover over a perfect field, the category of Rost cycle modules is equivalent to the heart of the category of Voevodsky motives in the homotopy  $t$ -structure (see Deglise modules de cycles et motifs mixtes)

**Theorem 3.12.5.** (*Properties of the Rost-Schmid complex*)

1. For  $X$  smooth, the Rost-Schmid complex provides an acyclic resolution in the Zariski or Nisnevich sites
2. The projection  $X \times \mathbb{A}^1 \rightarrow X$  induces a quasi-isomorphism

$$C^*(X, M_n) \rightarrow C^*(X \times \mathbb{A}^1, M_n).$$

This is [Mor12, 5.38], and it is a bit out of order here – one proves this first for a strongly invariant sheaf of abelian groups, and leverages this to conclude that strongly implies strictly.

**Theorem 3.12.6.** (Rost ?) For any smooth  $X$ , we have that  $H^n(X, \mathbf{K}_n^{\text{M}}) = \text{CH}^n(X)$ .

*Proof.* At the tail end of the Rost-Schmid complex, we have

$$\cdots \rightarrow \bigoplus_{x \in X^{(n-1)}} K_1^{\text{M}}(\kappa(x)) \rightarrow \bigoplus_{x \in X^{(n)}} K_0^{\text{M}}(\kappa(x)) \rightarrow 0,$$

since a contraction of  $K_0^M$  is trivial. Note that  $K_1^M(\kappa(x)) = \kappa(x)^\times$  and  $K_0^M(\kappa(x)) = \mathbb{Z}$ . This differential can precisely be identified with the divisor function in the definition of Chow groups. The result follows immediately.  $\square$

**Corollary 3.12.7.** We have that

$$H^n(X, \mathbf{K}_n^M) = H^n(X, \mathbf{K}_n^Q) = \mathrm{CH}^n(X).$$

This is the so-called *Gersten conjecture*.

**Definition 3.12.8.** The sheaf cohomology groups

$$H^n(X, \mathbf{K}_n^{\mathrm{MW}}) = \widetilde{\mathrm{CH}}^n(X)$$

are called *Chow–Witt groups*. Note that negative Milnor–Witt  $K$ -theory sheaves are not identically zero like Milnor  $K$ -theory, so the Rost–Schmid complex doesn’t truncate. Nevertheless, the differential going from  $\mathbf{K}_1^{\mathrm{MW}}$  to  $\mathbf{K}_0^{\mathrm{MW}}$  is still an analogue of a divisor of a rational function, however equipped with some orientation data.

**Remark 3.12.9.** The multiplicative structure on cycle modules induce a multiplicative structure on the Rost–Schmid complexes, and therefore a multiplicative structure on the total cohomology.

### 3.12.1 Vanishing of $I^j$ -cohomology

**Proposition 3.12.10.** If  $\mathbf{I}^n \neq 0$  for some  $n$ , then  $\mathbf{I}^n \neq \mathbf{I}^{n+1}$ .

*Proof.* It suffices to check on fields, from which it was originally proven by Arason and Pfister [AP71, Korollar 2].  $\square$

**Corollary 3.12.11.** If  $\mathbf{I}^n/\mathbf{I}^{n+1} = 0$  then  $\mathbf{I}^n = 0$ .

**Theorem 3.12.12.** Let  $X$  be a smooth scheme of dimension  $d$  over an algebraically closed field  $k$ . Then  $\mathbf{I}^n$  is identically zero over  $X$  for  $n > d$ .

*Proof.* It suffices to check on the residue field of  $X$ , and suffices to check that  $\mathbf{I}^n/\mathbf{I}^{n+1} = 0$  by **Corollary 3.12.11**. By the Milnor conjecture, we have an isomorphism

$$\mathbf{I}^n/\mathbf{I}^{n+1} \cong \mathbf{K}_n^M/2,$$

so we have that  $I^n(k(X))/I^{n+1}(k(X)) = H_{\mathrm{Gal}}^n(k(X), \mu_2^{\otimes n})$ , which vanishes above the dimension of  $X$ .  $\square$

**Remark 3.12.13.** (*Vanishing of  $I^j$ -cohomology*)

1. Over an arbitrary field  $k$ , the same vanishing result will hold for  $n \geq d + r + 1$ , where  $r = \mathrm{cd}_2(k)$  is the 2-cohomological dimension of the field  $k$  [AF14a, 5.1].
2. An even strongly vanishing statement is proven in [AF14a, 5.2], using a Bloch–Ogus spectral sequence argument.

### 3.12.2 Contraction yoga

**Proposition 3.12.14.** If  $M \rightarrow N$  is a map of homotopy sheaves, which is an isomorphism after  $k$ -fold contraction, then

$$H^n(X, M) \cong H^n(X, N)$$

for  $n > k$ .

*Proof.* The map induces a map of Rost complexes and see that they are identical starting at the  $k$ th slot.  $\square$

We can look at some maps of sheaves we currently have and see how far we have to contract them to get an isomorphism. Recall we have a short exact sequence

$$0 \rightarrow \mathbf{I}^{j+1} \rightarrow \mathbf{K}_j^{\text{MW}} \rightarrow \mathbf{K}_j^{\text{M}} \rightarrow 0.$$

Since  $(\mathbf{K}_j^{\text{M}})_{-k} = 0$  if  $k > j$ , we have that  $\mathbf{I}^{j+1} \rightarrow \mathbf{K}_j^{\text{MW}}$  is an isomorphism after  $j+1$  contractions. This gives us

**Proposition 3.12.15.** We have that

$$H^n(X, \mathbf{I}^{j+1}) \cong H^n(X, \mathbf{K}_j^{\text{MW}})$$

for  $n > j+1$ .

**Proposition 3.12.16.** We have that

$$H^n(X, \mathbf{K}_j^{\text{M}}) = 0$$

for  $j < n$ .

What happens to  $\mathbf{I}^j$  and  $\mathbf{K}_j^{\text{MW}}$  when they are contracted below zero?

**Proposition 3.12.17.** We have that

$$\mathbf{K}_{-n}^{\text{MW}} \cong \mathbf{W}$$

is the Witt for each  $n \geq 1$ .

*Proof.* We first make the claim that  $\mathbf{K}_{-1}^{\text{MW}}$  is generated by elements of the form  $\eta \langle a \rangle$ . In other words, the map

$$\begin{aligned} \mathbf{K}_0^{\text{MW}} &\rightarrow \mathbf{K}_{-1}^{\text{MW}} \\ \langle a \rangle &\mapsto \eta \langle a \rangle \end{aligned}$$

is onto. This is analogous to the argument we did before (now we remove an  $\eta^{-1}$ , and get everything else in  $\text{GW}(k)$ ).

We first show the claim for  $n = 1$ . We claim that the map factors

$$\begin{array}{ccc} \mathbf{K}_0^{\text{MW}} & \xrightarrow{\cong} & \mathbf{GW} \\ \eta \downarrow & & \downarrow \text{mod } h \\ \mathbf{K}_{-1}^{\text{MW}} & \dashrightarrow & \mathbf{W}, \end{array}$$

where the bottom map sends  $\eta \langle a \rangle$  to  $\langle a \rangle$ . We have to check this is well-defined, i.e. that if we have  $\alpha, \beta \in \text{GW}(k)$ , and  $\eta\alpha = \eta\beta$  is it true that  $\alpha \equiv \beta \pmod{h}$ . This is clear since  $\eta$  will kill any hyperbolic elements. From this it is also clear it is bijective. To see it for  $n > 1$ , we observe that multiplication by  $\eta$  induces an isomorphism  $\mathbf{K}_{-n}^{\text{MW}} \xrightarrow{\sim} \mathbf{K}_{-n-1}^{\text{MW}}$ .  $\square$

**Corollary 3.12.18.** We have that  $\mathbf{I}_{-1} = \mathbf{W}$ .

**Corollary 3.12.19.** For  $j < n$  we have that

$$H^n(X, \mathbf{I}^{j+1}) \cong H^n(X, \mathbf{K}_j^{\text{MW}}).$$

### 3.13 The fundamental group of $\mathbb{P}^1$

Our goal is to compute  $\pi_1^{\mathbb{A}^1}(\mathbb{P}^1)$ . Recall we have

$$\text{HI}(k) \subseteq \text{Shv}_{\text{Nis}}(\text{Sm}_k; \text{Set}_*).$$

It makes sense for any  $F$  to ask for some *universal* approximation to it, lying in the category  $\mathbf{HI}(k)$ . Let  $G$  be a strongly invariant sheaf of groups, and  $S$  a sheaf of pointed sets. Then there is a bijection

$$[\Sigma S, BG]_{\mathrm{Spc}(k)_*} \cong [S, G]_{\mathrm{Shv}(\mathrm{Sm}_k; \mathrm{Set}_*)},$$

given by taking  $\pi_1^{\mathbb{A}^1}$ . In particular this gives the following definition.

**Definition 3.13.1.** [Mor12, 7.23] If  $S$  is a Nisnevich sheaf of pointed sets, we denote by

$$F_{\mathbb{A}^1}(S) := \pi_1^{\mathbb{A}^1}(\Sigma S)$$

the *free strongly invariant sheaf* of groups on  $S$ . This has the universal property that for any strongly invariant sheaf of groups  $G$ , we have

$$\begin{array}{ccc} S & \xrightarrow{\quad} & G \\ \downarrow & \nearrow \text{dashed} & \\ F_{\mathbb{A}^1}(S) & & \end{array}$$

**Definition 3.13.2.** If  $S$  is a sheaf of pointed sets, we can alternatively consider the free strongly invariant sheaf of *abelian groups* on  $S$ . We denote this by<sup>3</sup>

$$F_{\mathbb{A}^1}^{\mathrm{ab}}(S),$$

and it is the universal element in  $\mathbf{HI}(k)$  receiving a morphism of sheaves of pointed sets from  $S$ . We don't have an obvious way to compute this yet, unless  $F_{\mathbb{A}^1}(S)$  already happens to be abelian.

**Proposition 3.13.3.** For any  $n \geq 2$ , we have that

$$F_{\mathbb{A}^1}^{\mathrm{ab}}(S) \cong \pi_n(\Sigma^n S),$$

compatibly with suspension homomorphisms.

*Proof.* Let  $n \geq 2$ , and let  $G$  be any strictly invariant sheaf of abelian groups. Then by adjunction, we have a string of equivalences

$$[S, G]_* \cong [S, \Omega^n K(G, n)] \cong [\Sigma^n S, K(G, n)].$$

Since  $K(G, n)$  is  $n$ -connected, any map factors through its  $n$ -truncation  $\tau_{\leq n} \Sigma^n S$ , which is equal to  $K(\pi_n(\Sigma^n S), n)$  since  $\Sigma^n S$  is  $n$ -connected. Therefore by taking  $\pi_n$  we get a bijection

$$[\Sigma^n S, K(G, n)] \cong [K(\pi_n \Sigma^n S, n), K(G, n)] = \mathrm{Hom}_{\mathrm{Shv}(\mathrm{Grp})}(\pi_n(\Sigma^n S), G).$$

□

The independence on  $n$  can be thought of as a preliminary version of the  $S^1$ -Freudenthal suspension theorem.

**Remark 3.13.4.** There is a natural map

$$F_{\mathbb{A}^1}(S) \rightarrow F_{\mathbb{A}^1}^{\mathrm{ab}}(S),$$

which is not necessarily an isomorphism.

**Example 3.13.5.** We can consider  $\mathbb{G}_m$  as a sheaf of pointed sets. Then by definition, we have that

$$F_{\mathbb{A}^1}(\mathbb{G}_m) = \pi_1^{\mathbb{A}^1}(\mathbb{P}^1).$$

This is the sheaf of groups we want to compute.

**Notation 3.13.6.** For  $n \geq 1$ , we denote by

$$F_{\mathbb{A}^1}(n) := F_{\mathbb{A}^1}(\mathbb{G}_m^{\wedge n})$$

the free strongly invariant sheaf of groups on the smash product  $\mathbb{G}_m^{\wedge n}$ .

---

<sup>3</sup>The notation here is nonstandard.



We have that  $\mathbb{G}_m$ , as a motivic space, represents units of global sections. Since smash products are computed sectionwise, we have that  $\mathbb{G}_m^{\wedge n}$  represents the  $n$ -fold smash product of  $\mathcal{O}_X(X)^\times$  for any  $X$ . This is a sheaf of pointed sets, with no higher homotopy. Hence it gives rise to a map

$$\begin{aligned} (\mathcal{O}_X(X)^\times)^{\wedge n} &\rightarrow (k(X)^\times)^{\wedge n} \rightarrow K_n^{\text{MW}}(k(X)) \\ (a_1, \dots, a_n) &\mapsto [a_1, \dots, a_n]. \end{aligned}$$

By the yoga of  $\mathcal{F}_k$ -data, this bootstraps up to a *symbol map*, which we denote by

$$\sigma_n: \mathbb{G}_m^{\wedge n} \rightarrow \mathbf{K}_n^{\text{MW}}.$$

This factors through the universal strongly invariant presheaf of *abelian* groups:

$$\begin{array}{ccc} \mathbb{G}_m^{\wedge n} & \xrightarrow{\sigma_n} & \mathbf{K}_n^{\text{MW}} \\ \downarrow & \nearrow & \\ F_{\mathbb{A}^1}^{\text{ab}}(n) & & \end{array}$$

**Theorem 3.13.7.** [Mor12, 3.37] For  $n \geq 1$ , the induced symbol map

$$F_{\mathbb{A}^1}^{\text{ab}}(n) \rightarrow \mathbf{K}_n^{\text{MW}}$$

is an equivalence. In other words,  $\mathbf{K}_n^{\text{MW}}$  is the free strongly invariant sheaf of *abelian* groups on  $\mathbb{G}_m^{\wedge n}$ .

*Proof intuition.* Let  $M \in \text{HI}(k)$ . Then any map of pointed sheaves of sets  $\mathbb{G}_m^{\wedge n} \rightarrow M$  can be viewed as a map of pointed motivic spaces, and we see by adjunction:

$$\begin{aligned} \text{Hom}_{\text{Spc}(k)_*}(S^{n,n}, M) &= \text{Hom}_{\text{Spc}(k)_*}(S^0, \Omega^{n,n}M) \\ &= \text{Hom}_{\text{Spc}(k)_*}(S^0, M_{-n}) \\ &= M_{-n}(k). \end{aligned}$$

Since  $M_{-n} = \underline{\text{Hom}}_{\text{HI}}(\mathbf{K}_n^{\text{MW}}, M)$ , we might expect that this gives rise to a unique map  $\mathbf{K}_n^{\text{MW}} \rightarrow M$ . This ends up being true, and the difficulty is just in verifying that this is indeed a morphism of homotopy sheaves. In particular one must construct it from the symbol map and verify it is well-defined with respect to the relations for Milnor–Witt  $K$ -theory.  $\square$

**Proposition 3.13.8.** We have that

$$F_{\mathbb{A}^1}(2) \cong F_{\mathbb{A}^1}^{\text{ab}}(2) \cong \mathbf{K}_2^{\text{MW}}.$$

*Proof.* Recall by Proposition 3.3.11 we have

$$\text{SL}_2 \simeq \mathbb{A}^2 \setminus 0 \simeq S^{3,2} \simeq \Sigma(\mathbb{G}_m \wedge \mathbb{G}_m).$$

Since  $\text{SL}_2$  is a group object, its  $\pi_1$  is abelian by an Eckmann–Hilton argument, hence the result follows.  $\square$

### 3.13.1 The fundamental group of $\mathbb{P}^1$

Since  $\mathbb{A}^2 \setminus 0 \rightarrow \mathbb{P}^1$  is the total space of a  $\mathbb{G}_m$ -torsor, we get an  $\mathbb{A}^1$ -fiber sequence

$$\mathbb{G}_m \rightarrow \mathbb{A}^2 \setminus 0 \rightarrow \mathbb{P}^1,$$

which deloops to

$$\mathbb{A}^2 \setminus 0 \rightarrow \mathbb{P}^1 \rightarrow B\mathbb{G}_m.$$

Since both  $B\mathbb{G}_m$  and  $\mathbb{A}^2 \setminus 0$  are connected, the long exact sequence on homotopy becomes a short exact sequence of the form [Mor12, (7.4)]

$$0 \rightarrow \mathbf{K}_2^{\text{MW}} \rightarrow F_{\mathbb{A}^1}(1) \rightarrow \mathbb{G}_m \rightarrow 0.$$

This turns out to be a central extension, but  $F_{\mathbb{A}^1}(1)$  is not commutative [Mor12, 7.29].

### 3.14 The Brouwer degree

We are interesting in leveraging our computation of  $\pi_1^{\mathbb{A}^1}(\mathbb{P}^1)$  to say something about  $\text{End}_{\text{Spc}(k)_*}(\mathbb{P}^1)$ . By convention,  $\mathbb{P}^1$  is pointed at  $\infty$ . Given any pointed endomorphism of motivic spaces  $\mathbb{P}^1 \rightarrow \mathbb{P}^1$ , we can look at the induced map on  $\pi_1^{\mathbb{A}^1}$ , and we get an assignment

$$[\mathbb{P}^1, \mathbb{P}^1]_{\text{Spc}(k)_*} \rightarrow \text{End}(F_{\mathbb{A}^1}(1)). \quad (3.1)$$

We are going to argue that this is a bijection.

**Proposition 3.14.2.** There is a bijection

$$[\mathbb{P}^1, \mathbb{P}^1]_{\mathbb{A}^1} \cong [\mathbb{P}^1, B\pi_1^{\mathbb{A}^1}(\mathbb{P}^1)]_*.$$

*Proof.* We have that

$$[\mathbb{P}^1, \mathbb{P}^1]_* = [S^1, \Omega^{1,1}\mathbb{P}^1]_* = \pi_1(\Omega^{1,1}\mathbb{P}^1) = [S^1, B\pi_1(\Omega^{1,1}\mathbb{P}^1)]_*.$$

Since  $\pi_1(\mathbb{P}^1)_{-1}$  is strongly invariant, the above is isomorphic to

$$[S^1, \Omega^{1,1}B\pi_1(\mathbb{P}^1)]_* \cong [\mathbb{P}^1, B\pi_1(\mathbb{P}^1)]_*.$$

□

**Corollary 3.14.3.** The assignment Equation 3.1 is a bijection.

*Proof.* Since  $BF_{\mathbb{A}^1}(1)$  is 1-truncated, we have that any map  $\mathbb{P}^1 \rightarrow BF_{\mathbb{A}^1}(1)$  factors through  $L_{\text{mot}}\tau_{\leq 1}\mathbb{P}^1$ . Since  $\mathbb{P}^1$  is connected,<sup>4</sup> we have that  $\tau_{\leq 1}\mathbb{P}^1$  is equal to  $BF_{\mathbb{A}^1}(1)$ , which is  $\mathbb{A}^1$ -local. □

**Proposition 3.14.4.** The contraction of  $\mathbb{G}_m$  is  $\mathbb{Z}$ .

*Proof.* We first check that

$$\text{Hom}_{\text{Smk}}(\mathbb{G}_m \times U, \mathbb{G}_m) = \text{Hom}_{\text{Smk}}(U[t, t^{-1}], \mathbb{G}_m).$$

If  $U$  is reduced and irreducible, the units in  $U[t, t^{-1}]$  are all of the form  $ut^n$  for some  $u \in U^\times$  and  $n \in \mathbb{Z}$ . Hence we get an isomorphism

$$\mathbb{G}_m(\mathbb{G}_m \times U) \cong \mathbb{G}_m(U) \times \mathbb{Z}.$$

In the short exact sequence defining contraction, this gives us that  $(\mathbb{G}_m)_{-1}$  is the constant presheaf  $\mathbb{Z}$ . □

**Proposition 3.14.5.** The short exact sequence, obtained from the contraction above

$$0 \rightarrow \mathbf{K}_1^{\text{MW}} \rightarrow F_{\mathbb{A}^1}(1)_{-1} \rightarrow \mathbb{Z} \rightarrow 0$$

is canonically split exact, giving an isomorphism

$$F_{\mathbb{A}^1}(1)_{-1} \cong \mathbf{K}_1^{\text{MW}} \oplus \mathbb{Z}.$$

*Proof.* It suffices to see that  $\text{Ext}(\mathbb{Z}, \mathbf{K}_1^{\text{MW}}) = 0$ , which is clear since  $\mathbb{Z}$  is projective. □

**Proposition 3.14.6.** We have that  $\text{End}(F_{\mathbb{A}^1}(1))$  is a group, and moreover is an associative commutative ring.

**Proposition 3.14.7.** There is a canonical isomorphism

$$\text{End}(F_{\mathbb{A}^1}(1)) \cong F_{\mathbb{A}^1}(1)_{-1}(k).$$

---

4

*Proof.* Since  $F_{\mathbb{A}^1}(1)$  is by definition the free pointed sheaf of abelian groups on  $\mathbb{G}_m$ , there is a canonical isomorphism

$$\mathrm{Hom}_{\mathrm{Shv}(\mathrm{Grp})}(F_{\mathbb{A}^1}(1), F_{\mathbb{A}^1}(1)) = \mathrm{Hom}_{\mathrm{Shv}_*}(\mathbb{G}_m, F_{\mathbb{A}^1}(1)).$$

This is the contraction  $F_{\mathbb{A}^1}(1)_{-1}$ .  $\square$

Since  $F_{\mathbb{A}^1}(1)_{-1}(k) = \mathrm{End}(F_{\mathbb{A}^1}(1))$  is the group we want to compute, we can take global sections  $H_{\mathrm{Nis}}^0(\mathrm{Spec}(k), -)$  in the above SES to get

$$[\mathbb{P}^1, \mathbb{P}^1]_* = \mathrm{End}(F_{\mathbb{A}^1}(1)) \cong K_1^{\mathrm{MW}}(k) \oplus \mathbb{Z}. \quad (3.8)$$

Any endomorphism of  $\mathrm{End}(F_{\mathbb{A}^1}(1))$  induces an endomorphism of the contraction  $F_{\mathbb{A}^1}(1)_{-1}$  by functoriality, which restricts to an endomorphism of  $\mathbf{K}_1^{\mathrm{MW}}$ . All endomorphisms of  $\mathbf{K}_1^{\mathrm{MW}}$  are given by multiplication by some element in  $K^{\mathrm{MW}}_0 = \mathrm{GW}(k)$ . This gives a natural homomorphism called the  $\mathbb{A}^1$ -Brouwer degree:

$$[\mathbb{P}^1, \mathbb{P}^1]_* = \mathrm{End}(F_{\mathbb{A}^1}(1)) \xrightarrow{\deg^{\mathbb{A}^1}} \mathrm{End}(\mathbf{K}_1^{\mathrm{MW}}) = \mathrm{GW}(k).$$

Some facts are true about this:

**Proposition 3.14.9.**

1. The  $\mathbb{A}^1$ -Brouwer degree is an epimorphism
2. The following diagram commutes

$$\begin{array}{ccccccc} & & & & \mathrm{GW}(k) & & \\ & & & \nearrow \deg^{\mathbb{A}^1} & \downarrow \mathrm{rank} & & \\ 0 & \longrightarrow & K_1^{\mathrm{MW}}(k) & \longrightarrow & [\mathbb{P}^1, \mathbb{P}^1]_* & \longrightarrow & \mathbb{Z} \longrightarrow 0. \end{array}$$

*Proof.* todo  $\square$

Hence we think about  $\mathbb{Z}$  as recording *rank*.

**Question 3.14.10.** What is the kernel of the degree homomorphism

$$\deg^{\mathbb{A}^1}: [\mathbb{P}^1, \mathbb{P}^1]_* \rightarrow \mathrm{GW}(k).$$

To answer this, we can consider the following commutative diagram, which is just rewriting the one before:

$$\begin{array}{ccccccc} 0 & \longrightarrow & ? & \longrightarrow & [\mathbb{P}^1, \mathbb{P}^1]_* & \longrightarrow & \mathrm{GW}(k) \longrightarrow 0 \\ & & \downarrow & & \parallel & & \downarrow \\ 0 & \longrightarrow & K_1^{\mathrm{MW}} & \longrightarrow & [\mathbb{P}^1, \mathbb{P}^1]_* & \longrightarrow & \mathbb{Z} \longrightarrow 0. \end{array}$$

By the snake lemma, the cokernel of the map  $? \rightarrow K_1^{\mathrm{MW}}(k)$  is the kernel of  $\mathrm{GW}(k) \rightarrow \mathbb{Z}$ , which is  $I(k)$ .

**Proposition 3.14.11.** The connecting homomorphism  $K_1^{\mathrm{MW}}(k) \rightarrow I(k)$  is given by the right exact sequence

$$K_1^{\mathrm{MW}}(k) \xrightarrow{h} K_1^{\mathrm{MW}}(k) \rightarrow I(k) \rightarrow 0,$$

where  $[u] \mapsto \langle u \rangle - \langle 1 \rangle$ .

*Proof.* todo  $\square$

**Proposition 3.14.12.** The kernel of  $K_1^{\text{MW}}(k) \rightarrow I(k)$  is  $(k^\times)^2$ .

*Proof.* Recall the pullback diagram from [Corollary 3.8.14](#):

$$\begin{array}{ccc} K_1^{\text{MW}} & \longrightarrow & K_1^M(k) \\ \downarrow & \lrcorner & \downarrow \\ I(k) & \longrightarrow & K_1^M(k)/2. \end{array}$$

The fiber of the left vertical map is isomorphic to the fiber of the rightmost vertical map, which is  $2K_1^M(k)$ . Since addition on  $K_1^M(k) = k^\times$  is multiplication of field units, the result follows.  $\square$

Altogether we get the following result.

**Theorem 3.14.13.** [[Mor12](#), 7.36] We have a short exact sequence of abelian groups

$$0 \rightarrow (k^\times)^2 \rightarrow [\mathbb{P}^1, \mathbb{P}^1]_* \xrightarrow{\deg^{\mathbb{A}^1}} \text{GW}(k) \rightarrow 0. \quad (3.14)$$

**Remark 3.14.15.** (On [Theorem 3.14.13](#))

1. Morel writes  $(k^\times)/\pm 1$  instead of  $(k^\times)^2$  in his book, which I believe is an error?
2. The short exact sequence above can alternatively be repackaged as a pullback diagram (the sequence is obtained by taking horizontal fibers):

$$\begin{array}{ccc} [\mathbb{P}^1, \mathbb{P}^1]_* & \longrightarrow & \text{GW}(k) \\ \downarrow & \lrcorner & \downarrow \\ k^\times & \longrightarrow & k^\times / (k^\times)^2. \end{array}$$

This is the form that [Theorem 3.14.13](#) takes in various forms in the literature, where people write

$$[\mathbb{P}^1, \mathbb{P}^1]_* \cong \text{GW}(k) \times_{k^\times / (k^\times)^2} k^\times.$$

3. The ring structure on  $\text{GW}(k) \times_{k^\times / (k^\times)^2} k^\times$  is spelled out explicitly in [[Caz09](#)].
4. The units will go away after stabilizing, and we will see that  $\text{GW}(k)$  is the home for the *stable*  $\mathbb{A}^1$ -Brouwer degree, in other words it is the  $(0,0)$ th stable stem motivically.

The additive structure comes from the isomorphism  $\mathbb{P}^1 \simeq S^1 \wedge \mathbb{G}_m$ . We have a pinch map of simplicial sets

$$S^1 \rightarrow S^1 \vee S^1.$$

Smashing with  $\mathbb{G}_m$ , we use that smash products distribute over wedge products to get a comultiplication

$$\mathbb{P}^1 \rightarrow \mathbb{P}^1 \vee \mathbb{P}^1.$$

This cogroup structure induces an addition on  $[\mathbb{P}^1, \mathbb{P}^1]_{\mathbb{A}^1}$  which corresponds to addition in  $\text{GW}(k)$ .

### 3.14.1 Explicit computations

Given an endomorphism  $f/g: \mathbb{P}^1 \rightarrow \mathbb{P}^1$ , explicit formulas exist for computing the associated symmetric bilinear form in  $\text{GW}(k)$  [[Caz12](#)]. We spell these out now.

By convention  $\mathbb{P}^1$  is pointed at  $\infty$  as a motivic space, however we could pick any basepoint and get analogous formulas.

**Note 3.14.16.** We say a morphism of schemes  $f/g: \mathbb{P}^1 \rightarrow \mathbb{P}^1$  is *pointed* if it sends  $\infty$  to  $\infty$ . Explicitly this implies that  $\deg(g) < \deg(f)$ . WLOG we can assume  $f$  is monic, and we can dehomogenize and write

$$\begin{aligned} f(x) &= x^n + a_{n-1}x^{n-1} + \dots + a_0 \\ g(x) &= b_{n-1}x^{n-1} + b_{n-2}x^{n-2} + \dots + b_0. \end{aligned}$$

Then we require  $f$  and  $g$  to not be simultaneously zero in order to give a well-defined morphism. Taking affine space in the coefficients

$$\mathbb{A}_k^{2n} = \text{Spec } k[a_0, \dots, a_{n-1}, b_0, \dots, b_{n-1}],$$

the resultant is a hypersurface in here. So the data of  $f/g$  of leading degree  $n$  is the same as a rational point in the hypersurface complement.

**Definition 3.14.17.** Let  $f/g: \mathbb{P}_k^1 \rightarrow \mathbb{P}_k^1$  be a pointed morphism of degree  $\deg(f) = n$ . We introduce auxiliary variables  $X$  and  $Y$ , and consider the quantity

$$\frac{f(X)g(Y) - f(Y)g(X)}{X - Y} = \sum_{1 \leq i, j \leq n} c_{ij} X^{i-1} Y^{j-1}.$$

This is called the *Bézoutian*, and we denote by  $\text{Béz}(f/g)$  the  $n \times n$  matrix  $(c_{ij})_{i,j}$  over  $k$ .

**Theorem 3.14.18.** (Cazanave) The Bézoutian is a symmetric bilinear matrix, equal to the  $\mathbb{A}^1$ -degree of  $f/g$ , and the bijection

$$\begin{aligned} [\mathbb{P}^1, \mathbb{P}^1]_* &\rightarrow \text{GW}(k) \times_{k^\times / (k^\times)^2} k^\times \\ (f/g) &\mapsto \left( \text{Béz}(f/g), (-1)^{\frac{n(n-1)}{2}} \det \text{Béz}(f/g) \right) \end{aligned}$$

is an isomorphism of abelian groups.

**Example 3.14.19.** Consider  $f(x) = x^2$  and  $g(x) = 1$ . Then we have that

$$\frac{f(X)g(Y) - f(Y)g(X)}{X - Y} = \frac{X^2 - Y^2}{X - Y} = X + Y = \begin{pmatrix} X & Y \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} X \\ Y \end{pmatrix}$$

Hence

$$\deg^{\mathbb{A}^1}(x^2/1) = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} = h.$$

**Corollary 3.14.20.** An analogous argument shows that

$$\deg^{\mathbb{A}^1} \left( \frac{x^n}{1} \right) = \begin{cases} \frac{n}{2}h & n \text{ even} \\ \frac{n-1}{2}h + \langle 1 \rangle & n \text{ odd.} \end{cases}$$

### 3.15 Naive homotopies

We localized at  $\mathbb{A}^1$  in order to build the category of motivic spaces, but we can talk briefly about a more naive way to go about this.

**Definition 3.15.1.** Let  $X, Y \in \text{Sm}_k$ , and take two maps  $f, g: X \rightarrow Y$  between them in the category of schemes. A *naive  $\mathbb{A}^1$ -homotopy* is a morphism  $H: X \times \mathbb{A}^1 \rightarrow Y$  so that the diagram commutes

$$\begin{array}{ccc} X & & Y \\ & \searrow 0 \quad \quad \nearrow f & \\ & X \times \mathbb{A}^1 & \xrightarrow{H} \\ & \nearrow 1 \quad \quad \searrow g & \\ X & & Y \end{array}$$

**Example 3.15.2.** If  $X = \operatorname{Spec}(B)$  and  $Y = \operatorname{Spec}(A)$  are affine  $k$ -algebras, then a map

$$X \times_k \mathbb{A}_k^1 \rightarrow Y$$

is the same data as a  $k$ -algebra homomorphism  $B[t] \leftarrow A$ . A homotopy between  $f, g: A \rightarrow B$  is a map that restricts to  $f$  and  $g$  by setting  $t = 0$  and  $t = 1$ , respectively.

We define a relation  $\sim_N$  on  $\operatorname{Hom}_{\operatorname{Sm}_k}(X, Y)$ , where we identify two maps if there exists a naive homotopy between them.

**Note 3.15.3.**  $\sim_N$  is not an equivalence relation (it isn't transitive). However it admits a transitive closure, giving rise to an equivalence relation, which we also denote by  $\sim_N$  by abuse of notation. We use the notation

$$[X, Y]_N := \operatorname{Hom}_{\operatorname{Sm}_k}(X, Y) / \sim_N .$$

We call this *naive  $\mathbb{A}^1$ -homotopy classes of maps*.

**Proposition 3.15.4.** If  $X, Y \in \operatorname{Sm}_k$  we have that

$$[X, Y]_N = \pi_0(L_{\mathbb{A}^1}h_Y)(X).$$

*Proof.* We recall  $L_{\mathbb{A}^1}h_Y$  was the sifted colimit

$$L_{\mathbb{A}^1}h_Y = \operatorname{colim}_{[n] \in \Delta^{\operatorname{op}}} h_Y(\mathbb{A}^n \times -).$$

Since  $h_Y$  is a discrete sheaf, it is equivalent to its own  $\pi_0$ , so we can pass  $\pi_0$  through the colimit. Using that we are now valued in a 1-category we can use the cofinality of  $\Delta_{\leq 1}^{\operatorname{op}, \operatorname{inj}} \subseteq \Delta^{\operatorname{op}}$  [Example 2.3.7](#) to rewrite the diagram as

$$\begin{aligned} \pi_0(L_{\mathbb{A}^1}h_Y)(X) &= \operatorname{colim}(\pi_0(L_{\mathbb{A}^1}h_Y)(X \times \mathbb{A}^1) \rightrightarrows \pi_0(L_{\mathbb{A}^1}h_Y)(X)) \\ &= \operatorname{colim}(\operatorname{Hom}_{\operatorname{Sm}_k}(X \times \mathbb{A}^1, Y) \rightrightarrows \operatorname{Hom}_{\operatorname{Sm}_k}(X, Y)), \end{aligned}$$

so it is the coequalizer of evaluation at 0 and 1, which is precisely the quotient of  $\operatorname{Hom}_{\operatorname{Sm}_k}(X, Y)$  by the equivalence relation generated by naive homotopy.  $\square$

**Remark 3.15.5.** There is an analogous notion of a pointed naive homotopy which is the definition one would expect, and a pointed version of [Proposition 3.15.4](#) can be formulated. See e.g. [\[Bar+23, Appendix A\]](#).

There is a natural map

$$L_{\mathbb{A}^1}h_Y \rightarrow L_{\operatorname{mot}}h_Y,$$

which induces a map

$$[X, Y]_N = \pi_0(L_{\mathbb{A}^1}h_Y)(X) \rightarrow \pi_0(L_{\operatorname{mot}}h_Y)(X) = [X, Y]_{\mathbb{A}^1}.$$

Here the latter equality is by [Example 3.4.3](#).

We remark that if  $L_{\mathbb{A}^1}h_Y$  is already a Nisnevich sheaf, then we get that these two classes agree, however this rarely happens.

**Definition 3.15.6.** [\[MV99, §3.2.4\]](#) We say that  $Y \in \operatorname{Sm}_k$  is  $\mathbb{A}^1$ -rigid if  $h_Y$  is  $\mathbb{A}^1$ -invariant.

**Example 3.15.7.**  $\mathbb{G}_m$  is  $\mathbb{A}^1$ -rigid, as is any smooth curve of genus  $g \geq 1$  [\[MV99, §3.2.4\]](#).

**Proposition 3.15.8.** If  $Y$  is  $\mathbb{A}^1$ -rigid, then for any  $X \in \operatorname{Sm}_k$ , we get that

$$[X, Y]_N \cong [X, Y]_{\mathbb{A}^1}.$$

In general this is rare. Recently, it has been observed that it is much more common for representable presheaves (and motivic spaces) to be  $\mathbb{A}^1$ -invariant on affines.

**Definition 3.15.9.** We say that a simplicial presheaf  $F \in \text{PSh}(\text{Sm}_k)$  is  $\mathbb{A}^1$ -naive if the induced map

$$(L_{\mathbb{A}^1}F)(U) \rightarrow (L_{\text{mot}}F)(U)$$

is an equivalence for every affine  $U \in \text{Sm}_k^{\text{aff}}$ .

**Proposition 3.15.10.** If  $h_Y$  is  $\mathbb{A}^1$ -naive, then

$$[U, Y]_N \cong [U, Y]_{\mathbb{A}^1}$$

for every affine  $U \in \text{Sm}_k^{\text{aff}}$ .

**Proposition 3.15.11.** [AHW18, 2.1.3] A simplicial presheaf  $F$  is  $\mathbb{A}^1$ -naive if and only if  $L_{\mathbb{A}^1}F$  is a Nisnevich sheaf on  $\text{Sm}_k^{\text{aff}}$ .

**Example 3.15.12.** The following schemes are  $\mathbb{A}^1$ -naive over  $k$ :

1. The smooth quadric [AHW18, 4.2.1]

$$Q_{2n-1} = \left\{ \sum_{i=1}^n x_i y_i = 1 \right\} \subseteq \mathbb{A}_k^{2n}.$$

2. The projective line  $\mathbb{P}^1$ . [Bar+23, 127].
3. Any  $\mathbb{A}^1$ -rigid scheme.
4.  $\mathbb{A}^n \setminus 0$  [AHW18, 4.2.6]
5. Any isotropic reductive group scheme, assuming  $k$  is infinite [AHW18, 4.3.1]

### 3.15.1 Cazanave's theorem

In general naive and genuine  $\mathbb{A}^1$ -homotopy classes of maps are quite different, but making statements relating the two has been a topic of interest throughout the history of motivic homotopy theory.

As a particular example, we have seen that  $[\mathbb{P}^1, \mathbb{P}^1]_*$  is an abelian group.

**Theorem 3.15.13.** (Cazanave) The set  $[\mathbb{P}^1, \mathbb{P}^1]_N$  of pointed naive homotopy classes of maps admits a monoid structure, and the natural map

$$[\mathbb{P}^1, \mathbb{P}^1]_N \rightarrow [\mathbb{P}^1, \mathbb{P}^1]_{\mathbb{A}^1}$$

is a group completion.

Cazanave's monoidal structure is defined explicitly in terms of rational maps.

## 3.16 Unstable connectivity

We'll take the following theorem without proof.

**Theorem 3.16.1.** [MV99, §.3.22 (p.94)] (c.f. [AD09, 2.7]) For any  $F \in \text{Shv}_{\text{Nis}}(\text{Sm}_k)$ , the map to its singularization  $F \rightarrow \text{Sing}(F)$  induces a surjection on  $\pi_0$ .

This has an immediate corollary, which is that connectivity is preserved in passing from the sheaf topos to motivic spaces.

**Corollary 3.16.2.** (c.f. [Bac24, 1.2]) If  $F \in \text{Shv}_{\text{Nis}}(\text{Sm}_k)$  is connected, then so is  $L_{\text{mot}}F$ .

*Proof.* Homotopy sheaves commute with filtered colimits, hence connected objects are closed under filtered colimits. By cofinality (Remark 3.1.24) it suffices to argue that  $\pi_0 L_{\text{Nis}} L_{\mathbb{A}^1} F$  is trivial for every connected  $F \in \text{Shv}_{\text{Nis}}(\text{Sm}_k)$ . Since  $\pi_0(F) \rightarrow \pi_0(L_{\text{Nis}} L_{\mathbb{A}^1} F)$  is surjective by Theorem 3.16.1, the result follows.  $\square$

This is the  $n = 0$  of the *unstable connectivity theorem*.

**Theorem 3.16.3.** (Unstable connectivity theorem, Morel) If  $F$  is a simplicial sheaf which is  $n$ -connected, then  $L_{\text{mot}}F$  is  $n$ -connected.

The goal of this section is to prove this.

**Lemma 3.16.4.** [Bac24, 1.9] Let  $F \rightarrow E \rightarrow B$  be a fiber sequence in  $\text{Shv}_{\text{Nis}}(\text{Sm}_k)$ , and let  $B$  be connected and  $\mathbb{A}^1$ -local. Then  $F$  is  $\mathbb{A}^1$ -local if and only if  $E$  is  $\mathbb{A}^1$ -local. In particular if these conditions are satisfied, it is also a fiber sequence in  $\text{Spc}(k)$ .

**Proposition 3.16.5.** [Bac24, 1.8(2)] If  $X \in \text{Spc}(k)_*$  is a connected pointed motivic space, then for each  $n \geq 1$ , the truncation  $\tau_{\leq n}X \in \text{Shv}_{\text{Nis}}(\text{Sm}_k)$  is  $\mathbb{A}^1$ -local, and hence a motivic space.

*Proof.* We have fiber sequences

$$K(\pi_{n+1}X, n+1) \rightarrow X_{\leq n+1} \rightarrow X_{\leq n}.$$

Since  $X$  is a motivic space, its homotopy sheaves are strongly invariant, hence  $K(\pi_{n+1}X)$  is  $\mathbb{A}^1$ -local. So  $X_{\leq n}$  being  $\mathbb{A}^1$ -local will imply that  $X_{\leq n+1}$  is  $\mathbb{A}^1$ -local. This is the inductive step.

For the base case it suffices to check that  $X_{\leq 0}$  is  $\mathbb{A}^1$ -local. However we note that  $X_{\leq 0} = \pi_0X$ , which is trivial since  $X$  is connected, and hence  $\mathbb{A}^1$ -local.  $\square$

We can now prove the connectivity theorem, following [Bac24, 1.10].

*Proof of Theorem 3.16.3.* We already proved this when  $n = 0$  (Corollary 3.16.2), so we may assume  $n > 0$ . Consider the fiber sequence in the sheaf topos

$$0 \rightarrow (L_{\text{mot}}F)_{>n} \rightarrow L_{\text{mot}}F \rightarrow (L_{\text{mot}}F)_{\leq n} \rightarrow 0.$$

The middle space is  $\mathbb{A}^1$ -local, and  $L_{\text{mot}}F$  has  $\mathbb{A}^1$ -local truncation by Proposition 3.16.5. Now by Lemma 3.16.4 the fiber  $(L_{\text{mot}}F)_{>n}$  is  $\mathbb{A}^1$ -local as well. Consider the diagram

$$\begin{array}{ccccc} & & F & & \\ & \swarrow \text{dashed} & \downarrow & \searrow & \\ (L_{\text{mot}}F)_{>n} & \longrightarrow & L_{\text{mot}}F & \longrightarrow & (L_{\text{mot}}F)_{\leq n} \end{array}$$

By null homotopy of the composite map of the map to the bottom right we get a lift  $F \rightarrow (L_{\text{mot}}F)_{>n}$ . Since the target is local, the lift factors through  $L_{\text{mot}}F$ , giving a section  $s$ . Hence we have a retract:<sup>5</sup>

$$\begin{array}{ccc} L_{\text{mot}}F & \xrightarrow{s} & (L_{\text{mot}}F)_{>n} \\ & \searrow & \downarrow \\ & & L_{\text{mot}}F. \end{array}$$

The property of being  $n$ -connected is closed under retracts,<sup>6</sup> so the result follows.  $\square$

**Corollary 3.16.6.** [Bac24, 1.11] Let  $F \in \text{Shv}_{\text{Nis}}(\text{Sm}_k)$  be a connected sheaf. Then  $F$  is a motivic space if and only if  $\pi_i F$  is strongly  $\mathbb{A}^1$ -invariant for all  $i \geq 1$ .

<sup>5</sup>Letting  $r$  denote the map  $(L_{\text{mot}}F)_{>n} \rightarrow L_{\text{mot}}F$ , it is perhaps not obviously immediate why  $rs = \text{id}$ . It is true after precomposing with  $F \rightarrow L_{\text{mot}}F$ , then applying  $L_{\text{mot}}$  we have that  $L_{\text{mot}}(rs) = \text{id}$ . Since  $r$  and  $s$  are already maps between motivic spaces, the result follows.

<sup>6</sup>Apply  $\pi_i$  to the diagram and see that the identity on  $\pi_i L_{\text{mot}}F$  factors through zero.



*Proof.* The forward direction is clear. For the reverse, we can inductively prove that  $X_{\leq n}$  is local. Since  $\mathrm{Shv}_{\mathrm{Nis}}(\mathrm{Sm}_k)$  is Postnikov complete, we have that

$$X = \lim_n X_{\leq n},$$

and the result follows from observing limits of  $\mathbb{A}^1$ -invariant sheaves are still  $\mathbb{A}^1$ -invariant.  $\square$

**Proposition 3.16.7.** Suppose that  $X \in \mathrm{Shv}_{\mathrm{Nis}}(\mathrm{Sm}_k)$  has  $\pi_i X = 0$  for  $i < n$ . Then the map

$$\pi_n X \rightarrow \pi_n L_{\mathrm{mot}} X$$

is the initial map to a strongly invariant sheaf of groups (if  $n = 1$ ) and of abelian groups (if  $n \geq 2$ ).

*Proof.* If  $n = 1$ , let  $M$  be any strongly invariant sheaf of groups, and for  $n \geq 2$ , suppose it is a sheaf of abelian groups. Then take any map  $\pi_n X \rightarrow M$ . This is equivalent to the data of a map

$$K(\pi_n X, n) \rightarrow K(M, n).$$

By our assumptions on  $X$ , we have that  $\tau_{\leq n} X = K(\pi_n X, n)$ , and by adjunction, we get equivalences

$$[\pi_n(X), M] \cong [\tau_{\leq n} X, K(M, n)] = [X, K(M, n)].$$

By our hypotheses on  $M$ , the EM space  $K(M, n)$  is  $\mathbb{A}^1$ -local, hence any map  $X \rightarrow K(M, n)$  factors through  $L_{\mathrm{mot}} X$ . Applying  $\pi_n(-)$  to the factorization gives the universal property.  $\square$

**Remark 3.16.8.** This result can be improved, via induction on the Postnikov tower, to assume that  $\pi_0 X = 0$  and that  $\pi_i X$  is strongly invariant for  $0 < i < n$ , see [Mor12, 6.60].

### 3.17 Simplicial Freudenthal

**Recall:** A map  $f: X \rightarrow Y$  is said to be  $n$ -connective if its homotopy fiber is  $n$ -connective, i.e.  $\pi_i(\mathrm{fib}(f)) = 0$  for  $i < n$ .

▷  $n$ -connective means  $\pi_i = 0$  for  $i < n$

▷  $n$ -connected means  $\pi_i = 0$  for  $i \leq n$ .

**Theorem 3.17.1.** [Mor12, 6.61], [ABH24, 3.7] Let  $n \geq 0$  and  $X \in \mathrm{Spc}(k)_*$ , and suppose that  $X$  is  $\mathbb{A}^1$ - $n$ -connected. Then for any  $i \geq 1$  the map

$$X \rightarrow \Omega^{i,0} \Sigma^{i,0} X$$

has  $2n$ -connected fibers.<sup>7</sup>

*Proof.* Let  $F$  be the fiber, computed in the sheaf topos

$$F \rightarrow X \rightarrow \Omega \Sigma X.$$

Since  $L_{\mathrm{mot}}$  preserves this fiber sequence, and the middle and rightmost spaces were already local, then  $F$  is local as well. Since  $F$  is already  $2n$ -connected in the sheaf topos by classical connectivity results, we have by unstable connectivity (Theorem 3.16.3) that  $L_{\mathrm{mot}} F$  is as well.  $\square$

**Corollary 3.17.2.** If  $X$  is  $\mathbb{A}^1$ - $n$ -connected, then the suspension morphisms

$$\pi_i^{\mathbb{A}^1}(X) \rightarrow \pi_{i+1}^{\mathbb{A}^1}(\Sigma^{1,0} X)$$

are isomorphisms for  $i \leq 2n$  and an epimorphism for  $i = 2n + 1$ .

<sup>7</sup>Note that Morel initially assumed  $n \geq 2$ , but this hypothesis is removed in [ABH24].

### 3.18 Motivic homotopy groups of spheres

Considering  $\mathbb{G}_m^{\wedge n}$  as a sheaf of pointed sets, it is discrete, hence we have that

$$\pi_2(\Sigma^2 \mathbb{G}_m^{\wedge n}) \xrightarrow{\sim} \pi_3(\Sigma^3 \mathbb{G}_m^{\wedge n}) \xrightarrow{\sim} \dots$$

by Freudenthal. Note that we already sort of knew this (modulo the black boxing we've been doing), since we identified the free strongly invariant sheaf of abelian groups on a sheaf of pointed sets  $S$  with  $\pi_n(\Sigma^n S)$  for any  $n \geq 2$ . Identifying this with Milnor–Witt  $K$ -theory allows us to conclude that

$$\pi_{n,n} \mathbb{S} \cong \mathbf{K}_n^{\text{MW}}$$

for  $n \leq 0$ . We'll offer a slightly different perspective.

**Intuition 3.18.1.** In classical homotopy theory, the Freudenthal suspension theorem, together with the Hurewicz theorem, identifies the first nonvanishing homotopy group of  $S^n$  with the *free discrete abelian group* on  $S^0$ :

$$\pi_n(S^n) \cong H_n(S^n) \cong H_{n-1}(S^{n-1}) \cong \dots \cong H_0(S^0) = \mathbb{Z}[S^0].$$

In motivic homotopy theory, the first nonvanishing homotopy sheaf of  $\Sigma^{n-1} \mathbb{G}_m^{\wedge n} = S^{2n-1,n}$  is the free strongly invariant sheaf of abelian groups on the sheaf of pointed sets  $\mathbb{G}_m^{\wedge n}$  [Mor12, 1.23].

We can identify many groups now just using contraction:

**Corollary 3.18.2.** We have a canonical isomorphism

$$\pi_n(S^{n+i,i}) \cong \mathbf{K}_i^{\text{MW}},$$

for  $n \geq 2$  and  $i \geq 1$  [Mor12, p. 167].

The proof of the theorem follows the classical proof — we develop a notion of  $\mathbb{A}^1$ -homology and a Hurewicz theorem, which we are omitting in this class. Instead we'll look at the generators in  $\mathbf{K}_*^{\text{MW}}$  as maps of motivic spheres and verify all the relations hold.

Recall the universal symbol map

$$\sigma_n: \mathbb{G}_m^{\wedge n} \rightarrow \mathbf{K}_n^{\text{MW}}.$$

In degree 1 it send an element  $u \in k^\times$  to  $[u] \in \mathbf{K}_1^{\text{MW}}$ . Explicitly, by taking global sections  $H^0(\text{Spec}(k)_+, -)$  of the symbol map, we see that  $[u] \in \mathbf{K}_1^{\text{MW}}$  is represented by the map

$$S^0 \xrightarrow{[u]} \mathbb{G}_m,$$

sending the basepoint to the basepoint  $1 \in \mathbb{G}_m$ , and the other point to  $u \in \mathbb{G}_m$ . The element  $\eta$  is the *Hopf map*

$$\begin{aligned} \eta: \mathbb{A}^2 \setminus 0 &\rightarrow \mathbb{P}^1 \\ (x, y) &\mapsto [x : y]. \end{aligned}$$

This is a map  $S^{3,2} \rightarrow S^{2,1}$ , so it occurs in bidegree  $\pi_{-1,-1} \mathbb{1}$ .

**Proposition 3.18.3.** The following diagram commutes in  $\mathrm{Sm}_k$ :

$$\begin{array}{ccccc}
 \mathbb{G}_m \times \mathbb{G}_m & \xrightarrow{(\mathrm{id}, \mathrm{id})} & \mathbb{G}_m \times \mathbb{A}^1 & & \\
 \downarrow (\cdot) & \searrow \mu & \downarrow \mathrm{id} & \searrow \mu & \\
 & \mathbb{G}_m & \xrightarrow{\mathrm{id}} & \mathbb{A}^1 & \\
 & \downarrow & \downarrow & \downarrow & \\
 \mathbb{G}_m \times \mathbb{A}^1 & \xrightarrow{\quad} & \mathbb{A}^2 & \xrightarrow{\quad} & \mathbb{A}^1 \\
 & \searrow & \downarrow & \searrow \eta & \\
 & \mathbb{A}^1 & \xrightarrow{\quad} & \mathbb{P}^1 & 
 \end{array}$$

**Corollary 3.18.4.** Rearranging the diagram, we have that the Hopf map factors in  $\mathrm{Spc}(k)_*$  as

$$\mathbb{A}^2 \setminus 0 \rightarrow \Sigma(\mathbb{G}_m \times \mathbb{G}_m) \xrightarrow{\Sigma\mu} \Sigma\mathbb{G}_m. \quad (3.5)$$

We'll see a slightly different perspective on this soon.

### 3.18.1 The Steinberg relation

Recall the first relation **MW1** was  $[u][1-u] = 0$ . This is often called the *Steinberg relation*. The statement that it holds at the level of maps of motivic spheres was first claimed by Hu and Kriz, however their paper contained an error. It was proven by Druzhinin in the stable setting, and reproven in the unstable setting by Hoyois (see [Hoy18] for details).

Consider the pushout diagram

$$\begin{array}{ccc}
 \mathbb{A}^1 - \{0, 1\} & \longrightarrow & \mathbb{A}^1 - 1 \\
 \downarrow & \lrcorner & \downarrow \\
 \mathbb{A}^1 - 0 & \longrightarrow & \mathbb{A}^1.
 \end{array}$$

By taking the pullback of the bottom right cospan, we obtain a natural map of unpointed spaces, where each  $\mathbb{G}_m$  here is now  $\mathbb{A}^1 - 0$ :

$$\begin{aligned}
 \mathbb{A}^1 - \{0, 1\} &\rightarrow \mathbb{G}_m \times \mathbb{G}_m \\
 a &\mapsto (a, 1 - a).
 \end{aligned}$$

Given any unit  $u \in k^\times$ , the map  $\mathrm{Spec}(k) \xrightarrow{(u, 1-u)} \mathbb{G}_m \wedge \mathbb{G}_m$  factors through the unpointed map above

$$\begin{array}{ccc}
 \mathrm{Spec}(k) & \xrightarrow{\quad} & \mathbb{G}_m \wedge \mathbb{G}_m \\
 & \searrow & \nearrow \\
 & \mathbb{A}^1 - \{0, 1\} & 
 \end{array}$$

Note that  $\mathbb{G}_m \times \mathbb{G}_m$  is naturally pointed, but  $\mathbb{A}^1 - \{0, 1\}$  isn't, so we add a disjoint basepoint to get a diagram of pointed spaces

$$\begin{array}{ccc}
 S^0 & \xrightarrow{\quad} & \mathbb{G}_m \wedge \mathbb{G}_m \\
 & \searrow & \nearrow \mathrm{st} \\
 & (\mathbb{A}^1 - \{0, 1\})_+ & 
 \end{array}$$

Hence it suffices to show that the *pointed map*  $(\mathbb{A}^1 - \{0, 1\})_+ \rightarrow \mathbb{G}_m \wedge \mathbb{G}_m$  is null-homotopic after

suspension. The mistake in Hu-Kriz was not considering the image of the additional basepoint one must add.

**Theorem 3.18.6.** [Hoy18] The  $S^1$ -suspension of the Steinberg map

$$\mathrm{st}: (\mathbb{A}^1 - \{0, 1\})_+ \rightarrow \mathbb{G}_m \wedge \mathbb{G}_m$$

is null-homotopic.

*Sketch.* The main idea is to blow up  $\mathbb{A}^2$  at  $(1, 0)$  and  $(0, 1)$ , and show that the map above factors after 1-fold suspension through an induced map from the union of three lines, one connecting  $(0, 1)$  and  $(1, 1)$ , one connecting  $(0, 1)$  and  $(1, 0)$ , and the exceptional divisor over  $(0, 1)$ .  $\square$

### 3.18.2 Digression: the Hopf construction

Let  $X$  be a pointed space (motivic or topological). If  $X$  has a comultiplication  $X \rightarrow X \vee X$ , then  $\mathrm{Hom}(X, Y)$  has a monoid structure, induced by precomposition with the monoidal structure

$$\mathrm{Hom}(X, Y) \times \mathrm{Hom}(X, Y) \cong \mathrm{Hom}(X \vee X, Y) \rightarrow \mathrm{Hom}(X, Y).$$

We say that the comultiplication is *cocommutative* if the diagram commutes (up to homotopy)

$$\begin{array}{ccc} X & \longrightarrow & X \vee X \\ & \searrow & \downarrow \text{swap} \\ & & X \vee X. \end{array}$$

**Example 3.18.7.** The pinch maps

$$S^n \rightarrow S^n \vee S^n,$$

defined by collapsing a great circle containing the basepoint, define a coassociative comultiplication co- $H$ -space structure on the  $n$ -sphere. For  $n \geq 2$  this operation is cocommutative.

The comultiplication on  $S^n$  defines a group structure on  $[S^n, X]_*$ , and the cocommutative property is exactly the statement that  $\pi_n(X)$  is abelian for  $n \geq 2$ .

**Example 3.18.8.** We have that  $\Sigma X$  admits a comultiplication for any  $X \in \mathrm{Spc}_*$ , induced by the pinch map  $S^1 \rightarrow S^1 \vee S^1$ .

Suppose now that  $G \in \mathrm{Spc}_*$  is a group object. Then  $[X, G]_*$  admits a group structure, and an equation of the form  $f = f_1 + f_2$  in  $[X, G]_*$  means exactly that  $f$  factors as

$$\begin{array}{ccc} X & \xrightarrow{f_1, f_2} & G \times G \\ & \searrow f & \downarrow \mu \\ & & G. \end{array}$$

If  $G$  is a group object, this does *not* imply that  $\Sigma G$  is a group object, and in general this won't be true. So what happens to the equation  $f = f_1 + f_2$  when we suspend? In other words can we express  $\Sigma f$  in terms of  $\Sigma f_1$  and  $\Sigma f_2$  in  $[\Sigma X, \Sigma G]_*$ ?

### 3.18.3 Splitting cofiber sequences

The following we learned from [DI13, Appendix A].

Suppose we have a cofiber sequence in  $\mathrm{Spc}(k)_*$

$$A \xrightarrow{j} B \xrightarrow{p} B/A,$$

and suppose that  $j$  is split after suspension – in other words there exists some map  $f: \Sigma B \rightarrow \Sigma A$  so that the following diagram commutes up to homotopy

$$\begin{array}{ccc} \Sigma A & \longrightarrow & \Sigma B \\ & \searrow \text{id} & \downarrow f \\ & & \Sigma A. \end{array}$$

Applying  $[-, X]_*$ , we get a sequence of groups, which turns out to be short exact

$$1 \rightarrow [\Sigma(B/A), X]_* \rightarrow [\Sigma B, X]_* \rightarrow [\Sigma A, X]_*.$$

This is not necessarily split exact – if  $f$  were a suspension of a map, then it would be a map of  $H$ -spaces and induce a splitting, but the best we can say here is that the sequence is exact.

**Case:** Let's take  $X = \Sigma B$ . Then we get a short exact sequence

$$0 \rightarrow [\Sigma(B/A), \Sigma B] \rightarrow [\Sigma B, \Sigma B] \rightarrow [\Sigma A, \Sigma B].$$

Note that since  $\text{id}_{\Sigma B} = (\Sigma j)f$ , the map  $\text{id}_{\Sigma B} - (\Sigma j)f$  is in the kernel of the rightmost map, hence is uniquely in the image of  $[\Sigma(B/A), \Sigma B]$ .

Let  $\chi: \Sigma(B/A) \rightarrow \Sigma B$  be the unique map sent to  $\text{id}_{\Sigma B} - (\Sigma j)f$ .

**Lemma 3.18.9.** [DI13, A.3] We have that the diagram commutes

$$\begin{array}{ccc} \Sigma B/A & \xrightarrow{\chi} & \Sigma B \\ & \searrow & \downarrow \Sigma p \\ & & \Sigma B/A, \end{array}$$

and the composite

$$\Sigma B/A \xrightarrow{\chi} \Sigma B \xrightarrow{f} \Sigma A$$

is null-homotopic.

**Proposition 3.18.10.** For any pointed spaces  $X, Y \in \text{Spc}(k)_*$ , the cofiber sequence

$$X \vee Y \rightarrow X \times Y \rightarrow X \wedge Y$$

admits a canonical splitting after suspension:

$$\Sigma X \vee \Sigma Y \vee \Sigma(X \wedge Y) \cong \Sigma(X \times Y).$$

We claim further that this is an equivalence of co- $H$ -spaces, hence the identity on  $\Sigma(X \times Y)$  is a sum of the projection maps.

*Proof.* Suspension is a left adjoint and hence preserves the above coproduct. Since the suspension of anything is a co-group object (by the pinch map), we have an induced map

$$\Sigma(X \times Y) \rightarrow \Sigma(X \times Y) \vee \Sigma(X \times Y).$$

Post-composing with the suspensions of the two projections  $X \times Y \rightarrow X$  and  $X \times Y \rightarrow Y$  yields a map  $\Sigma(X \times Y) \rightarrow \Sigma(X) \vee \Sigma(Y) = \Sigma(X \vee Y)$  which we claim is left inverse to the map in the suspended cofiber sequence.  $\square$

**Corollary 3.18.11.** If  $G$  is a group object, we can suspend the multiplication  $G \times G \rightarrow G$ , and use the splitting above to get three maps

$$\Sigma G \vee \Sigma G \vee \Sigma(G^{\wedge 2}) \rightarrow \Sigma G.$$

The first two component functions are the identity, and the latter is the *Hopf construction*  $\Sigma(G \wedge G) \rightarrow \Sigma G$ .

**Example 3.18.12.** The Hopf construction  $\Sigma(S^1 \wedge S^1) \rightarrow \Sigma S^1$  can be identified with the Hopf map  $\eta: S^3 \rightarrow S^2$  classically.

**Proposition 3.18.13.** The preferred map  $\Sigma\mathbb{G}_m \wedge \mathbb{G}_m \rightarrow \Sigma\mathbb{G}_m$  from the multiplication on  $\mathbb{G}_m$  can be identified with the Hopf map  $\eta$  [Mor12, p. 71]. In particular since  $\Sigma(\mathbb{G}_m \wedge \mathbb{G}_m) = \mathbb{A}^2 \setminus 0$ , we have that the suspension of the natural map  $\mathbb{G}_m \times \mathbb{G}_m \rightarrow \mathbb{G}_m \wedge \mathbb{G}_m$  admits a splitting  $s$ , hence we obtain a composite

$$\Sigma(\mathbb{G}_m \wedge \mathbb{G}_m) \xrightarrow{s} \Sigma(\mathbb{G}_m \times \mathbb{G}_m) \xrightarrow{\Sigma\mu} \Sigma\mathbb{G}_m,$$

which is precisely the factorization of the Hopf map we found in Equation 3.5.

**Proposition 3.18.14.** The splitting

$$\Sigma(\mathbb{G}_m \times \mathbb{G}_m) \xrightarrow{\pi_1 \vee \pi_2} \Sigma(\mathbb{G}_m \vee \mathbb{G}_m) \vee \Sigma(\mathbb{G}_m \wedge \mathbb{G}_m)$$

has the property that the two projections sum to the identity on  $\Sigma(\mathbb{G}_m \times \mathbb{G}_m)$ .

We can now prove **MW2**:

**Proposition 3.18.15.** The following relation holds in  $\pi_{*,*}\mathbb{S}$ :  $[ab] = [a] + [b] + \eta[a][b]$ .

*Proof.* The map

$$S^0 \xrightarrow{[ab]} \mathbb{G}_m$$

factors as

$$S^0 \xrightarrow{[a],[b]} \mathbb{G}_m \times \mathbb{G}_m \xrightarrow{\mu} \mathbb{G}_m.$$

Suspending, and leveraging the splitting above, we get a commutative diagram

$$\begin{array}{ccccc} & & \Sigma[ab] & & \\ & \nearrow & & \searrow & \\ S^1 & \xrightarrow{\Sigma([a],[b])} & \Sigma(\mathbb{G}_m \times \mathbb{G}_m) & \xrightarrow{\Sigma\mu} & \Sigma\mathbb{G}_m \\ & \searrow & \downarrow \cong & \nearrow & \\ & & \Sigma(\mathbb{G}_m \wedge \mathbb{G}_m) \vee \Sigma\mathbb{G}_m \vee \Sigma\mathbb{G}_m & & \end{array}$$

$\Sigma[a] \vee \Sigma[b] \vee \eta[a][b]$

□

# Bibliography

- [ABH23] Aravind Asok, Tom Bachmann, and Michael J. Hopkins. *On  $\mathbb{P}^1$ -stabilization in unstable motivic homotopy theory*. 2023. arXiv: [2306.04631 \[math.AG\]](#).
- [ABH24] Aravind Asok, Tom Bachmann, and Michael J. Hopkins. *On the Whitehead theorem for nilpotent motivic spaces*. 2024. arXiv: [2210.05933 \[math.AG\]](#).
- [AD09] Aravind Asok and Brent Doran.  *$A^1$ -homotopy groups, excision, and solvable quotients*. 2009. arXiv: [0902.1564 \[math.AG\]](#).
- [AF14a] Aravind Asok and Jean Fasel. “A cohomological classification of vector bundles on smooth affine threefolds”. In: *Duke Math. J.* 163.14 (2014), pp. 2561–2601.
- [AF14b] Aravind Asok and Jean Fasel. “Algebraic vector bundles on spheres”. In: *Journal of Topology* 7.3 (Feb. 2014), pp. 894–926.
- [AF14c] Aravind Asok and Jean Fasel. “Algebraic vector bundles on spheres”. In: *J. Topol.* 7.3 (2014), pp. 894–926.
- [AHW18] Aravind Asok, Marc Hoyois, and Matthias Wendt. “Affine representability results in  $A^1$ -homotopy theory, II: Principal bundles and homogeneous spaces”. In: *Geom. Topol.* 22.2 (2018), pp. 1181–1225.
- [AHW20] Aravind Asok, Marc Hoyois, and Matthias Wendt. “Affine representability results in  $A^1$ -homotopy theory III: finite fields and complements”. In: *Algebr. Geom.* 7.5 (2020), pp. 634–644.
- [AO19] Aravind Asok and Paul Arne Østvær.  *$A^1$ -homotopy theory and contractible varieties: a survey*. 2019. arXiv: [1903.07851 \[math.AG\]](#).
- [AP71] Jón Kristinn Arason and Albrecht Pfister. “Beweis des Krullschen Durchschnittsatzes für den Witttring”. In: *Inventiones mathematicae* 12.2 (June 1971), pp. 173–176.
- [Aso21] Aravind Asok. *Algebraic geometry from an  $A^1$ -homotopic viewpoint*. 2021.
- [Bac] Tom Bachmann. *Algebraic K-theory from the viewpoint of motivic homotopy theory*.
- [Bac24] Tom Bachmann. *Strongly  $A^1$  – invariant sheaves (after F. Morel)*. 2024. arXiv: [2406.11526 \[math.AG\]](#).
- [Bar+23] Viktor Balch Barth et al. *Making the motivic group structure on the endomorphisms of the projective line explicit*. 2023. arXiv: [2306.00628 \[math.AG\]](#).
- [Car23] Robin Carlier. *Milnor-Witt K-theory and Witt K-theory of a field*. 2023. arXiv: [2306.16985 \[math.AG\]](#).
- [Caz09] Christophe Cazanave. “Théorie homotopique des schémas d’Atiyah et Hitchin”. Université Paris 13, 2009.

- [Caz12] Christophe Cazanave. “Algebraic Homotopy Classes of Rational Functions”. In: *Annales scientifiques de l’École normale supérieure* 45.4 (2012), pp. 511–534.
- [Dég07] Frédéric Déglise. “Finite correspondences and transfers over a regular base”. In: *Algebraic cycles and motives. Vol. 1*. Vol. 343. London Math. Soc. Lecture Note Ser. Cambridge Univ. Press, Cambridge, 2007, pp. 138–205.
- [Dég11] F. Déglise. “Modules homotopiques”. In: *Doc. Math.* 16 (2011), pp. 411–455.
- [Dég23] Frédéric Déglise. *Notes on Milnor-Witt K-theory*. 2023. arXiv: [2305.18609](#) [math.AG].
- [Del74] Pierre Deligne. “Théorie de Hodge. III”. In: *Inst. Hautes Études Sci. Publ. Math.* 44 (1974), pp. 5–77.
- [DI13] Daniel Dugger and Daniel C. Isaksen. “Motivic Hopf elements and relations”. In: *New York J. Math.* 19 (2013), pp. 823–871.
- [Dug01] Daniel Dugger. “Universal homotopy theories”. In: *Adv. Math.* 164.1 (2001), pp. 144–176.
- [EKM08] Richard Elman, Nikita Karpenko, and Alexander Merkurjev. *The algebraic and geometric theory of quadratic forms*. Vol. 56. American Mathematical Society Colloquium Publications. American Mathematical Society, Providence, RI, 2008, pp. viii+435.
- [Fel21] Niels Feld. “Milnor-Witt sheaves and modules”. PhD thesis. 2021.
- [Fri82] Eric M. Friedlander. *Étale homotopy of simplicial schemes*. Vol. No. 104. Annals of Mathematics Studies. Princeton University Press, Princeton, NJ; University of Tokyo Press, Tokyo, 1982, pp. vii+190.
- [Ful98] William Fulton. *Intersection theory*. Second. Vol. 2. Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]. Springer-Verlag, Berlin, 1998, pp. xiv+470.
- [GJ99] Paul G. Goerss and John F. Jardine. *Simplicial homotopy theory*. Vol. 174. Progress in Mathematics. Birkhäuser Verlag, Basel, 1999, pp. xvi+510.
- [Gro] Alexandre Grothendieck. “Torsion homologique et sections rationnelles”. fr. In: ().
- [GS17] Philippe Gille and Tamás Szamuely. *Central simple algebras and Galois cohomology*. Second. Vol. 165. Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, 2017, pp. xi+417.
- [Hal] Connor Halleck-Dubé. *Some remarks on Galois cohomology and linear algebraic groups*.
- [Har77] Robin Hartshorne. *Algebraic geometry*. Vol. No. 52. Graduate Texts in Mathematics. Springer-Verlag, New York-Heidelberg, 1977, pp. xvi+496.
- [Hol08] Sharon Hollander. “A homotopy theory for stacks”. In: *Israel J. Math.* 163 (2008), pp. 93–124.
- [Hoy14] Marc Hoyois. “A quadratic refinement of the Grothendieck-Lefschetz-Verdier trace formula”. In: *Algebr. Geom. Topol.* 14.6 (2014), pp. 3603–3658.
- [Hoy15] Marc Hoyois. “From algebraic cobordism to motivic cohomology”. In: *J. Reine Angew. Math.* 702 (2015), pp. 173–226.
- [Hoy17] Marc Hoyois. “The six operations in equivariant motivic homotopy theory”. In: *Adv. Math.* 305 (2017), pp. 197–279.
- [Hoy18] Marc Hoyois. *The Steinberg Relation*. 2018.



- [Lam05] T. Y. Lam. *Introduction to quadratic forms over fields*. Vol. 67. Graduate Studies in Mathematics. American Mathematical Society, Providence, RI, 2005, pp. xxii+550.
- [Lur] Jacob Lurie. *Spectral algebraic geometry*.
- [Lur09] Jacob Lurie. *Higher topos theory*. Vol. 170. Annals of Mathematics Studies. Princeton University Press, Princeton, NJ, 2009, pp. xviii+925.
- [Mil13] James S. Milne. *Lectures on Etale Cohomology (v2.21)*. Available at [www.jmilne.org/math/](http://www.jmilne.org/math/). 2013.
- [Mil15] James S. Milne. *Algebraic Groups (v2.00)*. Available at [www.jmilne.org/math/](http://www.jmilne.org/math/). 2015.
- [Mil70] John Milnor. “Algebraic  $K$ -theory and quadratic forms”. In: *Invent. Math.* 9 (1969/70), pp. 318–344.
- [Mor12] Fabien Morel. *A1-Algebraic Topology over a Field*. en. Vol. 2052. Lecture Notes in Mathematics. Berlin, Heidelberg: Springer Berlin Heidelberg, 2012.
- [MV99] Fabien Morel and Vladimir Voevodsky. “ $\mathbf{A}^1$ -homotopy theory of schemes”. In: *Inst. Hautes Études Sci. Publ. Math.* 90 (1999), pp. 45–143.
- [Rie14] Emily Riehl. *Categorical homotopy theory*. Vol. 24. New Mathematical Monographs. Cambridge University Press, Cambridge, 2014, pp. xviii+352.
- [Rob15] Marco Robalo. “ $K$ -theory and the bridge from motives to noncommutative motives”. In: *Adv. Math.* 269 (2015), pp. 399–550.
- [Sha23] Jay Shah. “Parametrized higher category theory”. In: *Algebraic Geometric Topology* 23.2 (May 2023), pp. 509–644.
- [Stacks] The Stacks Project Authors. *Stacks Project*. <https://stacks.math.columbia.edu>. 2018.
- [Tra70] Carlo Traverso. “Seminormality and Picard group”. In: *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (3)* 24 (1970), pp. 585–595.
- [Wei13] Charles A. Weibel. *The K-book*. Vol. 145. Graduate Studies in Mathematics. An introduction to algebraic  $K$ -theory. American Mathematical Society, Providence, RI, 2013, pp. xii+618.
- [Wei89] Charles A. Weibel. “Homotopy algebraic  $K$ -theory”. In: *Algebraic K-theory and algebraic number theory (Honolulu, HI, 1987)*. Vol. 83. Contemp. Math. Amer. Math. Soc., Providence, RI, 1989, pp. 461–488.
- [Wei94] Charles A. Weibel. *An introduction to homological algebra*. Vol. 38. Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, 1994, pp. xiv+450.
- [You] Alex Youcis. “A brief discussion of torsors”.