

NUMERICAL ANALYSIS

PROBLEM SET 4

ROB TOMPKINS

1. PROBLEM 1

1.a. We are deriving the 2-point open Newton-Coats formula

$$\int_a^b f(x)dx \approx I_{2,\text{open}}(f) = \frac{b-a}{3} \left(2f(a+h) - f(a+2h) + 2f(a+3h) \right)$$

for evenly spaced points with $h = \frac{b-a}{n+2}$. Now we set $n = 2$ so that $h = \frac{b-a}{4}$, and we need to sort out

$$(1) \quad I_{2,\text{open}}(f) = \int_a^b \sum_{i=0}^2 L_{2,i}(x) f(x_i) dx.$$

But we need to shift to an easier or simpler variable to work with. So we set $x = a + t \cdot h$, with $dx = h \cdot dt$. Now as x moves from a to b we see that ($b = a + 4h$), t moves from 0 to 4, and we have

$$\int_a^b f(x)dx = h \int_0^4 f(a+ht)dt,$$

which we interpolate $f(a+ht)$ at $t = 1, 2, 3$. We write out the Lagrangian polynomials $L_{2,0}, L_{2,1}, L_{2,2}$ for the sake of the problem. This follows as

$$L_{2,0}(x) = \frac{(t-2)(t-3)}{(1-2)(1-3)} = \frac{(t-2)(t-3)}{2} = \frac{1}{2}(t^2 - 5t + 6)$$

$$L_{2,1}(x) = \frac{(t-1)(t-3)}{(2-1)(2-3)} = -(t-1)(t-3) = -t^2 + 4t - 3$$

$$L_{2,2}(x) = \frac{(t-1)(t-2)}{(3-1)(3-2)} = \frac{(t-1)(t-2)}{2} = \frac{1}{2}(t^2 - 3t + 2).$$

And we then need to integrate these Lagrange interpolants, which we do as

$$\begin{aligned}\int_0^4 \frac{1}{2}(t^2 - 5t + 6)dt &= \left. \frac{t^3}{6} - \frac{5t^2}{4} + 3t \right|_0^4 = \frac{8}{3} \\ \int_0^4 -t^2 + 4t - 3dt &= \left. -\frac{t^3}{3} + 2t^2 - 3t \right|_0^4 = -\frac{4}{3} \\ \int_0^4 \frac{1}{2}(t^2 - 3t + 2)dt &= \left. \frac{t^3}{6} - \frac{3t^2}{2} + 2t \right|_0^4 = \frac{8}{3}\end{aligned}$$

Thus, when we plug this into equation (1), we get the following

$$\begin{aligned}\int_a^b f(x)dx &\approx h \left(\frac{8}{3}f(a+h) - \frac{4}{3}f(a+2h) + \frac{8}{3}f(a+3h) \right) \\ &\approx \frac{4h}{3} \left(2f(a+h) - f(a+2h) + 2f(a+3h) \right) \\ &\approx \frac{b-a}{3} \left(2f(a+h) - f(a+2h) + 2f(a+3h) \right).\end{aligned}$$

which solves our problem

1.b. We must show that our quadrature is exact for $1, x, x^2, x^3$ but not for x^4 . Again we rescale the interval in teh fashion

$$x = a + ht, \quad h = \frac{b-a}{4}, x \in [a, b] \rightarrow t \in [0, 4]$$

So $Q(g) = \frac{4}{3} \left(2g(1) - g(2) - 2g(3) \right)$. But, we need to check if $I(g) = Q(g)$ for $g(t) \in \{1, x, x^2, x^3, x^4\}$. So we write

$$\begin{aligned}\int_0^4 1dt &= 4 \stackrel{?}{=} \frac{4}{3}(2 - 1 + 2) = 4 \quad \checkmark \\ \int_0^4 tdt &= \left. \frac{t^2}{2} \right|_0^4 = 8 \stackrel{?}{=} \frac{4}{3}(2 \cdot 1 - 2 + 2 \cdot 3) = 8 \quad \checkmark \\ \int_0^4 t^2dt &= \left. \frac{t^3}{3} \right|_0^4 = \frac{64}{3} \stackrel{?}{=} \frac{4}{3}(2 \cdot 1 - 4 + 2 \cdot 9) = \frac{4 \cdot 16}{3} = \frac{64}{3} \quad \checkmark\end{aligned}$$