

ADVANCED LINEAR ALGEBRA HOMEWORK 1

ROB TOMPKINS

P.1.2 Is $[1 \ 0 \ 1]^\top \in \mathbb{R}^3$ in the span of $[-1 \ 3 \ 2]^\top$ and $[1 \ 1 \ -2]^\top$? Why?

Solution. We will show that $[1 \ 0 \ 1]^\top \notin \text{span}([-1 \ 3 \ 2]^\top, [1 \ 1 \ -2]^\top)$ by contradiction. So assume that there is some $a, b \in \mathbb{R}$ with

$$a \begin{bmatrix} -1 \\ 3 \\ 2 \end{bmatrix} + b \begin{bmatrix} 1 \\ 1 \\ -2 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}$$

Which if we expand we get the following equations:

$$-a + b = 1$$

$$3a + b = 0$$

$$2a - 2b = 1$$

If we simply try to solve the first and third equation we need that the matrix equation to hold:

$$\begin{bmatrix} -1 & 1 \\ 2 & -2 \end{bmatrix} \begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

For this to be the case we would need that $\begin{bmatrix} -1 & 1 \\ 2 & -2 \end{bmatrix}$ be invertible so its determinant would need to be non-zero. Let's calculate the determinant:

$$\det \begin{bmatrix} -1 & 1 \\ 2 & -2 \end{bmatrix} = (2 - 2) = 0.$$

So our matrix is not invertible, and we cannot find a unique a and b to solve our system of equations. $[1 \ 0 \ 1]^\top \notin \text{span}([-1 \ 3 \ 2]^\top, [1 \ 1 \ -2]^\top)$ $\square$

P.1.3 Is $[7 \ 8 \ 9]^\top \in \mathbb{R}^3$ in the span of $[1 \ 2 \ 3]^\top$ and $[4 \ 5 \ 6]^\top$? Why? Is $[10 \ 11 \ 12]^\top$?

Solution. In a similar fashion to problem **P.1.2**, we choose an $a, b \in \mathbb{R}$, and we set:

$$a \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} + b \begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix} = \begin{bmatrix} 7 \\ 8 \\ 9 \end{bmatrix}.$$

This reduces to the equations:

$$\begin{aligned}a + 4b &= 7 \\2a + 5b &= 8 \\3a + 6b &= 9\end{aligned}$$

So if we let $a = -4b + 7$, as given by the first equation and then substitute into the second equation, we see that

$$\begin{aligned}2(-4b + 7) + 5b &= 8 \\-8b + 14 + 5b &= 8 \\-3b &= -6 \\b &= 2\end{aligned}$$

Substituting back into our top equality we see that $a + 4(2) = 7$, thus $a = -1$. Furthermore if we plug these values back into our original linear combination, we get that

$$\begin{aligned}-1 \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} + 2 \begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix} &= \begin{bmatrix} 7 \\ 8 \\ 9 \end{bmatrix} \\ \begin{bmatrix} -1 \\ -2 \\ -3 \end{bmatrix} + \begin{bmatrix} 8 \\ 10 \\ 12 \end{bmatrix} &= \begin{bmatrix} 7 \\ 8 \\ 9 \end{bmatrix}\end{aligned}$$

If we were to do the same inspection for the vector $[10 \ 11 \ 12]^\top$ we write $a, b \in \mathbb{R}$ with

$$a \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} + b \begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix} = \begin{bmatrix} 10 \\ 11 \\ 12 \end{bmatrix},$$

We then expand to:

$$\begin{aligned}a + 4b &= 10 \\2a + 5b &= 11 \\3a + 6b &= 12\end{aligned}$$

So we have that $a = -4b + 10$ which if we plug into the second equation we get:

$$2(-4b + 10) + 5b = 11$$

$$-8b + 20 + 5b = 11$$

$$-3b = -9$$

$$b = 6$$

And if we plug b back in we get $a = -24 + 10 = -14$. So we test this out to see if it works:

$$-14 \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} + 6 \begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix} = \begin{bmatrix} 10 \\ 11 \\ 12 \end{bmatrix}$$

$$\begin{bmatrix} -14 \\ -28 \\ -42 \end{bmatrix} + \begin{bmatrix} 24 \\ 30 \\ 36 \end{bmatrix} = \begin{bmatrix} 10 \\ 2 \\ -6 \end{bmatrix} \neq \begin{bmatrix} 10 \\ 11 \\ 12 \end{bmatrix}$$

So $[10 \ 11 \ 12]^\top \notin \text{span}([1 \ 2 \ 3]^\top, [4 \ 5 \ 6]^\top)$

□

P.1.8 Consider the following lists of vectors in $\mathbb{R}^3$:

$$\beta = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \begin{bmatrix} 4 \\ s \\ 6 \end{bmatrix}, \begin{bmatrix} 7 \\ 8 \\ 9 \end{bmatrix},$$

and

$$\gamma = \begin{bmatrix} 4 \\ 2 \\ 6 \end{bmatrix}, \begin{bmatrix} 1 \\ t \\ -1 \end{bmatrix}, \begin{bmatrix} 2 \\ -2 \\ 3 \end{bmatrix},$$

For what values of s is β linearly dependent. For what values of t is γ linearly dependent.

Solution.

□

P.1.12 Show that the set of $n \times n$ Hermitian matrices is a real vector space.

Proof.

□

P.1.13 Let

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix}$$

(a) Choose two of the following and show that each is a factorization $A = XY$, in which the columns of X and rows of Y are each linearly independent:

$$\begin{bmatrix} 1 & 2 \\ 4 & 5 \\ 7 & 8 \end{bmatrix} \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 2 \end{bmatrix}, \begin{bmatrix} 2 & 3 \\ 5 & 6 \\ 8 & 9 \end{bmatrix} \begin{bmatrix} 2 & 0 & 1 \\ -1 & 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 3 \\ 4 & 6 \\ 7 & 9 \end{bmatrix} \begin{bmatrix} 1 & \frac{1}{2} & 0 \\ 0 & \frac{1}{2} & 1 \end{bmatrix}, \begin{bmatrix} -1 & 3 \\ -1 & 9 \\ -1 & 15 \end{bmatrix} \begin{bmatrix} \frac{1}{2} & -\frac{1}{2} & -\frac{3}{2} \\ \frac{1}{2} & \frac{1}{2} & \frac{1}{2} \end{bmatrix}$$

(b) Which of the decompositions is the one described by Theorem 1.7.5? Why?

Solution.

□

P.1.27 (a) Show that $\text{col}(A) = \mathbb{R}^3$ for

$$A = [\mathbf{a}_1 \quad \mathbf{a}_2 \quad \mathbf{a}_3 \quad \mathbf{a}_4] = \begin{bmatrix} 1 & 1 & 1 & 0 \\ 0 & 1 & 1 & 1 \\ 0 & 0 & 1 & 1 \end{bmatrix} \in \mathbf{M}_n(\mathbb{R})$$

(b) Find $i, j, k \in \{1, 2, 3, 4\}$ such that $\text{span}\{\mathbf{a}_i, \mathbf{a}_j, \mathbf{a}_k\} = \mathbb{R}^3$. And find distinct $i, j, k \in \{1, 2, 3, 4\}$ such that $\text{span}\{\mathbf{a}_i, \mathbf{a}_j, \mathbf{a}_k\}$.

Solution.

□