

## Axioms of Probability

1.  $0 \leq \Pr(A) \leq 1$
2.  $\Pr(S) = 1$
3. If  $A_1, A_2, \dots$  are mutually exclusive (disjoint),  
i.e.  $A_i \cap A_j = \emptyset$  when  $i \neq j$ , then  $\Pr(\cup_{i=1}^{\infty} A_i) = \sum_{i=1}^{\infty} \Pr(A_i)$   
In particular, if events  $A$  and  $B$  are mutually exclusive, then  $\Pr(A \cup B) = \Pr(A) + \Pr(B)$

## Properties of Probability

1.  $\Pr(\emptyset) = 0$
2. If  $A_1, A_2, \dots, A_n$  are mutually exclusive events, then  $\Pr(\cup_{i=1}^n A_i) = \sum_{i=1}^n \Pr(A_i)$
3.  $\Pr(A') = 1 - \Pr(A)$
4.  $\Pr(A) = \Pr(A \cap B) + \Pr(A \cap B')$
5.  $\Pr(A \cup B) = \Pr(A) + \Pr(B) - \Pr(A \cap B)$
6.  $\Pr(A \cup B \cup C) = \Pr(A) + \Pr(B) + \Pr(C) - \Pr(A \cap B) - \Pr(B \cap C) - \Pr(A \cap C) + \Pr(A \cap B \cap C)$

## Conditional Probability, $P(A | B)$

- $\Pr(A | B) = \frac{\Pr(A \cap B)}{\Pr(B)}$ , if  $\Pr(B) > 0$
- For fixed  $A$ ,  $\Pr(B | A)$  satisfies the postulates of probability.
- False positive:  $\Pr(+ | \text{condition})$

## Multiplication rule

- $\Pr(A \cap B) = \Pr(A) \Pr(B | A) = \Pr(B) \Pr(A | B)$ , providing  $\Pr(A) > 0, \Pr(B) > 0$
- $\Pr(A \cap B \cap C) = \Pr(A) \Pr(B | A) \Pr(C | A \cap B)$
- $\Pr(A_1 \cap \dots \cap A_n) = \Pr(A_1) \Pr(A_2 | A_1) \Pr(A_3 | A_1 \cap A_2) \dots \Pr(A_n | A_1 \cap \dots \cap A_{n-1})$

## The Law of Total Probability

- Let  $A_1, A_2, \dots, A_n$  be a partition of sample space  $S$  (mutually exclusive and exhaustive events s.t.  $A_i \cap A_j = \emptyset$  for  $i \neq j$  and  $\cup_{i=1}^n A_i = S$ ).
- Then  $\Pr(B) = \sum_{i=1}^n \Pr(B \cap A_i) = \sum_{i=1}^n \Pr(A_i) \Pr(B | A_i)$

## Bayes' Theorem

- Let  $A_1, A_2, \dots, A_n$  be a partition of  $S$
- $\Pr(A_k | B) = \frac{\Pr(A_k) \Pr(B | A_k)}{\sum_{i=1}^n \Pr(A_i) \Pr(B | A_i)} = \frac{\Pr(A_k) \Pr(B | A_k)}{\Pr(B)}$ ,  $k \in [1, n]$

## Independent Events

- Definition: iff  $\Pr(A \cap B) = \Pr(A) \Pr(B)$

## Properties

- Suppose  $\Pr(A) > 0, \Pr(B) > 0$ ,  $A$  and  $B$  are independent:
  - $\Pr(B | A) = \Pr(B)$  and  $\Pr(A | B) = \Pr(A)$
  - $A$  and  $B$  cannot be mutually exclusive (and vice versa)
- The sample space  $S$  and  $\emptyset$  are independent of any event
- If  $A \subset B$ , then  $A$  and  $B$  are dependent unless  $B = S$

## Theorem

If  $A, B$  are indep, then so are  $A$  and  $B'$ ,  $A'$  and  $B$ ,  $A'$  and  $B'$ .

## $n$ Independent Events

- **Pairwise Independent Events:**

Events  $A_1, A_2, \dots, A_n$  are pairwise indep

iff  $\Pr(A_i \cap A_j) = \Pr(A_i) \Pr(A_j)$

- **Mutually Independent:**

Events  $A_1, A_2, \dots, A_n$  are (mutually) independent iff for any subset  $\{A_{i_1}, A_{i_2}, \dots, A_{i_k}\}$  of  $A_1, A_2, \dots, A_n$ ,

$\Pr(A_{i_1} \cap A_{i_2} \cap \dots \cap A_{i_k}) = \Pr(A_{i_1}) \Pr(A_{i_2}) \dots \Pr(A_{i_k})$

## Remarks

- $A_1, A_2, \dots, A_n$  are mutually independent  $\Leftrightarrow$  for any pair of events  $A_j, A_k$  where  $j \neq k$ , the multiplication rule holds, for any 3 distinct events, the multiplication rule holds, and so on  $\Pr(A_1 \cap A_2 \cap \dots \cap A_n) = \Pr(A_1) \Pr(A_2) \dots \Pr(A_n)$ . In total there are  $2^n - n - 1$  different cases.
- Mutually indep  $\Rightarrow$  pairwise indep (not the converse)
- Suppose  $A_1, A_2, \dots, A_n$  are mutually indep events, let  $B_i = A_i$  or  $A'_i$ ,  $i \in [1, n]$ . Then  $B_1, B_2, \dots, B_n$  are also mutually indep events.

## Discrete Probability Distributions

### Discrete R.V.

Let  $X$  be an R.V. If  $R_X$  is **finite or countable infinite**,  $X$  is discrete R.V.

### Probability Function (p.f.) or Probability Mass Function (p.m.f.)

- For a discrete R.V., each value  $X$  has a certain probability  $f(x)$ . Such a function  $f(x)$  is called the p.f.
- The collection of pairs  $(x_i, f(x_i))$  is prob distribution of  $X$
- The probability of  $X = x_i$  denoted by  $f(x_i)$  must satisfy:
  1.  $f(x_i) \geq 0 \forall x_i$
  2.  $\sum_{i=1}^{\infty} f(x_i) = 1$

## Continuous Probability Distributions

### Continuous R.V.

Suppose that  $R_X$  is an **interval or a collection of intervals**, then  $X$  is a continuous R.V.

### Probability Density Function (p.d.f.)

- Let  $X$  be a continuous R.V.
- p.d.f.  $f(x)$  is a function satisfying:
  1.  $f(x) \geq 0 \forall x \in R_X$
  2.  $\int_{R_X} f(x) dx = 1$  or  $\int_{-\infty}^{\infty} f(x) dx = 1$  as  $f(x) = 0 \forall x \notin R_X$
  3.  $\forall c, d : c < d$  (i.e.  $(c, d) \subset R_X$ ),  $\Pr(c \leq X \leq d) = \int_c^d f(x) dx$

## Remarks

- $\Pr(c \leq X \leq d) = \int_c^d f(x) dx$  represents area under the graph of the p.d.f.  $f(x)$  between  $x = c$  and  $x = d$
- Let  $x_0$  be a fixed value,  $\Pr(X = x_0) = 0$
- $\leq$  and  $<$  can be used interchangeably in a prob statement.
- $\Pr(A) = 0$  does not necessarily imply  $A = \emptyset$
- $R_X \in [a, b] \Rightarrow f(x) = 0 \forall x \notin [a, b]$

## Cumulative Distribution Function (c.d.f.)

- Let  $X$  be an R.V., discrete or continuous.
- $F(x)$  is a c.d.f. of  $X$  where  $F(x) = \Pr(X \leq x)$

### c.d.f. for Discrete R.V.

- $F(x) = \sum_{t \leq x} f(t) = \sum_{t \leq x} \Pr(X = t)$
- c.d.f. of a discrete R.V. is a step function
- $\forall a, b$  s.t.  $a \leq b$ ,  $\Pr(a \leq X \leq b) = \Pr(X \leq b) - \Pr(X < a) = F(b) - F(a^-)$  where  $a^-$  is the largest possible value of  $X$  that is strictly less than  $a$
- $R_X \subset \mathbb{Z}, a, b \in \mathbb{Z} \Rightarrow$ 
  - $\Pr(a \leq X \leq b) = \Pr(X = a \text{ or } a + 1 \text{ or } \dots \text{ or } b) = F(b) - F(a - 1)$
  - Taking  $a = b$ ,  $\Pr(X = a) = F(a) - F(a - 1)$

### c.d.f. for Continuous R.V.

- $F(x) = \int_{-\infty}^{\infty} f(t) dt$
- $f(x) = \frac{dF(x)}{dx}$  if the derivative exists
- $\Pr(a \leq X \leq b) = \Pr(a < X < b) = F(b) - F(a)$
- $F(x)$  is a non-decreasing function:  $x_1 < x_2 \Rightarrow F(x_1) \leq F(x_2)$
- $0 \leq F(x) \leq 1$

## Mean and Variance of an R.V.

### Expected Value / Mean / Mathematical Expectation

- **Discrete:**  $E(X) = \mu_X = \sum_i x_i f(x_i) = \sum_x x f(x)$
- **Continuous:**  $E(X) = \mu_X = \int_{-\infty}^{\infty} x f(x) dx$
- **Remark:** The expected value exists provided the sum/integral exists

### Expectation of a function of an R.V.

$\forall g(X)$  with p.f.  $f_X(x)$

- **Discrete:**  $E[g(X)] = \sum_x g(x) f_X(x)$
- **Continuous:**  $E[g(X)] = \int_{-\infty}^{\infty} g(x) f_X(x) dx$
- Provided the sum/integral exists.

### Variance ( $\sigma_X^2 = V(X)$ )

- $g(x) = (x - \mu_X)^2$ , Let  $X$  be an R.V. with p.f.  $f(x)$
- $\sigma_X^2 = V(X) = E[(X - \mu_X)^2]$
- $E[(X - \mu_X)^2] = \begin{cases} \sum_x (x - \mu_X)^2 f_X(x) & \text{if } X \text{ is discrete} \\ \int_{-\infty}^{\infty} (x - \mu_X)^2 f_X(x) dx & \text{if } X \text{ is continuous} \end{cases}$
- $V(X) \geq 0$ ,  $V(X) = E(X^2) - [E(X)]^2$
- **Standard deviation**  $= \sigma_X = \sqrt{V(X)}$

## Properties of Expectation

1.  $E(aX + b) = aE(X) + b$
2.  $V(X) = E(X^2) - [E(X)]^2$
3.  $V(aX + b) = a^2 V(X)$