



Linear Algebra

Name: Aditya Vangala(Masters from IISc Bangalore)

Subjects: Engineering Maths for all branches and
Signals and Systems for ECE, EE and IN.

Reference Book: Matrix Analysis and applied Linear
Algebra by Carl D Meyer

Email Id: adityasrnvs@gmail.com



Contents

1. System of linear equations
2. Matrix Algebra
3. Eigen values and Eigen vectors

For ECE students vector space

For CSE students LU decomposition



Imagine there are three friends. They go to a market

Friend 1 $3\text{apples} + 2\text{bananas} + 1\text{orange} = 39$ — (1)

Friend 2 $2\text{apples} + 3\text{bananas} + 1\text{orange} = 26$ — (2)

Friend 3 $1\text{apples} + 2\text{bananas} + 3\text{orange} = 25$ — (3)

How to find cost of each fruit??

Solve (1) and (2) to eliminate x (4)

Solve (1) and (3) to eliminate x (5)

Solve (4) and (5) to eliminate y and get z

} Elimination of variables



Solve the following linear system of equations

$$2x + y + z = 1$$

$$6x + 2y + z = -1$$

$$-2x + 2y + z = 7$$

$$\begin{bmatrix} 2 & 1 & 1 \\ 6 & 2 & 1 \\ -2 & 2 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 \\ -1 \\ 7 \end{bmatrix}$$

$A \quad X = b$

$$[A|b] = \begin{pmatrix} 2 & 1 & 1 & 1 \\ 6 & 2 & 1 & -1 \\ -2 & 2 & 1 & 7 \end{pmatrix}$$

$$R_2 \rightarrow R_2 - 3R_1, R_3 \rightarrow R_3 + R_1$$

$$\begin{pmatrix} 2 & 1 & 1 & 1 \\ 0 & -1 & -2 & -4 \\ 0 & 3 & 2 & 8 \end{pmatrix}$$



$$\begin{pmatrix} 2 & 1 & 1 & 1 \\ 0 & -1 & -2 & -4 \\ 0 & 3 & 2 & 8 \end{pmatrix}$$

$$R_3 \rightarrow R_3 + 3R_2$$

$$\begin{pmatrix} 2 & 1 & 1 & 1 \\ 0 & -1 & -2 & -4 \\ 0 & 0 & -4 & -4 \end{pmatrix}$$

$$-4z = -4$$

$$z = 1$$

$$-y - 2z = -4$$

$$-y - 2 = -4$$

$$y = 2$$

$$2x + y + z = 1$$

$$2x + 2 + 1 = 1$$

$$x = -1$$

$$\begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} -1 \\ 2 \\ 1 \end{pmatrix}$$



Pivots elements

$$\begin{pmatrix} 2 & 1 & 1 & 1 \\ 0 & -1 & -2 & -4 \\ 0 & 0 & -4 & -4 \end{pmatrix} \Rightarrow \text{this is in Row echelon form.}$$

Columns containing pivot elements are called

Basic Columns.

Columns not containing pivot elements are called

Non Basic Columns.

$$\text{Rank}(A) = 3$$

$$\text{Rank}(A|b) = 3$$



Solve the following system of linear equations

$$x_1 + 2x_2 + x_3 + 3x_4 + 3x_5 = 5$$

$$2x_1 + 4x_2 + 4x_4 + 4x_5 = 6$$

$$x_1 + 2x_2 + 3x_3 + 5x_4 + 5x_5 = 9$$

$$2x_1 + 4x_2 + 4x_4 + 7x_5 = 9$$

$$[A|b] = \begin{bmatrix} 1 & 2 & 1 & 3 & 3 & 5 \\ 2 & 4 & 0 & 4 & 4 & 6 \\ 1 & 2 & 3 & 5 & 5 & 9 \\ 2 & 4 & 0 & 4 & 7 & 9 \end{bmatrix}$$

$$R_2 \rightarrow R_2 - 2R_1, R_3 \rightarrow R_3 - R_1 \\ R_4 \rightarrow R_4 - 2R_1$$

$$\begin{bmatrix} 1 & 2 & 1 & 3 & 3 & 5 \\ 0 & 0 & -2 & -2 & -2 & -4 \\ 0 & 0 & 2 & 2 & 2 & 4 \\ 0 & 0 & -2 & -2 & 1 & -1 \end{bmatrix}$$

$$R_3 \rightarrow R_3 + R_2, R_4 \rightarrow R_4 - R_2$$

$$\begin{bmatrix} 1 & 2 & 1 & 3 & 3 & 5 \\ 0 & 0 & -2 & -2 & -2 & -4 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 3 & 3 \end{bmatrix}$$



$$\begin{pmatrix} 1 & 2 & 1 & 3 & 3 & 5 \\ 0 & 0 & -2 & -2 & -2 & -4 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 3 & 3 \end{pmatrix}$$

$R_3 \rightarrow R_4$

$$\begin{pmatrix} \textcircled{1} & 2 & 1 & 3 & 3 & 5 \\ 0 & 0 & \textcircled{-2} & -2 & -2 & -4 \\ 0 & 0 & 0 & 0 & \textcircled{3} & 3 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

$$3x_5 = 3 \Rightarrow x_5 = 1$$

$$-2x_3 - 2x_4 - 2x_5 = -4$$

$$-2x_3 - 2x_4 - 2 = -4$$

$$-2x_3 - 2x_4 = -2$$

$$x_3 + x_4 = 1$$

Non basic columns of A are 2 and 4
 $\Rightarrow x_2$ and x_4 are called free variables.

$$\text{Let } x_4 = 1 \Rightarrow x_3 + 1 = 1$$

$$\Rightarrow x_3 = 0$$

$$x_1 + 2x_2 + x_3 + 3x_4 + 3x_5 = 5$$

$$x_1 + 2x_2 + 0 + 3 + 3 = 5$$



$$x_1 + 2x_2 = -1$$

Let $x_2 = 0 \Rightarrow x_1 = -1$

$$\begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \end{pmatrix} = \begin{pmatrix} -1 \\ 0 \\ 0 \\ 1 \\ 1 \end{pmatrix}$$

Let us general solution
 $x_5 = 1$

$$x_3 + x_4 = 1$$

$$\text{Let } x_4 = k_1 \Rightarrow x_3 = 1 - k_1$$

$$x_1 + 2x_2 + x_3 + 3x_4 + 3x_5 = 5$$

$$x_1 + 2x_2 + 1 - k_1 + 3k_1 + 3 = 5$$

$$x_1 + 2x_2 = 1 - 2k_1$$

$$\text{Let } x_2 = k_2 \Rightarrow x_1 = 1 - 2k_1 - 2k_2$$



$$\begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \end{pmatrix} = \begin{pmatrix} 1 & -2K_1 & -2K_2 \\ & K_2 \\ & 1-K_1 \\ & K_1 \\ & 1 \end{pmatrix} = K_1 \begin{pmatrix} -2 \\ 0 \\ -1 \\ 1 \\ 0 \end{pmatrix} + K_2 \begin{pmatrix} -2 \\ 1 \\ 0 \\ 0 \\ 0 \end{pmatrix} + \begin{pmatrix} 1 \\ 0 \\ 1 \\ 0 \\ 1 \end{pmatrix}$$

$K_1 = K_2 = 0$

$X = \begin{pmatrix} 1 \\ 0 \\ 1 \\ 0 \\ 1 \end{pmatrix},$

$K_1 = 1 \quad K_2 = 2$

X

The system has infinitely many solutions

Aditya Vangala



$$\begin{pmatrix} 1 & 2 & 1 & 3 & 3 & 5 \\ 0 & 0 & -2 & -2 & -2 & -4 \\ 0 & 0 & 0 & 0 & 3 & 3 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$



Solve the following system of linear equations

$$x_1 + 2x_2 + x_3 + 3x_4 + 3x_5 = 0$$

$$2x_1 + 4x_2 + 4x_4 + 4x_5 = 0$$

$$x_1 + 2x_2 + 3x_3 + 5x_4 + 5x_5 = 0$$

$$2x_1 + 4x_2 + 4x_4 + 7x_5 = 0$$

$$\begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \end{pmatrix} = k_1 \begin{pmatrix} -2 \\ 0 \\ -1 \\ 1 \\ 0 \end{pmatrix} + k_2 \begin{pmatrix} -2 \\ 1 \\ 0 \\ 0 \\ 0 \end{pmatrix} + \cancel{\begin{pmatrix} 1 \\ 0 \\ 1 \\ 0 \\ 1 \end{pmatrix}}$$

$$\begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \end{pmatrix} = k_1 \begin{pmatrix} -2 \\ 0 \\ -1 \\ 1 \\ 0 \end{pmatrix} + k_2 \begin{pmatrix} -2 \\ 1 \\ 0 \\ 0 \\ 0 \end{pmatrix}$$

① $k_1 = k_2 = 0$ $X = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}$ zero solution

② $k_1 = 1$ $k_2 = 1$ $X = \begin{pmatrix} -4 \\ 1 \\ -1 \\ -1 \\ 0 \end{pmatrix}$



Summary:

Ex 1

$$\begin{pmatrix} 2 & 1 & 1 & 1 \\ 0 & -1 & -2 & -4 \\ 0 & 0 & -4 & -4 \end{pmatrix}$$

0 0 0 x

$x \neq 0$

Number of variables = $n = 3$

Rank(A) = 3 $R(A|b) = 3$

Rank(A) = $r = 3$

Number of free variables = $n - r = 3 - 3 = 0$

Unique solution

b is non basic column



Ex 2

$$\begin{pmatrix} 1 & 2 & 1 & 3 & 3 & 5 \\ 0 & 0 & -2 & -2 & -2 & -4 \\ 0 & 0 & 0 & 0 & 3 & 3 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

Number of variables = $n = 5$

Rank(A) = 3 Rank(A|b) = 3 Rank(A) = $r = 3$

Number of free variables = $n - r = 5 - 3 = 2$

Infinitely many solutions.

b is non basic column

$$0 \ 0 \ 0 \ 0 \ 0 \ \alpha$$

$$\alpha \neq 0$$

Ex 3

$$\begin{pmatrix} 1 & 2 & 1 & 3 & 3 & 5 \\ 0 & 0 & -2 & -2 & -2 & -4 \\ 0 & 0 & 0 & 0 & 3 & 3 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

$$n = 5$$

$$\text{Rank}(A) = 3, \quad \text{Rank}(A|b) = 4$$

$$\text{Rank}(A) \neq \text{Rank}(A|b)$$

$$0x_1 + 0x_2 + 0x_3 + 0x_4 + 0x_5 \neq 1$$

NO solution

b is a basic
column

$$00000x \\ x \neq 0$$



m linear equations in n unknowns is given by

$$a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n = b_1 \quad E_1$$

$$a_{21}x_1 + a_{22}x_2 + \dots + a_{2n}x_n = b_2 \quad E_2$$

$$\vdots$$

$$a_{m1}x_1 + a_{m2}x_2 + \dots + a_{mn}x_n = b_m \quad E_m$$

$$A_{m \times n} X_{n \times 1} = b_{m \times 1}$$

The above equations in matrix form can be written as **$Ax = b$** .

$$A = \begin{pmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{pmatrix} \quad X = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} \quad b = \begin{pmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{pmatrix}$$

$A_{m \times n}$



Let $S = \begin{Bmatrix} E_1 \\ E_2 \\ \vdots \\ E_m \end{Bmatrix}$ be the linear system with m equations and n unknowns.

For a linear system S , each of the following three elementary operations results in an equivalent system S' .

(1) Interchange the i^{th} and j^{th} equations. That is, if

Type I

$$S = \begin{Bmatrix} E_1 \\ E_2 \\ \vdots \\ E_i \\ E_j \\ \vdots \\ E_m \end{Bmatrix}$$

$$S' = \begin{Bmatrix} E_1 \\ E_2 \\ \vdots \\ E_j \\ E_i \\ \vdots \\ E_m \end{Bmatrix}$$



(2) Replace the i^{th} equation by a nonzero multiple of itself. That is

$$S = \begin{bmatrix} E_1 \\ E_2 \\ \vdots \\ E_i \\ E_j \\ \vdots \\ E_m \end{bmatrix}$$

$$S' = \begin{bmatrix} E_1 \\ E_2 \\ \vdots \\ \alpha E_i \\ E_j \\ \vdots \\ E_m \end{bmatrix}$$

$$\alpha \neq 0$$

Type II



(3) Replace the j^{th} equation by a combination of itself plus a multiple of the i^{th} equation. That is,

$$S = \begin{Bmatrix} E_1 \\ E_2 \\ \vdots \\ E_i \\ E_j \\ \vdots \\ E_m \end{Bmatrix} \quad S' = \begin{Bmatrix} E_1 \\ E_2 \\ \vdots \\ E_i \\ E_j + \alpha E_i \\ \vdots \\ E_m \end{Bmatrix} \quad \text{Type III}$$



A typical structure for a matrix in row echelon form is illustrated below with the pivots circled.

$$\begin{bmatrix} \textcircled{*} & * & * & * & * & * & * & * \\ 0 & 0 & \textcircled{*} & * & * & * & * & * \\ 0 & 0 & 0 & \textcircled{*} & * & * & * & * \\ 0 & 0 & 0 & 0 & 0 & 0 & \textcircled{*} & * \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$



What is a Consistent System?

A system which has at least one solution is called Consistent system

Inconsistent System: A system with no solution is called Inconsistent system.



Consistency

Each of the following is equivalent to saying that $[\mathbf{A}|\mathbf{b}]$ is consistent.

- ✓ ▶ In row reducing $[\mathbf{A}|\mathbf{b}]$, a row of the following form never appears:

$$(0 \ 0 \ \dots \ 0 | \alpha) \quad \text{where } \alpha \neq 0$$

- ▶ \mathbf{b} is a nonbasic column in $[\mathbf{A}|\mathbf{b}]$.
- ▶ $\text{rank}[\mathbf{A}|\mathbf{b}] = \text{rank}(\mathbf{A})$.
- ▶ \mathbf{b} is a combination of the columns in \mathbf{A} .

$$0x_1 + 0x_2 + \dots + 0x_n \neq \alpha$$



b is a combination of the columns in **A**.

$$2x + y + z = 1$$

$$6x + 2y + z = -1$$

$$-2x + 2y + z = 7$$

$$-1 \begin{pmatrix} 2 \\ 6 \\ -2 \end{pmatrix} + 2 \begin{pmatrix} 1 \\ 2 \\ 2 \end{pmatrix} + 1 \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 1 \\ -1 \\ 7 \end{pmatrix}$$