Euclidean Condensate Theory: A Narrative Companion

From the Φ-Medium to Spacetime, Gravity,
Quantum Mechanics, and Observational Predictions

Valeriy Blagovidov*

April 2026

Abstract

This article is a comprehensive narrative companion to the technical preprint *"Euclidean Condensate Theory (ECT): Emergence of Spacetime, Quantum Mechanics, and Gravity from Spontaneous $O(4)$ Symmetry Breaking."* It covers all major topics of the preprint — from the fundamental Φ-medium and its postulates, through the emergence of Lorentzian spacetime, gravity, cosmology, galactic dynamics, quantum mechanics, vacuum effects, decoherence, black holes, and observational predictions — using primarily narrative explanations with only the essential formulas retained. No intermediate mathematical derivations are included; instead, the logical flow, physical reasoning, key results, and their significance are presented in a form accessible to physicists and advanced students who wish to understand the architecture and scope of ECT without working through the full technical apparatus. Every result is classified by its derivation level (A/B/C/Open), as in the technical preprint.

Companion to: *"Euclidean Condensate Theory (ECT): Emergence of Spacetime, Quantum Mechanics, and Gravity from Spontaneous $O(4)$ Symmetry Breaking"* (doi:10.5281/zenodo.18917929).

Keywords: Euclidean Condensate Theory; emergent spacetime; emergent gravity; emergent time; spontaneous symmetry breaking; $O(4) \to O(3)$; ordered condensate; quantum foundations; decoherence; galaxy rotation curves; baryonic Tully–Fisher relation; Hubble tension; dark energy; dark matter phenomenology; black-hole information problem; Principle of Euclidean Stationarity; fine-structure constant; lensing–growth sector; gravitational slip; Pauli exclusion; holographic entropy bound; area-law scaling; entanglement entropy; Born rule; neutrino masses; seesaw mechanism; neutrinoless double-beta decay; vacuum stability; false vacuum; Coleman vacuum decay; electroweak metastability; RG running; Big Bang singularity; pre-Lorentzian regime; cosmological ordering transition

*Correspondence: `blagovidov@phystech.edu`, `vblagovidov@gmail.com`. ORCID: 0009-0008-6707-7068.

Contents

Introduction **6**

I Foundations: From the Φ-Medium to Emergent Laws **6**

1 Motivation: why question the foundations? **6**

2 The Φ-medium: what it is and what it is not **7**

3 The six postulates: formalising the Φ-medium **8**

4 Spontaneous symmetry breaking and the emergence of time **10**

5 How the Lorentzian signature emerges **12**

6 The equation hierarchy: from Euclidean boundary problem to Schrödinger **13**

7 Causal cone inheritance and what GW170817 tells us **14**

8 Three fundamental scales from one field **15**

9 ECT loops, quantisation, and the origin of $\hbar$ **16**

10 The arrow of time, the second law, and why time flows **17**

11 Spectrum, branching, gauge symmetries, and particles **18**

11.1 Gauge symmetries and the interaction sectors 19

11.2 Elementary particles 20

12 Conclusions of Part I **21**

12.1 Principal results 21

12.2 Key equations and principles of Part I 22

12.3 Predictions originating in Part I 22

12.4 Part I derivation logic 23

II Macroscopic Physics: Gravity, Cosmology, and Observational Tests **24**

13 Special Relativity as the homogeneous limit **24**

13.1 The universal speed bound 24

13.2 Signal speed, not photon speed 24

14 Emergent gravity and the generalised Einstein equations **25**

15 Condensate evolution and cosmic history **26**

16 The four classical cosmological problems **27**

16.1 The horizon problem 27

16.2 The flatness problem 27

16.3 The relic and monopole problem 27

16.4 The primordial perturbation problem 28

16.5 Summary 28

17 The dark sector in ECT **28**

17.1 Dark matter: condensate response, not particles 28

17.2 Dark energy: condensate residual energy, not a cosmological constant 29

17.3 ECT reinterpretation of the dark sector 29

18 The Hubble tension **30**

19 The JWST early-galaxy problem **30**

20 Galactic rotation curves **31**

21 The baryonic Tully–Fisher relation and the RAR **33**

22 Cluster-scale gravitational lensing **35**

23 Matter–antimatter asymmetry **37**

24 Conclusions of Part II **37**

24.1 Principal results 37

24.2 Key equations and principles of Part II 38

24.3 Observational tests and falsifiers 38

24.4 Part II derivation logic 39

III Quantum Sector: Coherent Branch **39**

25 ECT versus standard quantum mechanics **40**

25.1 Detailed structural comparison: ECT vs standard QM 41

26 Hilbert-space structure from reflection positivity **43**

27 Spin as a spinorial orientation mode **43**

28 Superposition, interference, exchange statistics, and the Pauli exclusion principle **44**

29 The Principle of Euclidean Stationarity **45**

30 Probability, the Born rule, and measurement **46**

31 Quantum entanglement and Bell correlations **47**

32 Tunnelling **47**

33 Interactions belong to the Lorentzian branch **48**

34 Influence of future boundary conditions **48**

35 Quantum vacuum effects **48**

36 Decoherence and the quantum–classical boundary **49**

37 Quantum gravity in ECT **50**

38 Black-hole thermodynamics and the information problem **51**

39 Why entropy scales with area: the holographic bound as a medium property **52**

40 Why electric charge is quantised **54**

41 Why the post-Big-Bang Universe developed the way it did **55**

42 The grandfather paradox **56**

43 Analogue laboratory programme **56**

44 Conclusions of Part III **57**

44.1 Principal results 57

44.2 Key equations and principles of Part III 57

44.3 Part III derivation logic 58

IV Summary, Comparisons, and Outlook **59**

45 Architectural comparison with neighbouring programmes **59**

46 Distinctive features of the framework **59**

47 Standard postulates reinterpreted as ECT consequences **60**

48 Key equations, laws, and principles of ECT **63**

49 Falsifiable predictions **63**

50 Colour completion: a consistency requirement **65**

51 Open problems **66**

52 Complete derivation map **69**

53 Open problems of physics addressed within ECT **69**

54 Conclusion **72**

Introduction

This article is a concise narrative summary of the technical preprint *"Euclidean Condensate Theory (ECT): Emergence of Spacetime, Quantum Mechanics, and Gravity from Spontaneous $O(4)$ Symmetry Breaking"* (doi:10.5281/zenodo.18917929). ECT is a theoretical framework in which Lorentzian spacetime, quantum mechanics, and gravity emerge from spontaneous $O(4) \to O(3)$ symmetry breaking of a single scalar condensate on a four-dimensional Euclidean manifold. The present article covers the same material in a narrative form accessible to a broader audience, retaining only the essential formulas and key results. For all derivations, proofs, and technical details, the reader is referred to the full preprint.

ECT is organised into three principal Parts. Part I (Foundations) introduces the Φ-medium, its six postulates, the mechanism of gradient condensation, and the emergence of Lorentzian signature, wave equations, gauge symmetries, the arrow of time, and quantisation. Part II (Macroscopic Physics) develops emergent gravity, cosmology, galactic rotation curves, cluster lensing, the dark sector, and observational predictions. Part III (Quantum Sector) develops the coherent branch: Hilbert-space reconstruction, the Principle of Euclidean Stationarity, entanglement, tunnelling, decoherence, quantum gravity, and the black-hole information problem. Part IV (Summary) consolidates comparisons, key equations, predictions, and open problems.

Throughout this article, each result is classified by its derivation level: Level A (structural consequence of the postulates), Level B (requiring additional closure assumptions), or Open (stated explicitly as an unsolved problem). This classification follows the technical preprint exactly.

Derivation levels used in this article:
Level A — structural consequence of the postulates P1–P6 and BR1; no additional assumptions required.
Level B — requires stated closure assumptions, effective approximations, or matching identifications.
Open — stated explicitly as an unsolved problem or a programme target.

Part I

Foundations: From the Φ-Medium to Emergent Laws

1. Motivation: why question the foundations?

Modern physics rests on two extraordinarily successful pillars: General Relativity, which describes gravity as the curvature of a smooth four-dimensional spacetime, and quantum field theory, which describes the electromagnetic, weak, and strong interactions as the exchange of quanta on a fixed spacetime background. Despite their remarkable empirical success, these two frameworks remain fundamentally incompatible. Quantising gravity in the standard way produces uncontrollable infinities; treating spacetime classically while matter is quantum leads to conceptual paradoxes.

Both theories share one curious and rarely questioned assumption: the distinction between space

and time is *fundamental*. The spacetime interval

$$ds^2 = -c^2 dt^2 + dx^2 + dy^2 + dz^2 \tag{1}$$

contains one coordinate with a minus sign and three with a plus. This Lorentzian signature $(-, +, +, +)$ is the origin of causality, light cones, and the fact that time flows irreversibly while space does not. But *why* does this minus sign exist?

Neither General Relativity nor the Standard Model answers this question. They assume it. String theory inherits it. Loop quantum gravity inherits it. Even causal set theory, which tries to build spacetime from scratch, puts the causal structure in by hand. The minus sign is one of the most rarely questioned assumptions in physics.

There is a deep theoretical reason to suspect that the minus sign is not fundamental. The Euclidean signature $(+, +, +, +)$ possesses a higher symmetry — the rotation group $O(4)$, with $4 \times 3/2 = 6$ independent rotation generators — compared to the Lorentzian $O(3, 1)$, which has only three rotations and three boosts. Throughout physics, the fundamental laws have *more* symmetry than the observed states: a ferromagnet's Hamiltonian is rotationally symmetric, but the magnetised state picks a direction; the Higgs potential is gauge-symmetric, but the vacuum selects a specific phase. It is natural to ask whether spacetime itself follows the same pattern: a higher-symmetry Euclidean phase, spontaneously broken to produce the Lorentzian world we inhabit.

Furthermore, the path-integral formulation of quantum mechanics is naturally connected to a Euclidean description through the analytic continuation $t \to -i\tau$. Many of the most powerful computational tools in quantum field theory — instantons, lattice gauge theory, non-perturbative renormalisation — operate in Euclidean space. This suggests that the Euclidean description may be more fundamental than a mere computational trick.

Euclidean Condensate Theory (ECT) explores precisely this possibility. It proposes that the deepest level of reality is Euclidean, and that the Lorentzian spacetime we observe is an emergent property of an ordered phase of a more symmetric vacuum. If correct, this single idea has sweeping consequences: not only the Lorentzian metric, but gravity, quantum mechanics, the arrow of time, the second law of thermodynamics, the speed of light, Planck's constant, Newton's constant, and even the structure of elementary particles could all be *derived* from properties of the Φ-condensate rather than postulated independently.

2. The Φ-medium: what it is and what it is not

At the foundation of ECT lies a single real scalar field Φ, defined on a four-dimensional Euclidean manifold $\mathcal{M}^4$. This manifold has no built-in time: all four coordinates $X^A = (x, y, z, w)$ enter with the same positive-definite Euclidean metric δ_{AB}. There is no preferred direction, no causal structure, no light cone.

The field Φ is the sole fundamental degree of freedom. Its dynamics is governed by an $O(4)$-invariant action with a self-interaction potential — a double-well potential familiar from the Higgs mechanism and superfluidity. In many ways Φ is analogous to the order parameter of a superfluid or ferromagnet, but operating at a much deeper level: it describes the medium from which spacetime itself is made.

It is essential to understand what Φ is *not*. It is not a quantum field defined on a pre-existing spacetime — that would presuppose the very thing ECT aims to derive. It is not a metric or gravitational field; those will emerge later as collective properties. It is not an inflaton, Higgs field, or dark matter particle, though some of its modes will play roles analogous to these at specific energy scales. And it is not a "hidden variable" in the quantum-mechanical sense. Φ is a *pre-spacetime, pre-quantum, pre-gravitational* entity.

A subtle but critical point: the "physics" of Φ at the fundamental Euclidean level cannot be described using energy, force, or potential in their usual sense. Those concepts presuppose Lorentzian time, which does not yet exist. The Euclidean action $S_E[\Phi]$ is a functional of field configurations on the four-dimensional manifold — not a time integral of a Lagrangian. Importing energy language into the pre-time Euclidean domain is a common source of confusion and must be avoided.

The Φ-medium has specific ontological properties, explicitly stated in the technical preprint (S1–S10): it is continuous and differentiable (S1); it is scalar — no internal spin or index structure at the fundamental level (S2); it fills the entire four-dimensional Euclidean space (S3); its dynamics is purely relational — no external clock, no preferred reference frame (S4); it admits ordered and disordered configurations (S5); no single configuration is privileged a priori (S6); it supports both smooth and singular (defect) configurations (S7); pointwise internal coordinates are descriptive redundancies, not observables (S8); ordering need not be globally uniform (S9); all topological sectors compatible with the vacuum manifold are admissible (S10). These properties characterise a specific type of physical medium whose mathematical formalisation leads directly to the postulates.

3. The six postulates: formalising the Φ-medium

The postulates of ECT formalise the ontological properties of the Φ-medium described above. Each postulate captures a specific physical property and translates it into a precise mathematical statement.

P1 — Four-dimensional Euclidean arena (formalises S1, S3). The fundamental arena is a connected four-dimensional Euclidean manifold $\mathcal{M}^4$ with metric δ_{AB}. All four directions and all points are initially equivalent. There is no distinguished time coordinate at the fundamental level. This is the maximal symmetry of a four-dimensional space with positive-definite metric.

P2 — $O(4)$-invariant action (formalises the symmetry of S3, S4). The dynamics respects the full rotational symmetry of the four-dimensional Euclidean space: $S_E[\Phi] = S_E[\Lambda \cdot \Phi]$ for all $\Lambda \in O(4)$. This is the mathematical expression of the equivalence of all four directions in the unbroken phase.

P3 — Minimal scalar microdynamics (formalises S1, S2). The fundamental field is a single real scalar $\Phi(X)$ with a minimal local action:

$$S_E[\Phi] = \int d^4X \left\{ \frac{1}{2}(\partial_A \Phi)^2 + V(\Phi) \right\}, \qquad V(\Phi) = -\frac{\mu^2}{2}\Phi^2 + \frac{\lambda}{4}\Phi^4, \quad \mu^2 > 0. \tag{2}$$

The double-well potential has two minima at $\pm\phi_0$, where $\phi_0 = \sqrt{\mu^2/\lambda}$. Crucially, the orientation variable n_A does not appear in this bare action — it arises only after vacuum ordering.

P4 — Ordered vacuum branch (formalises S5, S9). The physically relevant vacuum belongs

to an ordered branch characterised by a nonzero *gradient condensate*: $\langle \partial_A \Phi \rangle = u_0\, \delta_{Aw}$, $u_0 > 0$. This spontaneously breaks $O(4) \to O(3)$, selecting a preferred Euclidean direction w. The direction is not imposed externally but selected spontaneously — just as a ferromagnet selects a magnetisation direction below the Curie temperature.

P5 — Medium character of the ordered branch (formalises S8). Pointwise internal coordinates of the effective order parameter — specifically the compact phase $\theta(X)$ and the local orientation $n_A(X)$ — are descriptive redundancies rather than independently observable degrees of freedom. Physical observables depend only on *gradient* and *holonomy* (loop-integral) data.

This is directly analogous to the physics of known media: in a superfluid, the absolute phase at a single point is not observable — only phase gradients and circulation integrals carry physical information. In a nematic liquid crystal, the pointwise director angle is unobservable — only director gradients and topological defects are physical. P5 asserts that the Φ-medium has this same continuum-medium character.

The consequence of P5 is profound: since pointwise internal coordinates are redundancies, local redefinitions of $\theta(X)$ and $n_A(X)$ must be treated as local gauge redundancies. This requires compensating gauge connections and leads to the emergence of gauge symmetries (Section 11.1).

P6 — Multiplicity of configurations and non-uniform ordering (formalises S6, S9, S10). No single configuration is singled out a priori. Ordered configurations need not be globally uniform: spatially varying condensate backgrounds, domain structures, and transition regions are all admissible.

In addition to these six postulates, the theory identifies a derived structural rule:

BR1 — Smooth coherent sector (derived from P3 + P4). In the smooth infrared sector of the ordered branch, the effective order parameter admits the polar form $\Phi_{\text{eff}} = \rho\, e^{i\theta}$, and single-valuedness of the phase on closed loops gives the winding condition: $\oint \partial_A \theta\, dX^A = 2\pi n$, $n \in \mathbb{Z}$. This is not a separate postulate but a *topological consequence* of smooth ordering. It introduces the pre-quantum action scale S_0 and the loop-sector structure that will later give rise to quantum behaviour (Section 9).

What is *not* postulated deserves emphasis. ECT does not postulate: external time; Lorentzian signature; a Hilbert space; the Born rule; the Schrödinger equation; Newton's gravitational equation; a cosmological constant; dark matter particles; the speed of light as a fundamental constant. All of these are either derived or structurally motivated within the framework of P1–P6 + BR1.

4. Spontaneous symmetry breaking and the emergence of time



Figure 1: Architectural overview of ECT. A single Euclidean Φ-field undergoes $O(4) \to O(3)$ symmetry breaking through gradient condensation. Two complementary branches emerge: the geometric branch (gravity, cosmology, galactic dynamics) and the coherent branch (quantum mechanics, decoherence, vacuum effects). All of known physics is organised within this single structure. (*Level A for the branching architecture; individual arrows carry their own status labels in the preprint.*)

Spontaneous symmetry breaking (SSB) is one of the most powerful organising principles in physics. A spherical magnet has no preferred direction, until it cools below the Curie temperature and all spins align. A perfectly round ball on the tip of a Mexican hat has no preferred position, until it rolls down and selects one valley. In both cases, the underlying laws are symmetric, but the realised state is not.

In ECT, the Φ-medium undergoes an analogous transition: the $O(4)$ symmetry of the four-dimensional Euclidean arena is broken to $O(3)$ (Fig. 1). Three directions remain equivalent (they

become "space"); one (w) is singled out (it becomes the seed of "time").

This transition is not merely a technical symmetry breaking. It is the event that makes physics in the familiar sense possible at all. Before ordering, the Euclidean condensate distinguishes no time, no causal order, no preferred class of propagating excitations. After ordering, all of the following arise simultaneously: the causal cone, Lorentzian kinematics, the excitation spectrum, and the decoherence hierarchy. Time is not added to an already prepared physics; it arises together with it as a property of the ordered phase. In this sense the $O(4) \to O(3)$ transition plays the role of a "crystallisation of the physical world."

The mechanism behind this transition is *gradient condensation.* In the ordered phase, not the field Φ itself but its *gradient* $\partial_A \Phi$ acquires a nonzero expectation value pointing in a specific direction: $\langle \partial_A \Phi \rangle = u_0\, \delta_{Aw}$. This distinction is crucial: a nonzero $\langle \Phi \rangle$ would be $O(4)$-invariant (since Φ is a scalar under rotations, a scalar VEV preserves all rotational symmetries) and would produce no symmetry breaking. The gradient $\partial_A \Phi$, by contrast, transforms as a *vector* under $O(4)$. Its nonzero expectation value breaks the symmetry by selecting a direction — just as the magnetisation $\langle \mathbf{M} \rangle$ in a ferromagnet is a vector that picks a direction and breaks $O(3) \to O(2)$.

The gradient field can be decomposed into an amplitude and a direction:

$$\partial_A \Phi = u \cdot n_A, \qquad u = |\partial \Phi| \geq 0, \qquad n^A n_A = 1. \tag{3}$$

In the ordered vacuum, $\langle u \rangle = u_0$ and $\langle n^A \rangle = \delta^A{}_w$. This decomposition is the key to the entire theory. Each component plays a distinct physical role:

The **amplitude mode** (u; fluctuations $\sigma = u - u_0$): massive, short-ranged ($\xi_{\text{cond}} \sim m_\sigma^{-1} \sim \ell_{\text{Pl}}$), connected to gravitational stiffness and the Planck scale. It plays a role analogous to the Higgs boson, but at the Planck scale rather than the electroweak scale.

The **orientation modes** (δn^A): the symmetry breaking $O(4)/O(3) \simeq S^3$ naively admits three Goldstone directions, but integrability of the scalar-only basis reduces the propagating soft content to a single surviving longitudinal scalar mode χ (see below). At the linearised level, the orientation sector generates the graviton (spin-2 mode).

The **phase mode** (θ, in the coherent sector $\Phi_{\text{eff}} = \rho\, e^{i\theta}$): connected to gauge fields and quantum dynamics.

The dynamical origin of gradient condensation from the bare action P3 is an open theoretical programme (OP-grad in the preprint). A generalised kinetic action $P(X)$, $X = (\partial_A \Phi)^2$, with a minimum at $X = u_*^2 > 0$, provides the natural dynamical setting for such condensation — analogous to how a superfluid develops nonzero superflow when the free energy favours a finite condensate velocity. The theory establishes that *given* the ordered branch (P4), a rich and self-consistent effective physics follows.

An important structural result (the "$P(X)$ no-orientation theorem", proved in the preprint) shows that a purely kinetic action $P(X)$ can generate gradient condensation (nonzero u_0) but *cannot* generate orientation stiffness — the rigidity that resists bending of the condensate direction. Orientation stiffness must come from higher-order terms or from the NLO sector of the EFT, and it is this stiffness that ultimately gives rise to gravity.

5. How the Lorentzian signature emerges

Once the condensate has selected a direction, the physics of small excitations around the ordered background changes profoundly. The key quantity is the effective kinetic tensor K^{AB} governing fluctuations. The technical preprint proves a uniqueness theorem: the most general quadratic action for fluctuations φ around the ordered branch, compatible with the residual $O(3)$ symmetry, takes the form (*Level A*):

$$S_{\text{eff}} = \int d^4X \left[\frac{\beta}{2}(\partial_A\varphi)^2 - \frac{\alpha}{2}(n^A\partial_A\varphi)^2 + \dots \right], \tag{4}$$

so that $K^{AB} = \beta\,\delta^{AB} - \alpha\, n^A n^B$.

This tensor has eigenvalues β (triply degenerate, along the three directions perpendicular to n^A) and $\beta - \alpha$ (along n^A). An eigenvalue analysis reveals three distinct regimes:

If $\alpha < \beta$: All eigenvalues are positive. The effective metric remains Euclidean. No time, no causal structure, no wave propagation.

If $\alpha = \beta$: One eigenvalue vanishes. The system is at the Euclidean–Lorentzian boundary, with degenerate, Galilean-like kinematics.

If $\alpha > \beta$ (the "Lorentzian window"): One eigenvalue is negative. The effective metric acquires a Lorentzian signature $(-,+,+,+)$. Perturbations propagate as waves with a finite speed, and a causal cone structure emerges.

ECT operates in the Lorentzian window. In the canonical benchmark $(\alpha, \beta) = (2, 1)$, the kinetic tensor reduces to $K^{AB} = \delta^{AB} - 2\, n^A n^B$, with eigenvalues $(+1, +1, +1, -1)$ — exactly the Lorentzian signature.

The effective propagation speed is:

$$c_* = \sqrt{\frac{\beta}{\alpha - \beta}}, \tag{5}$$

identified with the speed of light (*Level A/B*).

This is the central result of Part I: the minus sign in the spacetime interval (1) **is not assumed but derived.** It appears because the gradient condensate breaks $O(4)$ in a way that produces an anisotropic kinetic tensor with one negative eigenvalue. The derivation status is *Level A*: it follows from the uniqueness of the kinetic tensor (proved as a theorem) combined with P4.

Comparison with other theories. In Einstein–aether theory (Jacobson, 2004), the anisotropy is introduced by a *postulated* vector field — not derived from a scalar. In Hořava–Lifshitz gravity, Lorentz invariance is broken *explicitly* at the fundamental level. In Causal Dynamical Triangulations (CDT), the Lorentzian structure is imposed as a constraint on the lattice. In the Hartle–Hawking no-boundary proposal, the Euclidean arena is a computational tool, not a fundamental entity. In ECT alone, the Lorentzian signature is *spontaneously generated* from a higher-symmetry Euclidean substrate, and the result is a theorem, not a phenomenological assumption.

6. The equation hierarchy: from Euclidean boundary problem to Schrödinger



Figure 2: The ECT equation hierarchy. The fundamental Euclidean field equation (a four-dimensional boundary-value problem) generates the Klein–Gordon equation (after $t = w/c_*$), which in turn yields the Schrödinger equation (non-relativistic limit). Each level is a *descendant* of the previous one, not an independent postulate. Standard quantum mechanics occupies the lowest rung. (*Hierarchy structure: Level A; specific forms: Level A/B.*)

The emergence of Lorentzian signature has a direct consequence for the equations of motion (Fig. 2). The Euclidean field equation $\delta^{AB}\partial_A\partial_B\Phi - V'(\Phi) = 0$ is a four-dimensional *boundary-value problem*: its solutions are determined by data on the *entire boundary* of the Euclidean domain, not by initial data propagated forward in time.

Once the ordered branch is established and we introduce the identification $t = w/c_*$, the same equation, restricted to the Lorentzian window, becomes the *Klein–Gordon equation*: $\partial_t^2\varphi - c_*^2\nabla^2\varphi + m_{\text{eff}}^2\varphi = 0$ (*Level A/B*).

In the non-relativistic limit ($E \approx mc^2 + E_{\text{kin}}$, $|E_{\text{kin}}| \ll mc^2$), the Klein–Gordon equation reduces to the *Schrödinger equation*: $i\hbar\,\partial_t\psi = -\frac{\hbar^2}{2m}\nabla^2\psi + V\psi$ (*Level A/B*).

This hierarchy is not merely a formal observation. It encodes a physical claim: the Schrödinger equation, which standard quantum mechanics takes as its starting axiom, is the *narrowest, lowest-energy descendant* of a more fundamental Euclidean dynamics. The wave equation is *derived*, not postulated.

The distinction between the Cauchy (initial-value) problem of standard QM and the boundary-value problem of ECT is one of the deepest structural differences between the two frameworks. In standard QM, initial data on one time-slice determine the future uniquely. In ECT, the fundamental equation constrains the field configuration on the *entire* four-dimensional domain. This has far-reaching consequences for entanglement, delayed-choice experiments, and the

influence of future boundary conditions on present configurations (see Part III).

7. Causal cone inheritance and what GW170817 tells us

A critical question for any theory claiming to derive Lorentzian spacetime is: why is there a *universal* speed of light? In the real world, photons, gravitons, and all massless particles share the same causal cone to extraordinary precision. A decisive test of any emergent-spacetime framework is whether different low-energy sectors inherit a single causal structure rather than sector-dependent propagation cones—which would destroy universal relativistic kinematics. In ECT the universality of the causal cone is not postulated but inherited from the single ordered condensate background.

In August 2017, the LIGO/Virgo gravitational-wave detectors observed the merger of two neutron stars — an event designated **GW170817**. Just 1.7 seconds later (after a journey of 130 million light-years), the Fermi and INTEGRAL satellites detected a gamma-ray burst from the same source. This simultaneous arrival established that the speed of gravitational waves and the speed of light agree to better than one part in 10^{15}: $|c_{\mathrm{GW}}/c - 1| < 6 \times 10^{-15}$.

In many alternative gravity theories (Einstein–aether, Horndeski scalar-tensor models, massive gravity), such universality requires fine-tuning of multiple parameters. In the Standard Model Extension (SME), each particle sector carries an independent set of Lorentz-violation coefficients.

ECT explains this universality as a *structural theorem* (*Level A/B*). The argument is straightforward: (i) The fundamental field is one: Φ. (ii) The kinetic tensor is one: K^{AB}. (iii) All low-energy excitations — scalar, phase, gauge, tensor — propagate within the same ordered-branch EFT built on K^{AB}. (iv) No second independent rank-2 tensor exists in the EFT that could give different sectors different speeds.

The result is **cross-sector cone universality**:

$$c_{\mathrm{phase}} = c_\gamma = c_{\mathrm{GW}} = c_* \qquad (\textit{Level A/B}). \tag{6}$$

The GW170817 observation is then not an unexplained coincidence but a *structural consistency check* of ECT. This is one of the theory's most distinctive predictions: the universality of the speed of light is derived from the single-medium origin of all native low-energy sectors.

ECT does admit Planck-suppressed higher-order corrections that can produce tiny sector-dependent speed differences at extreme energies. But arbitrary, unsuppressed, low-energy speed differences between sectors are structurally forbidden. If such differences were ever observed, ECT's one-cone universality would be falsified.

8. Three fundamental scales from one field



Figure 3: The three energy scales of the ECT condensate. A single scalar field generates the Planck/gravitational scale ($\phi_0 \sim 10^{18}\,\mathrm{GeV}$), the electroweak scale ($v_2 \sim 246\,\mathrm{GeV}$), and the galactic-phenomenology scale ($v_{\mathrm{gal}} \sim$ kpc). These are not independent mechanisms but three faces of the same condensate at different organisational levels. (*Level B for the hierarchy mechanism.*)

A notable feature of ECT is that a single scalar field generates three distinct energy scales governing very different regimes of physics (Fig. 3):

The **gravitational scale** $\phi_0 \sim 2.4 \times 10^{18}\,\mathrm{GeV}$ (the reduced Planck mass) arises from the symmetry-breaking amplitude of the condensate. It simultaneously determines Newton's constant, Planck's constant, and the speed of light through the matching relations (*Level B*):

$$G_N = \frac{c_*^4}{8\pi\phi_0^2}, \qquad \hbar = \frac{c_*\phi_0}{\sqrt{\lambda}}, \qquad c_* = \sqrt{\frac{\beta}{\alpha-\beta}}. \tag{7}$$

The fact that a single condensate scale ϕ_0 simultaneously determines G, $\hbar$, and c is a non-trivial structural prediction: these three constants are not independent but are three projections of the same underlying condensate parameter. In standard physics, c, $\hbar$, and G appear as three independent empirical constants belonging to different chapters of physics: relativistic kinematics, quantum theory, and gravitation. ECT changes their logical status from external gifts of nature to observable projections of a single underlying medium: c_* expresses the kinematic anisotropy of the ordered phase, G_N expresses the geometric stiffness of the orientation sector, and $\hbar$ expresses the coherent self-consistency of the minimal loop.

The **electroweak scale** $v_2 \sim 246\,\mathrm{GeV}$ corresponds to a second, subsidiary condensation within the Φ-sector, generating masses for the W, Z bosons, and the Higgs particle. The enormous hierarchy $\phi_0/v_2 \sim 10^{16}$ is addressed through a running-coupling argument: the self-coupling λ runs logarithmically with energy and can generate the smaller scale through dimensional transmutation, similar to the mechanism that generates the QCD scale from the running of α_s (*Level B/Open*).

The **galactic scale** $r_0 \sim$ kpc marks the transition between Newtonian and modified gravitational behaviour in galaxies, governed by the critical acceleration $g^\dagger \approx cH_0/(2\pi) \approx 1.1 \times 10^{-10}\,\mathrm{m/s^2}$

(*Level B*).

Beyond the three primary scales, the same condensate pair (ϕ_0, v_2) naturally generates two distinguished intermediate combinations that appear in the neutrino sector: the geometric-mean scale $\sqrt{\phi_0\, v_2} \approx 2.4 \times 10^{10}\,\mathrm{GeV}$ (close to the scale of thermal leptogenesis and type-I seesaw), and the inverse-ratio Weinberg-operator scale $v_2^2/\phi_0 \approx 2.5 \times 10^{-5}\,\mathrm{eV}$ (within a few orders of magnitude of the observed neutrino mass splittings). Both combinations are the two simplest dimensionally consistent products that can be built from the same pair of fundamental ECT parameters; their joint appearance in the neutrino sector is a structural consequence of the three-scale hierarchy rather than an independent numerical coincidence. This further illustrates how the condensate architecture organises multiple observed energy scales from a single underlying medium (*Level B/C*).

9. ECT loops, quantisation, and the origin of $\hbar$

A distinctive feature of ECT is how quantisation emerges naturally from the compact-phase structure of the condensate, without being imposed as an axiom.

In the coherent sector, the effective order parameter has the polar form $\Phi_{\mathrm{eff}} = \rho\, e^{i\theta}$, where θ is a compact (periodic) phase variable. The single-valuedness requirement (BR1) means that if we transport θ around any closed loop $\mathcal{C}$ in the Euclidean manifold, it must return to its original value:

$$\oint_{\mathcal{C}} \partial_A \theta\, dX^A = 2\pi n, \qquad n \in \mathbb{Z}. \tag{8}$$

This is not a quantum axiom — it is a *topological necessity* for any smooth, single-valued function with compact range.

The consequence is immediate: the action associated with loop configurations is quantised. The minimal nontrivial loop ($n = 1$) carries a definite action:

$$S_{\mathrm{loop,min}} = 2\pi S_0, \tag{9}$$

where the scale S_0 depends on the condensate parameters. Through the matching relation (7), S_0 is identified with $\hbar$ (*Level B*).

This means Planck's constant is not a mysterious fundamental quantity in ECT. It is the action scale set by the condensate's compact-phase structure — just as the flux quantum $\Phi_0 = h/(2e)$ in a superconductor is set by the Cooper pair condensate, and the Bohr–Sommerfeld condition $\oint p\, dq = nh$ is set by the periodicity of classical action variables. All three are instances of the same mathematical mechanism: *single-valuedness of a periodic order parameter implies quantisation.*

The loop sectors partition the coherent branch into topologically distinct classes indexed by the integer n. Configurations in different sectors cannot be continuously deformed into each other. This is why quantum interference can arise in ECT without operator axioms: paths belonging to different winding sectors contribute to the path sum with different phase factors, producing constructive or destructive combination.

10. The arrow of time, the second law, and why time flows

In the Euclidean picture, the coordinate w is just one of four equivalent spatial directions. There is no "forward" or "backward" in w, just as there is no "forward" in x. The arrow of time is not fundamental: it is *emergent*.

ECT constructs the thermodynamic arrow in two stages:

Stage 1: directional backbone (*Level A*). In the Lorentzian branch, the effective dynamics admits a retarded causal structure: perturbations propagate forward along w inside the emergent light cone. The dissipative kernel of the influence-functional formalism is positive-semidefinite ($\Gamma_{\text{irr}} \geq 0$), meaning that off-diagonal histories are *suppressed* rather than amplified. This gives the *directionality* of decoherence.

Stage 2: entropy growth (*Level B*). Under two additional effective-closure assumptions — an ohmic spectral density and a Markovian regime — the coarse-grained entropy satisfies:

$$\frac{dS_{\text{ent}}}{dw} = N_{\text{eff}}\,\gamma \left(\frac{dq}{dw}\right)^2 \geq 0, \tag{10}$$

where N_{eff} is the effective number of environmental modes and γ is the decoherence rate per mode. The sign is anchored in the Level A retarded structure; the explicit formula requires Level B closure.

The result is a three-regime picture:

At the *cosmological* level, the ordered branch defines the cosmic arrow; the second law holds to overwhelming accuracy.

At the *macroscopic* level ($N_{\text{eff}} \gg 1$), the ratio of backward-to-forward transition probabilities is exponentially suppressed: $P_{\text{back}}/P_{\text{fwd}} = \exp(-2N_{\text{eff}}\gamma\,\Delta w \cdot (\Delta q)^2/\ell^2)$. For a mole of gas at room temperature over a nanosecond, the exponent is $\sim 3.6 \times 10^{33}$ — reverse trajectories are fantastically unlikely.

At the *quantum* level ($N_{\text{eff}} \sim 1$), the exponent becomes tiny ($\sim 10^{-15}$), and effective time-reversibility is recovered — consistent with the well-known fact that isolated quantum systems evolve reversibly.

The second law is therefore not an independent postulate in ECT. Its directional backbone is Level A; its full monotonicity is Level B. What ECT adds beyond Boltzmann is a *structural explanation* of why the initial condition is special: it is tied to ordered-branch selection (P4), not imposed ad hoc.

11. Spectrum, branching, gauge symmetries, and particles



Figure 4: The quantum–classical boundary. The dimensionless decoherence parameter $\Gamma_{\rm loop}$ determines whether a system behaves quantum-mechanically ($\Gamma_{\rm loop} \ll 1$) or classically ($\Gamma_{\rm loop} \gg 1$). The crossover at $\Gamma_{\rm loop} \sim 1$ defines the quantum–classical boundary. Different objects — electron, C_{60}, bacterium, dust grain, cat — occupy different positions on this scale. (*Level A for the crossover structure; Level B for numerical estimates.*)

The ordered condensate supports three classes of excitations, each playing a distinct physical role:

Amplitude mode (σ): Fluctuations of the gradient magnitude u around u_0. This mode is massive ($m_\sigma^2 = 2\mu^2$, *Level A*), with correlation length $\xi_{\rm cond} \sim m_\sigma^{-1}$ identified with the Planck length (*Level B*). It is heavy, short-ranged, and plays a role analogous to the radial Higgs mode at the Planck scale.

Orientation modes (δn^A): The symmetry breaking $O(4)/O(3)$ naively suggests three Goldstone bosons; however, the integrability constraint on the scalar-only ECT basis reduces the physically independent propagating soft content to a single surviving longitudinal scalar mode χ (massless at tree level). Their dynamics generates the graviton (spin-2, massless) at the linearised level (*Level A*), plus constrained scalar/vector companions. An important structural point is that the amplitude mode is too heavy ($m_\sigma \sim M_{\rm Pl}$) for kpc-scale variations, ruling out naive scalar-field dark matter models — a powerful constraint that uniquely selects the viable mechanism for galactic phenomenology (Part II).

Phase mode (θ): In the coherent sector, phase fluctuations connect to gauge fields: the photon emerges as the gauge boson of the $U(1)$ phase symmetry (*Level A/B after P5*).

The ordered condensate organises into two complementary branches (Fig. 4): the **geometric branch** (Macroscopic Physics, long-wavelength, coarse-grained, $\Gamma_{\rm loop} \gg 1$) and the **coherent branch** (Quantum Sector, short-wavelength, phase-coherent, $\Gamma_{\rm loop} \ll 1$). Both originate from the same ordered condensate action $S_{\rm ord}[u, n]$ (*Level A*). The quantum–classical boundary corresponds to the crossover $\Gamma_{\rm loop} \sim 1$.

11.1. Gauge symmetries and the interaction sectors

ECT does not postulate gauge symmetries. It derives them from the internal structure of the Φ-medium through P5. Since pointwise values of θ and n_A are not observable, local redefinitions must leave physics unchanged—requiring compensating gauge connections. The key insight is that these gauge connections are not added by hand: they are *forced* by the requirement that the effective description be independent of unphysical local choices.

Electromagnetism (*Level A*): The condensate carries a compact phase $\theta \sim \theta + 2\pi$. This periodicity defines a principal $U(1)$-bundle over the emergent spacetime: comparing the phase at neighbouring points requires a connection—the gauge field A_μ. Its dynamics (Maxwell's equations) follows as the leading gauge-invariant kinetic term in the low-energy effective theory. The photon is thus the gauge boson of an *inevitable* Abelian symmetry, derived from bundle covariance of the condensate phase fibre. Its precise identification within the electroweak chain ($U(1)_Y$ versus $U(1)_{\mathrm{em}}$) is fixed only after electroweak symmetry breaking.

Weak interaction (*Level B+*): The residual rotation group $O(3)$ after $O(4) \to O(3)$ SSB is algebraically isomorphic to $SU(2)$ (as Lie algebras: $\mathfrak{so}(3) \cong \mathfrak{su}(2)$). The oriented condensate—through the sign of the four-volume form—selects one of the two $SU(2)$ factors of $Spin(4) \simeq SU(2)_+ \times SU(2)_-$, structurally selecting a single *chiral* $SU(2)$ sector. This provides a structural route to weak-sector parity asymmetry—a feature postulated in the Standard Model acquires a geometric origin in ECT through the orientation of the ordered branch. The exactly maximally parity-violating Standard-Model pattern, however, remains an open derivation problem. The full derivation of the weak sector as an independent internal gauge field, however, remains open. The phenomenological realisation occurs through a secondary condensate scale at $v_2 \simeq 246\,\mathrm{GeV}$, producing $W^\pm$, Z bosons and electroweak symmetry breaking.

Strong interaction (*Level B*): $SU(3)$ colour *cannot* arise from the minimal $O(4) \to O(3)$ pattern: the algebra $\mathfrak{su}(3)$ has dimension 8, while $\mathfrak{so}(4)$ has dimension 6, so $\mathfrak{su}(3)$ is simply too large to fit inside $\mathfrak{so}(4)$. This is not merely a technical inconvenience. It means that the colour algebra cannot arise as a direct subgroup or residual factor of the minimal $O(4) \to O(3)$ pattern. The electroweak-type $U(1) \times SU(2)$ architecture is the *maximal* gauge structure derivable from the minimal condensate symmetry breaking. Any colour sector therefore requires additional internal degrees of freedom: a local complex rank-3 module with $SU(3)$-structure. A deeper geometric candidate (G_2 internal fibre) is also identified. Full QCD derivation remains open.

Gravity (*Level A/B*): Gravity arises from orientation stiffness—the condensate's resistance to bending of n^A (Part II).

Director coupling (*Level A/B*): Beyond the four known interactions, ECT identifies a fifth interaction channel: the *director coupling* $\mu_5 \bar{\Psi} \gamma^A n_A \Psi$—the lowest-dimension fermion–condensate operator permitted by the ordered-branch symmetry. Its name derives from the analogy with the director field of nematic liquid crystals. Possible signatures include spin precession in the approximate range $\omega_{\mathrm{dir}} \sim 10^{-10}$–$10^{-15}\,\mathrm{rad/s}$ (depending on the local condensate profile) and composition-dependent effects; the $\eta \sim 10^{-15}$ window is a target sensitivity rather than a sharp prediction.

The colour asymmetry. Among these sectors, colour is the only one that cannot be accessed from the minimal $O(4) \to O(3)$ pattern alone. This qualitative asymmetry motivates the dedicated discussion of colour completion below (Section 50).

Common pre-geometric origin. ECT provides a common pre-geometric framework for all known interaction sectors. Gravity arises from the ordered condensate geometry; the Abelian gauge sector follows from the compact phase fibre; a preferred chiral $SU(2)$ sector is structurally selected by the oriented broken phase; and the colour sector requires additional internal condensate structure beyond the minimal $O(4) \to O(3)$ pattern. Together with the generic director coupling, this yields a unified condensate-based organisation of the interaction sectors. In this sense ECT is broader than conventional GUT schemes, while its notion of unification remains structural rather than simple-group theoretic.

11.2. Elementary particles

The bosonic content of ECT is best organised hierarchically, by structural level. The description below records the bosonic sectors retained in the present formulation at successive levels of condensate structure; it should not be read as a final unique census of all possible emergent excitations. At the minimal level (Level 0: real scalar condensate with $O(4) \to O(3)$ SSB), ECT contains the graviton (spin-2, *Level A/B*), the surviving Goldstone mode χ (ultra-light condensate scalar, *Level A/B*), and the radial mode σ (heavy, *Level A/B*)—no gauge bosons. At Level 1, the compact phase fibre produces the photon ($U(1)$ gauge boson, *Level A*). At Level 2, the electroweak condensate at $v_2 = 246\,\mathrm{GeV}$ generates $W^{\pm}$, Z, and the Higgs boson (*Level B+*). At Level 3, extended internal structure produces the eight gluons (*Level B*). Fermions appear as spinorial excitations of the ordered branch; in the emergent Lorentzian regime their dynamics is captured by a reconstructed Dirac equation, while the chiral structure of the weak sector is tied to the preferred orientation selected by the condensate (*Level A/B*), and topological sectors yield an instanton mass pattern $m_n = n \times 1.6\,\mathrm{GeV}/c^2$ (*Level B*).

Neutrino sector. The geometrically one-chiral weak route that follows from the condensate symmetry breaking structurally selects one chiral branch of the fermion sector (*A/B*). This constrains the type of viable neutrino mass mechanism: a pure Planck-suppressed Majorana route undershoots the observed atmospheric neutrino scale by three orders of magnitude (*B*), making an embedded seesaw mechanism with a heavy sterile partner the structurally preferred completion (*C*). The two simplest dimensionally consistent combinations of ϕ_0 and v_2 populate the neutrino sector: the geometric-mean seesaw scale $M_R^{\mathrm{geom}} = \sqrt{\phi_0\, v_2} \approx 2.4 \times 10^{10}\,\mathrm{GeV}$ and the inverse-ratio Weinberg-operator scale $v_2^2/\phi_0 \approx 2.5 \times 10^{-5}\,\mathrm{eV}$; both are structural scale observations, not predictions (*C*). Standard PMNS/MSW oscillation phenomenology remains intact at leading order: the ECT preferred-direction correction is suppressed by approximately eight orders of magnitude relative to the standard oscillation Hamiltonian (*B*). Neutrinoless double-beta decay is the central experimental discriminant of the preferred Majorana/seesaw branch. The full flavour closure (generations, textures, PMNS angles, mass ordering) remains open.

Relative to the minimal Standard-Model field content, the present ECT closure retains an additional condensate scalar sector, represented by the ultra-light χ-mode and the heavy radial mode σ. The detailed dark-sector role of χ remains model-dependent in the current formulation.

What remains open (*Level Open*): the origin of three fermion generations; the Yukawa coupling hierarchy; the fine-structure constant $\alpha = 1/137$. Qualitative structural routes exist (e.g. generations may be related to topological sectors of S^3), but rigorous quantitative derivations do not.

Three logical classes of excitations. ECT does not place all particles on the same logical footing—a fundamental difference from standard QFT. Category (i), the *required condensate excitations* (graviton, radial mode σ, soft mode χ), are analogous to phonons in a crystal: they are inevitable consequences of an ordered medium with broken symmetry. Category (ii), the *structurally embedded Standard Model fields* (photon, $W^{\pm}$, Z, Higgs, fermions, gluons), are analogous to impurity atoms in a lattice: compatible with the medium but not derived from the minimal scalar action. Category (iii), the *topological sectors* (textures from $\pi_3(S^3) = \mathbb{Z}$; anti-predictions from $\pi_0 = \pi_1 = \pi_2 = 0$: no monopoles, no domain walls, no cosmic strings), are analogous to dislocations: determined by the topology of the vacuum manifold. In standard QFT all fields are equally fundamental postulates; in ECT they occupy different logical tiers.

Multi-particle description and the status of Fock space. Ordinary quantum mechanics works with a fixed number of particles N. Processes with variable N (emission, annihilation, pair creation) require quantum field theory, where the Fock space and creation/annihilation operators are additional postulates. In ECT, changing particle number is a reconfiguration of one condensate medium—analogous to vortices appearing and disappearing in a superfluid. ECT does not require Fock space as a fundamental postulate; however, an explicit emergent sector decomposition with variable excitation number remains to be constructed from condensate dynamics (Open; OP-Q11). This is a nontrivial reconstruction problem, not a foundational gap.

Multi-particle description is necessary even in the coherent branch (where no event-like Lorentzian interaction is available) for four reasons: (a) entanglement as non-factorisable common-medium correlation; (b) exchange statistics ($\pi_1(\mathcal{M}_2) = \mathbb{Z}_2$); (c) the decoherence threshold, which depends on N_{eff}; (d) processes with variable excitation number.

Figure 5 summarises the derivation logic of Part I.

12. Conclusions of Part I

12.1. Principal results

From six postulates (P1–P6) and one derived branch rule (BR1), the following structures are obtained:

1. Lorentzian signature $(-,+,+,+)$ from $O(4) \to O(3)$ SSB (A).
2. Universal causal cone: all species share $c_* = c$ (A/B).
3. Equation hierarchy: Euclidean $\to$ KG $\to$ Schrödinger (A/B).
4. Arrow of time from monotonic decoherence (A).
5. Second law: $dS_{\text{ent}}/dw \geq 0$ (A/B).
6. Branching into geometric and coherent sectors (A).
7. $U(1)$ electromagnetism from P5 (A/B).
8. $SU(2)$ weak sector (B); $SU(3)$ colour ($B/Open$).
9. Graviton as spin-2 orientation mode (A).

10. $S_0 = \hbar$ from compact-phase winding (B).

11. Quantisation from loop-sector topology (A).

12. Fifth-force coupling $\beta_5 \sim m_f/\bar{M}_{\rm Pl}$ (B).

13. Neutrino sector: geometric chirality selection (A/B); embedded seesaw as preferred completion with $M_R^{\rm geom} = \sqrt{\phi_0\, v_2}$ (C); PMNS/MSW phenomenology intact (B); $0\nu\beta\beta$ as central discriminant.

12.2. Key equations and principles of Part I

Effective kinetic tensor: $K^{AB} = \beta\,\delta^{AB} - \alpha\, n^A n^B$ (uniqueness theorem).

Propagation speed: $c_* = \sqrt{\beta/(\alpha-\beta)}$.

Matching relations: $G_N = c_*^4/(8\pi\phi_0^2)$, $\hbar = c_*\phi_0/\sqrt{\lambda}$.

Loop quantisation: $\oint \partial_A\theta\, dX^A = 2\pi n$; $S_{\rm loop,min} = 2\pi\hbar$.

Cross-sector universality: $c_{\rm phase} = c_\gamma = c_{\rm GW} = c_*$.

Second law: $dS_{\rm ent}/dw = N_{\rm eff}\gamma(dq/dw)^2 \geq 0$.

12.3. Predictions originating in Part I

GW170817 consistency: $|c_{\rm GW}/c - 1| < 6\times 10^{-15}$ (satisfied).

Fifth force: $\omega_5 \sim 10^{-10}\,$rad/s, $\eta \sim 10^{-15}$ (MICROSCOPE-2).

LIV: $\Delta t_{\rm GW} \sim 10^{-64}\,$s (Planck-suppressed, currently unobservable).

Casimir boundary-sensitive vacuum response (conditional; soft-mode counting provisional).

PMNS/MSW oscillation intactness: ECT preferred-direction correction suppressed by $\sim 10^{-8}$ relative to standard oscillation scale at $E \sim 1\,$GeV (B).

12.4. Part I derivation logic



Figure 5: Part I derivation logic. From six postulates and BR1, the theory derives: Lorentzian signature, universal causal cone, equation hierarchy, arrow of time, second law, branching, gauge symmetries, particles, quantisation. Colours indicate Level A, B, and Open status.

Figure 5 shows the complete derivation architecture of Part I.

Part II

Macroscopic Physics: Gravity, Cosmology, and Observational Tests

13. Special Relativity as the homogeneous limit

In the homogeneous ordered phase, where the condensate amplitude and direction are uniform throughout space, the effective metric reduces to the Minkowski form: $ds^2 = -c_*^2 dt^2 + dx^2 + dy^2 + dz^2$. All familiar consequences of special relativity follow as theorems: Lorentz transformations arise as the invariance group of the causal cone; time dilation and length contraction are kinematic consequences; $E^2 = p^2c^2 + m^2c^4$ follows from the effective wave equation. In ECT, special relativity is a *derived limit* (*Level A*), not an independent postulate.

13.1. The universal speed bound

One of the most familiar facts of physics — that nothing can travel faster than light — is, in standard special relativity, a direct consequence of Einstein's second postulate. In ECT, this fact is not postulated: it is a theorem.

The argument rests not on any particular type of excitation but on the *characteristic cone* of the ordered-branch field equation. The principal symbol of the equation of motion for Φ defines a unique causal cone with speed c_*. Through the symmetric-hyperbolic structure of the dynamics and the Leray–Choquet-Bruhat theorem, ECT establishes at Level A that the domain of dependence of any point lies strictly inside its past causal cone, and that no ordered-branch disturbance — linear or nonlinear — propagates outside the cone. This is a theorem about the full nonlinear scalar dynamics, not merely about linearised waves.

For linear plane waves, the bound takes a concrete form. A massive mode has a group velocity strictly less than c_*, with equality only in the massless limit: massive excitations travel inside the cone, massless excitations on its boundary.

An important subtlety deserves mention. The *phase velocity* of massive modes formally *exceeds* c_*. This is a well-known feature of relativistic wave equations and is *not* a violation of causality: phase velocity does not carry information. The causal bound is set by the characteristic cone — the front speed — not by the phase velocity. This distinction highlights the correct ECT formulation: the speed limit is a bound on *causal propagation*, not on every quantity one can label a "velocity".

13.2. Signal speed, not photon speed

A deeper point emerges from the ECT derivation. The fundamental limiting velocity is not "the speed of light" as a photon-specific constant. It is the *signal speed* of the ordered-branch causal cone — the maximum speed of causal influence, front propagation, and information transfer.

The photon merely *realises* this boundary speed. It does so because the emergent Abelian gauge boson is massless (the relevant $U(1)$ symmetry remains unbroken, and gauge invariance forbids

a mass term) and because the gauge sector inherits the same causal cone as the scalar branch. Gravitational waves likewise inherit the same cone in the minimal tensor route. The near-equality confirmed by GW170817 ($|c_{\mathrm{GW}}/c - 1| < 6 \times 10^{-15}$) is therefore a structural consistency check, not a coincidence.

Because inertial coordinate changes in the homogeneous ordered branch are precisely the transformations preserving one and the same null cone, the same vacuum signal speed is measured in every inertial frame. What Einstein postulated as the universality of the vacuum light speed is, in ECT, reduced to a structural consequence of the unique ordered-branch causal cone (*Level A*) together with cone inheritance in the minimal gauge route (*Level A/B*).

In this sense, the vacuum speed of light is not fundamental by itself; it is the experimentally accessible manifestation of the deeper ordered-branch causal speed.

14. Emergent gravity and the generalised Einstein equations

In ECT geometry is not the primitive ontology. The effective metric is a derived bookkeeping object encoding how excitations propagate on the ordered condensate background. Gravity is not a fundamental force but the long-wavelength geometric response of the ordered condensate to deformations of its orientation structure n^A. The graviton is not a quantum of fundamental geometry but a low-energy excitation of the condensate order. This inverts the conventional logic of quantum gravity: ECT replaces the question "how to quantise an existing geometry?" with the question "how do collective condensate excitations produce a geometric description?".

The mechanism is analogous to elastic stresses in a crystal lattice. When matter bends n^A away from equilibrium, "orientation stress" acts as an effective gravitational field.

The derivation proceeds through stages. The orientation stiffness $\kappa_n \sim u_0^2/m_\sigma^2 \sim \phi_0^2$ determines the energy cost of bending n^A. Linearising around the flat background, transverse-traceless fluctuations satisfy a spin-2 wave equation identical to linearised GR (*Level A/B*).

At the nonlinear level, ECT derives **generalised Einstein equations (GEE)**:

$$G_{\mu\nu} + \Lambda_{\mathrm{eff}}\, g_{\mu\nu} = 8\pi G_{\mathrm{eff}}\, T_{\mu\nu} + \Theta_{\mu\nu}[n], \tag{11}$$

where $\Theta_{\mu\nu}[n]$ is the orientation-stress tensor (*Level B*). This term has no analogue in standard GR. It vanishes in homogeneous configurations but becomes dynamically important in three distinct observational domains:

On *galactic scales*, $\Theta_{\mu\nu}$ provides the additional acceleration that explains flat rotation curves without particle dark matter (Section 20).

On *cluster scales*, it modifies gravitational lensing morphology, reproducing the Bullet Cluster offset without dark matter halos (Section 22).

In *cosmology*, the evolving G_{eff} and $\Theta_{\mu\nu}$ jointly address the Hubble tension and the JWST problem (Sections 18–19).

Gravitational waves propagate at c_* with two transverse-traceless modes (*Level A/B*), consistent with LIGO/Virgo. Weak-field and post-Newtonian limits reproduce Newtonian gravity and are compatible with Solar System tests (PPN parameters).

15. Condensate evolution and cosmic history



Figure 6: Cosmic timeline comparison: ΛCDM (top) and ECT (bottom). The earliest epoch in ECT is a timeless Euclidean phase; the transition $O(4) \to O(3)$ creates time itself. The lower panel shows the condensate amplitude $u_0(t)$, scale factor $a(t)$, and effective gravitational coupling $G_{\text{eff}}(t)$. (*Level A for the pre-Lorentzian regime; Level B for the detailed timeline.*)

In ΛCDM, the universe begins at $t = 0$ (the Big Bang singularity) and proceeds through inflation, nucleosynthesis, recombination, structure formation, and dark-energy domination. ECT tells a fundamentally different story (Fig. 6).

Before the $O(4) \to O(3)$ transition, there is no time at all. The universe exists as a four-dimensional Euclidean medium with full $O(4)$ symmetry. The concept of "before" has no meaning in the usual temporal sense. The Big Bang singularity is replaced by the ordering transition, at which the condensate amplitude u_0 grows from zero to its equilibrium value $\phi_0 \sim \bar{M}_{\text{Pl}}$. This growth naturally produces a period of rapid ordered-branch expansion, replacing the standard inflationary postulate with a pre-Lorentzian Euclidean ordering picture.

After the ordering is established, the evolution shares familiar milestones (nucleosynthesis, recombination, galaxy formation), but two features distinguish ECT from ΛCDM: the gravitational coupling G_{eff} evolves slowly with redshift, and the dark-energy component arises from residual condensate energy rather than from a cosmological constant.

A notable consequence is the **age of the Lorentzian branch**. In ECT, "the age of the universe" refers to the duration of the ordered Lorentzian branch, not the time since a fundamental singularity. Because $G_{\text{eff}}(z) > G_N$ at high redshift, the Hubble rate $H(z)$ is larger over part of

cosmic history, which reduces the age integral $t_0 = \int_0^\infty dz/[(1+z)H(z)]$. Benchmark estimates give $t_0^{\rm ECT} \approx 13.0$–$13.5\,$Gyr, modestly shorter than the ΛCDM value of $\sim 13.8\,$Gyr (*Level B*). This reduction arises directly from the enhanced early-time gravity and is not a separate assumption. It has consequences for the JWST problem discussed below.

At late times, ECT admits two scenarios: Scenario A ($u_0 \to$ const, eternal expansion) and Scenario B ($u_0 \to 0$, recollapse on a timescale $\sim 10^{100}$ years). Distinguishing these requires data not yet available.

16. The four classical cosmological problems

The standard Big Bang model faces four well-known puzzles. The conventional resolution invokes cosmic inflation — a brief period of exponential expansion driven by a hypothetical inflaton field. While inflation is remarkably effective, it introduces its own unresolved issues: the inflaton has never been directly observed; inflation requires specific initial conditions to start (the initial homogeneity problem — inflation itself needs a sufficiently uniform region to begin, yet it was introduced to *explain* homogeneity); the measure problem makes probabilistic predictions ill-defined in the context of eternal inflation; and the trans-Planckian problem arises because CMB modes observed today had sub-Planckian wavelengths at the onset of inflation, potentially probing unknown physics.

ECT resolves all four puzzles at the level of the ordering transition itself, without introducing an inflaton field and without encountering any of these difficulties.

16.1. The horizon problem

The cosmic microwave background (CMB) — relic radiation from 380 000 years after the Big Bang — is uniform to one part in 10^5 across the entire sky. In ΛCDM, opposite sides of the sky were never in causal contact at that epoch: their past light cones do not overlap. The uniformity is unexplained without inflation.

In ECT, the resolution is structural (*Level A for the pre-ordered regime; Level B for its cosmological realisation*). Before the $O(4) \to O(3)$ transition, the Euclidean phase has no causal structure at all: there are no light cones, no causal horizons, no "regions out of contact." All points of $\mathcal{M}^4$ are connected by the Euclidean metric. Uniformity across the CMB is inherited from the symmetric pre-Lorentzian phase.

16.2. The flatness problem

Observations show $|\Omega_k| < 0.002$ (Planck 2018). In ΛCDM, this requires fine-tuning to $\sim 10^{-60}$.

In ECT: the Euclidean manifold $\mathcal{M}^4$ in P1 is flat by construction (δ_{AB}). Spatial flatness of the emergent Lorentzian geometry is a structural consequence of the flat substrate (*Level A*).

16.3. The relic and monopole problem

Grand Unified Theories predict massive topological defects (magnetic monopoles). None observed. In ECT: the vacuum manifold S^3 has $\pi_2(S^3) = 0$ — *no stable monopoles are topologically possible*

(*Level A*).

16.4. The primordial perturbation problem

The CMB contains tiny fluctuations ($\sim 10^{-5}$) with a nearly scale-invariant spectrum ($n_s = 0.9649 \pm 0.0042$, Planck 2018). In ECT, the ordered branch carries exactly one independent scalar perturbation mode at leading order (*Level A*). ECT-native critical O(4) fluctuation estimates yield $n_s \sim 0.96$–0.97, consistent with observations. The full first-principles derivation of n_s, r, and A_s remains *Level B/Open*.

16.5. Summary

All four problems are resolved without introducing an inflaton field, without fine-tuning initial conditions, and without the measure, trans-Planckian, and initial-homogeneity difficulties that inflation carries. The pre-Lorentzian Euclidean phase provides a natural starting point from which the Lorentzian ordered branch inherits uniformity, flatness, topological simplicity, and a single-mode perturbation sector. Phenomenological introduction of inflation is unnecessary.

17. The dark sector in ECT

The terms "dark matter" and "dark energy" were introduced phenomenologically to account for observations that standard gravity with visible matter alone cannot explain: flat rotation curves, gravitational lensing, large-scale structure, accelerated expansion. These phenomena are real and well-established. However, the entities postulated to explain them — invisible particles (WIMPs, axions, sterile neutrinos) and a cosmological constant Λ — have never been directly confirmed despite decades of dedicated searches. ECT provides a unified structural explanation for the entire dark sector from the condensate dynamics, without introducing any new fundamental entities.

17.1. Dark matter: condensate response, not particles

In ΛCDM, the flat rotation curves of galaxies, gravitational lensing in clusters, and the large-scale structure of the universe are attributed to an invisible component — dark matter particles — constituting $\sim 27\%$ of the energy budget. Despite extensive experimental programmes spanning direct detection (LUX, XENON, PandaX, LZ), indirect detection (Fermi-LAT, AMS-02), and collider searches (LHC), no dark matter particle has ever been observed. The parameter space for WIMPs — once the leading candidate — is almost entirely excluded.

In ECT, all observational phenomena attributed to dark matter arise from the nonlinear response of the condensate amplitude ϕ in weak gravitational fields (*Level B*). The orientation-stress tensor $\Theta_{\mu\nu}[n]$ in the generalised Einstein equations (11) provides additional effective gravitational acceleration in the regimes where dark matter is conventionally invoked: galaxy outskirts, cluster environments, and cosmological scales. Flat rotation curves, the BTFR, the RAR, and cluster lensing offsets are all reproduced without a single dark matter particle (Sections 20–22).

17.2. Dark energy: condensate residual energy, not a cosmological constant

In ΛCDM, the accelerated expansion of the universe is attributed to a cosmological constant Λ — a fixed parameter responsible for $\sim 68\%$ of the total energy budget. No fundamental theory explains its value, which differs from quantum field theory estimates by approximately 120 orders of magnitude (the cosmological constant problem).

In ECT, what appears as dark energy is the *residual amplitude energy* of the ordered condensate background — the same medium that produces gravity and quantum mechanics. No independent cosmological constant or dark-energy substance is postulated. The equation-of-state parameter satisfies $w_0 \neq -1$, unlike the fixed $w = -1$ of a cosmological constant; ECT predicts $w_0 \approx -0.83$, compatible with DESI 2024 results which found evidence for dynamical dark energy at $\sim 2.5\sigma$ significance. If $w = -1$ is confirmed with high precision by future surveys (DESI, Euclid), the ECT late-time acceleration closure would be falsified. Conversely, if dynamical dark energy is confirmed, ECT provides a structural explanation while ΛCDM does not.

A structural advantage of the ECT reframing: loop contributions to the vacuum energy are finite (the physical cutoff is $m_\sigma \sim M_{\mathrm{Pl}}$, not infinity), and the effective cosmological contribution is a dynamical functional of the condensate background rather than a rigid constant. This does not yet solve the 120-orders-of-magnitude fine-tuning problem; it reframes it as a question about condensate dynamics rather than a divergent sum over zero-point energies (*Open: OP-Λ3*).

17.3. ECT reinterpretation of the dark sector

Within the minimal ECT benchmark developed here, the phenomena usually attributed to dark matter particles and to a cosmological constant are reinterpreted as condensate-response effects: condensate-induced gravitational response for galactic mass-discrepancy phenomenology, and ordered-branch background dynamics for late-time accelerated expansion. The terms "dark matter" and "dark energy" are retained as labels for observational phenomena for comparison with the literature; they do not refer to independent physical substances in ECT. The present benchmark does not require dark matter particles or an independent cosmological constant to account for the observational data. This does not yet exclude every possible ECT extension with additional weakly coupled condensate excitations; it states only that the present minimal framework does not need them.

Important interpretive caveat. The statement "ECT does not need dark matter particles in its minimal benchmark" is not equivalent to "ECT proves that no particle beyond the Standard Model can ever exist." The first is a statement about the present minimal closure. The second would be an overclaim that the theory does not make. Collective condensate excitations and topological sectors remain structurally possible as optional extensions; they are not required by the core benchmark fit.

18. The Hubble tension



Figure 7: The ECT Hubble-tension alleviation mechanism. The evolving effective gravitational coupling $G_{\rm eff}(z) > G_N$ at high redshift shifts the inferred Hubble constant upward by $\sim 3\,$km/s/Mpc. (*Sign: Level A structural prediction; magnitude: Level B.*)

The Hubble constant H_0 measures the current expansion rate of the universe. Two independent methods give discrepant results: the Planck satellite, analysing the CMB assuming ΛCDM, infers $H_0 = 67.4 \pm 0.5\,$km/s/Mpc; the SH0ES team, using the cosmic distance ladder (Cepheids + Type Ia supernovae), measures $H_0 = 73.0 \pm 1.0\,$km/s/Mpc. The discrepancy exceeds 5σ and is known as the *Hubble tension* — one of the most pressing open problems in cosmology.

ECT offers a structural alleviation mechanism (Fig. 7). At high redshift, the condensate was slightly less screened, meaning $G_{\rm eff}(z) > G_N$. Specifically, $G_{\rm eff}(z) \approx G_N(1+z)^{2\varepsilon}$ with $\varepsilon \approx 0.01$. The enhanced gravity reduces the angular diameter distance to the last scattering surface; a CMB-calibrated analysis then interprets this as a higher H_0, shifted upward by $\sim 3\,$km/s/Mpc — the direction and approximate magnitude needed.

The *sign* of the shift is a structural prediction (*Level A*): any ordered-branch background with $G_{\rm eff} > G_N$ at high z necessarily shifts H_0 upward. The magnitude is closure-dependent (*Level B*). The same condensate background simultaneously affects the age budget and early structure formation, producing correlated predictions rather than independent parameters.

19. The JWST early-galaxy problem

The James Webb Space Telescope (JWST), launched in December 2021, has detected galaxies at redshifts $z > 10$, when the universe was less than 500 million years old. Several are unexpectedly massive and mature — containing old stellar populations, organised disk structures, and chemical

enrichment that seem to require more formation time than ΛCDM allows. Standard structure formation with fixed G_N cannot produce such objects quickly enough.

ECT addresses this through two complementary mechanisms, both arising from the same ordered-branch condensate dynamics:

First, the evolving $G_{\text{eff}}(z) > G_N$ at high z means gravitational collapse was *more efficient* in the early universe.

Second, the GEE (11) include $\Theta_{\mu\nu}[n]$, which provides additional effective attraction beyond a simple scalar-tensor model.

It is important to note that the shorter ECT age (Section 15) means these high-redshift galaxies had *even less* formation time available than in ΛCDM — making the problem nominally worse. However, the two enhancement mechanisms (stronger gravity + orientation-stress attraction) more than compensate: the accelerated collapse rate outweighs the reduced time budget. These are not independent tuning parameters: the same condensate background that shifts H_0 upward also enhances early structure formation.

20. Galactic rotation curves



Figure 8: Milky Way rotation curve. ECT (solid) fits the data with one parameter $r_0 = 14\,\text{kpc}$: $\chi^2/N = 2.95$. Newtonian prediction from baryons alone (dashed) falls off at large radii. (*Level B.*)

When astronomers measure the orbital speeds of stars in spiral galaxies, they find that stars in the outskirts orbit much faster than predicted by the visible matter alone. In a Newtonian calculation, the rotation speed should decrease as $v \propto 1/\sqrt{R}$ beyond the luminous disk, but observations show flat rotation curves extending to many disk radii. The standard explanation invokes a halo of dark matter particles surrounding each galaxy.

Despite decades of direct detection experiments (LUX, XENON, PandaX, LZ, and others), no

dark matter particle has ever been observed. The parameter space for WIMPs — once the leading candidate — is almost entirely excluded. This persistent non-detection motivates consideration of alternative explanations.

ECT provides one. The nonlinear dynamics of the macroscopic amplitude variable $\phi = \frac{1}{\beta}\ln(u/u_\infty)$, which measures the local "degree of Lorentzianness" of the vacuum, reproduces MOND-like dynamics on galactic scales without particle dark matter (*Level B*). In dense environments (Solar System, inner galaxy), ϕ is strongly screened and standard gravity is recovered. On galactic outskirts, the ϕ-sector enters a nonlinear regime whose scale-invariant form gives:

$$g(R) = \tfrac{1}{2}(g_N + \sqrt{g_N^2 + 4\, g_N\, g^\dagger}), \tag{12}$$

with $g^\dagger \approx cH_0/(2\pi)$. In the deep-field limit ($g_N \ll g^\dagger$): $g \approx \sqrt{g_N g^\dagger}$ (flat rotation curves). In the strong-field limit: $g \approx g_N$ (Newton recovered).



Figure 9: Rotation-curve fits for a sample of SPARC galaxies, spanning a wide range of morphologies: from dwarf irregulars to massive spirals. ECT (solid) with one parameter per galaxy. (*Level B.*)

This has been tested against **165 galaxies from the SPARC database** (Fig. 8, 9). ECT fits each galaxy with a single parameter (r_0 or equivalently $g^\dagger_{\text{eff}}$), achieving median $\chi^2/N \approx 2.95$ — substantially better than ΛCDM with NFW dark matter halos ($\chi^2/N \approx 6.55$), which requires more parameters.

21. The baryonic Tully–Fisher relation and the RAR



Figure 10: The baryonic Tully–Fisher relation (BTFR) from the SPARC sample. ECT predicts the exact algebraic slope 4 (solid line): $v_{\rm flat}^4 = G_N M_{\rm bar}\, g^\dagger$. (*Slope: Level A; $g^\dagger$ value: Level B.*)

The BTFR — $M_{\rm bar} \propto v_{\rm flat}^4$ — is one of the tightest empirical correlations in extragalactic astronomy (Fig. 10). In ΛCDM, reproducing it requires fine-tuning the dark matter halo for each galaxy. In ECT, the BTFR is an *exact algebraic consequence* of the ϕ-closure: $v_{\rm flat}^4 = G_N M_{\rm bar}\, g^\dagger$ with slope exactly 4 (*Level A*).



Figure 11: Radial acceleration relation (RAR) diagnostics from the SPARC sample. ECT reproduces the tight observed correlation between total and baryonic gravitational acceleration, with $g^\dagger \approx 1.1 \times 10^{-10}$ m/s^2 matching the McGaugh et al. (2016) empirical value. (*Level B.*)

The RAR (Fig. 11), discovered by McGaugh et al. (2016), is a direct prediction of the ECT interpolation law (12). Unlike standard MOND where $g^\dagger$ is a universal constant, in ECT $g^\dagger$ depends on the local degree of Lorentzianness (*Level B*), leading to a testable environmental modulation.

22. Cluster-scale gravitational lensing



Filled circles = BCG (collisionless); open circles = X-ray gas (collisional)

Figure 12: Gravitational lensing in the Bullet Cluster (1E 0657-56). The lensing signal (contours) is offset from the X-ray gas (colour), conventionally attributed to collisionless dark matter. ECT reproduces this offset through the condensate ϕ-closure without dark matter particles. (*Level B.*)



Cluster	Peak	$d_*^{\rm ECT}$	$d_*^{\rm obs}$	A_0	$A_{\rm fin}$
Bullet	**BCG**	37 kpc	<25 kpc	0.42	0.53
MACS J0025	**BCG**	10 kpc	<30 kpc	0.68	0.86
El Gordo	**BCG**	1 kpc	<35 kpc	0.43	0.56
Abell 520	*gas*	361 kpc	gas core	0.71	0.87

Figure 13: Cluster-merger gravitational lensing in ECT. Results for the Bullet Cluster, El Gordo, MACS J0025, and Abell 520, including lensing mass ratios and morphology classification. In Abell 520, the lensing peak falls on the gas core — consistent with ECT but a direct tension point for collisionless dark matter models. (*Level B.*)

Galaxy-cluster mergers provide a critical dark-sector test. In a collision, the hot intracluster gas decelerates (electromagnetic interaction), while galaxies and any collisionless dark matter should pass through. In the Bullet Cluster, the lensing signal is offset from the gas toward the galaxies — conventionally interpreted as evidence for collisionless dark matter halos.

ECT addresses cluster lensing through the ϕ-closure with chameleon screening (Fig. 13). The $\Theta_{\mu\nu}$ contribution generates gravitational slip, producing lensing morphology consistent with observations (*Level B*).

The case of **Abell 520** is particularly significant. In this system, the dominant lensing peak coincides with the *gas-rich core*, not with the galaxy concentrations. This is a *direct contradiction* to collisionless dark matter models, which predict that the lensing signal should track the collisionless component (galaxies). In ECT, this is the expected result: the gas core is locally more compact, so the ϕ-enhancement is strongest there. A single rule generates galaxy-peaked lensing in Bullet/El Gordo/MACS J0025 and *gas-peaked* lensing in Abell 520 — a unified prediction absent in models requiring a fixed collisionless species.

23. Matter–antimatter asymmetry

The observable universe contains far more matter than antimatter: $\eta_B^{\rm obs} \approx 6 \times 10^{-10}$. This asymmetry requires baryon number violation, C/CP violation, and departure from thermal equilibrium (the three Sakharov conditions, 1967). The Standard Model satisfies these conditions in principle but cannot quantitatively produce the observed asymmetry.

ECT provides a structural framework in which all three Sakharov conditions are satisfied by native ordered-medium mechanisms (*Level A* for the structural possibility). The $O(4) \to O(3)$ transition provides out-of-equilibrium conditions; the fifth coupling $\mathcal{L}_5$ supplies CP violation; defect-mediated topology changes in the condensate allow baryon number violation. The specific leptogenesis mechanism and the estimated $\eta_B \sim 9 \times 10^{-10}$ (factor 1.5 from observation) are *Level B*.

24. Conclusions of Part II

24.1. Principal results

From the ordered-branch EFT, the geometric branch produces:

1. Special Relativity as the homogeneous limit (A).
2. GEE with $\Theta_{\mu\nu}[n]$ (B).
3. Four cosmological problems resolved without inflation (A/B).
4. Dynamical dark energy $w_0 \approx -0.83$ (B).
5. Hubble tension: $\Delta H_0 \sim +3\,\mathrm{km/s/Mpc}$ (*sign: A; magnitude: B*).
6. JWST early galaxies via enhanced $G_{\rm eff} + \Theta_{\mu\nu}$ (B).
7. 165 SPARC galaxies: $\chi^2/N \approx 2.95$ vs. ΛCDM 6.55 (B).
8. BTFR slope exactly 4 (A).
9. RAR: $g^\dagger \approx cH_0/(2\pi)$ (B).
10. Cluster lensing incl. Abell 520 gas-peaked (B).

11. Baryon asymmetry: Sakharov conditions natively (A/B).

12. Dark matter and dark energy eliminated (B).

24.2. Key equations and principles of Part II

Generalised Einstein equations: $G_{\mu\nu} + \Lambda_{\text{eff}} g_{\mu\nu} = 8\pi G_{\text{eff}} T_{\mu\nu} + \Theta_{\mu\nu}[n]$.

Galactic interpolation: $g(R) = \frac{1}{2}(g_N + \sqrt{g_N^2 + 4 g_N g^\dagger})$.

BTFR: $v_{\text{flat}}^4 = G_N M_{\text{bar}} g^\dagger$ (slope 4, algebraic).

$G_{\text{eff}}(z) \approx G_N (1+z)^{2\varepsilon}$, $\varepsilon \approx 0.01$ (Hubble shift).

24.3. Observational tests and falsifiers

ECT is falsifiable. Key predictions of the macroscopic sector:

- BTFR slope = 4 (A); $g^\dagger \propto cH_0$ (B).
- Fifth force $\eta \sim 10^{-15}$ (B); dark energy $w_0 \approx -0.83$ (B).
- Environmental $g^\dagger$ modulation (B); graviton polarisation: two TT modes (B).
- CMB quadrupole deficit (*B/Open*).

Full list with explanations is given in Part IV, Section 49.



Figure 14: ECT cosmological predictions and comparison with observations and ΛCDM. ECT provides correlated predictions for dark energy, the Hubble tension, the JWST problem, and galactic phenomenology from the same ordered-branch dynamics. (*Level B.*)

24.4. Part II derivation logic



Figure 15: Part II derivation logic. From the ordered-branch EFT, the geometric branch produces all macroscopic results listed above. Colours indicate derivation levels.

The full derivation logic of Part II is shown in Fig. 15. The cosmological predictions are summarised in Fig. 14.

Part III

Quantum Sector: Coherent Branch

Why does quantum mechanics exist? In ECT, the answer has three layers.

First: the Euclidean functional integral. Because the fundamental arena is a four-dimensional Euclidean manifold with no built-in time, the natural mathematical object is the functional integral $\int \mathcal{D}\Phi\, e^{-S_E[\Phi]}$, which sums over *all* field configurations — including "closed histories" where the field returns to itself along the fourth coordinate. There are no events, no temporal ordering, no cause-effect sequences at this level. By the Osterwalder–Schrader reconstruction theorem, this Euclidean functional integral is mathematically equivalent to a unitary quantum

theory on a Hilbert space (*Level A/B*).

Second: the action scale $S_0 = \hbar$. The compact phase θ of the condensate ($\Phi_{\text{eff}} = \rho\, e^{i\theta}$) imposes single-valuedness on closed loops: $\oint \partial_A \theta \, dX^A = 2\pi n$. This topological constraint sets the minimal loop action $S_{\text{loop,min}} = 2\pi S_0$, where S_0 is identified with $\hbar$ through the condensate matching relation (*Level B*).

Third: the physical interpretation of the Euclidean–Lorentzian boundary. In the Euclidean regime, all configurations contribute coherently to the functional integral — including closed histories that would be "forbidden" in the Lorentzian sense. After the emergence of the Lorentzian causal structure, these closed histories continue to contribute to quantum amplitudes: they are the source of interference, superposition, and tunnelling. For macroscopic systems with $N_{\text{eff}} \gg 1$, decoherence exponentially suppresses the coherent contribution of these closed Euclidean histories, and classical behaviour emerges. For microscopic systems with $N_{\text{eff}} \sim 1$, the closed histories remain coherent — this is quantum behaviour. The quantum–classical boundary is therefore not a separate postulate but a consequence of the number of environmental modes coupled to the system (*Level A/B*).

25. ECT versus standard quantum mechanics



Figure 16: Structural comparison of standard quantum mechanics (left) and ECT's coherent branch (right). Standard QM starts from axioms (Hilbert space, operators, Born rule, Schrödinger equation) and computes predictions. ECT derives these structures from the Euclidean condensate dynamics. The central column shows the structural origins that ECT provides for each axiom. (*Level A/B for the structural routes; Level Open for the fully closed Born theorem.*)

Standard quantum mechanics answers the question "how?": given a Hamiltonian and an initial state, it prescribes how to compute probabilities. ECT aims to answer the additional question "why?": why do Hilbert spaces, operators, superposition, the Born rule, and decoherence exist at all? (Fig. 16).

The deepest structural distinction lies in the type of mathematical problem. In standard QM, the fundamental problem is the *Cauchy problem* (initial-value problem): $i\hbar\,\partial_t\psi = H\psi$. Given the state at $t = 0$, time evolution uniquely propagates it forward. In ECT, the fundamental

problem is a *boundary-value problem* on the Euclidean manifold: $\delta^{AB}\partial_A\partial_B\Phi - V'(\Phi) = 0$. The solution is constrained by data on the *entire boundary* of the four-dimensional domain, not just on one time-slice.

This has profound consequences. The Cauchy problem generates *one-directional* evolution: past determines future, but not vice versa. The boundary-value problem generates a *four-dimensionally self-consistent* configuration: future boundary conditions constrain the present just as past boundary conditions do. This is not retrocausality in the Lorentzian sense — it is the normal functioning of a boundary-value problem. The probabilistic character of quantum mechanics, in ECT's view, arises not from fundamental indeterminacy but from coarse-graining: when untracked environmental modes are integrated out, the effective description of a subsystem becomes inherently statistical.

25.1. Detailed structural comparison: ECT vs standard QM

The following comparison illustrates how ECT's coherent branch differs from standard quantum mechanics on specific physical examples.

The hydrogen atom. In standard QM, one solves $H\psi = E\psi$ and obtains $E_n = -13.6/n^2\,\mathrm{eV}$ together with the probability densities $|\psi_{nlm}|^2$. ECT gives a structurally richer picture of the same system.

The atom exists not in a fundamentally given Lorentzian spacetime but in the symmetry-broken condensate background that supports the emergent spacetime and quantum dynamics (Layer 1: background). The electron is described as a spinorial field compatible with the condensate geometry and the emergent Lorentzian sector; it is not yet derived as a collective excitation of the bare bosonic condensate (Layer 2: electron). The proton is treated as a localised composite source in the emergent gauge sector; its full microscopic internal origin (colour structure) is not yet derived from first principles (Layer 3: proton). The electromagnetic binding interaction is interpreted within the emergent $U(1)$ gauge sector, rather than being introduced in ECT as an independent fundamental ingredient external to the condensate framework (Layer 4: electromagnetic bond). The atom as a whole is a self-consistent bound configuration involving the protonic source, the electronic spinorial field, the $U(1)$ gauge connection, and the surrounding condensate-supported spacetime and quantum background (Layer 5: bound state).

The ground state $1s$ is the lowest regular, node-free, spherically symmetric bound mode consistent with these structural ingredients. At the structural level, discrete levels may be understood as arising because only certain regular, phase-consistent, single-valued solutions are admissible—in harmony with the winding/single-valuedness conditions of the coherent branch: $\oint dw\cdot p_w = 2\pi S_0\, n$ (*Level B*).

From Dirac to Schrödinger. The full structural description of hydrogen is first spinorial and gauge-coupled. The Schrödinger orbital, with its familiar $|\psi_{nlm}|^2$ probability densities, is the non-relativistic envelope limit—the regime where velocities, binding energies, and spin-dependent relativistic corrections are small compared with the rest scale. Within ECT, the familiar Schrödinger description is therefore treated as a low-energy effective level of description rather than the deepest structural level.

What remains open. The current framework does not yet derive hydrogenic fine structure, the Lamb shift, hyperfine splitting, or the numerical value of $\alpha_{\mathrm{fs}} = 1/137$. These are open derivational

targets (*Level Open*).

The double-slit experiment. In standard QM, $\psi = \psi_1 + \psi_2$ and $P = |\psi_1 + \psi_2|^2$. In ECT, the interference pattern is interpreted as arising from a connected solution of the underlying elliptic boundary-value problem, whose effective Lorentzian projection contains contributions associated with both slit channels. The interference is a property of the four-dimensional field configuration, not merely of the effective probability-amplitude description (*Level A/B*). The slit arrangement constrains the families of admissible phase-coherent condensate configurations; the detected pattern emerges from their coherent recombination. On this reading, the wavefunction is the effective envelope of a deeper phase-coherent condensate response.

Loss of interference and decoherence. When which-path information is coupled to the environment, the relevant phase coherence is lost: the influence functional from the environmental modes suppresses the off-diagonal (cross-slit) terms. In the ECT language, this is precisely the transition $\Gamma_{\text{loop}} \ll 1 \to \Gamma_{\text{loop}} \gtrsim 1$ described in the decoherence section—interference vanishes not because of a mysterious collapse, but because the condensate-plus-environment response ceases to remain phase-coherent (*Level A/B*).

The Aharonov–Bohm effect. In standard QM, a charged particle acquires an observable phase shift $\Delta\varphi = (e/\hbar) \oint A_\mu \, dx^\mu$ even in regions where the electromagnetic field strength vanishes. In the observational regime this is the standard $\hbar$-based form; within the ECT coherent branch it corresponds to the same phase-holonomy structure after the identification $S_0 = \hbar$. In ECT, this is especially natural: the compact phase $\theta(X)$ is already a primary structural variable of the coherent branch, and the gauge connection A_A arises as its local compensating field. The AB phase shift is therefore a direct manifestation of a nontrivial holonomy of the emergent $U(1)$ connection around a closed loop. Phase and loops are fundamental ingredients of the ECT quantum sector, so the Aharonov–Bohm effect is not an odd addition to the theory but exactly the kind of phenomenon one should expect when quantum dynamics emerges from phase-sensitive loop structure (*Level B: interpretive*).

Measurement. In standard QM, collapse is an extra postulate. In ECT, the detector belongs to the full boundary-value problem. At the reduced-description level, effective collapse is the subsystem manifestation of a Euclidean-to-Lorentzian transition under strong environmental locking ($\Gamma_{\text{loop}} \gtrsim 1$). Multiple winding sectors coexist in the coherent regime; environmental coupling drives decoherence; when $\Gamma_{\text{loop}} \gg 1$, interference is suppressed and a single classical outcome is selected (*Level A/B*).

Determinism. The fundamental Euclidean equation is deterministic: one specifies boundary conditions and obtains one global field configuration. No probabilities, no collapse, no stochasticity at the deepest level. Statistical quantum mechanics appears because realistic observers never have access to the full condensate state. When environmental modes are integrated out, the subsystem acquires influence functionals, reduced density matrices, and effective probabilities. ECT stands to standard QM as microscopic Newtonian mechanics stands to statistical mechanics: deterministic underneath, probabilistic after coarse-graining (*Level A*).

"Future affects past." In standard QM, this does not arise because the theory is formulated as a forward-time Cauchy problem. In ECT, it is natural: for an elliptic equation, changing the boundary condition on one side changes the solution throughout the domain, including what the Lorentzian reading would call the "past" side. This is the structural reason why delayed-choice phenomena are naturally accommodated (*Level A/B*).

Unified origin of coherent and geometric branches. A central structural claim of ECT: gravity and quantum coherence are not separate ingredients forced together, but two sectors of one ordered condensate. The long-wavelength sector is read geometrically (gravity); the coherent shorter-wavelength sector is read wave-mechanically (quantum). This is not ad hoc stitching — it follows from the mode decomposition of the same condensate around the same ordered background (*Level A*).

What ECT already reproduces: (i) Schrödinger equation as non-relativistic limit; (ii) Klein–Gordon from ordered branch; (iii) action quantisation through S_0 and phase compactness; (iv) decoherence and quantum–classical boundary; (v) Euclidean-to-Lorentzian measurement reading; (vi) spin discretisation from $O(3)$; (vii) PES for persistent-state selection; (viii) gravitational decoherence; (ix) unified tunnelling and entanglement; (x) cross-sector S_0 universality.

What remains open: Born rule from first principles; complete measurement theorem; Bell correlators from condensate theory; three generations; $\alpha_{\text{fs}} = 1/137$; full Yukawa hierarchy. These are open derivational targets, not failures of the framework.

26. Hilbert-space structure from reflection positivity

One of the central results of Part III is the reconstruction of Hilbert-space structure from Euclidean data, using the mathematical tool of **reflection positivity** (RP). If the Euclidean two-point function satisfies RP (a property proved for the ordered-branch correlator), the Osterwalder–Schrader reconstruction theorem (1975) guarantees the existence of a Hilbert space $\mathcal{H}$ with a positive-definite inner product, a self-adjoint Hamiltonian generating unitary time evolution, and canonical commutation relations $[q, p] = i\hbar$ (*Level A/B*).

Standard QM uses the Osterwalder–Schrader theorem in the *opposite* direction: starting from a quantum theory, it checks that the Euclidean continuation is well-defined. ECT uses it *constructively*: starting from the Euclidean condensate, it *builds* the Hilbert space. The quantum formalism is constructed, not assumed.

27. Spin as a spinorial orientation mode

In standard quantum mechanics, spin is an intrinsic quantum number defined by the representations of the Lorentz group. It is usually introduced as an axiom: some particles carry half-integer spin, period.

ECT offers a deeper geometric backdrop. Before symmetry breaking, the condensate is invariant under the four-dimensional Euclidean rotation group $O(4)$, whose connected covering group is $Spin(4) \simeq SU(2)_L \times SU(2)_R$. This group admits spinorial representations—two-component objects that require a 4π rotation rather than 2π for identity return. After $O(4) \to O(3)$ symmetry breaking and emergence of the Lorentzian sector, these spinorial representations reduce to the standard Dirac spinors of relativistic quantum mechanics.

On this reading, spin is not the literal mechanical rotation of a tiny ball in ordinary three-dimensional space. It is the spinorial orientation degree of freedom of a fermionic field compatible with the condensate symmetry structure—a geometric inheritance from the pre-Lorentzian Euclidean rotational symmetry of the ordered medium. The magnetic moment is then the response of this spinorial degree of freedom to the emergent $U(1)$ gauge sector.

Important caveat. In the current formulation, fermions are introduced as fields compatible with the condensate symmetry, not yet derived as collective condensate excitations. The electron g-factor, the fine-structure constant α_{fs}, and the hydrogenic fine structure are not yet calculated from condensate microdynamics. These remain open derivational targets (*Level Open*).

Superfluid analogy. A useful but limited analogy: in superfluid ^{3}He-A (Volovik 2003), Weyl fermions emerge as low-energy excitations near topological Fermi points—the spinorial character is a property of the quasiparticle spectrum, not an external input. ECT does not yet reach this level of derivation for fermions. The structural compatibility route developed in the main text should be understood as the present limit of what the framework can establish. Nonetheless, the superfluid analogy illustrates how an ordered background can naturally support spinorial quasiparticle structures, and therefore helps motivate why such a route is conceptually plausible for ECT even though the derivation is not yet complete.

28. Superposition, interference, exchange statistics, and the Pauli exclusion principle

Superposition (*Level A/B*): The linearised fluctuation equation in the coherent branch is linear. Solutions can be added, producing superposition without any axiom.

Interference (*Level A*): Arises from the compact-phase structure. Paths with different phase accumulations contribute to the functional integral with different factors $e^{iS/\hbar}$, producing constructive or destructive combination — the same mechanism that produces interference in a superfluid.

Indistinguishability of same-species particles: In ECT, particles of the same species are quanta of the same emergent mode sector of the same ordered medium. There is no hidden individuation label; exchange reassigns occupation rather than permuting individuated objects. The many-particle configuration space is therefore already the quotient by exchange.

Exchange statistics (*Level A/B*): Connected to the topology of path configurations in the Euclidean manifold. Paths that exchange identical particles acquire a factor $+1$ (bosons) or -1 (fermions) depending on the topological class of the exchange. The spin-statistics connection (integer spin $\leftrightarrow$ bosons, half-integer $\leftrightarrow$ fermions) is structurally motivated (*Level A/B*); a fully rigorous derivation from bare P3 remains open.

Pauli exclusion principle (*Level A/B*): For fermions, the -1 exchange factor means that configurations with two identical fermions in the same state contribute with opposite signs and cancel. Pauli exclusion is a consequence of exchange topology, not a separate axiom.

Smooth backgrounds do not weaken Pauli exclusion in ECT (*Level B / provisional no-go*): Because ECT aims to derive, rather than simply assume, the fermionic sector, one might naively expect strong gravity or dense astrophysical backgrounds to induce small violations of exclusion. The opposite is true in the smooth coherent branch. On a smooth coherent ordered branch with a local first-order fermionic description, any condensate-dependent prefactor in the kinetic term can be absorbed into a *local canonical normalisation of the fermionic field*, leaving the underlying exclusion algebra unchanged. Condensate-dependent masses, the ECT-specific preferred-direction coupling $\mu_5\bar{\psi}\gamma^A n_A\psi$, and local fermionic interaction terms all modify the dynamics of fermionic modes without altering the canonical anticommutation relation that enforces Pauli exclusion. Consequently, if the ECT fermionic reconstruction eventually closes to

the standard local Grassmann/CAR form, Pauli exclusion remains exact throughout ordinary smooth regions of the ordered branch, including the interiors of compact objects where the same local fermionic EFT still applies. Any genuinely nonstandard fermionic behaviour that might eventually arise in ECT is therefore expected to come not from ordinary strong gravity, but from more exceptional settings where the coherent-branch description itself may cease to apply: condensate branch transitions, topological or defect sectors of the ordered phase, and nonlocal sector mixing beyond the ordinary coherent branch. A physical consequence worth emphasising is that an equivalence-principle argument already constrains the form of any such residual effect: uniform acceleration can be locally removed by a change of frame, so the physically irreducible parameter is the *tidal curvature* $\mathcal{E} \sim GM/R^3$, not the acceleration itself. Even as a coarse-grained packet-level mismatch estimate (which is not itself a Pauli-violation observable), the resulting scale $(\mathcal{E}\, L^2/c_*^2)^2$ with L the fermionic overlap size remains enormously suppressed in all regimes of interest, including compact-object interiors. At the present stage the no-go statement is conditional on the completion of the ECT fermionic reconstruction programme from the exchange-topological and representation-theoretic sector to the full operator-level canonical anticommutation structure; this is tracked as an open problem in the main text (OP-Q21).

In standard QM, all four of these properties are either postulated directly or follow from operator axioms that are themselves postulated. In ECT, they arise from the geometry and topology of the ordered condensate.

29. The Principle of Euclidean Stationarity



Figure 17: The Principle of Euclidean Stationarity (PES). Three conditions must be simultaneously satisfied: the configuration is an extremal of the Euclidean action (stationary), it is resistant to environmental decoherence ($\Gamma_{\text{loop}} \ll 1$), and it belongs to a well-defined phase-topological class (compact phase with integer winding). Their intersection defines the persistent configurations — the ECT counterpart of observable quantum states. (*Level B.*)

The Principle of Euclidean Stationarity (PES) is a key organising principle of ECT (Fig. 17). It is not a new postulate but an organising principle that emerges as a consequence of three already-established ingredients: the Euclidean-first ontology (P1), the compact-phase topology (BR1), and the influence-functional decoherence framework.

Statement. The physically persistent and observable configurations are those that simultaneously: (i) are stationary or quasi-stationary extremals of the Euclidean action; (ii) maintain $\Gamma_{\text{loop}} \ll 1$ (resist decoherence); (iii) belong to an admissible phase-topological class (integer winding). Configurations failing any of these conditions decohere and are absorbed into the classical branch.

Physical consequences. PES explains *why* quantum systems settle into energy eigenstates: these are precisely the Euclidean-stationary configurations that survive environmental coupling. After the Wick rotation $w \to -ic_* t$, a Euclidean-stationary configuration with action S_E corresponds to an energy eigenstate with energy $E = S_E/\Delta w$. The ground state corresponds to the lowest-action Euclidean extremal.

Relation to the principle of least action. PES is more fundamental than the classical principle of least action. The latter is the $\Gamma_{\text{loop}} \gg 1$ limit of PES: when decoherence is strong, only the single lowest-action path survives, and classical mechanics emerges. PES thus *contains* the principle of least action as a limiting case while also explaining quantum behaviour.

Relation to discreteness. The discreteness of atomic spectra, quantised energy levels, and all integer quantum numbers follow from PES: only integer-winding, action-stationary, decoherence-resistant configurations survive. This replaces the Bohr postulate (integer quantum numbers) with a structural selection mechanism.

30. Probability, the Born rule, and measurement

In standard quantum mechanics, the Born rule $P = |\psi|^2$ is a foundational axiom. In ECT it is not postulated but traced to the reflection-positivity construction of the physical inner product: the Euclidean condensate measure, combined with the OS reconstruction, produces a Hilbert space whose inner product is quadratic in the amplitudes by construction. Once decoherence suppresses interference between branches, the remaining branches behave as classical alternatives whose weights inherit this quadratic structure. Conditional uniqueness then follows via Gleason's theorem (*Level B*): no other probability assignment is compatible with the Hilbert/projector structure.

More broadly, ECT does not merely reinterpret the axioms of quantum mechanics; it changes their logical status from foundational postulates to reconstruction targets: Hilbert space is constructed (not postulated), unitarity follows from reflection positivity, the Born rule is motivated by decoherence, and "collapse" is identified with the Euclidean-to-Lorentzian transition at $\Gamma_{\text{loop}} \sim 1$— no mystical observer is invoked.

The measurement problem — why does a quantum superposition appear to "collapse" upon measurement? — is addressed through decoherence (*Level A/B*). The interaction between the measured system, the apparatus, and the condensate environment rapidly suppresses off-diagonal terms in the density matrix, producing effective collapse. No additional collapse postulate is needed. What remains open is a fully closed mathematical theorem deriving the Born rule without any extra closure input (*Level Open*).

31. Quantum entanglement and Bell correlations

In standard QM, entanglement is a primitive property of the tensor-product Hilbert space. In ECT, it is a structural feature of the Euclidean condensate (*Level A/B*).

In the coherent branch, subsystems are not fundamental isolates — they are reduced sectors of a single ordered medium. Two subsystems sharing a common condensate with $\Gamma_{\text{loop}} \ll 1$ are correlated through the Euclidean substrate without requiring Lorentzian signalling. Four-dimensional Euclidean coherence produces non-local (in the 3D Lorentzian sense) correlations: correlations that would be surprising if one thinks in terms of 3D local causality, but are entirely natural as correlations within a single 4D Euclidean field configuration.

ECT does not violate Bell's theorem: it is not a local hidden-variable theory. The "non-locality" of entanglement is reinterpreted as common-medium correlation in the Euclidean arena, consistent with the no-signalling theorem. Two subsystems are entangled not because a hidden superluminal signal connects them, but because both are partial descriptions of one and the same coherent condensate configuration. Measurement of one part does not "send an influence" to the other; it converts the local subsystem into a decohered branch, while the other was already part of the same global configuration. The observed correlation reflects prior inseparability, not signal transfer. A full first-principles derivation of Bell-inequality-violating correlators and Tsirelson bounds remains open (*Level Open*).

Delayed-choice experiments (Wheeler) receive a natural interpretation. In the boundary-value problem of ECT, future detector settings are part of the boundary data that constrain the Euclidean field configuration globally. This is not retrocausality — it is the standard behaviour of a boundary-value problem.

32. Tunnelling

In standard QM, quantum tunnelling — the ability of a particle to traverse a classically forbidden barrier — is a consequence of wave mechanics, but its physical mechanism is mysterious. In ECT, tunnelling has a clear geometric interpretation (*Level A/B*).

In the Euclidean picture, a "classically forbidden" region corresponds to a region where the Lorentzian kinetic energy would be negative. But in the Euclidean domain, all four directions are equivalent — there is no distinction between "classically allowed" and "classically forbidden." The field configuration simply passes through the barrier in the four-dimensional Euclidean space. After the Wick rotation, this Euclidean passage manifests as exponentially suppressed but nonzero amplitude on the other side of the barrier — tunnelling.

This is the same logic used in the standard instanton calculation, but ECT gives it an ontological reading: tunnelling is not a "mysterious quantum effect" but a natural consequence of the fact that the fundamental arena is Euclidean, and classically forbidden regions are an artefact of the Lorentzian projection. At the present level, the standard WKB/instanton suppression is quantitatively recovered; the gain is primarily interpretive and geometric, not yet an established distinct numerical deviation.

33. Interactions belong to the Lorentzian branch

A conceptually important distinction in ECT is that *interaction* — in the sense of one physical system exchanging energy, momentum, or information with another — is fundamentally a Lorentzian concept (*Level A*).

In the coherent branch ($\Gamma_{\text{loop}} \ll 1$), what exists is not interaction but *common-medium correlation*: subsystems share a coherent condensate configuration, and their effective descriptions are correlated through the Euclidean substrate. No energy exchange occurs; no signal is transmitted; no "event" in the Lorentzian sense takes place.

Interaction requires decoherence: one subsystem must couple to another strongly enough that Γ_{loop} crosses the threshold ~ 1. At that point, the coherent correlations are converted into classical, Lorentzian, causal events — energy is exchanged, momentum is transferred, information is communicated. Measurement, in this picture, is the process that converts Euclidean common-medium correlation into Lorentzian interaction.

This does not mean there is no coupling at all. The influence of the medium enters through the influence functional; the decoherence kernel, medium response, and suppression of off-diagonal elements are all consequences of the medium, not of "forces" in the Lorentzian sense. The correct distinction is not "interaction versus no interaction" but "decohered Lorentzian event-like interaction versus Euclidean common-medium correlation."

This separation explains why quantum systems can exhibit non-local correlations (entanglement) without violating causality: the correlations belong to the coherent Euclidean branch, while causal signalling belongs to the decoherent Lorentzian branch.

34. Influence of future boundary conditions

Because the fundamental ECT equation is a boundary-value problem on the Euclidean manifold (not a Cauchy problem), boundary data on *all* sides of the domain constrain the solution. After the Wick rotation to Lorentzian signature, this means that future boundary conditions can constrain the present configuration.

This is not backward-in-time causation in the Lorentzian sense. It is the standard mathematical property of boundary-value problems: the solution at any interior point depends on data everywhere on the boundary. Global Euclidean boundary-value dependence is not retro-signalling within Lorentzian spacetime; no Lorentzian causality violation is implied. In ECT, this property provides a natural framework for understanding delayed-choice experiments, quantum eraser phenomena, and the apparent "retrocausal" features of the two-state vector formalism (*Level A/B*).

35. Quantum vacuum effects

The ordered condensate is not an empty backdrop but a physically active medium. Its response to boundaries, acceleration, and cosmological expansion produces three classes of observable effects:

The Casimir effect (*Level B, conditional*): In standard QED, the Casimir force between

conducting plates arises from modified vacuum fluctuation modes. In ECT, the ordered vacuum can in principle respond to boundaries that reflect its soft orientation sector, not just the photon sector. Within a provisional multi-component soft-orientation EFT with idealised condensate-reflecting Dirichlet boundaries, a naive counting argument yields an enhancement factor of 3/2 relative to QED. However, this provisional factor depends on boundary conditions that standard metallic plates do not implement (the neutral soft sector couples to matter only gravitationally), and on a three-component coordinate counting that the integrability constraint of the scalar-only basis reduces at the physical level ($N_G^{\text{phys}} = 1$ at linear order). The 3/2 ratio is therefore conditional, boundary-sensitive, and model-dependent—not a universal laboratory prediction. What is structurally robust (Level A) is the claim that a nontrivial ordered vacuum must be boundary-sensitive.

The Unruh effect (*Level A/B*): An accelerated observer in the ECT vacuum perceives a thermal bath with temperature $T = \hbar a/(2\pi c k_B)$, identical to the standard prediction. The derivation traces this to the analytic continuation from the Euclidean frame to the Rindler frame.

Cosmological particle production (*Level B*): At the $O(4) \to O(3)$ phase transition, the abrupt change in the condensate background creates real particle–antiparticle pairs. In ECT this is not the mysterious "appearance of something from nothing": modes that were "quiet" in the symmetric phase cease to be quiet in the ordered phase because the background against which "silence" is defined has changed—analogous to the production of phonons when a superfluid is rapidly cooled.

36. Decoherence and the quantum–classical boundary



Figure 18: Decoherence regimes in ECT. The information content of a quantum system evolves differently depending on the environmental coupling strength $N_{\text{eff}}\gamma$. For weak coupling (quantum regime), coherence is preserved. For strong coupling (classical regime), off-diagonal density matrix elements are exponentially suppressed. (*Level A/B.*)

ECT provides a unified influence-functional framework for decoherence (Fig. 18), treating both environmental and gravitational channels through the same mathematical structure (*Level A/B*).

Environmental decoherence: interaction with thermal photons, air molecules, phonons, and other environmental degrees of freedom.

Gravitational decoherence: interaction with the condensate orientation field n^A. This is an ECT-specific prediction: the condensate itself acts as a universal decoherence bath, even in the absence of ordinary matter environments.

The decoherence time is $\tau_{\rm dec} = 1/(N_{\rm eff}\gamma)$, where $N_{\rm eff}$ is the number of effective environmental modes and γ is the coupling rate. Numerical estimates: free electron in vacuum, $\tau_{\rm dec} \gg 10^{10}$ yr; fullerene C_{60}, $\tau_{\rm dec} \sim 10^{-6}$ s (consistent with observed interference); dust grain in air, $\tau_{\rm dec} \sim 10^{-15}$ s.

Physical picture. In ECT, decoherence is not a postulate about observers destroying wave functions. It is a physical process: the subsystem loses its coherent integrity because it couples to a large number of condensate degrees of freedom that it cannot track. No separate collapse postulate is needed; its physical content is already covered by medium-induced branch selection.

Recoherence. The reverse process—partial reassembly of coherent integrity—is not forbidden in principle but exponentially suppressed for macroscopic systems: $P_{\rm back}/P_{\rm fwd} = e^{-\Gamma_{\rm loop}}$. For mesoscopic or engineered systems, partial recovery of coherence is a physically meaningful possibility, consistent with echo-type and quantum-control protocols.

37. Quantum gravity in ECT

In standard approaches, quantum gravity — the unification of GR and QM — is one of the central unsolved problems of physics. Loop quantum gravity, string theory, causal dynamical triangulations, and other programmes attempt to quantise the gravitational field on a pre-existing spacetime or to construct spacetime from more fundamental elements.

ECT approaches quantum gravity from a fundamentally different direction (*Level A/B for the conceptual framework*). Since both gravity (geometric branch) and quantum mechanics (coherent branch) are emergent from the same ordered condensate, their unification is not a separate problem to be solved — it is built into the architecture. The graviton (spin-2 orientation mode) and the quantum amplitude (coherent phase mode) originate from the same field Φ and propagate on the same kinetic tensor K^{AB}. There is no conceptual barrier to treating them consistently, because they are not fundamentally different entities.

A concrete structural advantage over standard perturbative quantum gravity already follows from the condensate architecture. In perturbative Einstein gravity, Newton's constant has dimension $[G_N] = M^{-2}$, forcing each loop order to generate new curvature counterterms (R^2, R^3, ...) that cannot be absorbed into a finite parameter set — the theory is non-renormalizable. In ECT, G_N is a condensate parameter, and the natural dimensionless gravitational coupling $g(\mu) = \lambda(\mu)/(8\pi) \approx 4 \times 10^{-5}$ at the Planck scale remains parametrically small throughout the ordered-branch EFT. The Landau pole of the scalar closure lies at $\mu_{\rm LP} \approx M_{\rm Pl}\, e^{5.26\times 10^4}$ — far above the physical EFT threshold $m_\sigma \sim M_{\rm Pl}$ where the low-energy description already ceases to apply. Above that threshold, the correct variables are the microscopic condensate degrees of freedom, not a graviton EFT extrapolated to infinite energy. This is a structurally different starting point from quantizing a pre-existing Lorentzian metric.

The remaining open tasks: explicit graviton-loop power counting (OP-UV1), two-loop RG of the condensate–graviton system (OP-UV3), and establishing reflection positivity for the spin-2

orientation sector (OP-OS1). If the last is achieved, graviton states appear in the Hilbert space automatically from the Osterwalder–Schrader reconstruction — no separate quantisation act needed. These are open problems, not solved theorems (*Level Open*). But the conceptual problem — "how do you put gravity and quantum mechanics together?" — is dissolved by the single-medium architecture.

38. Black-hole thermodynamics and the information problem



Figure 19: The ECT picture of a black hole. The classical event horizon is replaced by a critical shell where the condensate amplitude reaches a critical value and the effective geometry transitions between regimes. Information is encoded in shell correlations. (*Structurally fixed at postulate level: no fundamental information loss. Level B: shell closure and near-horizon construction.*)

ECT reframes the black-hole horizon not as a fundamental singularity of causal structure but as an emergent interface in the condensate medium. This provides a microscopic language for why semiclassical Hawking thermality may survive while allowing controlled corrections. The present section organises the mechanism and its observable channels; it does not yet provide a complete first-principles microtheory of black holes.

If information falls into a black hole and the black hole evaporates via Hawking radiation, is the information lost? This is the information paradox — one of the deepest problems in theoretical physics.

ECT addresses it at two levels (Fig. 19).

Level A (structural): The total condensate state is never fundamentally mixed. In ECT, the apparent mixedness (thermal character) of Hawking radiation is a *reduced-state* phenomenon: the external observer's density matrix $\rho_{\text{ext}} = \text{Tr}_{\text{shell,int}}|\Psi\rangle\langle\Psi|$ is mixed because it traces over the shell and interior degrees of freedom, not because information has been destroyed. A conditional theorem establishes that exact thermality of the radiation is incompatible with

unitary evaporation: if the condensate evolves unitarily (as posited), the radiation *must* carry subtle correlations encoding the infalling information. This is a structural consequence of the postulate architecture: fundamental information destruction is not part of the basic law structure of the theory.

Level B (shell closure): The event horizon is replaced by a **critical shell** — a surface where the condensate amplitude reaches a critical value and the effective ordered-phase geometry transitions between regimes. Shell entropy scales with area: $S_{\text{shell}} \sim A/\ell_*^2$ (structural area law; the exact coefficient 1/4 is an open matching condition, not yet derived from first principles). A finite-memory shell model reproduces a Page-curve-like behaviour. These results are model-based (Level B), not yet a full first-principles derivation of black-hole evaporation. The classical singularity at $r = 0$ is replaced by a Euclidean core where the condensate transitions from the Lorentzian ordered phase back to the Euclidean disordered phase.

Three-regime structure. The near-horizon condensate separates into three regimes: (i) the exterior ordered Lorentzian regime, where standard effective geometry is valid and deviations from GR are parametrically small ($\rho_c/r_s \sim \ell_{\text{Pl}}/r_s \ll 1$); (ii) the critical shell / interface regime, where the ordered-branch amplitude transitions from its vacuum value to zero; and (iii) the pre-Lorentzian core, where the metric language ceases to be fundamental.

Topological consistency. The vacuum manifold $O(4)/O(3) \cong S^3$ has $\pi_2(S^3) = 0$, so the critical shell carries no topological charge and can form and dissolve without topological obstruction. The nontrivial sector $\pi_3(S^3) = \mathbb{Z}$ admits possible topological configurations, but the present work does not derive a stabilising Skyrme-type term or a finite-energy gravitating topological black-hole solution.

What is not yet derived. The present shell analysis does not yet derive the full interior profile, first-principles horizon thermodynamic coefficients, or a constructive Page curve.

39. Why entropy scales with area: the holographic bound as a medium property

One of the deepest lessons of black-hole physics is that the maximum information content of a region of space scales not with its volume but with the area of its boundary. This observation, first made by Bekenstein and later elevated by 't Hooft and Susskind into the *holographic principle*, has shaped much of modern theoretical physics. In string theory it appears as the AdS/CFT duality; in loop quantum gravity it emerges from a counting of quantum-geometric states. In both cases, however, area scaling is tied to specific features of the gravitational theory.

ECT offers a different perspective. In the condensate picture, area scaling is not a gravity-specific property at all. It is a **structural property of any local medium with finite microscopic resolution**.

The physical mechanism. Imagine dividing a region of the ordered condensate into "inside" and "outside" by drawing a closed surface Σ. An observer outside Σ cannot directly access the interior degrees of freedom. All that reaches the observer is mediated through a thin *boundary layer* of condensate cells straddling Σ.

Why only a boundary layer? Because the condensate is local: each cell talks only to its immediate neighbours, and the massive radial mode σ ensures that correlations decay exponentially over a

distance $\xi_{\text{cond}} \sim \ell_* \sim m_\sigma^{-1}$ (the condensate correlation length, comparable to the Planck length under the matching hypothesis). Cells deep inside the bulk have no direct line of communication with the exterior—their influence is screened by the intervening medium.

The number of independent cells in this boundary layer scales with the area of Σ divided by ℓ_*^2. Since each cell carries at most a finite amount of information, the total accessible information is bounded by an area-type formula:

$$S_{\text{red}}(\Sigma) \;\leq\; c_0 \, \frac{A(\Sigma)}{\ell_*^2}, \qquad c_0 = O(1). \tag{13}$$

This is the **general boundary-layer area bound** of ECT (*Level A for the scaling; Level B for the exact coefficient*).

Why this is not just a restatement of the QFT result. In ordinary quantum field theory, entanglement entropy across a surface also scales with area—but with an ultraviolet-divergent coefficient that depends on a regulator. The result is mathematically correct but physically incomplete: the cutoff is artificial and its value is arbitrary.

ECT upgrades this. The UV regulator is replaced by the *physical* condensate threshold $\ell_* \sim m_\sigma^{-1}$, which is determined by the medium itself. The area law therefore becomes a finite, physically meaningful entropy bound rather than a formal divergence.

Black-hole entropy as a special case. The shell-entropy result discussed earlier, $S_{\text{shell}} \sim A \ln q/\ell_*^2$, now appears as a specific instance of the general area bound. The shell is just the boundary layer closest to the horizon, and its entropy counts the condensate cells at that interface. The exact Bekenstein–Hawking coefficient $1/(4\ell_{\text{Pl}}^2)$ represents a matching condition between condensate parameters and gravitational units—not yet derived from first principles, but consistent.

Strengthening the Page-curve programme. The general area bound has an important consequence for the information paradox. If the shell entropy is area-bounded, then the shell carries only a *finite* information reservoir. This finiteness was previously assumed in the conditional no-information-loss theorem; now it follows from the general area-law mechanism. The theorem's conclusion—that exact Hawking thermality is incompatible with unitary evaporation—is therefore on firmer structural ground. A complete Page-curve derivation still requires additional dynamical ingredients (information return mechanism, purification dynamics), which remain open.

Connection to decoherence. The same boundary-layer logic that bounds entropy also underlies decoherence. When a subsystem is embedded in the condensate medium, the inaccessible degrees of freedom beyond the interface act as an effective environment. Both the entanglement entropy and the decoherence rate are controlled by the same interfacial coupling layer. This is not a coincidence but a structural feature of the reduced-description framework: entropy and decoherence are two manifestations of the same boundary-mediated tracing-out.

From area law to holography. It is important not to overclaim. ECT establishes that reduced accessible entropy scales with area rather than volume—a holographic-*style* entropy bound. What ECT does not yet establish is:

- a covariant entropy bound analogous to the Bousso bound for dynamical horizons;
- a full bulk-boundary reconstruction (a gravity/boundary duality of the AdS/CFT type);
- a complete holographic principle in the strongest reconstruction sense.

The correct summary is:

ECT provides a medium-native structural route to holographic-style entropy bounds without postulating holography as a separate principle. Full holography in the strong reconstruction sense remains open.

How ECT differs from other approaches. In AdS/CFT, area scaling arises from a duality between a gravitational theory in the bulk and a conformal field theory on its boundary. In loop quantum gravity, it arises from counting discrete quantum-geometric states. In ECT, area scaling arises from a much simpler mechanism: boundary-layer locality in a continuous ordered medium with a physical ultraviolet threshold. No extra dimensions, no gravity/gauge duality, no discrete quantum geometry. ECT suggests that the holographic character of nature is not a deep mystery of quantum gravity but a generic interface property of a local ordered medium with finite microscopic resolution.

What remains open. The exact microscopic entropy capacity per UV cell, the detailed treatment of soft and gauge-edge sectors, and the covariant generalisation of the bound to null hypersurfaces all remain open. These are important targets for the programme, but they do not weaken the already established structural area-scaling mechanism.

40. Why electric charge is quantised

One of the oldest puzzles in fundamental physics is why electric charge comes in discrete units. In standard electrodynamics, the gauge group $U(1)$ can in principle be taken either as compact (the circle) or noncompact (the real line). Only the compact choice forces charges onto a discrete lattice; the noncompact choice allows a continuous range of charges. Within the standard framework, which option nature realises is simply an empirical fact.

ECT removes this ambiguity structurally. The Abelian gauge sector is not postulated independently: it arises from the compact phase of the condensate,

$$\theta \sim \theta + 2\pi.$$

There is no consistent way to obtain a noncompact $\mathbb{R}$ gauge group from a field whose phase is topologically a circle. Compactness is therefore not an extra assumption; it is built into the condensate architecture.

The consequence is immediate. Any matter excitation transforming under this compact $U(1)$ must carry a charge that is an integer multiple of the elementary condensate charge unit e_0:

$$q = n\, e_0, \qquad n \in \mathbb{Z}.$$

Continuous charges are simply not representable. In this sense, Abelian charge quantisation is a structural consequence of the compact-phase origin of the gauge sector (*Level A*).

A second, independent route reaches the same conclusion through the canonical structure: the charge operator conjugate to the compact phase has a discrete spectrum for the same topological reason (*Level A/B*).

What this does not yet explain. This compact-$U(1)$ result gives the Abelian charge lattice $q \in \mathbb{Z} \cdot e_0$, but the *observed* charge pattern of the Standard Model is richer: quarks carry charges $\pm e/3$ and $\pm 2e/3$, while leptons carry integer multiples of e. These values arise only after electroweak embedding,

$$Q = T_3 + \frac{Y}{2},$$

combined with the colour sector. Deriving the full observed charge spectrum therefore still requires hypercharge quantisation, anomaly-consistent matter representations, and the global structure of the full gauge group. ECT already provides the Abelian foundation; the full Standard-Model charge pattern remains part of the broader gauge/flavour programme.

41. Why the post-Big-Bang Universe developed the way it did

ECT already offers a new answer to one part of the Big-Bang question: the Big Bang is not the beginning of all reality but the boundary at which an ordered Lorentzian branch emerges from a pre-temporal Euclidean regime. Before that boundary there is no physical time in the usual sense.

But this still leaves a second question: once the ordered branch has formed, why does the Universe then evolve in the way we observe?

ECT does not yet solve the full transition dynamics. In particular, it does not yet derive from first principles why the expanding branch is selected rather than a contracting one, nor how the branch boundary is completed into a fully thermal hot Big-Bang history.

What ECT *can* already do is classify the kinds of post-transition histories that are allowed or excluded once the ordered branch exists. The observed history belongs to a particularly natural class:

- ordinary radiation and matter dominate early;
- the condensate background stays nearly frozen during the early epochs, preserving standard cosmology;
- the residual condensate energy is positive and becomes important only late;
- the effective equation of state stays at or above $w = -1$, so the late acceleration is moderate.

In condensed-matter systems, when a disordered phase transitions to an ordered one, ordered domains generically *grow* — they expand into the surrounding disordered medium. In ECT the ordered Lorentzian branch emerging from the disordered Euclidean medium is structurally analogous to such a growing ordered domain. This makes an expanding ordered branch physically natural once ordered-domain formation begins, though it does not yet amount to a first-principles derivation of expanding-branch selection over the contracting alternative.

Other branches are also possible in principle. A contracting branch is kinematically admissible but physically disfavoured by the growth analogy. A future recollapse is possible if the condensate

support weakens. Strong phantom-like evolution ($w < -1$) is excluded by the no-phantom bound of the healthy ordered branch.

The theory therefore already reduces the question to a much sharper one: instead of asking only "why did the Big Bang happen?", it asks which ordered-branch cosmologies are structurally natural, which are fine-tuned, and which are excluded. The observed history sits in the "structurally natural" class. In other words, ECT already offers a geometric origin of the Big-Bang branch, but not yet a complete thermal origin of the hot radiation bath that must follow it.

42. The grandfather paradox

ECT's emergent view of time dissolves the grandfather paradox structurally.

Classical causal paradoxes of time travel arise from trying to maintain two incompatible assumptions: that one can move backward along the physical time axis, and that causal chains remain linearly irreversible. The contradiction appears because the traveller is imagined as re-entering an already fixed causal sequence from within that sequence.

In the Euclidean picture, the coordinate w is not a fundamental causal axis — it is a spatial coordinate of the underlying substrate. Moving "backward in w" is not literal motion into an already-existing past; it is motion in the deeper Euclidean medium. The segment of w that the agent leaves no longer contains that agent as an independent continuation, and worldlines can cross in the Euclidean picture without producing a contradiction of the classical type. Causal order belongs to the Lorentzian branch, where it emerges through the cone structure and where effectively irreversible macroscopic histories appear when $\Gamma_{\text{loop}} \gg 1$.

The paradox is not "solved" by imposing a rule such as "the universe forbids inconsistencies." It is dissolved by recognising that Lorentzian causal intuitions have a limited domain of validity: they apply within the emergent branch, not at the fundamental Euclidean level.

This is a conceptual statement about the ontology of time in ECT, not an operational claim that macroscopic backward time travel is physically available within the emergent Lorentzian branch.

43. Analogue laboratory programme

If ECT is correct, then superfluid liquids, Bose–Einstein condensates, and other condensed-matter systems are not merely "analogies" but realisations of the same condensate physics at a different scale: ordered medium $\to$ symmetry breaking $\to$ emergent Lorentzian kinematics $\to$ vacuum effects. Experiments in these systems are therefore tests of condensate vacuum physics at accessible scales. Key platforms include:

Superfluid ^{4}He and Bose–Einstein condensates: phase-winding quantisation, vortex dynamics, Goldstone spectrum, and the condensate-mediated correlation structure.

Acoustic black holes ("dumb holes") in BECs: analogue Hawking effect and critical-shell structure.

Condensate-reflecting boundary experiments: if a system can be designed whose boundaries genuinely reflect the soft condensate sector, it would test the boundary-sensitive vacuum response predicted by ECT. Standard metallic-plate Casimir experiments do not probe this channel.

Atom interferometry: gravitational decoherence channel predicted by ECT.

44. Conclusions of Part III

44.1. Principal results

From the coherent branch, ECT derives or structurally motivates:

1. Hilbert-space structure via reflection positivity (A/B).
2. Schrödinger and KG equations as descendants (A/B).
3. Superposition, interference, Pauli exclusion (A/B).
4. Quantisation from compact-phase winding (A).
5. Abelian charge lattice from compact $U(1)$ gauge origin (A).
6. PES as organising principle (B).
7. Born rule from RP inner product (B).
8. Measurement through decoherence (A/B).
9. Entanglement as common-medium correlation (A/B).
10. Tunnelling as Euclidean passage (A/B).
11. Casimir boundary-sensitive vacuum response (*B, conditional*).
12. Unruh temperature (A/B).
13. Gravitational decoherence channel (B).
14. No fundamental information loss in BH (*structural, postulate-level*).
15. BH entropy $\propto$ area; exact coefficient open (B).
16. General boundary-layer area-law bound (*A for scaling; B for coefficient*).
17. Page curve from shell model (B).
18. Quantum gravity dissolved by architecture (A/B).

44.2. Key equations and principles of Part III

Winding condition: $\oint \partial_A \theta \, dX^A = 2\pi n$.

Elementary loop action: $S_{\text{loop,min}} = 2\pi\hbar$.

PES: persistent configurations = extremal $\cap$ decoherence-resistant $\cap$ phase-compact.

Decoherence time: $\tau_{\text{dec}} = 1/(N_{\text{eff}}\gamma)$.

Casimir: boundary-sensitive vacuum response; provisional 3/2 factor conditional on soft-mode counting and condensate-reflecting boundaries.

BH entropy: $S = A/(4\ell_{\text{Pl}}^2)$.

General area-law bound: $S_{\text{red}}(\Sigma) \leq c_0 \, A(\Sigma)/\ell_*^2$, $c_0 = O(1)$.

44.3. Part III derivation logic



Figure 20: Part III derivation logic. The coherent branch produces: Hilbert-space structure, wave equations, Born rule, decoherence, PES, vacuum effects, BH thermodynamics, and measurement theory. Colours indicate derivation levels.

Figure 20 shows the complete derivation architecture of Part III.

Part IV

Summary, Comparisons, and Outlook

45. Architectural comparison with neighbouring programmes

ECT is not the only attempt to address the foundational questions of physics. The following comparison locates it among its closest architectural neighbours.

General Relativity + Standard Model: Time and Lorentzian structure are fundamental. Gauge group is postulated. Quantum axioms are primitive. Dark sector requires extra particles + Λ. BH information paradox is open.

Einstein–aether theory (Jacobson, 2004): Lorentzian structure is fundamental; a preferred frame is added via a postulated unit vector u^A. In ECT, the orientation n^A is derived from $\partial_A \Phi$, not postulated.

Hořava–Lifshitz gravity: Lorentz invariance is broken *explicitly* at the fundamental level. In ECT, it is broken *spontaneously* from a higher symmetry.

Volovik's "Universe in a Helium Droplet": Spacetime emerges from a condensate, but the analogy remains conceptual. ECT proposes that the condensate mechanism is the actual physical mechanism.

Causal Dynamical Triangulations: Lorentzian structure is imposed as a lattice constraint. ECT derives it from continuous SSB.

Hartle–Hawking no-boundary proposal: The Euclidean arena is a computational tool for defining path integrals. In ECT, it is the fundamental ontological entity.

MOND (Milgrom, 1983): The critical acceleration $g^\dagger$ is a universal free constant. In ECT, it is derived from condensate dynamics and is environment-dependent.

Superfluid dark matter models: Spacetime is pre-existing; a superfluid component is added as dark matter. In ECT, spacetime and the "superfluid" dynamics arise from the same condensate.

In all cases, ECT distinguishes itself by deriving structures that other theories postulate, and by unifying the dark, gravitational, and quantum sectors within a single condensate framework.

46. Distinctive features of the framework

Several features distinguish ECT from all other approaches:

Single-field origin: Everything — spacetime, gravity, quantum mechanics, gauge interactions, dark sector phenomena — emerges from one scalar field Φ.

No external time: Time is derived from $O(4) \to O(3)$ SSB, not assumed. This is stricter than most quantum gravity approaches.

"How" and "why": ECT reproduces quantum equations *and* explains why they take the form they do (compact phase structure, reflection positivity, boundary-value vs. Cauchy problem).

Honest epistemology: Every result is classified as Level A, B, or Open. No overclaiming.

Cross-sector universality of S_0: The same action scale controls phase weighting, wave kinematics, canonical normalisation, and Born weights simultaneously. This is falsifiable: if different quantum layers required different scales, ECT would be refuted.

Unified dark sector: Both "dark matter" and "dark energy" effects arise from condensate dynamics without new entities.

Common pre-geometric origin of interactions: All known interaction sectors—gravity, electromagnetism, the weak and strong forces, and the predicted director coupling—emerge from one condensate substrate. The Abelian gauge sector is derived from bundle covariance (Level A); a preferred chiral $SU(2)$ sector is structurally selected by the oriented condensate (Level B+); the colour sector requires extended internal structure. This is broader than conventional Grand Unification, which never includes gravity. The weak-sector parity asymmetry acquires a structural route from the condensate orientation, but the exact maximally parity-violating Standard-Model pattern remains an open derivation problem.

Director coupling: A fifth interaction channel—the lowest-dimension fermion–condensate operator permitted by the symmetry content—is a generic consequence of the ordered medium. It provides testable predictions (spin precession, composition-dependent effects) absent from the Standard Model.

Minimal parameter count: The entire theory is controlled by four parameters: μ^2, λ, α, β. In the canonical benchmark $(\alpha, \beta) = (2, 1)$, only **two parameters** remain: μ^2 and λ. From these two numbers, ECT derives G_N, $\hbar$, c, the entire equation hierarchy, the gauge structure, the gravitational sector, galaxy rotation curves, cosmological evolution, and the quantum formalism — without introducing any additional free parameters, phenomenological fields, or ad hoc assumptions. No other current framework achieves comparable scope with comparable economy.

47. Standard postulates reinterpreted as ECT consequences

A large fraction of structures that standard physics treats as independent axioms, postulates, empirical inputs, or fundamental constants arise in ECT as derived or structurally grounded consequences of a single scalar condensate Φ on a four-dimensional Euclidean manifold, its spontaneous $O(4) \to O(3)$ symmetry breaking, and the postulates P1–P6. The following compact table collects the most representative cases. A full version with all entries and section references is given in the technical preprint (§47 corresponds to the preprint's "Comprehensive map"; see there for the extended seven-group classification). Status labels: A = strictly derived; B = derived under stated assumptions or closures; C = conditional/target; Open = not yet established.

Standard postulates vs. ECT consequences (compact version).

Standard physics	Status there	ECT origin	Level
Foundations and spacetime			
Lorentzian signature $(-,+,+,+)$	Fundamental input	SSB $O(4) \to O(3)$ with $\alpha > \beta$ selects the hyperbolic ordered branch	A

Standard physics (cont.)	Status there	ECT origin	Level
Time as a distinguished direction	Fundamental	Emergent direction w fixed by the gradient-ordered vacuum	A
Arrow of time / second law	Open / empirical	Retarded propagation + $\Gamma_{\text{irr}} \geq 0$ from influence-functional kernel	B
Speed of light c	Fundamental constant	$c_*^2 = \beta/(\alpha - \beta)$ from ordered-branch EFT	A
Newton's constant G_N	Fundamental constant	EFT matching $G_N = c_*^4/(8\pi M_G^2)$	B
Planck's constant $\hbar$	Fundamental constant	Elementary-loop action scale S_0; identification $S_0 = \hbar$	B
Quantum mechanics postulates			
Complex Hilbert space / superposition	Axioms	Reflection-positive OS reconstruction on Gaussian coherent subsectors	A
Schrödinger equation	Axiom	Non-relativistic limit of Klein–Gordon from coherent branch	B
Canonical commutator $[\hat{q}, \hat{p}] = i\hbar$	Postulate	Compact phase + coherent-branch Poisson brackets, normalised by S_0	B
Wavefunction collapse	Postulate	Decoherence at $\Gamma_{\text{loop}} \gg 1$	B
Born rule $P = \lvert\psi\rvert^2$	Postulate	Quadratic structure from RP inner product; conditional Gleason uniqueness under (C1)–(C3) + decoherence	B
Pauli exclusion principle	Postulate	Exchange topology $\pi_1(\mathcal{M}_2) = \mathbb{Z}_2$ + OS reflection positivity	A / B
Dirac equation	Postulated fermion dynamics	Minimal first-order closure on Spin(3) spinors of the ordered branch	B
Gauge sector			
Photon as $U(1)$ gauge boson	Postulated gauge principle	Compact phase fibre; bundle covariance forces an Abelian connection and Maxwell-type leading closure	A
Preferred chiral $SU(2)$ sector	Postulated weak chirality	Structural selection from the oriented broken phase; phenomenological realisation through the electroweak condensate doublet	B+

Standard physics (cont.)	Status there	ECT origin	Level
Neutrino masses and mixing	Seesaw / Dirac / Majorana (model-dependent)	Geometric chirality selection (A/B); embedded seesaw preferred (C); $M_R^{\text{geom}} = \sqrt{\phi_0 v_2}$ as scale anchor; PMNS intact	A/B–C
$SU(3)$ colour sector	Postulated colour symmetry	Not accessible from the minimal $O(4) \to O(3)$ pattern; requires extended internal condensate structure	B
Gravity, cosmology, vacuum			
Einstein equations	Postulate	Unique rank-2 tensor compatible with broken $O(4) \to O(3)$; linearised Fierz–Pauli to Einstein–Hilbert	A/B
Anisotropic term $\Theta_{\mu\nu}[n]$	Absent in GR	Derived from ordered-branch direction-dependent closure	A(lin.)/B
Late-time acceleration (Λ)	Postulated constant	Ordered-branch background energy, no separate dark-energy substance	B
Particle dark matter (galactic)	Postulated halo	Critical ϕ-branch closure reproduces rotation curves and BTFR slope 4	B
Casimir / Unruh / Hawking sector	QFT postulates	Unified coherent-vacuum response programme	A/B
Long-standing mysteries and targets			
Fine-structure constant $\alpha_{\text{fs}} = 1/137$	Fundamental constant	Low-energy $\alpha_{\text{fs}} = q^2/(4\pi Z_A)$; UV recast as electroweak (Z_B, Z_W, θ_W) target	B/C
Three fermion generations	Empirical input	Bound-state spectral route at $y = \mathcal{O}(1)$	B/C
Proton stability	Symmetry-protected fact	Pattern III benchmark $\tau_p \sim 10^{41}$–10^{42} yr, conditional	C
Baryon asymmetry η_B	Sakharov + model	Ordering-transition + benchmark leptogenesis closure	B
Quantum gravity (UV completion)	Open problem	Graviton as condensate excitation governed by same S_0	B/C
Measurement problem	Open problem	Γ_{loop} criterion + PES + conditional Born rule	B

The compact table above is a selective extraction: it highlights the most structurally clean cases. The full preprint table adds reflection-positivity details, Fermi–Dirac/Bose–Einstein statistics, graviton structure, Klein–Gordon derivation, Crooks fluctuation theorem, Hubble evolution, S_8

lensing–growth sector, neutrino geometric masses, and cosmological-constant fine-tuning as an acknowledged open problem. The reading of the table is not that every standard postulate is already fully derived in ECT, but that each one has been reclassified from "input" to "derived or structurally targeted within a single ordered-condensate framework", with its rigour level explicitly labelled. This is the distinctive scope of the ECT programme.

48. Key equations, laws, and principles of ECT

The following equations and principles define ECT not only qualitatively but in its applied, quantitative aspects:

Foundational action: $S_E[\Phi] = \int d^4X\{\frac{1}{2}(\partial_A\Phi)^2 + V(\Phi)\}$, $V = -\frac{\mu^2}{2}\Phi^2 + \frac{\lambda}{4}\Phi^4$.

Gradient condensate: $\langle\partial_A\Phi\rangle = u_0\,\delta_{Aw}$; breaks $O(4) \to O(3)$.

Effective kinetic tensor: $K^{AB} = \beta\,\delta^{AB} - \alpha\,n^A n^B$ (uniqueness theorem).

Propagation speed: $c_* = \sqrt{\beta/(\alpha-\beta)}$ (= speed of light).

Matching relations: $G_N = c_*^4/(8\pi\phi_0^2)$; $\quad \hbar = c_*\phi_0/\sqrt{\lambda}$.

Cross-sector cone universality: $c_{\text{phase}} = c_\gamma = c_{\text{GW}} = c_*$.

Loop quantisation: $\oint \partial_A\theta\, dX^A = 2\pi n$; $\quad S_{\text{loop,min}} = 2\pi\hbar$.

Second law: $dS_{\text{ent}}/dw = N_{\text{eff}}\gamma(dq/dw)^2 \geq 0$.

Generalised Einstein equations: $G_{\mu\nu} + \Lambda_{\text{eff}}g_{\mu\nu} = 8\pi G_{\text{eff}}T_{\mu\nu} + \Theta_{\mu\nu}[n]$.

Galactic interpolation: $g(R) = \frac{1}{2}(g_N + \sqrt{g_N^2 + 4g_N g^\dagger})$.

BTFR: $v_{\text{flat}}^4 = G_N M_{\text{bar}}\, g^\dagger$ (slope 4, algebraic).

Hubble evolution: $G_{\text{eff}}(z) \approx G_N(1+z)^{2\varepsilon}$, $\varepsilon \approx 0.01$.

PES: Persistent configurations = extremal $\cap$ decoherence-resistant $\cap$ phase-compact.

Decoherence time: $\tau_{\text{dec}} = 1/(N_{\text{eff}}\gamma)$.

BH entropy: $S = A/(4\ell_{\text{Pl}}^2)$.

Casimir (conditional): $F_{\text{ECT}} = \frac{3}{2}F_{\text{QED}}$ in the provisional three-component soft-orientation EFT with condensate-reflecting boundaries; not a universal laboratory prediction.

49. Falsifiable predictions

Each prediction below is specific, quantitative, and testable:

1. **BTFR slope = 4** (A). Algebraic consequence of the ϕ-closure. A confirmed deviation would rule out the ECT galactic sector.
2. $g^\dagger \propto cH_0$ (B). Links galactic and cosmological scales. Testable by comparing $g^\dagger$ from galaxies with independently measured H_0.
3. **Fifth force:** $\eta \sim 10^{-15}$ (B). Within reach of MICROSCOPE-2 (current: $\eta < 10^{-14}$).

4. **Dark energy:** $w_0 \approx -0.83$ (B). Distinguishable from $w = -1$ by DESI/Euclid. Confirmed $w = -1$ falsifies ECT dark energy.

5. **Environmental modulation of $g^\dagger$** (B). Field vs. cluster galaxies show different $g^\dagger$ — absent in MOND.

6. **Casimir boundary-sensitive vacuum response** (B, *conditional*). The ordered vacuum must respond to boundaries that reflect its soft sector. The provisional 3/2 factor depends on naive Goldstone-coordinate counting and idealised condensate-reflecting boundaries; it is not a universal laboratory prediction for standard metallic plates.

7. **Graviton polarisation: two TT modes only** (B). Linearised condensate perturbation has no additional scalar or vector polarisations. Testable via polarisation decomposition of the stochastic GW background.

8. **CMB quadrupole deficit** (B/*Open*). A structural argument for horizon-scale suppression exists, but the quantitative prediction of $\delta C_2/C_2$ remains open.

9. **No dark matter particles** (B). Detection of a DM particle falsifies this prediction.

10. $w \neq -1$ (B). Dark energy is dynamical. Static Λ at high precision rules out the mechanism.

11. **LIV:** $\Delta t_{\rm GW} \sim 10^{-64}$ **s** (A/B). Planck-suppressed. GW170817 bound satisfied.

12. **Proton stability** (C). Conditional on the gauge/flavour completion: three selection-rule patterns are possible — full low-energy protection (proton stable), partial protection ($\Delta B = 1$ forbidden, n–$\bar{n}$ oscillations possible), or generic Planck-suppressed $\Delta B = 1$ decay ($\tau_p \sim 10^{41}$–10^{42} yr). All consistent with Super-K limit $\tau_p > 2.4 \times 10^{34}$ yr. This is a Level C closure discriminant, not a strict-core prediction.

Near-future observational discriminants. Beyond the predictions already testable with current data, several observational programmes expected to deliver results within the next decade open especially powerful discriminating channels for ECT. *(i)* Full-sky CMB polarisation missions (LiteBIRD, launch expected in the early 2030s) will test whether the primordial tensor spectrum is compatible with any viable condensate-based early-Universe completion of ECT. *(ii)* Precision ground-based CMB lensing and growth measurements (Simons Observatory, Euclid, Rubin/LSST) can probe whether the condensate-induced modifications through $G_{\rm eff}(z)$ and the anisotropic response $\Theta_{\mu\nu}[n]$ leave a detectable signature in the lensing–growth consistency relation. *(iii)* Space-based gravitational-wave detection (LISA, launch planned for 2035) will test the ECT single-cone prediction for GW propagation, search for additional GW polarisation modes beyond the two transverse-traceless modes expected in the linearised ECT graviton sector, and provide independent standard-siren determinations of H_0 that can test the benchmark condensate-driven Hubble shift. A detailed table of near-future discriminants, with explicit ECT statements and falsification verdicts, is given in the technical preprint.

ISW diagnostic. Late-time evolution of the condensate response modifies the time dependence of the gravitational potentials and hence the integrated Sachs–Wolfe (ISW) channel. Using the same benchmark parameter $\varepsilon = 0.01$ as in the Hubble-tension discussion, a semi-analytic ISW–tracer overlap proxy yields a modest enhancement of about 6% in amplitude ($A_{\rm proxy} = 1.060$)

relative to ΛCDM. An internal source-power diagnostic gives $\int \mathcal{F}^2_{\rm ECT}\, da / \int \mathcal{F}^2_{\Lambda\rm CDM}\, da = 1.108$, corresponding to an $\sim 11\%$ enhancement of the semi-analytic ISW source kernel. This benchmark is not obviously in tension with the broad range of published ISW amplitude measurements, but the comparison remains approximate: the present treatment uses an overlap proxy rather than a full C_ℓ^{Tg} observable, and a non-Limber Boltzmann-level calculation with realistic tracer windows remains future work. Any additional slip contribution from $\Theta_{\mu\nu}[n]$ is illustrated only through phenomenological ansätze and is not yet a derived ECT prediction. The full derivation, diagnostic figures, benchmark values, and limitations are documented in the ISW subsection and ISW appendix of the technical preprint.

50. Colour completion: a consistency requirement

The current ECT derivations impose a reverse-engineered constraint on the medium: any viable completion must contain an exact unbroken colour sector. A strict obstruction excludes colour-carrying condensates: any condensate transforming nontrivially under an internal group G breaks G to its stabilizer. Therefore the condensate must remain a colour singlet, and colour must reside in a separate internal sector.

A complementary algebraic obstruction reinforces this conclusion: $\dim \mathfrak{su}(3) = 8 > 6 = \dim \mathfrak{so}(4)$, so the colour algebra cannot even fit inside $\mathfrak{so}(4)$ as a subalgebra. The electroweak-type $U(1) \times SU(2)$ is the maximal gauge structure derivable from the minimal $O(4) \to O(3)$ pattern.

The primary candidate completion assumes the medium carries a local complex rank-3 internal module with $SU(3)$-structure: a Hermitian metric and a distinguished antisymmetric trilinear form ε_{abc}. Local non-rigidity of this module (extending the P5 logic) yields $SU(3)$ gauge redundancy. The preserved ε_{abc} has a direct counterpart in observed physics: it is the tensor from which colour-neutral baryonic channels are constructed.

A deeper alternative is an internal G_2-structured fibre, where $SU(3)$ arises as the stabilizer of a unit internal direction: $\mathrm{Stab}_{G_2}(u) = SU(3)$. This would explain *why* the colour group is exactly $SU(3)$, but requires exceptional internal geometry not otherwise forced by the postulates.

Neither completion yet derives full QCD: coupling g_s, confinement, quark representations, and $\theta_{\rm QCD}$ remain open.

Table 2: Key status changes under the primary colour completion.

Item	Before	After	Still open
Colour symmetry	Missing	$SU(3)$ available	Dynamical realisation
Gauge bosons	4 of 12	$8+3+1=12$ at symmetry level	Full dynamics
Quark/lepton pattern	Inaccessible	Structurally compatible	Chirality, reps
Meson/baryon channels	Inaccessible	Singlet channels available	Bound-state dynamics
α_s, Λ_{QCD}	Undefined	Definable targets	Not derived
Proton mass ($\sim 938\,\mathrm{MeV}$)	Inaccessible	Accessible in principle	Binding dynamics
α_{fs} normalisation	Quark loops undefined	Z_A strengthened	Full calculation open
Baryogenesis (η_B)	Baryon number incomplete	Colour defines baryon number	Transport open
Proton stability (τ_p)	Selection rules not formulable	Three patterns (I/II/III) formulable	Pattern identification, channels open

51. Open problems

These are stated explicitly as unsolved targets:

1. **Three fermion generations.** Qualitative routes exist (topological sectors of S^3); rigorous derivation of exactly three is not achieved.
2. **Yukawa hierarchy.** Mass differences from $m_e \sim 0.5\,\mathrm{MeV}$ to $m_t \sim 173\,\mathrm{GeV}$ not derived.
3. $\alpha = 1/137$**.** No first-principles prediction.
4. **Renormalisability.** Probable on fixed Euclidean background; not proven.
5. **Nonlinear GEE.** Structurally motivated (Level B); full mathematical rigour not achieved.
6. **Born rule.** Motivated via RP inner product (Level B); not a first-principles theorem.
7. **Bell correlators.** Medium-based entanglement established; explicit correlators not derived.
8. **BH evaporation.** Page curve from toy model (Level B); full dynamics open.
9. **Full observed electric-charge pattern.** ECT now supports an Abelian charge lattice from the compact condensate phase, but the derivation of the full Standard-Model charge spectrum, hypercharge quantisation, anomaly-consistent representations, and the global gauge-group quotient remains open.
10. **Holographic bound and strong holography.** ECT supports a medium-native area-law entropy bound (Level A for scaling; Level B for coefficient), but the exact microscopic

coefficient, the treatment of soft/gauge-edge sectors, a covariant Bousso-type formulation, and full bulk-boundary reconstruction remain open.

11. **High-z background.** Self-consistent $\phi_b(z)$ at $z \gg 1$ not derived.

12. **Gradient condensation (OP-grad).** Dynamical origin of ordered branch from bare P3: the deepest open problem.

13. **Graviton-loop power counting (OP-UV1).** Explicit one-loop graviton power counting in the ECT condensate background. Structurally expected to be tame (Landau pole far above EFT threshold), but not yet computed.

14. **Two-loop RG of the condensate–graviton system (OP-UV3).** Full RG flow including condensate–graviton mixing; needed to establish the UV behaviour beyond the scalar closure.

15. **OS reconstruction for spin-2 sector (OP-OS1).** Reflection positivity for the orientation (tensor) sector. If established, graviton states appear automatically in the Hilbert space without a separate quantisation act.

16. **Cosmological-constant fine-tuning (OP-Λ3).** The tree-level condensate minimum gives $V(\phi_0) = -\mu^4/(4\lambda)$, a large negative contribution. ECT reframes (finite loop corrections, dynamical Λ_{eff}) but does not yet eliminate the residual fine-tuning of $\sim 10^{-120}$.

17. **Proton stability and baryon-number selection rules (OP-proton).** Derivation of which baryon-number selection-rule pattern (full protection / partial $\Delta B = 2$ only / generic $\Delta B = 1$) is realised in the completed gauge/flavour sector; if generic violation, identification of decay channels and coefficient structure.

18. **Origin of neutrino masses (OP-neutrino).** The present formulation does not derive the absolute neutrino mass scale, the Dirac/Majorana character, the PMNS matrix, or the mass ordering. Two ECT-specific scale observations (M_R^{geom} and m_ν^{geom}) show that the condensate hierarchy naturally provides scale combinations close to the observed neutrino range, but neither is a first-principles derivation. Any future derivation of neutrino masses from condensate dynamics would simultaneously constrain the leptogenesis benchmark and the electroweak input status of v_2.

19. **Gauge-sector completion.** The Abelian gauge sector is already derived from the compact phase fibre. What remains open is the full internal derivation of the weak sector, the construction of the colour sector from extended internal condensate structure, and the derivation of the observed matter representations.

20. **Charge quantisation (OP-GUT3).** If the global electroweak structure group is $U(2) = (SU(2) \times U(1))/\mathbb{Z}_2$, representations satisfy $2T + Y \in 2\mathbb{Z}$. The full observed charge assignments of quarks and leptons remain open.

21. **Anomaly cancellation (OP-GUT6).** Any chiral $SU(2)$ sector must be checked for both local triangle anomalies and the global Witten $\mathbb{Z}_2$ anomaly. These become sharp consistency tests once the fermion spectrum is derived. Besides the usual local triangle anomalies, any chiral $SU(2)$ completion must also pass the global Witten $\mathbb{Z}_2$ anomaly test, which excludes an odd number of left-handed $SU(2)$ doublets—a condition that becomes a sharp consistency constraint once the emergent fermion spectrum is fixed.

22. **Weinberg angle (OP-GUT7).** $\sin^2\theta_W$ from condensate parameters: not yet derived.

23. **Director coupling coefficient (OP-GUT8).** Microscopic derivation of μ_5 from condensate microdynamics and establishment of radiative stability.

24. **Full weak-gauge derivation (OP-GRAV-WEAK).** Whether the oriented connection sector contains independent spin-1 modes beyond those fixed by the metric, which could underlie the weak interaction as a geometric rather than purely internal gauge field.

52. Complete derivation map



Figure 21: Complete ECT derivation map. The full logical architecture from the Φ-medium and the six postulates through all four Parts of the theory. Colours indicate postulates (dark), phenomenological inputs (grey), Level A (light grey), Level B (white), and Open (dashed).

Figure 21 shows the complete derivation architecture of ECT.

53. Open problems of physics addressed within ECT

Within its framework, ECT addresses several well-known open problems of modern physics:

1. **Unification of quantum mechanics and gravity.** Both emerge from the same ordered condensate. The unification is architectural, not forced (A/B).

2. **The origin of time.** Why does spacetime have signature $(-,+,+,+)$? Derived from $O(4) \to O(3)$ SSB (A).

3. **The dark matter problem.** No DM particle has been detected despite decades of searches. ECT reinterprets all observational phenomena as condensate dynamics; the minimal benchmark does not require DM particles (B).

4. **The cosmological constant problem.** QFT predicts a vacuum scale that is enormously too large. ECT replaces a rigid external Λ with dynamical condensate energy and thereby reframes part of the problem, but the residual 10^{-120} vacuum-scale fine-tuning is still open (*B/Open*).

5. **The Hubble tension.** $> 5\sigma$ discrepancy. ECT provides structural alleviation through evolving G_{eff} (*sign: A; magnitude: B*).

6. **JWST early galaxies.** Enhanced early gravity + $\Theta_{\mu\nu}$ accelerate structure formation (B).

7. **Baryon asymmetry.** A benchmark leptogenesis closure yields $\eta_B \sim 9 \times 10^{-10}$ versus the observed $\sim 6 \times 10^{-10}$; this is a closure-level result, not yet a strict first-principles theorem (B).

8. **Black-hole information paradox.** Fundamental information loss structurally forbidden (A); Page curve reproduced (B); general area-law entropy bound established (A/B).

9. **The measurement problem.** Decoherence and PES remove the need for a fundamental collapse postulate, but the full Born-rule closure remains conditional (B; unconditional closure open).

10. **The origin of quantum mechanics.** ECT provides a structural route to Hilbert space, superposition, and quantum dynamics from the condensate; the Born rule is conditionally unique via the RP/Gleason route but not yet fully first-principles derived (A/B).

11. **The arrow of time.** Derived from condensate ordering and decoherence monotonicity (A/B).

12. **The need for inflation.** Four cosmological problems resolved without inflaton, avoiding inflation's paradoxes (A/B).

13. **The hierarchy problem.** $M_{\text{Pl}}/M_{\text{EW}} \sim 10^{16}$ addressed through running coupling (*B/Open*).

14. **Why $3+1$ dimensions and why no fundamental anyons?** The $O(4) \to O(3)$ transition singles out exactly three spatial directions and one temporal seed. In $d = 3$ the exchange topology permits only the ordinary boson/fermion split; fundamental anyons are excluded in the primary branch. Standard physics postulates $d = 3+1$ without explanation (A).

15. **Why no magnetic monopoles?** $\pi_2(S^3) = 0$: topological anti-prediction. GUT frameworks generically predict monopoles; ECT excludes them (A).

16. **Vacuum stability.** In the primary condensate sector, the ordered branch is locally stable at tree level, and the standard one-loop scalar running closes the Standard-Model-type high-field metastability channel in the primary amplitude sector: here $\beta_\lambda > 0$, unlike the Higgs sector of the Standard Model where the top-Yukawa contribution drives metastability. The symmetric point $\Phi = 0$ is a hilltop, not a metastable minimum, so Coleman false-vacuum decay does not apply to the minimal primary potential. The full stability of the complete ordered branch architecture, and the question of whether a secondary embedded electroweak sector inherits this stability, remain open (*B/Open*).

17. **Quantum gravity.** ECT offers a structural route in which gravity appears as a condensate response and the graviton as an excitation of the ordered medium on a Euclidean background; the full quantum-gravity closure remains open (*A/B*).

18. **Cosmological coincidence question.** The "why now" question — why Ω_m and Ω_Λ are of comparable order today — is debated in the literature, since in pure ΛCDM the order-unity window of ρ_m/ρ_Λ already spans a sizeable low-redshift range, and its force depends on measure and observer-selection assumptions. In ECT dark energy is supplied by the slow-rolling amplitude of the same ordered condensate that generates G_N, c_* and $\hbar$, which makes the matter-to-dark-energy ratio at least a dynamically defined quantity rather than a strictly rigid external constant. In the current calibrated thawing benchmark, however, the condensate dark-energy density varies by only ≈11% between the present epoch and the asymptotic past, and the corresponding order-unity window in ρ_m/ρ_ϕ is only marginally wider than in pure ΛCDM. We therefore do not list coincidence alleviation among current ECT results; it remains an open programmatic question, distinct from the separate 10^{-120} vacuum-scale fine-tuning (*Level B/C-Open*).

19. **Massless graviton.** The spin-2 mode emerges from SSB with $m_g = 0$ protected by gauge structure, consistent with the LIGO bound $m_g < 1.2 \times 10^{-22}$ eV (*A/B*).

20. **Why are c, G_N, and $\hbar$ not independent primitives?** In ECT these constants are tied to the same condensate architecture rather than introduced as unrelated fundamental inputs (*A/B*).

21. **Why do Casimir, Unruh, Hawking, and particle-production effects belong to one family?** ECT treats them as different manifestations of a single ordered-vacuum response logic rather than as unrelated effects (*A/B*).

22. **Why does smooth gravity not weaken Pauli exclusion?** In the smooth coherent branch, condensate inhomogeneity changes local fermionic dynamics but not the exclusion algebra after canonical normalisation; any non-standard statistics would have to come from more exceptional sectors (*B*).

23. **Fine-structure constant.** After colour completion the fine-structure problem is recast as a concrete electroweak kinetic/mixing target rather than a purely formal route: external one-loop benchmark running points to moderate ultraviolet electroweak gauge-sector normalisations, structurally tied to the same condensate architecture that also underlies G_N and $\hbar$, while a full first-principles derivation remains open (*B/C*).

24. **S_8 / lensing–growth sector.** ECT has a natural route to the weak-lensing / growth sector through the tensor/orientation stress of the ordered condensate, which can produce a

mismatch between the dynamical and lensing gravitational potentials. The relevant future target is the completed set of effective functions $\mu(k, z)$, $\eta(k, z)$, and $\Sigma(k, z)$. Whether this route actually alleviates the presently mixed lensing–growth discrepancy remains open; it is a high-priority discriminator rather than a current ECT result (*Level B/C*).

The breadth of problems addressed from a single starting point is a distinctive feature of the framework. The technical preprint provides a comprehensive table of about seventy structures—from superposition, the Pauli principle, and Born-rule probability to Einstein's equations, the Casimir effect, dark energy, and rotation-curve phenomenology—that are treated as independent postulates or empirical inputs in standard physics but arise as derived or structurally grounded consequences within ECT, together with a complementary table of ECT-specific predictions and observational handles with no direct analogue in standard physics in the same form.

54. Conclusion

ECT is an attempt to place spacetime, gravity, and quantum behaviour within a single pre-spacetime framework. In its current form, some parts of this programme are structural results, some are closure-dependent effective constructions, and some remain open.

Common pre-geometric origin of interactions. ECT now supports a hierarchical picture of the interaction sectors: an inevitable Abelian gauge sector from the compact phase fibre; a structurally selected chiral $SU(2)$ sector tied to the oriented broken phase (providing a structural route to weak-sector parity asymmetry, while the exact maximally parity-violating Standard-Model pattern remains open); and a colour sector that, being algebraically inaccessible from $O(4)$ alone ($\dim \mathfrak{su}(3) > \dim \mathfrak{so}(4)$), requires extended internal condensate structure. Together with emergent gravity and the generic director coupling, ECT provides a common pre-geometric origin for all known interaction sectors—a structural notion of unification that, unlike conventional GUT programmes, also attempts to include gravity within the same condensate framework. The full Standard-Model matching (matter representations, charge assignments, anomaly cancellation, Weinberg angle) remains an open programme.

Discrete-symmetry architecture. ECT identifies a layered discrete-symmetry structure of the emergent matter sector: a geometric backbone in which Euclidean reflections of $O(4)$ play the role of emergent parity and the Euclidean antecedent of time reversal (Level A); a conditional CPT route for those local sectors where Osterwalder–Schrader reconstruction is available (Level A/B for free/quadratic sectors, Open for the full nonlinear theory); and a branch-sensitive phenomenological layer in which the preferred-direction operator $\mathcal{L}_5$ is re-classified as a CPT-odd, CP-odd, C-odd, P-even, T-even operator. Three notions that are sometimes conflated are kept explicitly distinct in ECT: the fundamental anti-unitary T on the reconstructed Hilbert space, the effective branch-level T-violation carried by $\mathcal{L}_5$, and the thermodynamic arrow of time emerging through decoherence. The antipodal branch-pair structure of the S^3 vacuum manifold admits a speculative cosmological reading (CPT-symmetric pair with matter-dominated and antimatter-dominated partners) that is recorded as a future direction (Level C) rather than as an established result. A useful structural payoff of the discrete-symmetry analysis is that the preferred-direction coupling $\mathcal{L}_5$, being a vector bilinear, does not contribute to chiral anomaly coefficients and is therefore compatible with the standard anomaly-consistency requirements of the emergent chiral gauge sector.

What is established structurally (Level A): Lorentzian signature from $O(4) \to O(3)$ SSB; universal causal cone ($c_* = c$); the equation hierarchy (Euclidean $\to$ KG $\to$ Schrödinger); arrow of time from retarded decoherence; BTFR slope exactly 4; resolution of the horizon, flatness, and monopole problems; structural impossibility of fundamental information loss in black holes.

What is obtained at effective/closure level (Level B): Newton's constant, Planck's constant, and c as condensate matching relations; one-loop vacuum stability of the primary amplitude sector (no Standard-Model-type metastability; no Coleman false vacuum in the minimal potential); generalised Einstein equations with $\Theta_{\mu\nu}$; dark energy as condensate residual energy ($w_0 \approx -0.83$); Hubble tension alleviation ($\Delta H_0 \sim +3\,\mathrm{km/s/Mpc}$); galaxy rotation curves for 165 SPARC galaxies ($\chi^2/N \approx 2.95$); reinterpretation of the dark sector as condensate dynamics; decoherence and the quantum–classical boundary; BH area entropy and Page-curve-like behaviour from the shell model; PES; boundary-sensitive Casimir vacuum response (conditional).

What remains an open programme: Three fermion generations; Yukawa hierarchy; $\alpha = 1/137$; renormalisability; Born rule as first-principles theorem; Bell correlators; complete BH evaporation; gradient condensation from bare P3; proton stability as a UV discriminant (three selection-rule patterns conditional on gauge/flavour completion). This also includes the full interacting closure of the conditional CPT route and the exact derivation of the maximally parity-violating weak-sector pattern.

Distinctive conceptual achievements (not available in standard physics): The effective metric is derived, not fundamental—the condensate is primary, geometry is secondary. Three logically distinct classes of excitations replace the uniform field-postulate structure of QFT. The axioms of QM are relocated from postulates to reconstruction targets. Event-like interaction is emergent; in the coherent branch, subsystems are related by common-medium correlation. Gravitational decoherence is derived without the Diósi–Penrose postulate (nanosphere prediction: $\tau \sim 38\,\mathrm{ms}$). A recoherence law $P_{\mathrm{back}}/P_{\mathrm{fwd}} = e^{-\Gamma_{\mathrm{loop}}}$ is predicted; standard QM has no analogue. Fock space is a reconstruction target, not a foundational axiom.

Table 3: Distinctive ECT outputs and foundational questions addressed.

Problem / Achievement	ECT mechanism	Level	Unique?
Conceptual achievements			
Metric derived	Condensate primary, geometry secondary	A/B	Yes
Three particle sectors	Condensate scalar sector / gauge-sector realisations / topological sectors	A/B	Yes
QM axioms → targets	Reconstructed, not postulated	A/B	Yes
Interaction = Lorentzian	Coherent branch: correlation only	A	Yes
Fock as reconstruction target	Variable N = condensate reconfiguration	Open	Yes (conceptual)
Foundational questions			
Why 3+1 dimensions?	$O(4) \to O(3)$	A	Yes
No monopoles	$\pi_2(S^3) = 0$	A	Anti-prediction
Vacuum stability	Tree-level local stability; one-loop primary amplitude-sector stability; no Coleman false vacuum in minimal potential	B/Open	Yes
Quantum gravity	No metric quantisation	A/B	Architecture
Massless graviton	Spin-2 from ordered branch, gauge-protected	A/B	Structural
Quantitative predictions			
Grav. decoherence	$\tau \sim 38\,\mathrm{ms}$ (nanosph.)	B	No DP postulate
Recoherence law	$P_{\mathrm{back}}/P_{\mathrm{fwd}} = e^{-\Gamma}$	B	Yes
$w_0 \approx -0.83$	Condensate residual	B	DESI/Euclid
Director coupling $\eta \sim 10^{-15}$	Fermion–condensate (director coupling)	B	MICROSCOPE-2

In its minimal closure, ECT does not require dark-matter particles, a fundamental cosmological constant, an inflaton field, separate quantum axioms, or an ad hoc collapse postulate as independent starting ingredients. With at most four bare parameters — reducible to two in the canonical benchmark — ECT addresses a scope that typically requires many independent inputs in the standard framework.

Whether ECT ultimately becomes a viable theory of nature or remains a productive conceptual laboratory, it does something valuable: it demonstrates that spacetime, gravity, and quantum mechanics *can* be derived from a common substrate, and it identifies the specific structures — gradient condensation, compact-phase topology, orientation stiffness, and the Principle of Euclidean Stationarity — that make this possible.

If even partly correct, the deepest truth about reality is not that the universe *contains* time, but that under the right conditions, the universe *creates* it.

Note. This article is a narrative companion to: *"Euclidean Condensate Theory (ECT): Emergence of Spacetime, Quantum Mechanics, and Gravity from Spontaneous $O(4)$ Symmetry Breaking"* by Valeriy Blagovidov (2026). Full preprint: doi:10.5281/zenodo.18917929.

`https://github.com/chufelo/ECT-preprint-code`

Zenodo: `https://doi.org/10.5281/zenodo.18917929`