

CME 307 / MS&E 311: Optimization

Introduction

Professor Udell

Management Science and Engineering
Stanford

April 2, 2023

Outline

What topics and why?

Grading

Quadratic optimization

Nonlinear optimization

Conic optimization

Integer programming

Convex optimization

Who's here?

What is an optimization problem?

optimization problem: nonlinear form

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, \quad i = 1, \dots, m_1 \\ & h_i(x) = 0, \quad i = 1, \dots, m_2\end{array}$$

- ▶ objective f_0
- ▶ inequality constraints f_i
- ▶ equality constraints h_i

Modularity in optimization

how to optimize:

1. model problem as a mathematical optimization problem
2. identify the properties of the problem
3. use an appropriate solver (or write a new one)

... and iterate:

- ▶ approximate the problem to make it easier
- ▶ solve a sequence of approximated problems that converge to solve the original problem
- ▶ or initialize (“warm-start”) a solver for the original problem with a solution to the approximated problem

How will you use optimization?

- ▶ model problem to match solver capabilities
- ▶ use solvers as building blocks in larger algorithms
- ▶ write your own solver for a new problem class
- ▶ write a specialized solver for a particular problem

What will you need to know to use (or develop) optimization?

- ▶ convexity
- ▶ geometry
- ▶ numerical stability
- ▶ complexity
- ▶ reliability
- ▶ stopping conditions

Optimization courses

compared to

- ▶ EE364a: more algos, less modeling
- ▶ EE364b: more nonconvex, different mix of algos
- ▶ ...

my goal: $< 30\%$ overlap with either

Topics

let's look at topics on google sheets:

https://docs.google.com/spreadsheets/d/1PXv_sFkhz5jNAA765kgHanSPoILm2fHzzRNuOOJzpHM

- ▶ what other topics should we add?

Outline

What topics and why?

Grading

Quadratic optimization

Nonlinear optimization

Conic optimization

Integer programming

Convex optimization

Who's here?

Grading

let's look at the course website:

<https://stanford-cme-307.github.io/website-2023/>

- ▶ for PhD students, the grading is designed to help (not distract from) your research
- ▶ for UG/Masters, this course may push you hard. . .

Optimization problems

important optimization problem classes:

- ▶ quadratic
- ▶ unconstrained
- ▶ finite-sum
- ▶ linear
- ▶ conic
- ▶ convex
- ▶ nonlinear (with linear or nonlinear constraints)
- ▶ mixed-integer linear programs
- ▶ black-box with (0, 1, or 2)-order oracle

Outline

What topics and why?

Grading

Quadratic optimization

Nonlinear optimization

Conic optimization

Integer programming

Convex optimization

Who's here?

Quadratic optimization

a **quadratic optimization** problem is written as

$$\begin{array}{ll} \text{minimize} & \frac{1}{2} \|Ax - b\|^2 := f_0(x) \\ \text{variable} & x \in \mathbf{R}^n \end{array}$$

where

- ▶ $A \in \mathbf{R}^{m \times n}$: matrix
- ▶ $b \in \mathbf{R}^m$: vector

how to solve?

Quadratic optimization

a **quadratic optimization** problem is written as

$$\begin{array}{ll} \text{minimize} & \frac{1}{2} \|Ax - b\|^2 := f_0(x) \\ \text{variable} & x \in \mathbf{R}^n \end{array}$$

where

- ▶ $A \in \mathbf{R}^{m \times n}$: matrix
- ▶ $b \in \mathbf{R}^m$: vector

how to solve? take gradient and set to 0:

$$\nabla f_0(x) = A^T(Ax - b) = 0$$

\implies linear system solvers also solve quadratic optimization problems

Quadratic program

a **quadratic program** is written as

$$\begin{array}{ll}\text{minimize} & \frac{1}{2}x^T Qx + c^T x \\ \text{subject to} & Ax = b \\ \text{variable} & x \in \mathbf{R}^n\end{array}$$

where

- ▶ $Q \in \mathbf{R}^{n \times n}$: symmetric positive semidefinite matrix
- ▶ $c \in \mathbf{R}^n$: vector

how to solve?

Quadratic program

a **quadratic program** is written as

$$\begin{array}{ll}\text{minimize} & \frac{1}{2}x^T Qx + c^T x \\ \text{subject to} & Ax = b \\ \text{variable} & x \in \mathbf{R}^n\end{array}$$

where

- ▶ $Q \in \mathbf{R}^{n \times n}$: symmetric positive semidefinite matrix
- ▶ $c \in \mathbf{R}^n$: vector

how to solve? reduce to quadratic optimization problem:

- ▶ (explicit) form solution set
 $\{x : Ax = b\} = \{x_0 + Vz \mid z \in \mathbf{R}^{n-m}\}$ by computing a solution $Ax_0 = b$ and a basis V for the null space of A
- ▶ (implicit) use duality to recast problem as larger linear (KKT) system

Symmetric positive semidefinite matrices

Definition

a symmetric matrix $Q \in \mathbf{R}^{n \times n}$ is **positive semidefinite** (psd) if $x^T Q x \geq 0$ for all $x \in \mathbf{R}^n$.

these matrices are so important that there are many ways to write them! for $Q \in \mathbf{R}^{n \times n}$,

$$Q \in \mathbf{S}_+^n \iff Q \succeq 0 \iff Q = Q^T, \lambda_{\min}(Q) \geq 0$$

Symmetric positive semidefinite matrices

Definition

a symmetric matrix $Q \in \mathbf{R}^{n \times n}$ is **positive semidefinite** (psd) if $x^T Q x \geq 0$ for all $x \in \mathbf{R}^n$.

these matrices are so important that there are many ways to write them! for $Q \in \mathbf{R}^{n \times n}$,

$$Q \in \mathbf{S}_+^n \iff Q \succeq 0 \iff Q = Q^T, \lambda_{\min}(Q) \geq 0$$

$Q \in \mathbf{S}_+^n$ is **symmetric positive definite** (spd) ($Q \succ 0$) if $x^T Q x > 0$ for all $x \in \mathbf{R}^n$.

Symmetric positive semidefinite matrices

Definition

a symmetric matrix $Q \in \mathbf{R}^{n \times n}$ is **positive semidefinite** (psd) if $x^T Q x \geq 0$ for all $x \in \mathbf{R}^n$.

these matrices are so important that there are many ways to write them! for $Q \in \mathbf{R}^{n \times n}$,

$$Q \in \mathbf{S}_+^n \iff Q \succeq 0 \iff Q = Q^T, \lambda_{\min}(Q) \geq 0$$

$Q \in \mathbf{S}_+^n$ is **symmetric positive definite** (spd) ($Q \succ 0$) if $x^T Q x > 0$ for all $x \in \mathbf{R}^n$.

why care about psd matrices Q ?

- ▶ least-squares objective has a psd $Q = A^T A$
- ▶ level sets of $x^T Q x$ are (bounded) ellipsoids
- ▶ the quadratic form $x^T Q x$ is a metric iff $Q \succ 0$
- ▶ eigenvalue decomp and svd coincide for psd matrices

Outline

What topics and why?

Grading

Quadratic optimization

Nonlinear optimization

Conic optimization

Integer programming

Convex optimization

Who's here?

Unconstrained smooth optimization

for $f : \mathbf{R}^n \rightarrow \mathbf{R}$ continuously differentiable,

$$\begin{array}{ll} \text{minimize} & f(x) \\ \text{variable} & x \in \mathbf{R}^n \end{array}$$

how to solve?

Unconstrained smooth optimization

for $f : \mathbf{R}^n \rightarrow \mathbf{R}$ continuously differentiable,

$$\begin{array}{ll}\text{minimize} & f(x) \\ \text{variable} & x \in \mathbf{R}^n\end{array}$$

how to solve? approximate as a quadratic problem

$$f(x) \approx f(x_0) + \nabla f(x_0)^T (x - x_0) + \frac{1}{2}(x - x_0)^T H(x_0)(x - x_0)$$

and find solution x_{quad} to the quadratic problem.

then set $x_0 \leftarrow x_{\text{quad}}$ and repeat.

Finite sum

finite sum optimization problem

$$\begin{array}{ll} \text{minimize} & \sum_{i=1}^m f_i(x) \\ \text{variable} & x \in \mathbf{R}^n \end{array}$$

key fact: can approximate gradient using gradient on **minibatch** $S \subseteq \{1, \dots, m\}$:

$$\nabla f(x) \approx \frac{1}{|S|} \sum_{i \in S} \nabla f_i(x)$$

examples:

- ▶ statistical learning (logistic regression, SVM)
- ▶ deep learning

Logistic regression

(regularized) **logistic regression** minimizes the **finite sum**

$$\begin{array}{ll} \text{minimize} & \sum_{i=1}^m \log(1 + \exp(-b_i a_i^T x)) + r(x) \\ \text{variable} & x \in \mathbf{R}^n \end{array}$$

where

- ▶ $b_i \in \{-1, 1\}$, $a_i \in \mathbf{R}^n$
- ▶ $r : \mathbf{R}^n \rightarrow \mathbf{R}$ is a **regularizer**, e.g., $\|x\|^2$ or $\|x\|_1$

Support vector machine

support vector machine (SVM) minimizes the **finite sum**

$$\begin{array}{ll} \text{minimize} & \sum_{i=1}^m \max(0, 1 - b_i a_i^T x) + \gamma \|x\|^2 \\ \text{variable} & x \in \mathbf{R}^n \end{array}$$

where $b_i \in \{-1, 1\}$ and $a_i \in \mathbf{R}^n$.

Support vector machine

support vector machine (SVM) minimizes the **finite sum**

$$\begin{array}{ll} \text{minimize} & \sum_{i=1}^m \max(0, 1 - b_i a_i^T x) + \gamma \|x\|^2 \\ \text{variable} & x \in \mathbf{R}^n \end{array}$$

where $b_i \in \{-1, 1\}$ and $a_i \in \mathbf{R}^n$. not differentiable!

Support vector machine

support vector machine (SVM) minimizes the **finite sum**

$$\begin{array}{ll} \text{minimize} & \sum_{i=1}^m \max(0, 1 - b_i a_i^T x) + \gamma \|x\|^2 \\ \text{variable} & x \in \mathbf{R}^n \end{array}$$

where $b_i \in \{-1, 1\}$ and $a_i \in \mathbf{R}^n$. not differentiable!

how to solve?

Support vector machine

support vector machine (SVM) minimizes the **finite sum**

$$\begin{array}{ll} \text{minimize} & \sum_{i=1}^m \max(0, 1 - b_i a_i^T x) + \gamma \|x\|^2 \\ \text{variable} & x \in \mathbf{R}^n \end{array}$$

where $b_i \in \{-1, 1\}$ and $a_i \in \mathbf{R}^n$. not differentiable!

how to solve?

- ▶ use **subgradient** method
- ▶ transform to **conic form**
- ▶ **smooth** the objective

Nonlinear optimization

optimization problem: nonlinear form

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq b_i, \quad i = 1, \dots, m_1 \\ & h(x) = 0 \\ \text{variable} & x \in \mathbf{R}^n\end{array}$$

- ▶ $x = (x_1, \dots, x_n)$: optimization variables
- ▶ $f_0 : \mathbf{R}^n \rightarrow \mathbf{R}$: objective function
- ▶ $f_i : \mathbf{R}^n \rightarrow \mathbf{R}, i = 1, \dots, m$: constraint functions

special case: **unconstrained optimization**

Example: process control

You are the process engineer for a desalination plant that produces drinking water. The plant has a variety of knobs, collected in vector x , that you can turn to control the process. These control, e.g., how much water is pumped into the plant, how much pressure is used to force the water through filters, and how much of each chemical is added to the water.

- ▶ $f_0(x)$: cost of water produced
- ▶ $f_i(x)$: level of each measured impurity in the water
- ▶ b_i : maximum allowable level of each impurity

Given a setting of the knobs, you can observe the cost of water produced and the levels of impurities.

What is the optimal setting of the knobs?

Oracles

an optimization **oracle** is your interface for accessing the problem data:

e.g., an oracle for $f : \mathbf{R}^n \rightarrow \mathbf{R}$ can evaluate for any $x \in \mathbf{R}^n$:

- ▶ **zero-order:** $f_0(x)$
- ▶ **first-order:** $f_0(x)$ and $\nabla f_0(x)$
- ▶ **second-order:** $f_0(x)$, $\nabla f_0(x)$, and $\nabla^2 f_0(x)$

why oracles?

- ▶ can optimize real systems based on observed output (not just models)
- ▶ can use and extend old or complex but trusted code (e.g., NASA, PDE simulations, ...)
- ▶ can prove lower bounds on the oracle complexity of a problem class

source: Nesterov 2004 "Introductory Lectures on Convex Optimization"

Nonlinear optimization: how to solve?

depends on the oracle:

- ▶ first- or second-order: approximate by a sequence of quadratic problems
- ▶ zero-order: harder, lots of methods
 - ▶ simulated annealing
 - ▶ Bayesian optimization
 - ▶ pseudo-higher-order methods, e.g., compute approximate gradient

Outline

What topics and why?

Grading

Quadratic optimization

Nonlinear optimization

Conic optimization

Integer programming

Convex optimization

Who's here?

Linear program

a **linear program** is written as

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & b - Ax \geq 0 \\ \text{variable} & x \in \mathbf{R}^n\end{array}$$

where

- ▶ $A \in \mathbf{R}^{m \times n}$: matrix
- ▶ $b \in \mathbf{R}^m$: vector
- ▶ $c \in \mathbf{R}^n$: vector

how to solve?

Linear program

a **linear program** is written as

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & b - Ax \geq 0 \\ \text{variable} & x \in \mathbf{R}^n\end{array}$$

where

- ▶ $A \in \mathbf{R}^{m \times n}$: matrix
- ▶ $b \in \mathbf{R}^m$: vector
- ▶ $c \in \mathbf{R}^n$: vector

how to solve?

- ▶ use the simplex method
- ▶ use a conic solver

Conic form

conic form optimization problem generalizes LP:

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & b - Ax \in \mathcal{K},\end{array}$$

where \mathcal{K} is a **convex cone**:

$$x \in \mathcal{K} \iff rx \in \mathcal{K} \text{ for any } r > 0.$$

examples:

- ▶ zero cone $\mathcal{K}_0 = \{0\}$
- ▶ positive orthant $\mathcal{K}_+ = \{x : x_i \geq 0, i = 1, \dots, n\}$
- ▶ second order cone $\mathcal{K}_{\text{SOC}} = \{(x, t) : \|x\|_2 \leq t\}$
- ▶ positive semidefinite (PSD) cone
 $\mathcal{K}_{\text{SDP}} = \{X : X = X^T, v^T X v \geq 0, \forall v \in \mathbf{R}^n\}$
- ▶ cartesian products of cones

Conic form: how to solve?

Morally, conic problems are solved by reducing to a nonlinear optimization problem

- ▶ barrier methods (e.g., interior point methods)
 - ▶ add a barrier term to the objective that goes to infinity when constraints are violated
- ▶ penalty methods (e.g., augmented Lagrangian methods, ADMM, ...)
 - ▶ add a penalty term to the objective that depends on a dual variable
 - ▶ adjust the dual variable to enforce constraints

Conic form example: nonnegative least squares

$$\begin{array}{ll}\text{minimize} & \|Ax - b\| \\ \text{subject to} & x \geq 0\end{array}$$



$$\begin{array}{ll}\text{minimize} & t \\ \text{subject to} & x \in \mathcal{K}_+ \\ & (Ax - b, t) \in \mathcal{K}_{\text{soc}}\end{array}$$

Conic form example: SVM

$$\begin{array}{ll} \text{minimize} & \sum_{i=1}^m \max(0, 1 - b_i a_i^T x) + \|x\|^2 \\ \text{variable} & x \in \mathbf{R}^n \end{array}$$
$$\Updownarrow$$

$$\begin{array}{ll} \text{minimize} & \sum_i s_i + t \\ \text{subject to} & s \geq \mathbf{diag}(b)Ax - 1 \\ & s \geq 0 \\ & t \geq \|x\|^2 \end{array}$$
$$\Updownarrow$$

$$\begin{array}{ll} \text{minimize} & \sum_i s_i + t \\ \text{subject to} & s + \mathbf{diag}(b)Ax - 1 \in \mathcal{K}_+ \\ & s \in \mathcal{K}_+ \\ & [t \ x^T; x \ 0_n] \in \mathcal{K}_{\text{SDP}} \end{array}$$

Conic form example: semidefinite programming

$$\begin{array}{ll}\text{minimize} & \lambda_{\max}(X) + y^T X^{-1} y \\ \text{subject to} & X \succeq 0\end{array}$$



Conic form example: semidefinite programming

$$\begin{array}{ll}\text{minimize} & \lambda_{\max}(X) + y^T X^{-1} y \\ \text{subject to} & X \succeq 0\end{array}$$

$$\Updownarrow$$

$$\begin{array}{ll}\text{minimize} & t_1 + t_2 \\ \text{subject to} & t_1 I - X \in \mathcal{K}_{\text{SDP}} \\ & \begin{bmatrix} t_2 & y^T \\ y & X \end{bmatrix} \in \mathcal{K}_{\text{SDP}}\end{array}$$

Outline

What topics and why?

Grading

Quadratic optimization

Nonlinear optimization

Conic optimization

Integer programming

Convex optimization

Who's here?

Integer programming

integer linear programming generalizes linear programming:

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & b - Ax \geq 0 \\ \text{variable} & x \in \mathbf{Z}^n\end{array}$$

variants:

- ▶ **mixed integer linear programming (MILP):**
 $x \in \mathbf{Z}^{n-m} \cup \mathbf{R}^m$
- ▶ **mixed integer nonlinear programming (MINLP):**
 $x \in \mathbf{Z}^{n-m} \cup \mathbf{R}^m$ and nonlinear objective or constraints

how to solve?

Integer programming

integer linear programming generalizes linear programming:

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & b - Ax \geq 0 \\ \text{variable} & x \in \mathbf{Z}^n\end{array}$$

variants:

- ▶ **mixed integer linear programming (MILP):**
 $x \in \mathbf{Z}^{n-m} \cup \mathbf{R}^m$
- ▶ **mixed integer nonlinear programming (MINLP):**
 $x \in \mathbf{Z}^{n-m} \cup \mathbf{R}^m$ and nonlinear objective or constraints

how to solve?

- ▶ use Gurobi, CPLEX, ...
- ▶ branch and bound and cut (*i.e.*, a sequence of LPs)
- ▶ use duality to decompose into a sequence of simpler LPs

Outline

What topics and why?

Grading

Quadratic optimization

Nonlinear optimization

Conic optimization

Integer programming

Convex optimization

Who's here?

Convex sets

Definition

A set $S \subseteq \mathbf{R}^n$ is convex if it contains every chord: for all $\theta \in [0, 1]$, $w, v \in S$,

$$\theta w + (1 - \theta)v \in S$$

Convex sets

Definition

A set $S \subseteq \mathbf{R}^n$ is convex if it contains every chord: for all $\theta \in [0, 1]$, $w, v \in S$,

$$\theta w + (1 - \theta)v \in S$$

Q: Which of these are convex?

ellipsoid, half moon

Convex functions

a function $f : \mathbf{R}^n \rightarrow \mathbf{R}$ is convex iff

Convex functions

a function $f : \mathbf{R}^n \rightarrow \mathbf{R}$ is convex iff

- **Chords.** it never lies above its chord: $\forall \theta \in [0, 1], w, v \in \mathbf{R}^n$

$$f(\theta w + (1 - \theta)v) \leq \theta f(w) + (1 - \theta)f(v)$$

Convex functions

a function $f : \mathbf{R}^n \rightarrow \mathbf{R}$ is convex iff

- ▶ **Chords.** it never lies above its chord: $\forall \theta \in [0, 1], w, v \in \mathbf{R}^n$

$$f(\theta w + (1 - \theta)v) \leq \theta f(w) + (1 - \theta)f(v)$$

- ▶ **Epigraph.** $\text{epi}(f) = \{(x, t) : t \geq f(x)\}$ is convex

Convex functions

a function $f : \mathbf{R}^n \rightarrow \mathbf{R}$ is convex iff

- ▶ **Chords.** it never lies above its chord: $\forall \theta \in [0, 1], w, v \in \mathbf{R}^n$

$$f(\theta w + (1 - \theta)v) \leq \theta f(w) + (1 - \theta)f(v)$$

- ▶ **Epigraph.** $\text{epi}(f) = \{(x, t) : t \geq f(x)\}$ is convex
- ▶ **First order condition.** if f is differentiable,

$$f(v) - f(w) \geq \nabla f(w)^\top (v - w) \quad \forall w, v \in \mathbf{R}^n$$

Convex functions

a function $f : \mathbf{R}^n \rightarrow \mathbf{R}$ is convex iff

- ▶ **Chords.** it never lies above its chord: $\forall \theta \in [0, 1], w, v \in \mathbf{R}^n$

$$f(\theta w + (1 - \theta)v) \leq \theta f(w) + (1 - \theta)f(v)$$

- ▶ **Epigraph.** $\text{epi}(f) = \{(x, t) : t \geq f(x)\}$ is convex
- ▶ **First order condition.** if f is differentiable,

$$f(v) - f(w) \geq \nabla f(w)^\top (v - w) \quad \forall w, v \in \mathbf{R}^n$$

- ▶ **Second order condition.** If f is twice differentiable, its Hessian is always psd:

$$\lambda_{\min}(\nabla^2 f(x)) \geq 0 \quad \text{for all } x \in \mathbf{R}^n$$

Convex functions

a function $f : \mathbf{R}^n \rightarrow \mathbf{R}$ is convex iff

- ▶ **Chords.** it never lies above its chord: $\forall \theta \in [0, 1], w, v \in \mathbf{R}^n$

$$f(\theta w + (1 - \theta)v) \leq \theta f(w) + (1 - \theta)f(v)$$

- ▶ **Epigraph.** $\text{epi}(f) = \{(x, t) : t \geq f(x)\}$ is convex
- ▶ **First order condition.** if f is differentiable,

$$f(v) - f(w) \geq \nabla f(w)^\top (v - w) \quad \forall w, v \in \mathbf{R}^n$$

- ▶ **Second order condition.** If f is twice differentiable, its Hessian is always psd:

$$\lambda_{\min}(\nabla^2 f(x)) \geq 0 \quad \text{for all } x \in \mathbf{R}^n$$

Q: Which of these are convex?

quadratic, l1, pwl, step, jump, logistic, logistic loss

Convex optimization

an optimization problem is convex if:

- ▶ **Geometrically:** the feasible set and the epigraph of the objective are convex
- ▶ **NLP:** the objective and inequality constraints are convex functions, and the equality constraints are affine
- ▶ **Conic:** all the cones are convex cones

Convex optimization

an optimization problem is convex if:

- ▶ **Geometrically:** the feasible set and the epigraph of the objective are convex
- ▶ **NLP:** the objective and inequality constraints are convex functions, and the equality constraints are affine
- ▶ **Conic:** all the cones are convex cones

why convex optimization?

- ▶ relatively complete theory
- ▶ efficient solvers
- ▶ conceptual tools that generalize

duality, stopping conditions, ...

Convex optimization

an optimization problem is convex if:

- ▶ **Geometrically:** the feasible set and the epigraph of the objective are convex
- ▶ **NLP:** the objective and inequality constraints are convex functions, and the equality constraints are affine
- ▶ **Conic:** all the cones are convex cones

why convex optimization?

- ▶ relatively complete theory
- ▶ efficient solvers
- ▶ conceptual tools that generalize

duality, stopping conditions, ...

- ▶ a function f is concave if $-f$ is convex
- ▶ concave maximization results in a **convex** optimization problem

Modern solvers

- ▶ algebraic modeling languages, *e.g.*
 - ▶ JuMP facilitates nonlinear and mixed integer optimization
 - ▶ CVX* (CVX, CVXPY, Convex.jl, ...) transform a problem into conic form
- ▶ and modern solvers

Optimization modeling

- ▶ Rocket control
- ▶ Power systems
- ▶ AML

Outline

What topics and why?

Grading

Quadratic optimization

Nonlinear optimization

Conic optimization

Integer programming

Convex optimization

Who's here?

Questions for you

respond at pollev.com/madeleineudell824

- ▶ program / major
- ▶ year of program
- ▶ schedule conflicts
- ▶ what other optimization courses have you taken?
- ▶ what optimization techniques do you want to learn?
- ▶ what's your favorite algorithm (or algorithmic trick)