

MS&E 125: Intro to Applied Statistics

Inference and confidence intervals

Professor Udell

Management Science and Engineering
Stanford

March 23, 2023

Announcements

Models and samples


a **statistical model** says how data is generated

example: we model a coin flip as a Bernoulli random variable with parameter θ

we can **sample** from that model to create a dataset

example: $X_1, \dots, X_n \sim \text{Bernoulli}(\theta)$

Application: process control



Inference

inference goes backwards: we use the data to make statements about the model

- ▶ also called **learning** the model or distribution

example: we can learn the parameter θ from the data

one important kind of inference is **estimation**: we use the data to estimate some parameter of the model

- ▶ e.g., a mean or variance
- ▶ **point estimate**: a single value
- ▶ **confidence interval**: a range of values likely to contain the parameter

Q: how to estimate θ from X_1, \dots, X_n ?

Inference

inference goes backwards: we use the data to make statements about the model

- ▶ also called **learning** the model or distribution

example: we can learn the parameter θ from the data

one important kind of inference is **estimation**: we use the data to estimate some parameter of the model

- ▶ e.g., a mean or variance
- ▶ **point estimate**: a single value
- ▶ **confidence interval**: a range of values likely to contain the parameter

Q: how to estimate θ from X_1, \dots, X_n ? **A:** $\hat{\theta} = \frac{1}{n} \sum_{i=1}^n X_i$

Bias of an estimator

the **bias** of estimator $\hat{\theta}$ is $\mathbb{E}[\hat{\theta}] - \theta$

- ▶ the estimator is **unbiased** if $\mathbb{E}[\hat{\theta}] = \theta$

Bias of an estimator

the **bias** of estimator $\hat{\theta}$ is $\mathbb{E}[\hat{\theta}] - \theta$

► the estimator is **unbiased** if $\mathbb{E}[\hat{\theta}] = \theta$

Q: What is the bias of $\hat{\theta}$?

Bias of an estimator

the **bias** of estimator $\hat{\theta}$ is $\mathbb{E}[\hat{\theta}] - \theta$

► the estimator is **unbiased** if $\mathbb{E}[\hat{\theta}] = \theta$

Q: What is the bias of $\hat{\theta}$?

$$\begin{aligned} \mathbf{A:} \quad \mathbb{E}[\hat{\theta}] - \theta &= \frac{1}{n} \sum_{i=1}^n X_i - \theta = \frac{1}{n} \sum_{i=1}^n (X_i - \theta) = \\ &= \frac{1}{n} \sum_{i=1}^n (X_i - \mathbb{E}[X_i]) = \frac{1}{n} \sum_{i=1}^n (X_i - \theta) = 0 \end{aligned}$$

Bias of an estimator

the **bias** of estimator $\hat{\theta}$ is $\mathbb{E}[\hat{\theta}] - \theta$

► the estimator is **unbiased** if $\mathbb{E}[\hat{\theta}] = \theta$

Q: What is the bias of $\hat{\theta}$?

A:
$$\mathbb{E}[\hat{\theta}] - \theta = \frac{1}{n} \sum_{i=1}^n X_i - \theta = \frac{1}{n} \sum_{i=1}^n (X_i - \theta) = \frac{1}{n} \sum_{i=1}^n (X_i - \mathbb{E}[X_i]) = \frac{1}{n} \sum_{i=1}^n (X_i - \theta) = 0$$

Q: Give another example of an unbiased estimator for θ .

Bias of an estimator

the **bias** of estimator $\hat{\theta}$ is $\mathbb{E}[\hat{\theta}] - \theta$

► the estimator is **unbiased** if $\mathbb{E}[\hat{\theta}] = \theta$

Q: What is the bias of $\hat{\theta}$?

$$\begin{aligned}\mathbf{A:} \quad \mathbb{E}[\hat{\theta}] - \theta &= \frac{1}{n} \sum_{i=1}^n X_i - \theta = \frac{1}{n} \sum_{i=1}^n (X_i - \theta) = \\ &= \frac{1}{n} \sum_{i=1}^n (X_i - \mathbb{E}[X_i]) = \frac{1}{n} \sum_{i=1}^n (X_i - \theta) = 0\end{aligned}$$

Q: Give another example of an unbiased estimator for θ .

$$\mathbf{A:} \quad \hat{\theta} = X_1$$

Consistency of an estimator

estimator $\hat{\theta}$ is **consistent** if $\hat{\theta} \rightarrow \theta$ as $n \rightarrow \infty$

Consistency of an estimator

estimator $\hat{\theta}$ is **consistent** if $\hat{\theta} \rightarrow \theta$ as $n \rightarrow \infty$

Q: Is the estimator $\hat{\theta} = \frac{1}{n} \sum_{i=1}^n X_i$ consistent?

Consistency of an estimator

estimator $\hat{\theta}$ is **consistent** if $\hat{\theta} \rightarrow \theta$ as $n \rightarrow \infty$

Q: Is the estimator $\hat{\theta} = \frac{1}{n} \sum_{i=1}^n X_i$ consistent?

A: The sample mean is consistent because $\frac{1}{n} n\theta \rightarrow \theta$.

Consistency of an estimator

estimator $\hat{\theta}$ is **consistent** if $\hat{\theta} \rightarrow \theta$ as $n \rightarrow \infty$

Q: Is the estimator $\hat{\theta} = \frac{1}{n} \sum_{i=1}^n X_i$ consistent?

A: The sample mean is consistent because $\frac{1}{n} n\theta \rightarrow \theta$.

Q: What about $\hat{\theta} = X_1$?

Consistency of an estimator

estimator $\hat{\theta}$ is **consistent** if $\hat{\theta} \rightarrow \theta$ as $n \rightarrow \infty$

Q: Is the estimator $\hat{\theta} = \frac{1}{n} \sum_{i=1}^n X_i$ consistent?

A: The sample mean is consistent because $\frac{1}{n} n\theta \rightarrow \theta$.

Q: What about $\hat{\theta} = X_1$?

A: The first entry is either 1 or 0 and so is not consistent unless $\theta = 0$ or $\theta = 1$.

Standard error

the standard deviation of $\hat{\theta}$ is called the **standard error**

$$\text{se}(\hat{\theta}) = \sqrt{\text{Var}[\hat{\theta}]}$$

we call the estimated standard error $\hat{\text{se}}$

Standard error

the standard deviation of $\hat{\theta}$ is called the **standard error**

$$\text{se}(\hat{\theta}) = \sqrt{\text{Var}[\hat{\theta}]}$$

we call the estimated standard error $\hat{\text{se}}$

example:

- ▶ if $\hat{\theta} = \frac{1}{n} \sum_{i=1}^n X_i$, then $\text{se} = \frac{\text{varX}}{n}$
proof: $\text{var}[\hat{\theta}] = \text{var}\left[\frac{1}{n} \sum_{i=1}^n X_i\right] = \frac{1}{n^2} \sum_{i=1}^n \text{var}[X_i] =$
 $\frac{1}{n^2} n \text{varX} = \frac{\text{varX}}{n}$

Standard error

the standard deviation of $\hat{\theta}$ is called the **standard error**

$$\text{se}(\hat{\theta}) = \sqrt{\text{Var}[\hat{\theta}]}$$

we call the estimated standard error $\hat{\text{se}}$

example:

- ▶ if $\hat{\theta} = \frac{1}{n} \sum_{i=1}^n X_i$, then $\text{se} = \frac{\text{var}\mathbf{X}}{n}$
proof: $\text{var}[\hat{\theta}] = \text{var}\left[\frac{1}{n} \sum_{i=1}^n \mathbf{X}_i\right] = \frac{1}{n^2} \sum_{i=1}^n \text{var}[\mathbf{X}_i] = \frac{1}{n^2} n \text{var}\mathbf{X} = \frac{\text{var}\mathbf{X}}{n}$
- ▶ for our coin flip model, $\hat{\text{se}} = \sqrt{\frac{\theta(1-\theta)}{n}}$
proof: $\text{var}\mathbf{X} = \theta(1 - \theta)$, so $\text{var}[\hat{\theta}] = \frac{\theta(1-\theta)}{n}$

Demo

`https://colab.research.google.com/github/
stanford-mse-125/demos/blob/main/inference.ipynb`

Outline

Normal approximation

Confidence intervals

Central limit theorem

the **central limit theorem** says that the distribution of $\hat{\theta} = \frac{1}{n} \sum_{i=1}^n X_i$ is approximately normal with mean θ and variance $\text{var}\mathbf{X}/\mathbf{n} = \text{se}(\theta)^2$

$$\frac{\hat{\theta} - \theta}{\text{se}} \rightarrow \mathcal{N}(0, 1)$$

- ▶ the distribution of $\hat{\theta}$ is approximately normal with mean θ and standard deviation se
- ▶ also true if the standard error se is replaced by the estimated standard error $\hat{\text{se}}$

assumptions:

- ▶ $\mathbb{E}X = \theta$
- ▶ $\text{var}\mathbf{X}$ is finite

Central limit theorem

the **central limit theorem** says that the distribution of $\hat{\theta} = \frac{1}{n} \sum_{i=1}^n X_i$ is approximately normal with mean θ and variance $\text{var}\mathbf{X}/\mathbf{n} = \text{se}(\theta)^2$

$$\frac{\hat{\theta} - \theta}{\text{se}} \rightarrow \mathcal{N}(0, 1)$$

- ▶ the distribution of $\hat{\theta}$ is approximately normal with mean θ and standard deviation se
- ▶ also true if the standard error se is replaced by the estimated standard error $\hat{\text{se}}$

assumptions:

- ▶ $\mathbb{E}X = \theta$
- ▶ $\text{var}\mathbf{X}$ is finite

example: for our coin flip model, $\hat{\theta} \sim \mathcal{N}(\theta, \frac{\theta(1-\theta)}{n})$

Why use a normal approximation?

- ▶ normal distribution has just two parameters
- ▶ can estimate those parameters from data
- ▶ we can use those parameters to reason about tails of distribution

define the **z-score**: the number of standard deviations away from the mean

$$z = \frac{\hat{\theta} - \theta}{\text{se}}$$

Demo

`https://colab.research.google.com/github/
stanford-mse-125/demos/blob/main/inference.ipynb`

Outline

Normal approximation

Confidence intervals

Confidence interval

a **confidence interval** is an interval C likely to contain the parameter e.g. the $(1 - \alpha)$ confidence interval satisfies

$$\mathbb{P}[\theta \in C] \geq 1 - \alpha$$

- ▶ C is a random variable: it depends on the data X_1, \dots, X_n
- ▶ θ is fixed

Confidence interval

a **confidence interval** is an interval C likely to contain the parameter e.g. the $(1 - \alpha)$ confidence interval satisfies

$$\mathbb{P}[\theta \in C] \geq 1 - \alpha$$

- ▶ C is a random variable: it depends on the data X_1, \dots, X_n
- ▶ θ is fixed

two interpretations (e.g., for 95% confidence interval C):

- ▶ if we repeat the experiment, we expect C to contain θ 100(1 - α)% of the time
- ▶ if we do a bunch of different experiments, we expect the 95% confidence interval to contain the true value of θ for 95% of the experiments

Confidence intervals: examples

opinion polls:

- ▶ $49\% \pm 3\%$ think U.S. should lift Cuba embargo.
- ▶ $38\% \pm 3\%$ think U.S. should build more nuclear power plants.
- ▶ $16\% \pm 4\%$ think St. Louis Cardinals will win the World Series.

demographic surveys:

- ▶ The average height of adult males in the United States is between 5 feet 7 inches and 5 feet 10 inches
- ▶ The average salary of software engineers in San Francisco is between \$120,000 and \$140,000

medical research:

- ▶ average weight loss of participants in a weight loss program is between 10 and 15 pounds

operations management:

How to construct confidence interval?

- ▶ use a normal approximation with analytic formula for standard error
- ▶ use a normal approximation with bootstrap estimate for standard error
- ▶ use bootstrap quantiles

Normal approximation for confidence interval

Suppose $\hat{\theta} \approx N(\theta, \text{se}^2)$. Then

$$C = \left[\hat{\theta} - z_{\alpha/2} \hat{\text{se}}, \hat{\theta} + z_{\alpha/2} \hat{\text{se}} \right]$$

is an approximate $(1 - \alpha)$ confidence interval for θ , where $z_{\alpha/2}$ is the $(1 - \alpha/2)$ quantile of the standard normal distribution.

Confidence interval for coin flip

example: for our coin flip model, we can construct a $100(1 - \alpha)\%$ confidence interval for θ as

$$\hat{\theta} \pm z_{\alpha/2} \hat{s}_e$$

where $z_{\alpha/2}$ is the $(1 - \alpha/2)$ quantile of the standard normal distribution

e.g., for $\alpha = 0.05$, we use $z_{0.025} = 1.96$

Calibration

A $(1 - \alpha)$ confidence interval is called **calibrated** if

$$\mathbb{P}[\theta \in C] \approx 1 - \alpha$$

- ▶ if confidence interval is too large, it's useless
- ▶ if confidence interval is too small, it's wrong

Proof that normal confidence interval is calibrated

Proof:

$$\begin{aligned}\Pr(\theta \in C_n) &= \Pr(\hat{\theta}_n - z_{\alpha/2}\hat{s}e \leq \theta \leq \hat{\theta}_n + z_{\alpha/2}\hat{s}e) \\ &= \Pr(-z_{\alpha/2}\hat{s}e \leq \theta - \hat{\theta}_n \leq z_{\alpha/2}\hat{s}e) \\ &= \Pr\left(-z_{\alpha/2} \leq \frac{\theta - \hat{\theta}_n}{\hat{s}e} \leq z_{\alpha/2}\right) \\ &\approx \Pr(-z_{\alpha/2} \leq Z \leq z_{\alpha/2}) \\ &= 1 - \alpha\end{aligned}$$

- ▶ $\hat{s}e$ approximates the standard deviation of $\hat{\theta}$
- ▶ the central limit theorem says that $\hat{\theta}$ is approximately normal, so the standard deviation controls the tails of the distribution

\implies CI is calibrated if number of samples n is large enough to justify approximations