

# List 12

Michał Balcerek  
Computer Simulations of Stochastic Processes  
06.2022

For a stochastic process  $(X_t)_{t \geq 0}$  starting in  $x \in \mathbb{R}$ , let us define the exit time from interval  $(a, b)$  ( $a \leq x \leq b$ ) as

$$\tau_{(a,b)}^x \stackrel{\text{def}}{=} \inf\{t \geq 0 : X_t \notin (a, b)\}. \quad (1)$$

**Problem 1.** For  $x \in (a, b)$ , estimate the expected time to exit

$$x \mapsto \mathbb{E}\tau_{(a,b)}^x$$

for the following processes:

- a) Brownian motion
- b) fractional Brownian motion with  $H = 0.3, 0.7$
- c) Lévy stable motion with  $\alpha \in \{0.5, 1, 1.5, 2\}$  and  $\beta \in \{-1, -0.5, 0, 0.5, 1\}$ .

**Hint** First focus on symmetric processes.

**Problem 2.** Calculate the probability that the exit happened through  $b$ , i.e.

$$\mathbb{P}\left(X_{\tau_{(a,b)}^x} \geq b\right),$$

(again, for various  $x \in (a, b)$ ).

**Problem 3.** (might be hard) Fit the appropriate function to  $x \mapsto \mathbb{E}\tau_{(a,b)}^x$  and find the appropriate distribution for  $x \mapsto \mathbb{P}\left(X_{\tau_{(a,b)}^x} \geq b\right)$