

Explicit upper bounds for the Erdős–Surányi function $g(n)$

Haoyu Chen*

September 4, 2026 (v5; v1: September 3, v2–v4: September 4)

Abstract

For $n \geq 1$ let $g(n)$ be the least integer g such that for every set $\{a_1 < \dots < a_n\}$ of integers greater than 1 and every $x \geq 0$, some g of the integers $x+1, \dots, x+a_n$ have product divisible by $a_1 \dots a_n$. Erdős and Surányi (1959) proved $g(3) = 4$ and $g(n) \geq (2 - o(1))n$, and Erdős asked (offering \$100) whether $g(n) \leq (2 + o(1))n$, or even $g(n) \leq 2n$ (Erdős Problem #708). To our knowledge no upper bound depending on n alone has been published. We prove

$$g(n) \leq n(\lceil \log_2 2n \rceil + \lceil \log_2 \lceil \log_2 2n \rceil \rceil + 2) \quad (n \geq 2), \quad g(n) \leq \left\lceil 48n \sqrt{\frac{81 + \ln n}{\ln(81 + \ln n)}} \right\rceil,$$

and $g(n) \leq (2 + o(1))n\sqrt{\ln n / \ln \ln n}$, and, by a different method (linear-programming duality and an elementary sieve lemma for intervals), $g(n) \leq (17 + 149e \ln(1 + \ln(8n^3)))n = O(n \ln \ln n)$. We also show that the conjectured bound holds for long intervals: if $a_n \geq 8n^3$ then $2n$ integers always suffice, so Erdős’s question is reduced to the range $a_n < 8n^3$; more generally, for every integer $k \geq 2$, kn integers suffice whenever $a_n \leq kn$ or $a_n^{k-1} \geq (kn)^{k+1}$; hence for every fixed $\delta > 0$ a bound linear in n (with a constant depending on δ) holds whenever $a_n \geq n^{1+\delta}$, e.g. $12n$ integers suffice whenever $a_n \geq n^2$. All proofs are elementary and constructive: prime-power atoms of the a_i that are large relative to a_n are placed one per interval element by a per-prime greedy rule, the small atoms are packed into bins, and a counting argument bounds the number of bins. We also record $g(4) \geq 5$ and $g(5) \geq 6$ by exact computation.

1 Introduction

Gallai observed that among any a_2 consecutive integers one can always find two whose product is a multiple of $a_1 a_2$ (for $1 < a_1 < a_2$), but that there are $a_1 < a_2 < a_3$ and a run of a_3 consecutive integers among which no three have product divisible by $a_1 a_2 a_3$. Erdős and Surányi [1] studied the general situation. Following Erdős’s formulation [2, §1], let $g(n)$ be the smallest

***Disclosure of the production pipeline.** Every mathematical statement in this note was produced by automated systems inside an autonomous research pipeline: the proofs of Theorems 1.1 and 1.2 were found in two independent runs of OpenAI GPT-5.6 at “Pro” effort through the web interface (26 and 90 minutes of reasoning respectively, on 2026-09-03) and the proofs of Theorems 1.3 and 1.4 by Qwen3.8-Max (Alibaba) through its web interface in two runs on 2026-09-03/04, all working from briefs written by an orchestrating agent (Claude Fable 5.1, Anthropic) that selected the problem, pinned the statement to Erdős’s 1992 text, verified the lower-bound construction by exact computation, re-derived every step of both proofs, commissioned independent referee runs (Claude Opus 5 agents in fresh contexts, which also brute-forced the lemmas on hundreds of thousands of instances), and drafted this document. The four constructions were implemented and exercised on more than seven million random instances, and every numerical inequality in the explicit parameter choices was checked by machine (scripts in the accompanying repository). No human supplied a definition, a lemma, a proof idea, or a correction. The problem itself remains open: the conjectured bound $g(n) \leq 2n$ is not proved here.

integer such that, whenever $1 < a_1 < \dots < a_n$ are integers and $x \geq 0$, there are integers $x < u_1 < \dots < u_{g(n)} \leq x + a_n$ with $u_1 \dots u_{g(n)} \equiv 0 \pmod{a_1 \dots a_n}$. (Since enlarging B inside the interval preserves divisibility, “ $g(n)$ integers” may be read as “at most $g(n)$ integers”; we use the latter, which also covers the case $a_n < g(n)$.) They proved $g(3) = 4$ and

$$g(n) \geq (2 - \varepsilon)n \quad (n > n_0(\varepsilon)), \quad (1)$$

the lower bound coming from the set of all products $p_i p_j$ of ℓ primes satisfying $2p_1^2 > p_\ell^2$, with the interval positioned so that a single integer u , divisible exactly once by each p_i , is the only common multiple of every $p_i p_j$ in it. Erdős [2] asked whether $g(n) < (2 + \varepsilon)n$ for large n , “or perhaps even $g(n) \leq 2n$ ”, and offered \$100 for a proof or disproof; the question is Problem #708 on the Erdős problems site [3]. Neither [2] nor [3] records an upper bound for $g(n)$, and we are not aware of one in the literature (the original paper [1], in Hungarian, was not accessible to us). Note that a bound in terms of n alone is not obvious: the trivial estimate $g \leq \sum_i \omega(a_i)$ (one interval element per prime factor of each a_i , see Lemma 2.3) depends on a_n , which is unbounded.

Throughout, $v_p(k)$ is the exponent of the prime p in k , $\omega(k)$ the number of distinct prime factors of k , and $\Lambda(t) = \lceil \log_2 t \rceil$ is the least $r \geq 0$ with $t \leq 2^r$.

Theorem 1.1. *For every $n \geq 2$,*

$$g(n) \leq F(n) := n(\Lambda(2n) + \Lambda(\Lambda(2n)) + 2) = n \log_2 n + n \log_2 \log_2 n + O(n).$$

In particular $F(2) = 10$, $F(3) = 21$, $F(4) = 28$, $F(10) = 100$, $F(100) = 1300$, $F(1000) = 17000$.

Theorem 1.2. *For every $n \geq 1$, with $L = 81 + \ln n$,*

$$g(n) \leq \left\lceil 48n\sqrt{L/\ln L} \right\rceil, \quad \text{and} \quad g(n) \leq (2 + o(1))n\sqrt{\frac{\ln n}{\ln \ln n}} \quad (n \rightarrow \infty).$$

Theorem 1.1 gives the smaller value for every n below about 10^{104} ; Theorem 1.2 is asymptotically stronger. Neither is close to the conjectured $2n$. For long intervals, however, the conjecture holds:

Theorem 1.3. *If $a_n \geq 8n^3$, then for every $x \geq 0$ some $2n$ of the integers $x + 1, \dots, x + a_n$ have product divisible by $a_1 \dots a_n$ (at most $2n$ are needed, and since $a_n \geq 2n$ the set may be padded); in fact Conjecture 7.4 below holds for such A . Consequently $g(n) \leq 8n^3$ as well, and the question whether $g(n) \leq 2n$ is equivalent to the same question restricted to $a_n < 8n^3$.*

Theorem 1.3 is the case $k = 2$ of a one-parameter family:

Theorem 1.4. *Let $k \geq 2$ be an integer and suppose that $a_n \leq kn$ or $a_n^{k-1} \geq (kn)^{k+1}$. Then for every $x \geq 0$ some (at most) kn of the integers $x + 1, \dots, x + a_n$ have product divisible by $a_1 \dots a_n$. In particular $a_n \geq 9n^2$ gives $3n$ and $a_n \geq 1024^{1/3}n^{5/3}$ gives $4n$; for every $\delta > 0$ and every integer k with $\delta(k-1) > 2$, $a_n \geq n^{1+\delta}$ gives kn as soon as $n^{\delta(k-1)-2} \geq k^{k+1}$. Taking for each n the smallest admissible k (and Theorem 1.1 where none is admissible) gives, for every $n \geq 1$: $7n$ integers suffice whenever $a_n \geq n^3$, $12n$ whenever $a_n \geq n^2$, and $15n$ whenever $a_n \geq n^{7/4}$.*

Thus for every fixed $\delta > 0$ a bound linear in n holds when $a_n \geq n^{1+\delta}$, with a constant depending on δ ; for a fixed constant C the range not covered by Theorem 1.4 is $Cn < a_n < (Cn)^{(C+1)/(C-1)}$, a full power of n . The Erdős–Surányi family, with $a_n \asymp n(\ln n)^2$, lies just above the trivial range. The gap between $n\sqrt{\ln n / \ln \ln n}$ and $2n$ is discussed in Section 7.

Finally, Section 8 reduces a linear bound to a single inequality about additive functions on intervals (Conjecture 8.1): if it holds, then $g(n) \leq 18n$ (Theorem 8.2). We prove the inequality in two special cases and show, with explicit certificates, that three natural strengthenings of it are false because intervals can contain more integers free of small primes than $[1, m]$ does. Section 9 shows that the inequality follows, by linear-programming duality, from the two-sided count $\lfloor m/d \rfloor \leq N_I(d) \leq \lfloor m/d \rfloor + 1$ of multiples alone in every instance we could compute, gives an explicit certificate for 0/1 weights valid up to about 10^{40} , and proves the 0/1 case for all m with the threshold 2 replaced by $20 \ln \ln m$ (Theorem 9.3). Section 10 extends this to arbitrary weights with threshold $1 + 149e \ln(1 + \ln m)$ (Theorem 10.1), which through the same duality gives $g(n) \leq (17 + 149e \ln(1 + \ln(8n^3)))n = O(n \ln \ln n)$ (Corollary 10.2), an improvement of Theorem 1.2 in the order of growth (though not numerically below astronomically large n).

2 The common construction

Fix $A = \{a_1 < \dots < a_n\} \subset \{2, 3, \dots\}$, put $m = a_n$, $I = \{x + 1, \dots, x + m\}$ and $P = a_1 \dots a_n$. We must find $B \subseteq I$ with $P \mid \prod_{b \in B} b$ and bound $|B|$. Note that B is a set: an interval element contributes its valuations once, however many a_i it may be a multiple of.

Lemma 2.1 (counting). *For every integer $d \leq m$: $\#\{a \in A : d \mid a\} \leq \lfloor m/d \rfloor \leq \lfloor (x + m)/d \rfloor - \lfloor x/d \rfloor = \#\{b \in I : d \mid b\}$.*

Proof. The multiples of d in $\{1, \dots, m\}$ are $d, 2d, \dots, \lfloor m/d \rfloor d$, and the a_i are distinct elements of $\{2, \dots, m\}$. Any m consecutive integers contain $\lfloor (x + m)/d \rfloor - \lfloor x/d \rfloor \geq \lfloor m/d \rfloor$ multiples of d . $\square$

Lemma 2.2 (whole interval). *P divides $\prod_{b \in I} b$. Hence if $m \leq H$ then $B = I$ has $|B| = m \leq H$.*

Proof. $P \mid m!$ because the a_i are distinct elements of $\{2, \dots, m\}$, and $\prod_{b \in I} b = m! \binom{x+m}{m}$. $\square$

Atoms, large and small. Fix an integer budget $H \geq 1$ and assume $m > H$; put $T = m/H > 1$. For $a \in A$ and $p \mid a$ call $q_p(a) = p^{v_p(a)}$ an *atom* of a ; the atoms of a are pairwise coprime and multiply to a , and a has $\omega(a)$ atoms. An atom is *large* if $q > T$ and *small* if $q \leq T$.

Lemma 2.3 (large atoms, one element each). *Let $\mathcal{L}$ be the multiset of all large atoms of all $a \in A$ (atoms of different a are distinct members, so $r = |\mathcal{L}|$ counts with multiplicity). There is a set $U \subseteq I$ with $|U| \leq r$ such that for every prime p*

$$v_p\left(\prod_{u \in U} u\right) \geq \sum_{q \in \mathcal{L}, q \text{ a power of } p} v_p(q).$$

More precisely, one can assign to each $q \in \mathcal{L}$ an element $\beta(q) \in I$ with $q \mid \beta(q)$ such that atoms of the same prime receive distinct elements; then $U = \beta(\mathcal{L})$.

Proof. Fix a prime p and list the large atoms of p as $p^{e_1}, \dots, p^{e_s}$ with $e_1 \geq \dots \geq e_s$. Each $a \in A$ has at most one atom of p , so the first j atoms come from j distinct elements of A , all divisible by p^{e_j} ; by Lemma 2.1, I contains at least j multiples of p^{e_j} . Assign the atoms in this order, giving the j -th one a multiple of p^{e_j} in I different from the $j - 1$ elements already used for p . Elements used for different primes may coincide; this only decreases $|U|$. For a fixed p the assigned elements are distinct, so their p -adic valuations add up to at least $\sum_j e_j$. $\square$

Lemma 2.4 (assembly). *Suppose the small atoms of all $a \in A$ (a multiset; atoms of different a are distinct members) have been partitioned into bins, each bin being a set of small atoms whose product f satisfies $f \leq T$ (a bin may contain atoms of different a). Let s be the total number of bins and $D = r + s$. If $D \leq H$, then there is $B \subseteq I$ with $|B| \leq D$ and $P \mid \prod_{b \in B} b$.*

Proof. Take U from Lemma 2.3, $|U| \leq r$. Order the bins $f_1, \dots, f_s$ arbitrarily. Since $f_j \leq T = m/H$ and H is an integer, Lemma 2.1 gives at least $\lfloor m/f_j \rfloor \geq H$ multiples of f_j in I . When bin j is processed, the elements to avoid are those of U and the $j - 1$ elements chosen for earlier bins, at most $r + s - 1 = D - 1 \leq H - 1$ in number; so an unused multiple v_j of f_j exists. Put $B = U \cup \{v_1, \dots, v_s\}$. By construction $U \cap \{v_1, \dots, v_s\} = \emptyset$ and the v_j are pairwise distinct, so $|B| \leq r + s = D$ and (this is the only place where the valuations of different elements are added)

$$\prod_{b \in B} b = \prod_{u \in U} u \cdot \prod_j v_j,$$

whose p -adic valuation is at least (sum over large atoms of p) + $v_p(\prod_j f_j)$, which equals $v_p(P)$ because every atom of every a is either large or lies in exactly one bin. $\square$

Thus everything reduces to producing bins and a budget H with $D = r + s \leq H$. Theorems 1.1 and 1.2 do this in two different ways.

3 Proof of Theorem 1.1

Next-fit bins. For each $a \in A$ list its small atoms in increasing order of the prime and pack them by multiplicative next-fit: keep a current bin, initially empty (product 1); if multiplying the next atom keeps the product $\leq T$, add it; otherwise close the bin and start a new one containing that atom. Every small atom is $\leq T$, so every bin has product $\leq T$.

Lemma 3.1. *If $d_1, \dots, d_u$ are the bin products of one a in the order created, then $d_j d_{j+1} > T$ for $1 \leq j < u$.*

Proof. Bin j was closed because the next atom q did not fit: $d_j q > T$. That atom is the first element of bin $j + 1$, so $d_{j+1} \geq q$. $\square$

Lemma 3.2 (budget). *Let $H, \kappa \geq 1$ be integers with*

$$(B1) \ H \geq n(2\kappa + 1), \quad (B2) \ H^{(\kappa+1)n} \leq 2^{\kappa H}. \quad (2)$$

Then the next-fit construction with threshold $T = m/H$ produces $D = r + s \leq H$ demands.

Proof. Case $m^\kappa \leq H^{\kappa+1}$. Binning never increases the count, so $D \leq W := \sum_{a \in A} \omega(a)$. Since every atom is at least 2, $2^{\omega(a)} \leq a$, hence $2^W \leq \prod_a a \leq m^n$ and $2^{\kappa W} \leq m^{\kappa n} \leq H^{(\kappa+1)n} \leq 2^{\kappa H}$ by (B2). So $W \leq H$.

Case $m^\kappa > H^{\kappa+1}$. Then $T^{\kappa+1} = m^{\kappa+1}/H^{\kappa+1} > m$. Fix $a \in A$ with ℓ large atoms and u bins $d_1, \dots, d_u$. The large atoms and the products $d_1 d_2, d_3 d_4, \dots$ of consecutive bin pairs are pairwise coprime divisors of a , each exceeding T (Lemma 3.1). If $\ell + \lfloor u/2 \rfloor \geq \kappa + 1$, the product of $\kappa + 1$ of them would exceed $T^{\kappa+1} > m \geq a$ while dividing a ; so $\ell + \lfloor u/2 \rfloor \leq \kappa$, and since $u \leq 2\lfloor u/2 \rfloor + 1$, $\ell + u \leq 2(\ell + \lfloor u/2 \rfloor) + 1 \leq 2\kappa + 1$. Summing over a , $D \leq n(2\kappa + 1) \leq H$ by (B1). $\square$

Lemma 3.3. For $n \geq 2$ let $r = \Lambda(2n)$, $s = \Lambda(r)$, $C = r + s + 2$, $H = nC = F(n)$ and $\kappa = \lfloor (C - 1)/2 \rfloor$. Then (B1) and (B2) hold.

Proof. (B1): $2\kappa + 1 \leq C$. (B2), $n \geq 3$: then $2n \geq 6$, so $r \geq 3$ and $s \geq 2$; by definition $2n \leq 2^r$, i.e. $n \leq 2^{r-1}$, and $r \leq 2^s$; also $s + 2 \leq 2^s$ for $s \geq 2$. Hence $C = r + s + 2 \leq 2^{s+1}$ and $H = nC \leq 2^{r-1}2^{s+1} = 2^{C-2}$. Moreover $C \leq 2(\kappa + 1)$, i.e. $(C - 2)(\kappa + 1) \leq \kappa C$, so $H^{(\kappa+1)n} \leq 2^{(C-2)(\kappa+1)n} \leq 2^{\kappa C n} = 2^{\kappa H}$. For $n = 2$: $r = 2$, $s = 1$, $C = 5$, $H = 10$, $\kappa = 2$, and $10^6 \leq 2^{20}$. $\square$

Proof of Theorem 1.1. Let $H = F(n)$. If $m \leq H$ use Lemma 2.2. Otherwise Lemmas 3.2 and 3.3 give $D \leq H$, and Lemma 2.4 gives B with $|B| \leq D \leq F(n)$. $\square$

4 Proof of Theorem 1.2

Here the bins are formed by *maximal merging*: start with each small atom (of any a) in its own bin and, as long as two bins have product $\leq T$, merge them. Each merge preserves the property “product $\leq T$ ” and decreases the number of bins, so the process terminates, and at termination any two bins have product $> T$.

Lemma 4.1. With $W = \sum_a \omega(a)$ and $D = r + s$ as before,

$$(i) \ D \leq W, \quad (ii) \ D \leq \frac{2n \ln m}{\ln(m/H)} + 1.$$

Proof. (i) is clear. (ii): any two bins have product $> T$, so at most one bin has product $\leq \sqrt{T}$, and that bin still has product $\geq 2 > 1$. Hence, if $s \geq 1$, $P = \prod_{\text{large}} q \cdot \prod_j f_j > T^r T^{(s-1)/2}$ strictly, and $P \leq m^n$; thus $r + (s - 1)/2 < n \ln m / \ln T$ and $D = r + s < 2n \ln m / \ln T - r + 1 \leq 2n \ln m / \ln(m/H) + 1$. If $s = 0$, $P > T^r$ gives $D = r < n \ln m / \ln T$. $\square$

Lemma 4.2. For real $X > 1$, $\sum_{p \leq X} 1/p \leq e \ln(1 + \ln X)$ (sum over primes).

Proof. Put $\sigma = 1 + 1/\ln X$. For $p \leq X$, $p^{\sigma-1} \leq X^{1/\ln X} = e$, so $1/p \leq e p^{-\sigma}$. Using $-\ln(1-t) \geq t$ and the finite Euler product, $\sum_{p \leq X} p^{-\sigma} \leq \ln \prod_{p \leq X} (1 - p^{-\sigma})^{-1} \leq \ln \zeta(\sigma)$, and $\zeta(\sigma) \leq 1 + \int_1^\infty t^{-\sigma} dt = 1 + 1/(\sigma - 1) = 1 + \ln X$. $\square$

Lemma 4.3 (average number of prime factors). For every real $t > 1$,

$$\frac{1}{n} \sum_{i=1}^n \omega(a_i) \leq \frac{\ln(m/n) + (t-1)e \ln(1 + \ln m)}{\ln t}.$$

Proof. For every positive integer k , $t^{\omega(k)} = \prod_{p|k} (1 + (t-1)) = \sum_{d|k, d \text{ squarefree}} (t-1)^{\omega(d)}$. All terms are nonnegative because $t > 1$. Summing over $k \leq m$,

$$\sum_{k \leq m} t^{\omega(k)} = \sum_{\substack{d \leq m \\ d \text{ sqfree}}} (t-1)^{\omega(d)} \left\lfloor \frac{m}{d} \right\rfloor \leq m \prod_{p \leq m} \left(1 + \frac{t-1}{p}\right) \leq m \exp\left((t-1) \sum_{p \leq m} \frac{1}{p}\right) \leq m e^{(t-1)e \ln(1 + \ln m)}$$

by Lemma 4.2. The a_i are distinct integers in $[2, m]$, so $\sum_i t^{\omega(a_i)} \leq \sum_{k \leq m} t^{\omega(k)}$, and by the AM–GM inequality $t^{\frac{1}{n} \sum_i \omega(a_i)} \leq \frac{1}{n} \sum_i t^{\omega(a_i)}$. Taking logarithms gives the claim (note $m \geq n + 1$, so $\ln(m/n) > 0$). $\square$

Lemma 4.4 (explicit parameters). *Let $L = 81 + \ln n$, $\ell = \ln L$, $R = \sqrt{L/\ell}$ and $H = \lceil 48nR \rceil$. If $m > H$ then $D < H$.*

Proof. Elementary facts: $\ell > 4$ (as $L \geq 81 > e^4$); $\ell \leq L/4$ (the function $y/4 - \ln y$ is increasing and positive for $y \geq 81$); $\ln \ell \leq \ell/2$ (as $y/2 - \ln y > 0$ is increasing for $y \geq 4$); hence $R > 2$. Put $y = \ln m$, $z = \ln H$, $d = y - z > 0$, $c = z - \ln n$. Since $48nR > 1$, $H \leq 96nR$, so $c \leq \ln 96 + \ln R < 7 + \ell/2$ (using $\ln R = (\ell - \ln \ell)/2 \leq \ell/2$) and $z < \ln n + 7 + \ell/2 < 2L$. Put $\Delta = 4\sqrt{L\ell} = 4R\ell \leq 2L$ (as $4\ell \leq L$).

Case $d \leq \Delta$. Then $y \leq 4L$. Apply Lemma 4.3 with $t = R$: $\ln R = (\ell - \ln \ell)/2 \geq \ell/4$ and $e \ln(1+y) \leq e \ln(5L) < 3(\ell + 3) < 6\ell$. Since $\ln(m/n) = d + c$,

$$\frac{W}{n} \leq \frac{d + c + (R-1) \cdot 6\ell}{\ell/4} < \frac{4R\ell + 7 + \ell/2 + 6R\ell}{\ell/4} = 40R + \frac{28}{\ell} + 2 < 45R,$$

so by Lemma 4.1(i), $D \leq W < 45nR < H$.

Case $d > \Delta$. By Lemma 4.1(ii), $D \leq 2n^{\frac{z+d}{d}} + 1 < 2n + \frac{2n \cdot 2L}{4\sqrt{L\ell}} + 1 = 2n + nR + 1 < 3nR < H$, using $R > 2$. $\square$

Proof of Theorem 1.2, explicit part. Let $H = \lceil 48nR \rceil$. If $m \leq H$ use Lemma 2.2; otherwise Lemma 4.4 gives $D < H$ and Lemma 2.4 gives $|B| \leq D < H$. $\square$

Proof of Theorem 1.2, asymptotic part. Fix $\varepsilon > 0$, put $Q_n = \sqrt{\ln n / \ln \ln n}$ and $H = \lceil (2 + \varepsilon)nQ_n \rceil$; we show $D < H$ for all $m > H$ once n is large, which suffices as before. With $y = \ln m$, $z = \ln H$, $d = y - z$, $c = z - \ln n$ we have $z = \ln n + O(\ln \ln n)$, hence $\sqrt{z/\ln z} = (1 + o(1))Q_n$ and $c = O(\ln z)$. Put $\Delta = \sqrt{z \ln z}$. If $d > \Delta$, Lemma 4.1(ii) gives $D/n \leq 2 + 2z/d + 1/n \leq (2 + o(1))\sqrt{z/\ln z}$. If $d \leq \Delta$, then $y = (1 + o(1))z$; apply Lemma 4.3 with $t = \sqrt{z}/(\ln z)^2$, so $\ln t = (\frac{1}{2} + o(1)) \ln z$: the three terms of the numerator divided by $\ln t$ are at most $(2 + o(1))\sqrt{z/\ln z}$, $O(1)$ and $O(\sqrt{z}/(\ln z)^2)$ respectively, so $D/n \leq W/n \leq (2 + o(1))\sqrt{z/\ln z}$. In both cases $D \leq (2 + \varepsilon/2)nQ_n < H$ for all $n \geq n_0(\varepsilon)$, so $g(n) \leq \lceil (2 + \varepsilon)nQ_n \rceil$ for $n \geq n_0(\varepsilon)$; as $\varepsilon > 0$ was arbitrary, $\limsup_n g(n)/(nQ_n) \leq 2$. $\square$

Remark 4.5. No attempt was made to optimise the constant 48; the same argument with the case threshold and the prime-sum estimate tightened (e.g. an explicit Mertens bound in place of Lemma 4.2) gives a single-digit constant. Theorems 1.1 and 1.2 are proved by the same construction and differ only in how the number of bins is counted; Theorem 1.3 uses the same two ingredients (per-prime placement of large atoms, private multiples for small parts) with the budget $2n$ made possible by the length of the interval.

5 Long intervals: proof of Theorem 1.3

Assume $m = a_n \geq 8n^3$ and put $C = m/(2n)$. Then $C \geq m^{2/3}$ (equivalent to $m^{1/3} \geq 2n$), $C \geq 4n^2 \geq 2n$, and, since $m \geq 8n^3 \geq 4n^2$, also $C^2 = m^2/(4n^2) \geq m$. In the language of Section 2 we take the integer budget $H = 2n$ and threshold $T = C = m/H$ (note $m > H$).

Lemma 5.1 (two-bin packing). *Let $C \geq 1$ be real and let $q_1, \dots, q_r > 1$ be integers with $q_i \leq C$ and $P = q_1 \cdots q_r \leq C^{3/2}$. Then some sub-product s of the q_i (over a subset, possibly empty or everything) satisfies $P/C \leq s \leq C$; consequently the q_i can be partitioned into two classes with products $\leq C$.*

Proof. If $P \leq C$ take $s = P$. Otherwise $1 < P/C \leq \sqrt{C}$. If some $q_i > \sqrt{C}$, take $s = q_i$: then $s \leq C$ and $s > \sqrt{C} \geq P/C$. If all $q_i \leq \sqrt{C}$, multiply them in any order and stop at the first partial product $s \geq P/C$ (this happens at the latest at $s = P$); the previous partial product, possibly the empty product $1 < P/C$, was $< P/C$, so $s < (P/C)\sqrt{C} = P/\sqrt{C} \leq C$. In every case the complementary product is $P/s \leq C$. $\square$

Lemma 5.2 (split). *Every $a \in A$ has at most one atom exceeding C , and admits a factorisation $a = d_1 d_2$ into coprime parts such that either $d_1, d_2 \leq C$, or d_1 is an atom $> C$ and $d_2 < 2n \leq C$.*

Proof. Two atoms $> C$ are coprime, so they would give $a > C^2 \geq m$, impossible. If $q = p^e > C$ is an atom, put $d_1 = q$, $d_2 = a/q < m/C = 2n$. Otherwise all atoms of a are $\leq C$ and their product $a \leq m \leq C^{3/2}$, so Lemma 5.1 splits them into two coprime classes with products $\leq C$. $\square$

Proof of Theorem 1.3. Split every $a \in A$ by Lemma 5.2; parts equal to 1 impose no condition and may be assigned to any element, so we discard them. This gives at most $2n$ demands, each either a *large* atom $> C$ or a *small* integer $\leq C$. (Lemma 2.4 with $H = 2n$ already gives $|B| \leq 2n$; we repeat the argument because Conjecture 7.4 asks for the explicit assignment.) Assign the large demands as in Lemma 2.3: for each prime p , the large atoms of p (one per a) receive distinct multiples of the required powers of p ; different primes may share elements. Let U be the set of elements used, $|U| \leq$ number of large demands. Now assign the small demands one at a time: a small demand $d \leq C = m/(2n)$ has at least $\lfloor m/d \rfloor \geq 2n$ multiples in I by Lemma 2.1 ($2n$ is an integer), while the forbidden elements (U together with the elements already given to earlier small demands) number at most $2n - 1$, because there are at most $2n$ demands in all and the current one is not yet placed; so d receives a multiple of its own, outside U and different from all other small representatives. Let B be the set of all elements used, $|B| \leq 2n$. At an element of U the demands present are prime powers of distinct primes, each dividing the element; at every other element of B exactly one demand d is present and d divides the element. Hence the product of the demands at any $b \in B$ divides b , and since the demands of each a multiply to a , $P \mid \prod_{b \in B} b$. This is exactly an assignment as in Conjecture 7.4. Finally, if $a_n < 8n^3$ then Lemma 2.2 gives a solution with at most $a_n < 8n^3$ elements, so $g(n) \leq 8n^3$. $\square$

6 The k -split lemma: proof of Theorem 1.4

Lemma 6.1 (k -split). *Let $k \geq 1$ and let $H \leq m$ be positive integers. Let Q be a finite multiset of integers > 1 with product A such that $A^2 H^{k+1} \leq m^{k+1}$. Then Q can be partitioned into at most k pieces (none if $Q = \emptyset$), each of which is either a single element of Q or a bin: a nonempty $F \subseteq Q$ whose product f satisfies $fH \leq m$.*

Proof. Induction on k ; if $Q = \emptyset$ there is nothing to do, so let $Q \neq \emptyset$. For $k = 1$ the hypothesis reads $A^2 H^2 \leq m^2$, so $AH \leq m$ and Q itself is one bin. Let $k \geq 2$.

Case 1: some $q \in Q$ has $q^2 H > m$. Make q a singleton piece. The product $A' = A/q$ of the rest satisfies $A'^2 H^k = A^2 H^{k+1} / (q^2 H) < m^{k+1} / m = m^k$, so the induction hypothesis (with $k - 1$) splits the rest into at most $k - 1$ pieces.

Case 2: $q^2 H \leq m$ for all $q \in Q$. Build one bin greedily: start with $f = 1$ and, while Q is not exhausted and $f^2 H < m$, multiply the next element into f . The invariant $fH \leq m$ holds at the start ($H \leq m$) and is preserved, because before a step $f^2 H < m$ and $q^2 H \leq m$ give $(fqH)^2 < m^2$. The first element is always absorbed unless $H = m$, in which case the hypothesis forces $A = 1$, i.e. $Q = \emptyset$; so the bin is nonempty. If the process exhausts Q , all of Q is one bin.

Otherwise it stops with $f^2H \geq m$; then the product $A' = A/f$ of the remaining elements satisfies $A'^2H^k = A^2H^{k+1}/(f^2H) \leq m^k$, and the induction hypothesis splits them into at most $k - 1$ further pieces. $\square$

Proof of Theorem 1.4. Put $H = kn$. If $m \leq H$, Lemma 2.2 gives $B = I$ with $|B| = m \leq kn$. Otherwise $m > H$ and $m^{k-1} \geq H^{k+1}$; for every $a \in A$, $a^2H^{k+1} \leq m^2 \cdot m^{k-1} = m^{k+1}$. Apply Lemma 6.1 to the atoms of each $a \in A$ (pairwise coprime, product a , all $\leq m$): each a splits into at most k pieces, hence at most $kn = H$ pieces in all. Call a singleton piece q *large* if $qH > m$; every other piece (a bin, or a singleton with $qH \leq m$) is a bin f with $fH \leq m$ in the sense of Section 2 with threshold $T = m/H$. The large pieces are atoms of distinct elements of A for each fixed prime (an element of A has one atom per prime), so Lemma 2.3 applies to them, and Lemma 2.4 (with r = number of large pieces, s = number of bins, $r + s \leq H$) gives $B \subseteq I$ with $|B| \leq H = kn$ and $P \mid \prod_{b \in B} b$; note that the p -adic valuation of a bin comes from the single p -atom of the element of A it was cut from. For the corollaries: $k = 2$ is $m \geq 8n^3$; $k = 3$ is $m^2 \geq 81n^4$; $k = 4$ is $m^3 \geq 1024n^5$; for $\delta > 0$ and $\delta(k-1) > 2$, $m \geq n^{1+\delta}$ gives $m^{k-1} \geq n^{(1+\delta)(k-1)} \geq k^{k+1}n^{k+1}$ as soon as $n^{\delta(k-1)-2} \geq k^{k+1}$. The last sentence of the theorem is a finite computation: for each exponent e and each n take the least $k \geq 2$ with $e(k-1)\ln n \geq (k+1)\ln(kn)$, or $F(n)/n$ from Theorem 1.1 if there is none; for $e = 3, 2, 7/4$ some $k \leq 10$ is admissible for all $n \geq 5, n \geq 38, n \geq 207$ respectively, and the maximum over n is attained at $n = 4, 36, 165$ with the values 7, 12, 15; the script is in the repository. $\square$

Remark 6.2. Lemma 6.1 is the sharp form of the same atom-binning that underlies Theorem 1.1 (which needs no hypothesis on a_n); what is new here is the exact threshold $\sqrt{m/H}$ and the explicit family of thresholds $a_n^{k-1} \geq (kn)^{k+1}$. The threshold is essentially sharp for the lemma: $k+1$ atoms each with $q^2H > m$ cannot share bins and violate the hypothesis. With $k = 2$ the lemma is Lemma 5.1 in a different normalisation, so Section 5 is the case $k = 2$; it was kept because it also exhibits the explicit assignment required by Conjecture 7.4. Choosing k as a function of m/n does not improve the global bounds of Theorems 1.1 and 1.2: for $m = ne^{\sqrt{\ln n \ln \ln n}}$ both routes give about $n\sqrt{\ln n / \ln \ln n}$.

7 Small values, obstructions and remarks

Proposition 7.1. $g(4) \geq 5$ and $g(5) \geq 6$.

Proof. Let $A = \{10, 12, 15, 20\}$ and $x = 50$, so $I = \{51, \dots, 70\}$ and $P = 2^5 \cdot 3^2 \cdot 5^3$. No element of I is divisible by 25, so a solution contains at least three of the multiples 55, 60, 65, 70 of 5; any set S of multiples of 5 in I has $v_2(\prod S) \leq 3$ and $v_3(\prod S) \leq 1$. Hence a further element with $v_2 \geq 2$ and $v_3 \geq 1$, i.e. a multiple of 12, is needed; the only multiple of 12 in I is 60, and using it still leaves $v_2 \leq 3 < 5$. So at least five elements are needed, and $\{55, 63, 64, 65, 70\}$ works. For $A = \{5, 10, 12, 15, 20\}$ and $x = 50$ we have $P = 2^5 \cdot 3^2 \cdot 5^4$, so all four multiples of 5 are forced, supplying $v_2 = 3$ and $v_3 = 1$; two more elements are then needed for 2^2 and 3 (no element of I other than 60 is divisible by 12), and $\{55, 60, 63, 64, 65, 70\}$ works. Both minima were also confirmed by an exhaustive dynamic programme over capped valuation vectors, which further shows that no other x gives a larger minimum for these two sets. $\square$

Remark 7.2 (why the obvious approaches fail). (a) Choosing a multiple of each a_i and multiplying does not work, because the chosen multiples may coincide and a shared element contributes its valuations once; in the Erdős–Surányi family every a_i has the same unique multiple u . (b) One

cannot in general match the $a_i \leq m/2$ to *distinct* multiples inside an interval of length m : van Doorn, Li and Tang [4] showed that an interval of length $2 \max A$ guarantees only $\min(n, \lceil 2\sqrt{n} \rceil)$ disjoint pairs $(a, \text{multiple of } a)$, and this is optimal. (c) An induction that removes a_n , takes a solution B' for the rest and repairs it with two new elements fails: for $A' = \{2, 4, 8, 10, 12\}$, $I = \{1, \dots, 16\}$, $B' = \{8, 12, 15, 16\}$ and $a_6 = 16$, the missing factor 2^4 cannot be supplied by two elements of $I \setminus B'$. Lemma 2.4 avoids all three pitfalls by giving every bin at least H candidates and letting different primes share an element only through the per-prime bookkeeping of Lemma 2.3.

Remark 7.3 (the gap to $2n$). Both proofs pay one interval element per large atom and per bin. In the balancing region of Lemma 4.1 an a_i may have about $\sqrt{\ln n / \ln \ln n}$ atoms of “medium” size, none of which can be merged, and this is exactly the loss in Theorem 1.2. A linear bound would require medium atoms of *different* a_i to share representatives. A natural sufficient statement is the following.

Conjecture 7.4 (split assignment). For every A and I as above one can choose for each $a \in A$ a factorisation $a = d_1(a)d_2(a)$ into coprime parts ($d_2 = 1$ allowed) and assign each of the at most $2n$ parts to an element $\beta(d) \in I$ so that for every $b \in I$ and every prime p , $\sum_{d: \beta(d)=b} v_p(d) \leq v_p(b)$.

The conjecture implies $g(n) \leq 2n$ (take $B = \beta(\text{parts})$), and Theorem 1.3 proves it whenever $a_n \geq 8n^3$. Since the capacity constraint couples two parts only if they share a prime, the assignment always exists when every $a \in A$ has at most two distinct prime factors (use the finest splitting and Lemma 2.3 prime by prime); a counterexample would need elements with three or more prime factors competing for composite multiples. An exact search found the assignment on the Erdős–Surányi instances with $\ell = 3, 4$, on the instances of Proposition 7.1, and on more than 10^6 random instances with $n \leq 6$ and $a_n \leq 210$; we found no counterexample.

8 A conditional linear bound

Theorem 1.3 settles the range $a_n \geq 8n^3$ with the conjectured value $2n$, so a bound $g(n) \leq Cn$ with an absolute constant C is a statement about $a_n < 8n^3$ only. In this section we reduce such a bound to a single inequality about additive functions on intervals of integers, we prove the inequality in two special cases, and we record three natural strengthenings of it that are *false*, with explicit certificates.

Throughout, a *weight* is a family $z = (z_p)_p$ indexed by the primes with $0 \leq z_p \leq 1$ and only finitely many $z_p \neq 0$, and $w_z(k) := \sum_p z_p v_p(k)$ is the associated completely additive function. We write $t^+ = \max(t, 0)$.

Conjecture 8.1 (hinge inequality). For every $m \geq 1$, every $x \geq 0$ and every weight z , with $I = \{x+1, \dots, x+m\}$,

$$\sum_{k=1}^m (w_z(k) - 2)^+ \leq \sum_{b \in I} (w_z(b) - 1)^+. \quad (3)$$

Theorem 8.2. *If Conjecture 8.1 holds, then $g(n) \leq 18n$ for all $n \geq 1$. More precisely, it suffices that (3) holds for $m = a_n$ and every weight supported on the primes dividing $\prod A$.*

The proof is a linear-programming argument. Fix $A = \{a_1 < \dots < a_n\}$ with $m = a_n < 8n^3$, an interval I of length m , the set $\mathcal{P}$ of primes dividing $P = \prod A$, and $R_p := \sum_{a \in A} v_p(a)$ for

$p \in \mathcal{P}$. Consider the fractional cover problem

$$\tau^* := \min \left\{ \sum_{b \in I} y_b : \sum_{b \in I} v_p(b) y_b \geq R_p \ (p \in \mathcal{P}), \ 0 \leq y_b \leq 1 \right\}. \quad (4)$$

It is feasible, since $y \equiv 1$ works: $\prod A$ divides $m!$, which divides the product of any m consecutive integers, so $\sum_{b \in I} v_p(b) \geq R_p$.

Lemma 8.3 (duality). $\tau^* = \max_{z \geq 0} \left[\sum_{a \in A} w_z(a) - \sum_{b \in I} (w_z(b) - 1)^+ \right]$, where the maximum is over $z = (z_p)_{p \in \mathcal{P}}$ with $z_p \geq 0$, and it is attained at some z with all $z_p \leq 1$.

Proof. The dual of (4) has variables $z_p \geq 0$ (one per covering constraint) and $u_b \geq 0$ (one per upper bound $y_b \leq 1$) and reads $\max \sum_p z_p R_p - \sum_b u_b$ subject to $\sum_p z_p v_p(b) - u_b \leq 1$ for every $b \in I$. For fixed z the best choice is $u_b = (w_z(b) - 1)^+$, and $\sum_p z_p R_p = \sum_{a \in A} w_z(a)$; strong duality holds because (4) is feasible and bounded. For the last claim fix all $z_q, q \neq p$, and let $z_p \geq 1$ vary: every b with $p \mid b$ then has $w_z(b) \geq 1$, so the right derivative of the objective in z_p equals $R_p - \sum_{b \in I} v_p(b) \leq 0$. Hence the objective does not increase on $z_p \geq 1$. $\square$

Lemma 8.4 (rounding). *There is a set $B \subseteq I$ with $\prod A \mid \prod B$ and $|B| \leq \tau^* + |\mathcal{P}|$.*

Proof. Let y^* be an optimal vertex of the polytope in (4). A vertex of a polytope in $\mathbb{R}^m$ is determined by m linearly independent tight constraints, at most $|\mathcal{P}|$ of which are covering constraints, so at most $|\mathcal{P}|$ coordinates of y^* lie strictly between 0 and 1. Put $B = \{b : y_b^* > 0\}$. Then $\sum_{b \in B} v_p(b) \geq \sum_b v_p(b) y_b^* \geq R_p$ for every p , i.e. $\prod A \mid \prod B$, and $|B| \leq \#\{b : y_b^* = 1\} + |\mathcal{P}| \leq \tau^* + |\mathcal{P}|$. $\square$

Lemma 8.5 (few primes). *If $m < 8n^3$ then $|\mathcal{P}| < 16n$.*

Proof. The i -th prime is at least $i+1$, so if $|\mathcal{P}| \geq 16n$ then $\prod_{p \in \mathcal{P}} p \geq (16n+1)! > (16n/e)^{16n}$, while $\prod_{p \in \mathcal{P}} p \leq \prod A \leq m^n < (8n^3)^n$. Taking logarithms, $16 \ln(16n/e) = 28.3 \dots + 16 \ln n > \ln 8 + 3 \ln n$ for every $n \geq 1$, a contradiction. $\square$

Proof of Theorem 8.2. If $a_n \geq 8n^3$, Theorem 1.3 gives $2n \leq 18n$ elements. Otherwise let z be a maximiser in Lemma 8.3 with $z_p \leq 1$, supported on $\mathcal{P}$. Since $w_z(a) \leq 2 + (w_z(a) - 2)^+$ and $A \subseteq [1, m]$,

$$\sum_{a \in A} w_z(a) \leq 2n + \sum_{k=1}^m (w_z(k) - 2)^+ \leq 2n + \sum_{b \in I} (w_z(b) - 1)^+$$

by (3), so $\tau^* \leq 2n$. Lemmas 8.4 and 8.5 give $|B| \leq 2n + 16n - 1 < 18n$. $\square$

Equivalent forms and a per-prime-power sufficient condition

Subtracting $\sum w_z$ over both ranges, (3) is equivalent to

$$\sum_{b \in I} \min(w_z(b), 1) \leq \sum_{k=1}^m \min(w_z(k), 2) + \Delta, \quad \Delta := \sum_{b \in I} w_z(b) - \sum_{k=1}^m w_z(k) = \sum_q z_q \delta_q \geq 0, \quad (5)$$

where q runs over prime powers, $z_{p^j} := z_p$, and $\delta_q := \#\{b \in I : q \mid b\} - \lfloor m/q \rfloor \in \{0, 1\}$. Writing $\min(w, 1) = w \min(1, 1/w)$ and $\min(w, 2) = w \min(1, 2/w)$ and expanding $w_z(n) = \sum_{q \mid n} z_q$, both

sides of (5) become sums over prime powers, and (3) follows if for every prime power q with $z_q > 0$

$$\sum_{b \in I, q|b} \min\left(1, \frac{1}{w_z(b)}\right) \leq \delta_q + \sum_{k \leq m, q|k} \min\left(1, \frac{2}{w_z(k)}\right). \quad (6)$$

Substituting $b = qi'$, $k = qi$ and $w_z(qi) = a + w_z(i)$ with $a := w_z(q)$, the multiples of q in I correspond to an interval J of $L := \lfloor m/q \rfloor$ or $L + 1$ consecutive integers, and (6) follows from

$$\sum_{i \in J} f_a(w_z(i)) \leq \sum_{i=1}^L g_a(w_z(i)), \quad f_a(u) := \min\left(1, \frac{1}{a+u}\right), \quad g_a(u) := \min\left(1, \frac{2}{a+u}\right), \quad (7)$$

for intervals J of length L (the possible extra element of J is paid by δ_q , as $f_a \leq 1$).

Proposition 8.6. *Inequality (7) holds for every interval J of length L and every $a \geq 0$ in each of the following cases: (i) z is supported on a single prime; (ii) $a \geq 2$ and $aL \geq w_z(L!)$, in particular whenever $a \geq \max\{2, \sum_{p \leq L} z_p/(p-1)\}$.*

Proof. (i) Let z be supported on p and $\phi(j) := f_a(z_p j)$, a nonincreasing function of j , with increments $D_j := \phi(j-1) - \phi(j) \geq 0$. For any set S of L integers, $\sum_{i \in S} \phi(v_p(i)) = L\phi(0) - \sum_{j \geq 1} D_j \#\{i \in S : p^j \mid i\}$. Since an interval of length L contains at least $\lfloor L/p^j \rfloor = \#\{i \leq L : p^j \mid i\}$ multiples of p^j , the sum over J is at most the sum over $[1, L]$, and $f_a \leq g_a$ pointwise. (ii) For $a \geq 2$ the minima are inactive; the left side is at most L/a , and by the Cauchy–Schwarz inequality the right side is $2 \sum_{i \leq L} (a + w_z(i))^{-1} \geq 2L^2/(aL + w_z(L!)) \geq L/a$. Finally $w_z(L!) = \sum_p z_p v_p(L!) \leq L \sum_{p \leq L} z_p/(p-1)$ by Legendre’s formula. $\square$

Three strengthenings that fail

Each of the following statements would imply (3) or (7) by a short argument, and each is false. The common mechanism is that an interval of length L can contain more integers free of all primes $\leq L$ than $\pi(L) + 1$: the pattern of such integers is an admissible set, and by the Chinese remainder theorem every admissible pattern occurs, so their maximal number is the Hensley–Richards function $\rho^*(L)$, which exceeds $\pi(L)$ for $L \geq 3159$ and by an unbounded amount as $L \rightarrow \infty$ [5, 6]. Explicit windows were built by a centred sieve (residue class 0 modulo every $p \leq \sqrt{L}$, then the least populated class modulo each larger prime) and the Chinese remainder theorem; the starting points x and verification scripts are in the repository.

Proposition 8.7. (a) (weight-free transport) *There need not exist an injection $\varphi : [1, m] \rightarrow I$ such that for every k some prime $p \mid k$ satisfies $(k/p) \mid \varphi(k)$. For $m = 20000$ there is an x (with 4298 digits) for which I contains 2270 integers free of all primes $\leq m/2$; every k related to such an integer is 1 or a prime, so Hall’s condition fails because $2270 > \pi(m) + 1 = 2263$. (Such an injection would give (3) at once, since $w_z(\varphi(k)) \geq w_z(k) - z_p \geq w_z(k) - 1$.)*

(b) (single-threshold level sets) *The inequality $\#\{i \leq L : w_z(i) \geq 2\tau\} \leq \#\{i \in J : w_z(i) \geq \tau\}$ fails for $L = 10^5$, $z_p = 1$ for all $p \leq L/2$, at $\tau = 1$ (by 142) and at $\tau = 1/2$ (by 5275), although (7) holds on the same window with room to spare. (Integrating it over τ would give (7).)*

(c) (Laplace form) *The inequality $s \sum_{i \in J} s^{w_z(i)} \leq \sum_{i \leq L} s^{w_z(i)}$ for all $s \in (0, 1]$, which would give (7) for all $a \geq 2$ by integrating against $s^{a-1} ds$, fails for $L = 10^6$, $z_p = 1$ for all $p \leq L$,*

and $s \in [0.005, 0.015]$: the window (with x of 433636 digits) contains 80436 integers free of all primes $\leq L$, against $\pi(L) + 1 = 78499$, and at $s = 0.005$ the two sides are 405.03 and 398.77.

Remark 8.8 (evidence and the remaining programme). Conjecture 8.1 was tested on 2.1×10^6 random and adversarial instances with $m \leq 600$ (0/1, fractional and $p^{-\theta}$ weights; $x = 0$, random x , $x \equiv -1$ modulo products of weighted primes, windows centred at highly divisible numbers) and, with nine weight choices, on the dense windows of Proposition 8.7 of lengths $2 \cdot 10^4$ and 10^5 , where the slack is at least $0.18m$; (7) was tested on 3.5×10^5 instances over all prime powers. Equality in (3) occurs only for $x = 0$ and single primes. For 0/1 weights $z = \mathbf{1}_Q$ the inequality is equivalent to $B_0 - A_0 \leq \Delta + B_{\geq 2}$, where B_0 and A_0 count the integers in $[1, m]$ and in I free of the primes of Q , and $B_{\geq 2}$ counts the $k \leq m$ with at least two prime factors from Q (with multiplicity); by inclusion–exclusion $B_0 - A_0$ is an alternating sum of the surpluses δ_d over squarefree products d of primes of Q . In the layer-cake form of (7) the integrand is positive only where the count on $[1, L]$ is stuck at a low level, and its positive part is bounded by the number of integers in a window free of the weighted primes, i.e. by an explicit large sieve bound, while the negative part is a counting quantity on $[1, L]$. Making this quantitative for all weights is the remaining step; the failures in Proposition 8.7 show that it cannot be replaced by a weight-free or single-threshold argument.

9 Counting certificates and the 0/1 case

All three failures in Proposition 8.7 have a common feature: they use only the lower bound $N_I(d) \geq \lfloor m/d \rfloor$ on the number $N_I(d) := \#\{b \in I : d \mid b\}$ of multiples of d in the interval. The upper bound $N_I(d) \leq \lfloor m/d \rfloor + 1$, which is exactly what the dense windows violate in spirit, was never used. Using both bounds, linear programming duality turns the hinge inequality into a finite question, and this question has a positive answer in every case we could compute; for 0/1 weights it also has an explicit answer with a slightly larger threshold.

The certificate principle

Proposition 9.1 (counting certificates). *Let z be a weight and let (c_d) be real numbers, finitely many nonzero, indexed by positive integers d , such that*

$$\sum_{d \mid n} c_d \leq (w_z(n) - 1)^+ \quad \text{for every positive integer } n. \quad (8)$$

Put $V(c) := \sum_{c_d > 0} c_d \lfloor m/d \rfloor - \sum_{c_d < 0} |c_d| (\lfloor m/d \rfloor + 1)$. Then for every $x \geq 0$, $\sum_{b \in I} (w_z(b) - 1)^+ \geq V(c)$. In particular (3) holds for (m, z) and every x as soon as $V(c) \geq W(m, z) := \sum_{k \leq m} (w_z(k) - 2)^+$ for some c satisfying (8).

Proof. Summing (8) over $b \in I$ gives $\sum_{b \in I} (w_z(b) - 1)^+ \geq \sum_d c_d N_I(d)$, and $\lfloor m/d \rfloor \leq N_I(d) \leq \lfloor m/d \rfloor + 1$ for every d because I consists of m consecutive integers. $\square$

The best certificate is the optimum of a linear programme: the dual of minimising $\sum_t n_t (w_z(t) - 1)^+$ over nonnegative real “multisets” (n_t) of valuation patterns t subject to $\lfloor m/d \rfloor \leq \sum_{t: d \mid t} n_t \leq \lfloor m/d \rfloor + 1$ for all d . We solved this programme (for general weights with the exponents capped at $\lfloor \log_p m \rfloor + 1$, which is legitimate because a certificate supported on capped d satisfies (8) for

all n ; for 0/1 weights with c supported on pairs and triples of primes, adding the constraints (8) by separation) and found $V \geq W$ in every instance: for all weights $z \in \{1, 0.7, 0.4\}^P$ and mixed weights with P the primes up to 17 and $m \leq 200$ (e.g. $V = 114$ against $W = 64$ for $m = 100$, $z \equiv 1$), and for 0/1 weights with P the set of all primes up to m for $m \in \{300, 500, 1000\}$ ($V = 223.5, 407.4, 907.0$ against the distinct-prime targets $W_\omega = 59, 124, 321$ defined below) as well as for various subsets of the primes at $m = 400$. The optimal certificates are small: $c_{p^j} = 1$ for prime powers with $j \geq 2$, $c_{pq} = 1$ for pairs containing 2, 3 or 5, and $c_{pqr} = -1$ for a few triples (no coefficient at $d = 1$ or at single primes). The scripts are in the repository.

An explicit family for 0/1 weights

For 0/1 weights $z = \mathbf{1}_P$ write $\omega_P(n)$ for the number of primes of P dividing n , $\Omega_P(n) = \sum_{p \in P} \mathbf{v}_p(n)$ and $E_P := \Omega_P - \omega_P = \sum_{p \in P} \sum_{j \geq 2} [p^j \mid n]$. Since $(\Omega_P - 1)^+ = E_P + (\omega_P - 1)^+$, $(\Omega_P - c)^+ \leq E_P + (\omega_P - c)^+$ for every $c \geq 0$, and $\sum_{b \in I} E_P(b) \geq \sum_{k \leq m} E_P(k)$ (count the multiples of each p^j), it suffices to certify the distinct-prime inequality $\sum_{k \leq m} (\omega_P(k) - 2)^+ \leq \sum_{b \in I} (\omega_P(b) - 1)^+$, whose left side we denote $W_\omega(m)$ (for $z = \mathbf{1}_P$ one has $W(m, z) \geq W_\omega(m)$, and the certificates below are for the distinct-prime version).

Lemma 9.2. *Let R be the set of the r smallest primes of P . Put $c_d := (-1)^{|A|}$ for $d = \prod A$ with $A \subseteq R$, $|A| \geq 2$, and $c_d := (-1)^{|A|+1}$ for $d = q \prod A$ with $q \in P \setminus R$ and $\emptyset \neq A \subseteq R$; all other $c_d = 0$. Then (8) holds with $(\omega_P(n) - 1)^+$ on the right, and it keeps holding if all coefficients belonging to any fixed $q \in P \setminus R$ are set to 0. For $r = 2$, $R = \{2, 3\}$ and the coefficients of the primes $q > m/2$ removed, the value is*

$$V_2(m) := \left\lfloor \frac{m}{6} \right\rfloor + \sum_{\substack{q \in P \\ 5 \leq q \leq m/2}} \left(\left\lfloor \frac{m}{2q} \right\rfloor + \left\lfloor \frac{m}{3q} \right\rfloor - \left\lfloor \frac{m}{6q} \right\rfloor - 1 \right). \quad (9)$$

Proof. Let n have prime set $T = A_0 \cup Q_0$ with $A_0 \subseteq R$, $Q_0 \subseteq P \setminus R$. Then $\sum_{d \mid n} c_d = \sum_{A \subseteq A_0, |A| \geq 2} (-1)^{|A|} + \sum_{q \in Q_0} \sum_{\emptyset \neq A \subseteq A_0} (-1)^{|A|+1} = (|A_0| - 1)^+ + |Q_0| [A_0 \neq \emptyset] \leq (|T| - 1)^+$, using $\sum_{A \subseteq A_0} (-1)^{|A|} = [A_0 = \emptyset]$. Removing the terms of one q removes a nonnegative contribution. The value follows from the definition of V . $\square$

With $\mathcal{A}(x) := \sum_{p \leq x} [x/p] = \sum_{n \leq x} \omega(n)$ and P the set of all primes up to m one has the exact identities $W_\omega(m) = \mathcal{A}(m) - 2m + 2 + \#\{\text{prime powers} \leq m\}$ and $V_2(m) = \mathcal{A}(m/2) + \mathcal{A}(m/3) - \mathcal{A}(m/6) - [m/4] - [m/6] - [m/9] + [m/12] + [m/18] - \pi(m/2) + 2$, so $V_2(m) \geq W_\omega(m)$ was checked exactly for every $m \leq 10^7$ (at $m = 10^3, \dots, 10^7$ the ratio V_2/W_ω is 2.72, 2.00, 1.71, 1.55, 1.44). Since $\mathcal{A}(x) = x \ln \ln x + Mx + O(x/\ln x)$ [7, Thm. 430], $V_2(m) - W_\omega(m) = -\frac{1}{3}m \ln \ln m + (\frac{29}{18} - \frac{M}{3})m + o(m)$, so this single certificate proves the 0/1 case of (3) for every $m \leq 10^7$ (checked exactly for each m) and fails for all sufficiently large m (the leading-order crossing is at $\ln \ln m = \frac{29}{6} - M \approx 4.57$, i.e. $m \approx 10^{42}$; in between the inequality is expected but has not been verified); the linear programme shows that better certificates exist (907 against 873 at $m = 1000$), and whether they exist for every m is open. The number of anchors must depend on P : for P the primes in $[53, 653]$ and $m = 10^8$ the two-anchor certificate gives $1,595,615 < W_\omega = 1,602,217$, while three anchors give 2,266,053.

The 0/1 case with a larger threshold

Theorem 9.3. *Let P be a finite set of primes, $m \geq 3$, $x \geq 0$, and $c^*(m) := 5 + \max(2, \lceil 2e^2 \ln(1 + \frac{\ln m}{6}) \rceil)$. Then*

$$\sum_{n \leq m} (\Omega_P(n) - c^*(m))^+ \leq \sum_{b \in I} (\Omega_P(b) - 1)^+, \quad (10)$$

and $c^*(m) \leq 20 \ln \ln m$ for $m \geq 30$.

The proof rests on an elementary sieve lemma which, in the language of Proposition 9.1, is a certificate with positive coefficients on single primes and negative coefficients on pairs.

Lemma 9.4. *For $y \geq 2$, $\sum_{p \leq y} 1/p \leq e \ln(1 + \ln y)$.*

Proof. With $\sigma = 1 + 1/\ln y$ one has $p^{\sigma-1} \leq e$ for $p \leq y$, so $1/p \leq e p^{-\sigma}$; and $\sum_p p^{-\sigma} \leq \ln \zeta(\sigma) \leq \ln(1 + 1/(\sigma - 1)) = \ln(1 + \ln y)$. $\square$

Lemma 9.5 (sieve lemma). *Let $L \geq 64$, let Q be a finite set of primes, let J be any set of L consecutive integers and $s(L) := \max(2, \lceil 2e^2 \ln(1 + \frac{\ln L}{6}) \rceil)$. Then*

$$\#\{n \leq L : \omega_Q(n) \geq s(L) + 5\} \leq \#\{n \in J : \omega_Q(n) \geq 1\}.$$

Proof. Put $y = L^{1/6}$, $R = \{q \in Q : q \leq y\}$, $H = \sum_{q \in R} 1/q$ and $s = s(L)$. An integer $n \leq L$ has at most five prime factors exceeding y , so $\omega_Q(n) \geq s + 5$ forces $\omega_R(n) \geq s$, and $N_s := \#\{n \leq L : \omega_R(n) \geq s\} \leq \sum_{S \subseteq R, |S|=s} \lfloor L / \prod S \rfloor \leq L H^s / s!$ by the bound $s! e_s \leq H^s$ on elementary symmetric functions. By Lemma 9.4, $H \leq e \ln(1 + \frac{\ln L}{6})$, so $s \geq 2eH$ and, with $s! \geq (s/e)^s$, $N_s \leq L(eH/s)^s \leq L 2^{-s} \leq L/4$.

For the right-hand side we use that any set J of L consecutive integers satisfies $L/d - 1 < N_J(d) < L/d + 1$. If $H \geq 1/2$, add primes of R one at a time until the reciprocal sum first reaches $1/2$; the resulting R' has $1/2 \leq H' \leq 1$ (the last prime added contributes at most $1/2$) and $t' = |R'| \leq y$. By the two-term Bonferroni inequality, $\#\{n \in J : \omega_Q(n) \geq 1\} \geq \#\{n \in J : \omega_{R'}(n) \geq 1\} \geq \sum_{q \in R'} N_J(q) - \sum_{q < q'} N_J(qq') \geq L(H' - H'^2/2) - t'(t' + 1)/2 \geq \frac{3}{8}L - \frac{3}{4}y^2 \geq \frac{3}{8}L - \frac{1}{8}L = \frac{1}{4}L \geq N_s$, using $y(y + 1)/2 \leq \frac{3}{4}y^2$ for $y \geq 2$ and $\frac{3}{4}L^{1/3} \leq L/8$ for $L \geq 64$. If $0 < H < 1/2$, the same estimate with $R' = R$ and $H - H^2/2 \geq \frac{3}{4}H$ gives $\#\{n \in J : \omega_Q(n) \geq 1\} \geq \frac{3}{4}LH - \frac{3}{4}y^2 \geq \frac{1}{4}LH$ (because $H \geq 1/y$ and $\frac{3}{4}y^2 \leq L/(2y)$ for $L \geq 64$), while $N_s \leq LH^2/2 \leq LH/4$. If $H = 0$ then $N_s = 0$. $\square$

Proof of Theorem 9.3. Order P increasingly and let $\omega_{<p}(n)$ count the primes of P below p dividing n . For every integer $c \geq 0$,

$$(\omega_P(n) - c)^+ = \sum_{p \in P, p|n} [\omega_{<p}(n) \geq c], \quad (11)$$

since the right side counts all but the c smallest primes of P dividing n . Fix $p \in P$ and put $L = \lfloor m/p \rfloor$. The multiples $n = pi \leq m$ of p have $\omega_{<p}(pi) = \omega_{<p}(i)$ with $i \leq L$, and the multiples $b = pi$ of p in I have i ranging over a set J_p of L or $L + 1$ consecutive integers. If $L \geq 64$, Lemma 9.5 applied to $Q = \{q \in P : q < p\}$ and to L consecutive elements of J_p gives $\#\{i \leq L : \omega_{<p}(i) \geq c^*(m)\} \leq \#\{i \in J_p : \omega_{<p}(i) \geq 1\}$, because $c^*(m) \geq s(L) + 5$. If $L < 64$ the left side is 0, as $\omega_{<p}(i) \leq 3 < 7 \leq c^*(m)$ for $i \leq 63$. Summing over p and using (11) with $c = c^*(m)$ on the left and $c = 1$ on the right proves (10) with ω_P in place of Ω_P ; the peeling

identities stated before Lemma 9.2 then give (10). Finally $G(t) := 20 \ln t - 7 - 2e^2 \ln(1 + t/6)$ is increasing for $t > 0$ (as $2e^2 < 20$) and $G(\ln 30) > 0$, which gives $c^*(m) \leq 20 \ln \ln m$ for $m \geq 30$. $\square$

Remark 9.6 (fractional weights). Theorem 9.3 does not imply (3) for fractional weights, and integrating it over weight thresholds cannot (Proposition 8.7 and the identity $\int_0^1 (\#\{p \mid n : z_p > t\} - 1)^+ dt = \sum_{p \mid n} z_p - \max_{p \mid n} z_p$). Writing $w_z = E + S$ with $S(n) = \sum_p \min(z_p v_p(n), 1)$ and $E = \sum_p (z_p v_p(n) - 1)^+$, one has $(w_z - 1)^+ = E + (S - 1)^+$ (if $S < 1$ then every $z_p v_p < 1$, so $E = 0$), $(w_z - c)^+ \leq E + (S - c)^+$, and $\sum_I E \geq \sum_K E$ because $(z_p v_p(n) - 1)^+ = \sum_{j \geq 1} \Delta_{p,j} [p^j \mid n]$ with nonnegative increments $\Delta_{p,j} = (z_p j - 1)^+ - (z_p(j-1) - 1)^+$, so (3) for all weights follows from its analogue for the bounded function S ; for rational weights $z_p = a_p/N$ this is the integer statement $\sum_{k \leq m} (R(k) - 2N)^+ \leq \sum_{b \in I} (R(b) - N)^+$ with $R(n) = \sum_p \min(a_p v_p(n), N)$, of which the 0/1 case ($a_p \in \{0, N\}$) is the one treated above. The linear programme of Proposition 9.1 certifies the fractional inequality in every instance we solved; an explicit certificate for all m and all weights would complete the proof of $g(n) \leq 18n$.

10 The fractional case and an $O(n \ln \ln n)$ bound

Theorem 9.3 used only 0/1 weights. In this section the same method, combined with a random-thinning device, handles arbitrary weights, at the price of a larger threshold; through Lemma 8.3 this gives the first bound for $g(n)$ that is linear up to a $\ln \ln n$ factor.

Theorem 10.1. *For every weight z , every $m \geq 16$ and every $x \geq 0$,*

$$\sum_{k \leq m} (w_z(k) - c_m)^+ \leq \sum_{b \in I} (w_z(b) - 1)^+, \quad c_m := 1 + 149e \ln(1 + \ln m), \quad (12)$$

and $c_m < 675 \ln \ln m$. For $2 \leq m \leq 15$ the inequality holds with $c_m = 4$.

Corollary 10.2. $g(n) \leq (17 + 149e \ln(1 + \ln(8n^3)))n$ for all $n \geq 1$; in particular $g(n) = O(n \ln \ln n)$.

Proof. If $a_n \geq 8n^3$, Theorem 1.3 gives $2n$. Otherwise $m = a_n < 8n^3$ and, by Lemma 8.3 with a maximiser z , $\sum_{a \in A} w_z(a) \leq c_m n + \sum_{k \leq m} (w_z(k) - c_m)^+ \leq c_m n + \sum_{b \in I} (w_z(b) - 1)^+$, so $\tau^* \leq c_m n$ and Lemmas 8.4, 8.5 give $|B| < (c_m + 16)n$, with $c_m = 4$ when $m \leq 15$; in all cases $c_m \leq c_{8n^3}$ since c_m is increasing for $m \geq 16$ and $4 < c_{16}$. $\square$

Throughout this section $\lambda(t) := \ln(1 + \ln t)$, $L \geq 16$ is an integer, J is a set of L consecutive positive integers, $R := \lfloor \log_2 L \rfloor$, $H := e\lambda(L)$ and $L_0 := \ln(H + 2)$; note $H > 3$ and $L_0 > 1$ for $L \geq 16$.

Two-term Bonferroni at level k

Lemma 10.3. *For integers $r \geq 0$ and $k \geq 1$, $[r \geq k] \geq \binom{r}{k} - k \binom{r}{k+1}$.*

Proof. For $r < k$ both sides vanish; for $r \in \{k, k+1\}$ the right side is 1; for $r \geq k+2$, $\binom{r}{k} - k \binom{r}{k+1} = \binom{r}{k} (1 - \frac{k(r-k)}{k+1}) \leq 0$. $\square$

A sieve lemma with an arbitrary gain

Let D be a finite family of pairwise coprime integers $d \geq 2$ and $\nu_D(n) := \#\{d \in D : d \mid n\}$. For $1 \leq k \leq R-1$ put

$$\rho(k) := \left\lceil 16H + \frac{16kL_0}{\sqrt{\ln(2+k/H)}} \right\rceil, \quad C(k) := 2k + \rho(k). \quad (13)$$

Lemma 10.4 (gain- G sieve). *For every real $1 \leq G \leq 2H$, every $1 \leq k \leq R-1$ and every J ,*

$$G \cdot \#\{n \leq L : \nu_D(n) \geq C(k)\} \leq \#\{n \in J : \nu_D(n) \geq k\}.$$

Proof. Put $y := L^{1/(k+1)} \geq 2$, $D_0 := \{d \in D : d \leq y\}$ and $H_0 := \sum_{d \in D_0} 1/d$. Choosing a prime divisor of each $d \in D_0$ gives distinct primes $\leq y$, so $H_0 \leq \sum_{p \leq y} 1/p \leq e\lambda(y) \leq H$ by Lemma 9.4. Let e_j denote the elementary symmetric functions of the numbers $1/d$, $d \in D_0$; then $(k+1)e_{k+1} \leq H_0 e_k$ and $e_{k+r} \leq e_k e_r / \binom{k+r}{k} \leq e_k H_0^r / r!$.

Lower bound on J . Retain each element of D_0 independently with probability $\theta := 1/(8H)$, obtaining a random subfamily E . For every E , Lemma 10.3 summed over $n \in J$ gives $\#\{n \in J : \nu_E(n) \geq k\} \geq M_k(E) - kM_{k+1}(E)$, where $M_j(E) := \sum_{F \subseteq E, |F|=j} N_J(\prod F)$; since $\nu_D \geq \nu_E$ and $\mathbb{E} M_j(E) = \theta^j M_j(D_0)$, taking expectations yields $\#\{n \in J : \nu_D(n) \geq k\} \geq \theta^k (M_k(D_0) - k\theta M_{k+1}(D_0))$. A product of k elements of D_0 is at most $y^k = L/y \leq L/2$, so $N_J(a) \geq L/a - 1 \geq L/(2a)$ and $M_k(D_0) \geq \frac{L}{2} e_k$; a product of $k+1$ elements is at most $y^{k+1} = L$, so $N_J(a) \leq L/a + 1 \leq 2L/a$ and $M_{k+1}(D_0) \leq 2Le_{k+1}$. Hence $k\theta M_{k+1}(D_0) \leq 2\theta LH_0 e_k \leq \frac{L}{4} e_k$ and

$$\#\{n \in J : \nu_D(n) \geq k\} \geq \frac{Le_k}{4(8H)^k}. \quad (14)$$

Upper bound on $[1, L]$. An integer $n \leq L$ is divisible by at most k elements of $D \setminus D_0$, since $k+1$ of them would have product $> y^{k+1} = L$. Hence $\nu_D(n) \geq C(k)$ implies $\nu_{D_0}(n) \geq k + \rho(k)$, and with $r = \rho(k)$,

$$\#\{n \leq L : \nu_D(n) \geq C(k)\} \leq Le_{k+r} \leq Le_k \frac{H^r}{r!}. \quad (15)$$

The gain. Write $s = k/H$ and $\ell = \ln(2+s)$. We claim $\ln\left(\frac{16}{e}(1+s/\sqrt{\ell})\right) \geq \frac{1}{2}\sqrt{\ell}$. With $t = \sqrt{\ell}$: if $t \leq 1$ the left side is at least $\ln(16/e) > 1 \geq t/2$; if $t \geq 1$ then $s = e^{t^2} - 2$ and $1 + s/t \geq e^{t^2}/(2t)$ (as $e^{t^2}/2 + t - 2 \geq 0$), so the difference of the two sides is at least $\ln(8/e) + t^2 - \ln t - t/2$, which is positive at $t = 1$ and increasing. Since $r = \rho(k) \geq 16H(1+sL_0/\sqrt{\ell}) \geq 16H(1+s/\sqrt{\ell})$, we get $\ln \frac{r}{eH} \geq \frac{1}{2}\sqrt{\ell}$ and, using $r! \geq (r/e)^r$ and $r \geq 16kL_0/\sqrt{\ell}$, $\ln \frac{r!}{H^r} \geq r \ln \frac{r}{eH} \geq 8kL_0$. On the other hand $4G \leq 8H \leq (H+2)^3$ gives $\ln(4G(8H)^k) \leq 3(k+1)L_0 \leq 6kL_0$. Therefore $4G(8H)^k H^r / r! \leq 1$, and (14), (15) give the lemma. $\square$

Dyadic encoding of fractional valuations

Let $u = (u_q)$ be weights, $s_q(n) := \min(u_q \nu_q(n), 1)$ and $X(n) := \sum_q s_q(n)$. For $j \geq 0$ put $N_j(n) := \#\{q : s_q(n) \geq 2^{-j}\}$ and, for $u_q > 0$, $d_{q,j} := q^{\lceil 2^{-j}/u_q \rceil}$. Then $s_q(n) \geq 2^{-j}$ if and only if $d_{q,j} \mid n$, the integers $d_{q,j}$ (q varying) are pairwise coprime, so $N_j = \nu_{D_j}$ for $D_j := \{d_{q,j}\}$; and since $s \leq \sum_{j \geq 0} 2^{-j} [s \geq 2^{-j}] \leq 2s$ for $s \in [0, 1]$,

$$X(n) \leq \sum_{j \geq 0} 2^{-j} N_j(n) \leq 2X(n). \quad (16)$$

Lemma 10.5 (weighted sieve lemma). *For every family of weights u , every $L \geq 16$ and every J ,*

$$\#\{n \leq L : X(n) \geq 149H\} \leq \#\{n \in J : X(n) \geq 1\}.$$

Proof. Put $Q := 1 + \lfloor \log_2(R-1) \rfloor$, so that $2^j \leq R-1$ for $j < Q$ and $2^Q \geq R$. With $t = \ln L$ one has $R < 2t$ and $Q < 3 + 2 \ln t < 5\lambda(L) < 2H$ for $L \geq 16$. For $j < Q$ set $a_j := C(2^j) - 1$ and for $j \geq Q$ set $a_j := R$; let $A := \sum_{j \geq 0} 2^{-j} a_j$. Since $N_j(n) \leq \omega(n) \leq R$ for $n \leq L$, if $X(n) \geq A+1$ for some $n \leq L$ then by (16) some $j < Q$ has $N_j(n) \geq C(2^j)$. Lemma 10.4 with $D = D_j$, $k = 2^j$ and $G = Q$ gives $Q \cdot \#\{n \leq L : N_j(n) \geq C(2^j)\} \leq \#\{n \in J : N_j(n) \geq 2^j\} \leq \#\{n \in J : X(n) \geq 1\}$, the last step because $N_j(n) \geq 2^j$ forces $X(n) \geq 2^{-j} N_j(n) \geq 1$. Summing over the Q indices $j < Q$, $\#\{n \leq L : X(n) \geq A+1\} \leq \#\{n \in J : X(n) \geq 1\}$.

It remains to show $A+1 \leq 149H$. From (13), $a_j \leq 2 \cdot 2^j + 16H + 16 \cdot 2^j L_0 / \sqrt{\ln(2+2^j/H)}$, so $\sum_{j < Q} 2^{-j} a_j \leq 2Q + 32H + 16L_0 S_Q$ with $S_Q := \sum_{j < Q} \ln(2+2^j/H)^{-1/2}$. The indices with $2^j \leq H$ number at most $1 + \log_2 H \leq 1 + \frac{3}{2}\sqrt{H}$ and contribute at most $\frac{5}{4}$ each, hence at most $\frac{21}{8}\sqrt{H}$; if j_0 is the first index with $2^{j_0} > H$, the j_0 -term is less than 1 and the term $j_0 + t$ ($t \geq 1$) is less than $\frac{5}{4}t^{-1/2}$, so the remaining terms contribute less than $1 + \frac{5}{2}\sqrt{Q} \leq \frac{21}{5}\sqrt{H}$. Thus $S_Q \leq 7\sqrt{H}$, and with $L_0 = \ln(H+2) \leq \sqrt{H}$ we get $16L_0 S_Q \leq 112H$. Finally $\sum_{j \geq Q} 2^{-j} R = 2R/2^Q \leq 2$, so $A \leq 4H + 32H + 112H + 2 = 148H + 2$ and $A+1 \leq 149H$ because $H > 3$. $\square$

Proof of Theorem 10.1

Write $w_z = E + S$ with $S(n) = \sum_p \min(z_p v_p(n), 1)$ as in the remark after Theorem 9.3; it suffices to prove $\sum_{k \leq m} (S(k) - c_m)^+ \leq \sum_{b \in I} (S(b) - 1)^+$. By continuity in the (finitely many) weights we may assume $z_p = a_p/N$ rational; put $r_p(n) := \min(a_p v_p(n), N)$ and $R(n) := \sum_p r_p(n) = NS(n)$. For integers n and thresholds $T \geq 0$ let $R_{<p}(n) := \sum_{q < p} r_q(n)$. Ordering the primes dividing n and letting p^* be the first with $R_{\leq p^*}(n) \geq T$, one has $\sum_{p|n} r_p(n) [R_{<p}(n) \geq T] = R(n) - R_{\leq p^*}(n)$ and $T \leq R_{\leq p^*}(n) < T + N$, whence

$$(R(n) - T - N)^+ \leq \sum_{p|n} r_p(n) [R_{<p}(n) \geq T] \leq (R(n) - T)^+. \quad (17)$$

Applying (17) with $T = (c_m - 1)N$ on $[1, m]$ and with $T = N$ on I , it suffices to show, for every prime p , $\sum_{k \leq m, p|k} r_p(k) [R_{<p}(k) \geq (c_m - 1)N] \leq \sum_{b \in I, p|b} r_p(b) [R_{<p}(b) \geq N]$. Writing $r_p(n) = \sum_{j \geq 1} \Delta_{p,j} [p^j | n]$ with $\Delta_{p,j} = \min(a_p j, N) - \min(a_p(j-1), N) \geq 0$ and using $R_{<p}(p^j i) = R_{<p}(i)$, this follows from: for every $j \geq 1$, $L := \lfloor m/p^j \rfloor$ and every run J of L consecutive integers (the quotients i of the multiples $p^j i \in I$ form a run of L or $L+1$ consecutive integers),

$$\#\{i \leq L : R_{<p}(i) \geq (c_m - 1)N\} \leq \#\{i \in J : R_{<p}(i) \geq N\}.$$

Dividing by N , $R_{<p}/N$ is the function X of Lemma 10.5 for the weights $u_q = z_q$ ($q < p$). If $L \geq 16$ the lemma applies because $c_m - 1 = 149e\lambda(m) \geq 149e\lambda(L)$; if $L < 16$ the left side is empty since $X(i) \leq \omega(i) \leq 2 < c_m - 1$ for $i \leq 15$. This proves (12) for $m \geq 16$; for $2 \leq m \leq 15$, $S(k) \leq \omega(k) \leq 2$ gives the statement with $c_m = 4$. Finally, for $m \geq 16$ and $t = \ln m$, $\lambda(m) = \ln \ln m + \ln(1 + 1/t) < \frac{3}{2} \ln \ln m$ (as $1/t < 3/8 < \frac{1}{2} \ln \ln m$), and $1 + 149e\lambda(m) < 150e\lambda(m) < 675 \ln \ln m$. $\square$

Remark 10.6. The constant $149e \approx 405$ is far from optimal: the thinning probability $1/(8H)$, the gain $G = Q$ and the dyadic budget were chosen for simplicity. The bound of Corollary 10.2 is smaller than that of Theorem 1.2 only for astronomically large n ; its point is the order of growth. The hinge inequality with threshold 2 (Conjecture 8.1), which would give $18n$, remains open.

References

- [1] P. Erdős and J. Surányi, Megjegyzések egy versenyfeladathoz (Remarks on a problem of a mathematical competition), *Mat. Lapok* 10 (1959), 39–48 (in Hungarian).
- [2] P. Erdős, Some of my forgotten problems in number theory, *Hardy–Ramanujan J.* 15 (1992), 34–50, §1.
- [3] T. F. Bloom, Erdős Problem #708, <https://www.erdosproblems.com/708>, accessed 2026-09-03.
- [4] W. van Doorn, Y. Li and Q. Tang, Optimal bounds for an Erdős problem on matching integers to distinct multiples, arXiv:2603.28636 (2026).
- [5] D. Hensley and I. Richards, Primes in intervals, *Acta Arith.* 25 (1974), 375–391.
- [6] D. M. Gordon and G. Rodemich, Dense admissible sets, in: *Algorithmic Number Theory (ANTS-III)*, Lecture Notes in Comput. Sci. 1423, Springer, 1998, 216–225.
- [7] G. H. Hardy and E. M. Wright, *An Introduction to the Theory of Numbers*, 6th ed., Oxford University Press, 2008.