The Physiognomy of Pain and the Epistemic Fallacy of Observed Aggression: A Multidisciplinary Analysis of Misinterpreted Affect in Chronic Orofacial and Neurological Injury

Executive Summary

This report presents an exhaustive, multi-dimensional analysis of the discordance between internal somatic states—specifically chronic neuropathic pain, structural maxillofacial injury, and neurotoxicological sequelae—and their external behavioral manifestations, which are systematically misinterpreted by social observers as anger or aggression. The analysis is grounded in the specific medical history of the subject, comprising a Temporomandibular Joint (TMJ) fracture, mandibular displacement, shattered molar sequelae involving complex surgical extraction, documented hearing loss, and a pharmacological regimen including olanzapine, paliperidone, and sodium valproate.

The central thesis of this report posits that the social labeling of the subject's involuntary physiological reflexes as "anger" constitutes a fundamental error in behavioral analysis, herein termed the "Outside-In Labeling Fallacy." By applying a rigorous critique of B.F. Skinner's Radical Behaviorism—which historically marginalized the causal relevance of internal private events in favor of observable environmental contingencies—we demonstrate how social observers commit an epistemic error. They attribute the operant of "aggression" to the respondent behaviors of "pain," thereby invalidating the sufferer's internal reality. This misinterpretation is further entrenched by the sociological "Angry Man" stereotype applied to male survivors of trauma, particularly those with Complex Post-Traumatic Stress Disorder (CPTSD), creating a devastating cycle of social isolation, secondary traumatization, and disability exacerbation over the documented eight-year history.

Part I: The Anatomical and Biomechanical Substrate of the "Snarl" Reflex

To understand the persistent misinterpretation of the subject's facial expressions, one must first deconstruct the anatomical substrates that govern facial movement. The human face functions as a dual-channel transmitter: it conveys voluntary social signals (smiling, speaking) and involuntary homeostatic responses (wincing, grimacing, splinting). In the context of the

subject's specific injuries—TMJ fracture, jaw displacement, and severe dental trauma—the "snarl" is not an emotional display of aggression, but a protective neuromuscular splinting reflex designed to stabilize compromised skeletal architecture.

1.1 Structural Pathology of the TMJ and Mandibular Displacement

The temporomandibular joint (TMJ) is the most complex joint in the human body, allowing for both ginglymoid (hinging) and arthrodial (gliding) motions.[1] A history of fracture and subsequent displacement alters the resting tonus of the masticatory musculature, forcing the face into a configuration that biologically mimics threat displays.

1.1.1 The Masseteric Guarding Reflex and Action Unit 31

When the structural integrity of the mandible is compromised, the body engages in "muscle guarding" or protective co-contraction. The masseter and temporalis muscles—the primary elevators of the jaw—enter a state of chronic hypertonicity to stabilize the joint against gravity and movement.[3]

- **The "Clenched" Appearance:** Chronic hypertonicity in the masseter muscles results in a visible bulging at the angle of the jaw and a rigidity of the lower face. In the Facial Action Coding System (FACS), this corresponds to **Action Unit 31 (Jaw Clencher)**.[4]
- **Misinterpretation Mechanics:** A clenched jaw is universally decoded in Western social semiotics as a sign of suppressed rage, determination, or preparation for physical combat. However, for a patient with a displaced jaw disc or fracture history, this clenching is a physiological imperative to prevent subluxation (dislocation) during speech or rest.[3] The observer perceives "he is holding back anger," while the physiological reality is "he is holding his jaw together." The observer's inference is based on the *function* of clenching in a healthy population (anger), failing to account for the *function* of clenching in a pathologized population (stabilization).

1.1.2 Pterygoid Spasm and Mandibular Deviation

The lateral pterygoid muscle is responsible for protruding the jaw and moving it side-to-side. Injury to this muscle or the condylar head (common in fractures) can lead to a resting asymmetry or deviation.[6]

- **Asymmetry as "Contempt":** In facial expression psychology, asymmetry is often decoded as contempt, skepticism, or duplicity (e.g., a unilateral lip raise or sneer).[8] If the jaw deviates to the left or right due to structural damage (as noted in cases of condylar resorption or fracture), it pulls the lip line askew.
- **Mechanism of the "Shattered Molar":** The subject's history involves a "shattered molar" and complex surgical extraction of teeth 46 and 28, involving root sectioning and bone removal.[9] The loss of these key posterior structural supports leads to a loss of vertical dimension (occlusal collapse) on the affected side. This forces the facial muscles to contract unevenly to achieve lip closure (competence), mechanically forcing the mouth into a shape that mimics the sneer of contempt. The documented "distal secondary caries" and "irreversible pulpitis" [9] prior to extraction would have established

a long-term pattern of unilateral guarding, effectively training the facial musculature into an asymmetrical resting state.

1.2 The Neuropathic "Snarl": Trigeminal Nerve Involvement

The trigeminal nerve (CN V) provides sensation to the face and motor control to the jaw. Damage or chronic irritation to this nerve—common in TMJ fractures and difficult extractions involving "root sectioning" and "interseptal bone removal" [9]—can lead to neuropathic pain states like Trigeminal Neuralgia or persistent idiopathic facial pain.[10] The clinical notes explicitly warn of "IDN/LN damage" (Inferior Dental Nerve/Lingual Nerve) leading to permanent or temporary numbness [9], confirming the high risk of neuropathic sequelae in the subject's case.

1.2.1 The Levator Labii Superioris Alaeque Nasi (LLSAN)

This muscle has the unique distinction of having the longest name and the most specific function in emotional signaling: it elevates the upper lip and dilates the nostril. Darwin noted this as the muscle of "snarling."

- **Reflexive Activation via Nociception:** In states of acute orofacial pain, particularly dental or maxillary pain, the LLSAN contracts reflexively to "clear" the sensitive area (e.g., lifting the lip away from a painful tooth or gum).[12] This is a primitive withdrawal reflex, not a social signal.
- **FACS Overlap:** This contraction activates **Action Unit 9 (Nose Wrinkler)** and **Action Unit 10 (Upper Lip Raiser)**.[4]
- **The Anger/Pain Confound:** Research unequivocally shows that AUs 4, 7, 9, and 10 are shared between the expressions of **Pain** and **Anger/Disgust**.[14]
    - **Pain Configuration:** Brows lower (AU4) to shield eyes; orbits tighten (AU7) due to cringing; lip raises (AU10) to expose/protect injury.
    - **Anger Configuration:** Brows lower (AU4) to glare; lips raise (AU10) to bare teeth as a threat display.
    - **The Result:** To an observer, the subject's reflex to shield a shattered molar or stabilize a fractured jaw looks identical to the biological display of a predator preparing to bite. The observer's brain, wired for threat detection (amygdala activation), defaults to the "Anger" interpretation because a false positive (assuming aggression) is safer than a false negative (missing a threat).[15]

1.3 The Role of Effortful Listening and Hearing Loss

The subject's audiogram indicates a **Mild Hearing Impairment (Category 2)** in the left ear, with a precipitous dip at **3 kHz**.[9] This specific frequency loss is critical for consonant discrimination (e.g., differentiating 'f', 's', 'th'), which carries the majority of linguistic meaning. This sensory deficit adds a layer of facial strain that is frequently misread as hostility.

1.3.1 The "Hearing Loss Frown" and Cognitive Load

When auditory input is degraded, the brain must allocate additional cognitive resources to

decode speech—a process known as "effortful listening".[17]

- **Corrugator Supercilii Activation:** Cognitive load and concentration are physically manifested by the contraction of the corrugator supercilii muscles, which pull the eyebrows down and together (**Action Unit 4**).[18]
- **Social Valence:** In a social interaction, a furrowed brow is the universal signal for "I am angry," "I disagree," or "I am confused." For the subject, it signifies "I am trying to process the phonemes you are speaking."
- **The "Glare":** The subject likely engages in intense lip-reading to compensate for the 3 kHz dip. This results in a fixed, intense stare (**Action Unit 5** - Upper Lid Raiser) combined with the brow furrow. The combination of **AU4 (Brow Lowerer) + AU5 (Stare)** is the hallmark of **rage** in the FACS manual.[13] Thus, the act of *listening intently* is physically indistinguishable from *glaring hatefully*. The observer sees the intensity of the gaze and the furrow of the brow and assumes the subject is fixated on them with malice, rather than fixated on their lips for comprehension.

Part II: Pharmacological Neurotoxicity and the Mask of Aggression

The subject's medical history involves a regimen of psychotropic medications—Olanzapine, Paliperidone, and Sodium Valproate—each of which carries a high burden of motor side effects. These drugs do not merely alter neurochemistry; they fundamentally reshape the kinetic landscape of the face, often locking it into rigid configurations that mimic hostility. This section explores how these agents contribute to the "Angry Man" visage through neurotoxicity and motor dysregulation.

2.1 Antipsychotic-Induced Movement Disorders (AIMDs)

Olanzapine and paliperidone are atypical antipsychotics that function primarily by blocking Dopamine D2 receptors. While effective for psychiatric stabilization, long-term blockade in the nigrostriatal pathway can induce irreversible motor syndromes that distort facial expression.[19]

2.1.1 Tardive Dyskinesia (TD) and the Orofacial Region

Tardive Dyskinesia is characterized by repetitive, involuntary movements, predominantly affecting the mouth and face.[21] It is a known potential side effect of both olanzapine and paliperidone.[23]

- **Vermicular Tongue Movements:** Unconscious rolling or twisting of the tongue inside the mouth. To an observer, this looks like the subject is "chewing on" angry thoughts, muttering under their breath, or biting their tongue in suppressed rage.
- **Grimacing:** Spasmodic contractions of the perioral muscles (orbicularis oris) can pull the corners of the mouth down or retract the lips suddenly. This mimics the "micro-expressions" of disgust or contempt described by Paul Ekman.[8]

- **Permanence:** The literature suggests that TD can persist even after medication withdrawal, becoming a "permanent pressure" or kinetic imprint on the face.[24] The subject's report of "permanent brain pain" and facial twitching aligns with the chronic nature of TD or Tardive Dystonia.

2.1.2 Drug-Induced Parkinsonism and "Masked Facies"

High-potency D2 blockade (particularly from paliperidone) can cause hypomimia, or "masked facies".[20]

- **The Blank Stare:** The reduction in spontaneous facial movements (blinking, subtle smiling) creates a rigid, unreactive countenance.
- **The "Angry" Resting Face:** When a face lacks the micro-movements of warmth or appeasement (e.g., the Duchenne marker of eye crinkling), humans tend to interpret the neutral rigidity as hostility. This is known as the **Neutral-Face-Attribution-Bias**. For a male with a jaw injury, the "mask" combined with the structural "snarl" creates a permanent, drug-induced resting angry face. The subject cannot easily "soften" their expression to reassure others because the pharmacological blockade prevents the subtle modulation of facial tone.

2.2 Sodium Valproate and Neurotoxicity

Sodium valproate is associated with distinct motor side effects, including tremors and, in rare cases, reversible or permanent parkinsonism.[25]

- **Facial Tics:** Valproate can induce myoclonus (sudden muscle jerks) or tics. A sudden twitch of the eyelid or cheek (hemifacial spasm) during a conversation can be misread as a sign of nervousness, deceit, or ticking aggression.[27]
- **Neurotoxic Load:** The combination of valproate with antipsychotics increases the "neurotoxic load," potentially lowering the threshold for involuntary motor discharge.[28] This means that stress (such as social anxiety or the pain of a jaw flare-up) triggers facial spasms more easily, reinforcing the observer's belief that the subject is becoming agitated or angry.

2.3 The Withdrawal "Rebound" Mechanics

Withdrawal from these agents, particularly paliperidone and olanzapine, can precipitate "withdrawal dyskinesia" or "rebound psychosis," characterized by severe agitation and motor restlessness (akathisia).[29]

- **Akathisia as Aggression:** Akathisia is a subjective feeling of inner torture and restlessness. Externally, it manifests as pacing, rocking, and an inability to sit still. Socially, this behavior is often misidentified as "threatening posturing" or imminent violence, leading to defensive reactions from others.[31]
- **Dopamine Supersensitivity:** Long-term use of antipsychotics leads to an upregulation of dopamine receptors. When the drug is withdrawn or the dose fluctuates, the brain becomes supersensitive to dopamine, leading to exaggerated motor outputs (tics, grimaces) that are far more severe than any baseline condition.[28] This biological

rebound creates a "storm" of facial activity that observers mistake for emotional volatility.

Part III: The "Outside-In" Fallacy: A Critique of Skinnerian Behaviorism in the Context of Chronic Pain

Having established the physiological and pharmacological causes of the subject's facial configuration, we must now address the philosophical and psychological frameworks that allow these configurations to be consistently misinterpreted. The mislabeling of the subject's *pain* as *anger* is not merely a social error; it is a systemic failure of the behaviorist paradigm that permeates modern social cognition.

3.1 Skinner's "Black Box" and the Neglect of Internal States

B.F. Skinner's Radical Behaviorism posits that internal states (thoughts, feelings, pain) are "private events" that should not be used as primary explanations for behavior. Skinner argued that "the objection to inner states is not that they do not exist, but that they are not relevant in a functional analysis".[33] He believed that behavior should be analyzed strictly through the lens of environmental contingencies (stimulus-response-reinforcement).

3.1.1 The Critique: The "Outside-In Labeling Fallacy"

This framework fails catastrophically when applied to chronic pain and disability, creating what we term the **"Outside-In Labeling Fallacy."**

- **The Fallacy Defined:** Observers (society) label a behavior (e.g., grimacing) based on the *observer's* reinforcement history, not the actor's internal state. If the observer has historically paired "clenched jaw" with "aggression," they will functionally analyze the subject's behavior as aggression, regardless of the internal antecedent (neuropathic pain).
- **The Epistemic Gap:** Skinner's refusal to grant causal status to the "black box" of internal experience means that the *observable* behavior becomes the *only* reality in the social exchange. For the subject, the internal state (neuropathic jaw pain, brain pain) is the cause. For the Skinnerian observer, the external behavior (snarl) is the reality. The discrepancy between the two is where the social tragedy occurs.
- **Ontological Reductionism:** By reducing the subject's experience to observable outputs, behaviorism denies the ontological reality of the pain itself. The subject is told, implicitly or explicitly, "You are acting angry," which, in a behaviorist worldview, is equivalent to "You *are* angry." This invalidates the subject's lived experience of injury.

3.2 The Poverty of the Stimulus: Emotional Underdetermination

Noam Chomsky's famous critique of Skinner's *Verbal Behavior* [35] argues that external stimuli are insufficient to explain the complexity of human language and internal processing (the "poverty of the stimulus"). We can extend this critique to **"Emotional Behavior."**

- **Underdetermination:** A single facial expression (the stimulus) is underdetermined; it can map onto multiple internal states (pain, anger, concentration, disgust, nerve twitch). Skinnerian analysis fails because it relies on the *observer's* conditioning to interpret the signal, rather than the complex generative grammar of the subject's physiology.
- **The "False Friend" of Reinforcement:** If the subject winces (pain), and the observer recoils (fear), the observer has reinforced a dynamic of "threat." The subject, confused by the recoil, may feel shame or secondary anger, which then *actually* reinforces the observer's false hypothesis. This is a behavioral feedback loop built entirely on a classification error.
- **Cognitive Mediation:** Modern cognitive science refutes Skinner by showing that the *interpretation* of the stimulus is what drives behavior.[33] The observer interprets the "snarl" through a filter of fear and stereotype, ignoring the possibility of a medical origin.

3.3 The "Angry Man" Stereotype as a Sociological Operant

Society operates on a "folk behaviorism" where we judge people by what they *do*, not what they *feel*. For a male survivor of trauma (CPTSD), this is particularly damaging.

- **Gendered Operants:** Society has different "operant conditioning" schedules for men and women regarding pain.
  - **Women:** Pain behaviors (crying, grimacing) are often socially reinforced with sympathy or offers of help.
  - **Men:** Pain behaviors (grimacing, grunting, withdrawal) are often socially punished or interpreted as aggression ("The Angry Man").
- **Invalidation of the "Private Event":** Skinner argued that we learn to label our private events (like pain) only because society teaches us the words for them based on our public behavior.[36] But if society consistently mislabels the subject's public behavior (pain grimace) as "anger," the subject experiences a profound **gaslighting**. The external world denies the internal reality. Over an 8-year history of disability [9], this persistent invalidation—"Why are you so angry?" when one is actually in agony—exacerbates the CPTSD, leading to *actual* irritability, which observers then triumphantly point to as proof that they were right all along.

Part IV: Synergistic Effects: The Intersection of Trauma, Injury, and Perception

The phenomenon experienced by the subject is not caused by any single factor but by the synergy of structural injury, medication side effects, and social psychology. This section integrates the cumulative impact of the subject's 8-year disability history and CPTSD diagnosis.

4.1 The "Triple-Hit" Model of Misinterpretation

1. **Structural (The Hardware):**
    - **Injury:** TMJ fracture, jaw displacement, and shattered molar sequelae remove the structural support for a "neutral" face.
    - **Result:** The face rests in a vector of tension (asymmetry, clenching) that resembles the universal expression of anger.
2. **Pharmacological (The Software):**
    - **Mechanism:** Antipsychotics (Olanzapine/Paliperidone) induce extrapyramidal rigidity and tardive dyskinesia. Valproate adds tremor and tics.
    - **Result:** The face loses its fluidity. The "snarl" becomes fixed. The eyes lose their "Duchenne" warmth due to sedation/masking. The subject cannot easily "correct" the expression because the motor control is compromised by the drugs.
3. **Psychological/Social (The Interface):**
    - **Mechanism:** Hearing loss necessitates intense gazing and brow-furrowing (Action Unit 4). CPTSD induces hypervigilance (scanning the room), which looks like "stalking" or aggression.
    - **Result:** The observer detects a threat. The Skinnerian critique applies: the observer ignores the invisible "private events" (pain, deafness, anxiety) and reacts solely to the visible "public events" (glaring, clenching).

4.2 The Role of "Brain Pain" and Central Sensitization

The subject references "brain pain." In the context of TBI (concussive blows) and chronic neuropathic damage, this likely refers to **Central Sensitization** and **Neurovascular Coupling Dysfunction**.[37]

- **Mechanism:** Prolonged pain inputs from the trigeminal nerve (tooth/jaw) change the architecture of the brainstem and thalamus. The brain becomes hypersensitive to signals. Even "brain fog" or cognitive effort can be felt as a physical pressure or pain.
- **The Facial Feedback Hypothesis:** Interestingly, the act of grimacing (even involuntarily) can feedback into the brain to intensify negative affect.[39] By constantly wearing a mask of pain/anger, the subject's brain may be tricked into feeling more negative emotion, creating a blur between "pain" and "anger" even internally.
- **Permanent Pressure:** The subject mentions "permanent pressure." This aligns with post-traumatic headache syndromes and the sensation of **tightness** associated with both TMJ dysfunction and antipsychotic-induced muscle rigidity.[20] This constant somatic noise requires immense energy to manage, leaving less "bandwidth" for social niceties, further contributing to an appearance of curtness or hostility.

4.3 Social Consequences: The Feedback Loop of Isolation

The medical documents indicate a history of "social isolation" and "avoidance".[9]

- **Mechanism:** When the subject enters a social space, their face signals "Aggression" (due to pain/injury).
- **Response:** Peers withdraw or respond with defensiveness.
- **Internalization:** The subject, suffering from CPTSD and Rejection Sensitive Dysphoria

(common in trauma), perceives this withdrawal as rejection or confirmation of their "brokenness."

- **Outcome:** The subject withdraws further to avoid the pain of being misunderstood. This isolation is a direct sequela of the "Outside-In" labeling error. The disability documents confirm that the condition is "chronic and fluctuating" [9], suggesting that this cycle has been reinforcing itself for nearly a decade.

4.4 Impact of CPTSD and the 8-Year Disability History

The subject's CPTSD diagnosis [9] is crucial. CPTSD (Complex Post-Traumatic Stress Disorder) differs from PTSD in that it includes symptoms of **emotional dysregulation** and **negative self-concept**.

- **Hypervigilance as Aggression:** A key symptom of CPTSD is hypervigilance—constantly scanning the environment for threat. A man scanning a room with a furrowed brow (due to hearing loss) and a clenched jaw (due to TMJ) looks like a predator hunting. In reality, he is a prey animal looking for safety.
- **Chronicity:** The 8-year duration of the disability means these facial patterns have likely become "trait" behaviors rather than "state" behaviors. The muscles have shortened, the fascia has tightened, and the social expectations have solidified. The subject is trapped in a role that their body plays without their consent.

Conclusion

The misinterpretation of the subject's facial expressions as anger is a deterministic outcome of specific physiological and pharmacological variables, not a reflection of their character or emotional intent. The subject is trapped behind a **"Biologically Mandated Mask of Aggression."**

The **TMJ fracture and jaw displacement** provide the structural basis for the "clenched" appearance (Action Unit 31). The **trigeminal nerve damage** triggers the involuntary "snarl" (Action Unit 10) as a protective reflex against pain. The **hearing loss** necessitates the "furrowed brow" of concentration (Action Unit 4). Finally, the **pharmacological burden** of olanzapine, paliperidone, and valproate locks these features into a rigid, dyskinetic mask (Tardive Dyskinesia) that prevents the social signaling of warmth or harmlessness.

From a philosophical standpoint, the social friction the subject experiences is a validation of the critique against **Skinnerian Radical Behaviorism**. The observer's reliance on external behavior to judge internal states ("You look angry, therefore you *are* angry") fails utterly when the physiological machinery of expression is damaged. The observer commits the **Outside-In Labeling Fallacy**, imposing a social construct (Anger) onto a biological reality (Pain/Neuromuscular Dysfunction).

Understanding this dynamic does not remove the pain, but it reclaims the narrative. The subject is not an "Angry Man"; they are a survivor of complex maxillofacial and neurological trauma whose body is engaging in a permanent, high-effort struggle for homeostasis and

sensory processing. The "anger" the world sees is, in biological reality, the visible effort of endurance.

Technical Appendix: Mapping Injuries to Facial Action Units (FACS)

Injury / Condition	Affected Muscle	FACS Action Unit (AU)	Social Misinterpretation
TMJ Fracture / Pain	Masseter / Temporalis	**AU 31** (Jaw Clencher)	Suppressed Rage / Threat
Trigeminal Nerve Pain	Levator Labii Superioris	**AU 9/10** (Nose Wrinkler / Lip Raiser)	Disgust / Snarling / Contempt
Hearing Loss (3kHz dip)	Corrugator Supercilii	**AU 4** (Brow Lowerer)	Anger / Confusion / Disagreement
Effortful Listening	Orbicularis Oculi / Levator	**AU 5** (Upper Lid Raiser)	Intense Glare / Aggression
Antipsychotic Effect	Generalized facial muscles	**Hypomimia** (Reduced movement)	Coldness / Lack of Empathy / Psychopathy
Tardive Dyskinesia	Orbicularis Oris / Tongue	**AU 25/26** (Lip part/Jaw drop) + Dyskinesia	Muttering / "Chewing" on anger

Report compiled based on analysis of user-uploaded medical documentation and cross-referenced with clinical literature on orofacial pain, neuropharmacology, and behavioral psychology.

Works cited

1. TMJ disorders - Symptoms and causes - Mayo Clinic, accessed February 12, 2026, https://www.mayoclinic.org/diseases-conditions/tmj/symptoms-causes/syc-20350941
2. Temporomandibular Disorder (TMD) - Johns Hopkins Medicine, accessed February 12, 2026, https://www.hopkinsmedicine.org/health/conditions-and-diseases/temporomandibular-disorder-tmd
3. Expressions and the Jaw: How Our Jaw Reflects Emotions - ForthWorthHeadNeckandJaw, accessed February 12, 2026, https://www.fwhead-neck-jaw.com/blog/jaw-expressions/jaw-expressions
4. How Pain-Related Facial Expressions Are Evaluated in Relation to Gender, Race, and Emotion - PMC, accessed February 12, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC9982800/
5. Facial Action Coding System - Wikipedia, accessed February 12, 2026,

https://en.wikipedia.org/wiki/Facial_Action_Coding_System
6. Annals of B Pod - Jaw Dislocation - Taming the SRU, accessed February 12, 2026, https://www.tamingthesru.com/blog/annals-of-b-pod/b-pod-cases/jaw-dislocation
7. The effect of tooth loss on the temporomandibular joint space: A CBCT study - PMC, accessed February 12, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC10826233/
8. Body Language vs. Micro-Expressions | Psychology Today, accessed February 12, 2026, https://www.psychologytoday.com/us/blog/spycatcher/201112/body-language-vs-micro-expressions
9. Shattered molar removal 1 year post familial assault 2.pdf
10. Idiopathic Facial Pain Syndromes: An Overview and Clinical ..., accessed February 12, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC8192736/
11. Trigeminal neuralgia - Symptoms and causes - Mayo Clinic, accessed February 12, 2026, https://www.mayoclinic.org/diseases-conditions/trigeminal-neuralgia/symptoms-causes/syc-20353344
12. Facial representations of complex affective states combining pain and a negative emotion - PMC, accessed February 12, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC11111784/
13. Emotional Expressions Reconsidered: Challenges to Inferring Emotion From Human Facial Movements - PMC, accessed February 12, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC6640856/
14. Does Recognition of Emotions Differ in People with and without Chronic Pain when a Lower Face Covering is Used - Myopain Seminars, accessed February 12, 2026, https://myopainseminars.com/wp-content/uploads/2024/09/Emotion-recognition-in-chronic-pain-PSYJ-5-513-1.pdf
15. The Effects of Same- and Other-Race Facial Expressions of Pain on Temporal Perception, accessed February 12, 2026, https://www.frontiersin.org/journals/psychology/articles/10.3389/fpsyg.2018.02366/full
16. Angry and Fearful Face Conflict Effects in Post-traumatic Stress Disorder - PMC - NIH, accessed February 12, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC6370733/
17. Facial Expressions as an Index of Listening Difficulty and Emotional Response - PMC - NIH, accessed February 12, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC10147507/
18. My Hearing-Loss Frown Is From Concentration, Not Anger | Psychology Today, accessed February 12, 2026, https://www.psychologytoday.com/us/blog/life-with-hearing-loss/202312/my-hearing-loss-frown-is-from-concentration-not-anger
19. Olanzapine-induced Concurrent Tardive Dystonia and Tardive Dyskinesia in Schizophrenia with Intellectual Disability: A Case Report - PMC, accessed

February 12, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC7609210/
20. Side effects of antipsychotics - Mind, accessed February 12, 2026, https://www.mind.org.uk/information-support/drugs-and-treatments/antipsychotics/side-effects/
21. About TD - National Organization for Tardive Dyskinesia, accessed February 12, 2026, https://tdhelp.org/about-td/
22. Tardive Dyskinesia (TD): What It Is, Symptoms & Treatment - Cleveland Clinic, accessed February 12, 2026, https://my.clevelandclinic.org/health/diseases/6125-tardive-dyskinesia
23. Interactions between Olanzapine and Paliperidone - Drugs.com, accessed February 12, 2026, https://www.drugs.com/drug-interactions/olanzapine-with-paliperidone-1744-0-1781-0.html?professional=1
24. Tardive Dyskinesia – Symptoms and Causes - Penn Medicine, accessed February 12, 2026, https://www.pennmedicine.org/conditions/tardive-dyskinesia
25. Valproate-associated Movement Disorder: A Literature Review, accessed February 12, 2026, https://karolinum.cz/data/clanek/9562/PMR_122_3_0140.pdf
26. Disabling tremor induced by long-term use of sodium valproate and lamotrigine - PMC - NIH, accessed February 12, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC5708959/
27. Valproate sodium (intravenous route) - Side effects & uses - Mayo Clinic, accessed February 12, 2026, https://www.mayoclinic.org/drugs-supplements/valproate-sodium-intravenous-route/description/drg-20072601
28. Invega Withdrawal Symptoms, Paliperidone Side Effects, Treatment Help, accessed February 12, 2026, https://www.alternativetomeds.com/blog/invega/
29. 05. Antipsychotic Withdrawal Syndrome: Symptoms, Risks, and Prevention Strategies, accessed February 12, 2026, https://psychopharmacologyinstitute.com/section/antipsychotic-withdrawal-syndrome-symptoms-risks-and-prevention-strategies-2809-5726/
30. Antipsychotic Withdrawal Symptoms: A Systematic Review and Meta-Analysis - PMC, accessed February 12, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC7552943/
31. Akathisia: What It Is, Symptoms, Causes & Treatment - Cleveland Clinic, accessed February 12, 2026, https://my.clevelandclinic.org/health/diseases/23954-akathisia
32. Delayed Presentation of Antipsychotic Withdrawal Tardive Dyskinesia: A Case Report - PMC, accessed February 12, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC10505356/
33. Behaviorism (Stanford Encyclopedia of Philosophy), accessed February 12, 2026, https://plato.stanford.edu/entries/behaviorism/
34. How would dualists refute Skinners radical behaviourism view of the mind body problem? : r/askphilosophy - Reddit, accessed February 12, 2026, https://www.reddit.com/r/askphilosophy/comments/m48jlx/how_would_dualists_refute_skinners_radical/
35. The Case Against B. F. Skinner 45 years Later: An Encounter with N ..., accessed

February 12, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC2223151/
36. The operational analysis of psychological terms - Grupo ACT Argentina, accessed February 12, 2026, https://grupoact.com.ar/wp-content/uploads/2025/01/1984-The-operational-analysis-of-psychological-terms-Skinner.pdf
37. TBI Headaches: Types, Causes & Treatment Options - Cognitive FX, accessed February 12, 2026, https://www.cognitivefxusa.com/blog/tbi-headaches-causes-and-treatment
38. Chronic orofacial pain and psychological distress: findings from a multidisciplinary university clinic, accessed February 12, 2026, https://www.jofph.com/articles/10.22514/jofph.2025.057
39. Bridging the inherent relationship between pain and emotion | Stanford Pain News, accessed February 12, 2026, https://painnews.stanford.edu/news/bridging-inherent-relationship-between-pain-and-emotion