# Forensic and Biomechanical Reconstruction of Cervicothoracic Trauma: Analysis of Manual Strangulation, Respiratory Arrest, and Delayed Spinal Sequelae

## 1. Introduction: The Biomechanics of Asymmetric Strangulation Vectors

The forensic reconstruction of non-fatal strangulation (NFS) requires a precise analysis of the forces applied relative to the anatomical position of the victim and the assailant. The scenario presented—a bimanual compression of the anterior neck targeting the laryngeal prominence (thyroid cartilage/Adam's apple), combined with a downward traction force applied by an assailant of shorter stature—constitutes a complex, multi-vector trauma. This specific mechanism differs significantly from static compression. The superimposition of a downward vector (45° angle) upon a compressive vector creates a shear force at the cervicothoracic junction (C7-T1) while simultaneously compromising the integrity of the laryngotracheal complex.[1]

This report evaluates the physiological plausibility of the survivor's account, specifically addressing the sensation of structural failure ("crack"), the phenomenology of respiratory paralysis ("brain trying to talk to lungs"), the restorative mechanics of the retching reflex, and the correlation between the described assault vector and the subsequent radiological finding of spinal degeneration at the C7-T1 level. The synthesis of biomechanical data, neurophysiology, and trauma literature supports a high probability that the described events represent a sequence of laryngeal fracture, phrenic nerve neurapraxia, and flexion-distraction injury to the lower cervical spine.

### 1.1 The Physics of the "Downward Pull"

When a shorter assailant applies force to the neck of a taller victim, the force vector is rarely purely horizontal (anterior-to-posterior). Instead, the shorter stature necessitates a vector that is directed posterior-inferiorly.

- **The Fulcrum Effect:** The cervical spine acts as a cantilever beam. A downward force applied at the level of the thyroid cartilage (C4-C5) creates a significant bending moment. The fulcrum of this lever arm is not the mid-neck, but the transition zone where the mobile cervical spine meets the rigid thoracic spine—specifically, the C7-T1 junction.[3]

- **Force Magnification:** The "intergenerational trauma" and acute stress response described in the assailant suggests the recruitment of sympathetic nervous system reserves (hysterical strength). This can elevate manual grip strength well beyond the 4.4–6 kg (roughly 10–13 lbs) required to occlude the jugular veins or the 15 kg (33 lbs) required to fracture the trachea.[5] The combination of this grip strength with the weight of the assailant hanging or pulling downward amplifies the traction force on the cervical ligaments.

# 2. Structural Trauma: The "Crack" and Laryngotracheal Fracture

The survivor reports a distinct sensation of a "crack" immediately followed by respiratory failure. In the context of "two thumbs into the Adam's apple," this is pathognomonic of a fracture to the thyroid cartilage or the hyoid-larynx complex.

## 2.1 Thyroid Cartilage Fracture Mechanics

The thyroid cartilage ("Adam's apple") houses the vocal cords and protects the airway. It is suspended from the hyoid bone by the thyrohyoid membrane.

- **Fracture Thresholds:** Biomechanical studies indicate that the thyroid cartilage is vulnerable to direct anterior compression, particularly in the calcified cartilage of adults. The thumbs, providing point-specific pressure, can exert force exceeding 30 psi, easily sufficient to fracture the superior cornua (horns) or the lamina of the thyroid cartilage.[5]
- **The "Crack" Sensation:** The reported "crack" likely corresponds to one of three events:
    1. **Fracture of the Superior Thyroid Horn:** This structure is susceptible to lateral compression and downward traction. A fracture here disrupts the laryngeal suspension.[7]
    2. **Cricoarytenoid Joint Subluxation:** The downward pull can dislocate the arytenoid cartilages from the cricoid ring, a profound injury that can mechanically lock the vocal cords in a closed position.[9]
    3. **Ligamentous Avulsion:** The "crack" may also represent the avulsion of the nuchal ligament or a spinous process at the posterior neck (Clay Shoveler's fracture) due to the violent flexion.[11]

The immediate cessation of breathing following this sound strongly suggests that the structural damage directly impinged upon the airway's functional integrity, transitioning the trauma from a structural injury to a functional respiratory arrest.

# 3. Neurophysiology of Respiratory Paralysis vs. Obstruction

The survivor's description of the respiratory failure is clinically distinct and highly specific: *"It felt like I could breathe, but I couldn't breathe because I tried to activate my lungs and nothing happened... brain was trying to talk to lungs but nothing worked."* This is *not* the sensation of

a mechanical obstruction (choking), where the victim heaves violently against a blockage. Rather, it describes a failure of respiratory drive or neuromuscular transmission.

## 3.1 Mechanism of Phrenic Nerve Neurapraxia

The most plausible explanation for the sensation of "paralysis" is a transient injury to the phrenic nerves.

- **Anatomical Vulnerability:** The phrenic nerves (C3-C5) run vertically across the anterior scalene muscles, deep to the sternocleidomastoid. They provide the *sole* motor innervation to the diaphragm.[13]
- **Mechanism of Injury:** A deep, digging thumb pressure into the "Adam's apple" (C4-C5 level) combined with a violent downward traction (stretching the neck) can compress or stretch the phrenic nerves against the cervical vertebrae.[15]
- **Neuropraxia (The "Stun"):** This trauma can cause neurapraxia—a temporary conduction block where the nerve is intact but fails to transmit signals. Bilateral phrenic nerve neurapraxia results in immediate diaphragmatic paralysis.[16]
- **Sensory Correlation:** In this state, the brain sends the signal to breathe, but the "pump" (diaphragm) does not respond. The accessory muscles (intercostals) may not be sufficient to generate tidal volume, leading to the terrifying sensation of a disconnect between volition and lung actuation.[18]

## 3.2 Laryngospasm and the "Silent Chest"

Concurrently, the trauma likely triggered a severe **laryngospasm**.

- **Reflex Arc:** Direct pressure on the larynx or the superior laryngeal nerve triggers a primitive protective reflex, clamping the vocal cords shut to prevent aspiration.[20]
- **The "Ball-Valve" Effect:** In severe laryngospasm, the airway is hermetically sealed. If the diaphragm is also compromised (phrenic stun), the chest wall may remain motionless, contributing to the sensation that "nothing was happening".[22]
- **Neurogenic Shock:** Strangulation can induce a vagal autonomic storm, potentially leading to bradycardia and a transient state of neurogenic shock, further depressing respiratory drive and altering consciousness.[23]

## Table 1: Differentiating Respiratory Failure Mechanisms in Strangulation

| **Clinical Feature** | **Mechanical Obstruction (Choking)** | **Laryngospasm** | **Phrenic Nerve Paralysis (Neuropraxia)** |
|---|---|---|---|
| **Airway Status** | Physically blocked by object/swelling | Glottis spastically closed | Patent (Open) or Closed |
| **Respiratory Effort** | Intense heaving *against* blockage | High effort (stridor) or silent | **Absent or ineffective diaphragmatic movement** |

| | | | |
|---|---|---|---|
| **Sensation** | "Something is stuck" | "Throat is locking up" | **"Brain talking to lungs but nothing happening"** |
| **Anatomical Target** | Trachea/Pharynx | Laryngeal Nerve/Vocal Cords | C3-C5 Nerve Roots/Scalene Muscles |
| **Reversibility** | Requires object removal | Requires positive pressure/reflex break | Requires nerve recovery/accessory muscle use |

# 4. The Mechanics of Survival: Emesis as a Resuscitative Event

The survivor questions whether it is possible to survive a lethal squeeze if they "puked or force air up the oesophagus." The answer is affirmatively positive based on respiratory physiology. The act of heaving (vomiting) acted as a biological override to the paralysis.

## 4.1 "Larson's Maneuver" and the Breaking of Spasm

Medical protocols for breaking laryngospasm often involve applying painful pressure to the "laryngospasm notch" (Larson's point).[25] However, the physiological forces generated during retching are far more potent.

- **The Pressure Wedge:** Retching generates massive intra-abdominal and intrathoracic pressure (up to 200 cmH2O). As the survivor "tried to heave her through my stomach," this pressure wave forced gastric content and air up the esophagus.[26]
- **The Abduction Reflex:** While the glottis initially closes during the build-up of a vomit, the *expulsive phase* requires the coordination of the upper esophageal sphincter and the airway. The violent blast of air and fluid from below can mechanically force the adducted vocal cords apart, breaking the negative-pressure lock of the laryngospasm.[27]

## 4.2 Bypass of Phrenic Paralysis via Accessory Muscles

If the diaphragm was paralyzed (phrenic injury), the survivor’s conscious decision to "heave" utilized a different neural pathway.

- **Neural Rerouting:** The muscles of vomiting (abdominal rectus, obliques) are innervated by the **thoracic spinal nerves (T6-L1)**, not the cervical phrenic nerves.[13]
- **Manual Ventilation:** By violently contracting the abdominal muscles to retch, the survivor manually compressed the abdominal viscera, pushing the paralyzed diaphragm upward into the chest. When the muscles relaxed, the diaphragm dropped back down due to gravity and elastic recoil, creating a passive inhalation. This "abdominal piston" effect is a known compensatory mechanism in high spinal cord injury patients.[29]
- **Survival Implication:** The "heave" effectively acted as a manual ventilator, jump-starting the respiratory cycle and breaking the neurological deadlock.

### 4.3 Reduction of Arytenoid Subluxation

If the "crack" was a subluxation (dislocation) of the arytenoid cartilage blocking the airway, the violent muscular contractions of the pharynx during vomiting could have essentially "popped" the cartilage back into place (reduction). The dilation of the pharynx required to pass vomitus is maximal, potentially resolving a mechanical lock of the cricoarytenoid joint.[9]

## 5. Hemorrhagic Sequelae: The Origin of the Blood

The expulsion of blood ("blood came out") upon the restoration of breathing is a critical indicator of internal trauma. Differentiating the source confirms the severity of the event.

### 5.1 Laryngeal Trauma (Hemoptysis)

Given the direct compression of the thyroid cartilage, the blood likely originated from a **mucosal laceration** of the larynx or trachea.

- **Mechanism:** As the cartilage fractured or deformed against the cervical spine, the delicate mucous membranes were crushed or torn. This is common in manual strangulation even without external signs.[32]
- **Significance:** If the blood was bright red and frothy, or appeared *with* the first breath, it indicates direct airway injury.

### 5.2 Mallory-Weiss Tear (Hematemesis)

Alternatively, the violence of the "heave" required to break the spasm may have caused a **Mallory-Weiss tear**.

- **Mechanism:** This is a longitudinal tear at the junction of the esophagus and stomach caused by severe retching. The sudden increase in intragastric pressure against a closed or spastic glottis/esophagus tears the lining.[34]
- **Correlation:** The survivor mentions the heave "went up my oesophagus and then I could breathe again and blood came out." This sequence allows for a Mallory-Weiss tear to be a secondary injury necessary for survival—a cost paid to restore respiration.

## 6. Long-Term Sequelae: The C7-T1 Spinal Degeneration

The survivor reports a later X-ray showing "spinal degeneration" specifically at the C7-T1 junction (where the neck meets the shoulders). This finding is highly consistent with the biomechanics of a **downward traction injury** and is likely a misidentified or healed traumatic lesion rather than age-related degeneration.

### 6.1 The "Clay Shoveler's" Mechanism

The described vector—a shorter person pulling the neck down at a 45° angle—replicates the exact forces involved in a **Clay Shoveler's Fracture**.

- **Avulsion Physics:** This fracture typically occurs when the head is forced into flexion

against the resistance of the supraspinous ligaments. The downward pull by the mother forced the survivor's chin toward the chest while the assailant's weight provided the shear force.[11]

- **The Injury:** This results in an avulsion fracture of the spinous process of C7 or T1. Because these fractures are stable (they do not cause immediate paralysis), they are often missed if the patient does not seek immediate trauma care. The survivor noted they got an X-ray "eventually later".[37]
- **Chronic Appearance:** An untreated avulsion fracture or severe ligamentous tear (whiplash-associated disorder) at C7-T1 will heal with calcification, osteophytes (bone spurs), and disc space narrowing. To a radiologist viewing this years later, this scarring looks indistinguishable from "degenerative disc disease" or spondylosis.[38]

## 6.2 The Cervicothoracic Fulcrum

The C7-T1 junction is the hardest part of the cervical spine to image on standard X-rays and is a common site for "occult" (hidden) injuries.

- **Traction-Distraction:** The assailant's grip on the thyroid cartilage (anterior) combined with the downward pull creates a distraction force (pulling apart) on the posterior elements of the C7-T1 vertebrae.[40]
- **Causal Link:** In a young or otherwise healthy individual, isolated degeneration at the C7-T1 level is statistically unlikely to be natural. It is a hallmark of prior trauma. The specific "45-degree pull down" is the smoking gun for this specific location, as it concentrates the moment arm of the lever exactly at the cervicothoracic junction.[4]

## Table 2: Biomechanical Correlation of Assault Vector and Injury

| Assault Component | Physical Force Vector | resulting Pathological Lesion | Long-Term Radiographic Finding |
|---|---|---|---|
| **"Two thumbs in Adam's Apple"** | Direct Anterior-Posterior Compression (>30 psi) | Thyroid Cartilage Fracture; Laryngospasm; Phrenic Neuropraxia | Healed laryngeal calcification; Voice changes |
| **"Shorter Mother" (Height Differential)** | Inferior (Downward) Traction | Depression of Hyoid/Larynx; Traction on Cervical Roots | C7-T1 Instability |
| **"Pulled down at 45° angle"** | Flexion-Distraction Moment | **Clay Shoveler's Fracture (C7/T1)**; Nuchal Ligament Tear | **"Spinal Degeneration" at C7-T1** (Osteophytes, Spondylosis) |
| **"Heave through stomach"** | Intra-abdominal Pressure Spike (>200 cmH2O) | Mechanical Abduction of Glottis; Mallory-Weiss Tear | Gastric/Esophageal scarring |

# 7. Conclusion: Synthesis of Probability

The sequence of events described—from the "crack" and specific sensation of paralytic suffocation to the restorative heave and subsequent spinal pathology—is not only possible but physiologically consistent with a high-energy manual strangulation event.

1. **Airway Collapse:** The "crack" indicates a fracture of the laryngotracheal complex, which precipitated an immediate **laryngospasm** and/or **phrenic nerve neurapraxia**. This explains the unique sensation of the "brain trying to talk to the lungs" without effect, differentiating it from simple obstruction.[17]
2. **Survival via Emesis:** The "heave" was a critical survival reflex. The recruitment of thoracic and abdominal muscles bypassed the paralyzed phrenic pathways and generated sufficient pressure to mechanically break the laryngospasm or reduce a laryngeal subluxation, restoring airway patency.[26]
3. **Spinal Evidence:** The "spinal degeneration" at C7-T1 is the chronic radiological footprint of the acute trauma. The downward, 45-degree traction applied by the shorter assailant exerted maximum shear force at the cervicothoracic junction, likely causing a **Clay Shoveler's variant fracture** or severe ligamentous tearing that healed as post-traumatic spondylosis.[11]

The presence of blood confirms breach of the mucosa (either airway or esophageal), serving as a biological marker of the extreme pressures involved. The survivor's account aligns with the known pathophysiology of "near-miss" fatal strangulation, where internal injuries are severe despite the victim's survival.

### Works cited


1. Strangulation, Domestic Violence, and Brain Injury: An Introduction to a Complex Topic, accessed February 14, 2026, https://biausa.org/public-affairs/media/strangulation-domestic-violence-and-brain-injury-an-introduction-to-a-complex-topic
2. Strangulation Injuries - StatPearls - NCBI Bookshelf - NIH, accessed February 14, 2026, https://www.ncbi.nlm.nih.gov/sites/books/NBK459192/
3. Cervicothoracic junction - posterior fixation for B2 Posterior tension band injury (bony capsuloligamentous, ligamentous) - AO Surgery Reference, accessed February 14, 2026, https://surgeryreference.aofoundation.org/spine/trauma/subaxial-cervical/b2/cervicothoracic-junction-posterior-fixation
4. CHAPTER 7: THE CERVICAL SPINE - Chiro.org, accessed February 14, 2026, https://chiro.org/ACAPress/Biomechanics_of_Cervical_Spine.html
5. Dear Doctor Q: Doctor Smock can you please tell me about the ..., accessed February 14, 2026, https://www.allianceforhope.org/file-downloads/574
6. Facts Victims of Choking (Strangulation) Need to Know! - pottstown.org, accessed February 14, 2026,

https://www.pottstown.org/DocumentCenter/View/211/Strangulation-Information-for-Victims?bidId=
7. Biomechanics of a previously unknown thyroid cartilage fracture in fatal strangulation, accessed February 14, 2026, https://pubmed.ncbi.nlm.nih.gov/34915339/
8. Anthropological analysis of trauma in throat bone and cartilage: A review - PMC - NIH, accessed February 14, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC7452381/
9. Arytenoid Subluxation - StatPearls - NCBI Bookshelf - NIH, accessed February 14, 2026, https://www.ncbi.nlm.nih.gov/books/NBK544264/
10. Arytenoid Dislocation: A Diagnostic Strategy - Laryngopedia, accessed February 14, 2026, https://laryngopedia.com/arytenoid-dislocation-a-diagnostic-strategy/
11. The clay-shoveler's fracture, a rare occupational injury in the modern era: a case report, accessed February 14, 2026, https://www.springermedizin.de/the-clay-shoveler-s-fracture-a-rare-occupational-injury-in-the-m/26985716
12. Clay shoveler's fracture - Medical News Today, accessed February 14, 2026, https://www.medicalnewstoday.com/articles/clay-shovelers-fracture
13. Diaphragmatic Palsy - PMC, accessed February 14, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC5871962/
14. What Does Phrenic Nerve Damage Feel Like? - The Institute for Advanced Reconstruction, accessed February 14, 2026, https://www.advancedreconstruction.com/blog/phrenic-nerve-damage-symptoms-what-it-feels-like-when-to-act
15. What is phrenic neuropathy? - Michigan Medicine, accessed February 14, 2026, https://www.michiganmedicine.org/health-lab/what-phrenic-neuropathy
16. Paralyzed Diaphragm | Cedars-Sinai, accessed February 14, 2026, https://www.cedars-sinai.org/health-library/diseases-and-conditions/p/paralyzed-diaphragm.html
17. What Are The Different Types of Phrenic Nerve Injury and Diaphragm Paralysis?, accessed February 14, 2026, https://www.advancedreconstruction.com/blog/types-of-phrenic-nerve-injury-and-diaphragm-paralysis
18. Paralyzed Diaphragm – Symptoms and Causes - Penn Medicine, accessed February 14, 2026, https://www.pennmedicine.org/conditions/paralyzed-diaphragm
19. Diaphragm Paralysis Overview: Symptoms, Diagnosis, & Treatments | IFAR, accessed February 14, 2026, https://www.advancedreconstruction.com/nerve-reconstruction/diaphragm-paralysis
20. 3-6. Laryngospasm and stridor - Right Decisions - NHS Scotland, accessed February 14, 2026, https://rightdecisions.scot.nhs.uk/sjh-theatres-and-anaesthetics/anaesthetic-emergencies/3-6-laryngospasm-and-stridor/
21. Update on the Management of Laryngospasm - SciELO, accessed February 14,

2026, https://www.scielo.org.mx/scielo.php?script=sci_arttext&pid=S2448-87712018000200012
22. Laryngospasm • LITFL Medical Blog • CCC Ventilation, accessed February 14, 2026, https://litfl.com/laryngospasm/
23. The Probability of Neck Organ Block Fractures From Hanging, Ligature and Manual Strangulation - Florida Gulf Coast University, accessed February 14, 2026, https://scholarscommons.fgcu.edu/esploro/outputs/graduate/The-Probability-of-Neck-Organ-Block/99383341659706570
24. Strangulation - WikEM, accessed February 14, 2026, https://wikem.org/wiki/Strangulation
25. Trick of the Trade: Laryngospasm notch maneuver - Academic Life in Emergency Medicine, accessed February 14, 2026, https://www.aliem.com/trick-of-trade-laryngospasm-notch/
26. Mechanisms of airway protection during retching, vomiting, and swallowing. Am J Physiol Gastrointest Liver Physiol 283:G529-G536 - ResearchGate, accessed February 14, 2026, https://www.researchgate.net/publication/11206540_Mechanisms_of_airway_protection_during_retching_vomiting_and_swallowing_Am_J_Physiol_Gastrointest_Liver_Physiol_283G529-G536
27. Mechanisms of airway protection during retching, vomiting, and swallowing - PubMed - NIH, accessed February 14, 2026, https://pubmed.ncbi.nlm.nih.gov/12181164/
28. Choking and laryngospasm: Exploring commonalities and treatment strategies - PMC - NIH, accessed February 14, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC11864063/
29. Cervical Spinal Cord Injury and Respiratory Function - Total Community Care, accessed February 14, 2026, https://totalcommunitycare.co.uk/cervical-spinal-cord-injury-and-respiratory-function/
30. Respiratory problems and management in people with spinal cord injury - PMC, accessed February 14, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC5335574/
31. How to: A Novel Five-Step Reduction Technique of Arytenoid Dislocation - ENTtoday, accessed February 14, 2026, https://www.enttoday.org/article/how-to-a-novel-five-step-reduction-technique-of-arytenoid-dislocation/
32. Unexpected delayed death after manual strangulation: need for careful examination in the emergency room, accessed February 14, 2026, https://www.monaldi-archives.org/macd/article/view/359
33. Laryngeal Injury - StatPearls - NCBI Bookshelf, accessed February 14, 2026, https://www.ncbi.nlm.nih.gov/books/NBK556150/
34. Mallory-Weiss Tear | Johns Hopkins Medicine, accessed February 14, 2026, https://www.hopkinsmedicine.org/health/conditions-and-diseases/malloryweiss-tear
35. Mallory-Weiss Syndrome - StatPearls - NCBI Bookshelf, accessed February 14,

2026, https://www.ncbi.nlm.nih.gov/books/NBK538190/
36. Clay Shoveler's Fracture: A Pain in the Neck - PubMed, accessed February 14, 2026, https://pubmed.ncbi.nlm.nih.gov/39935987/
37. Clay-Shoveler Fracture (Cervical Spinous Process Fractures) - Spine - Orthobullets, accessed February 14, 2026, https://www.orthobullets.com/spine/12284/clay-shoveler-fracture-cervical-spinous-process-fractures
38. Traumatic Low Back Disc Injuries on Imaging | Passen Powell Jenkins I Trial Lawyers, accessed February 14, 2026, https://www.passenpowell.com/traumatic-low-back-disc-injuries-on-imaging/
39. The Influence of Pre-Accident Degenerative Joint Disease - ChiroTrust, accessed February 14, 2026, https://chiro-trust.org/advanced/influence-pre-accident-degenerative-joint-disease/
40. Treatment of chronic traumatic C7–T1 grade III spondylolisthesis with mild neurological deficit: case report - PMC, accessed February 14, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC5386908/
41. Reduction of cervicothoracic spondyloptosis in an ambulatory patient: when traction fails, accessed February 14, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC10477183/
42. Traumatic Fracture: Dislocation of Cervicothoracic Junction—Grand Round Presentation of C7-T1 Instabilities and Different Instrumentation Techniques - NIH, accessed February 14, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC7349622/