

Universidade de São Paulo  
Instituto de Física

# Acoplamento Kondo-Majorana em pontos quânticos duplos

Jesús David Cifuentes Pardo

Orientador: Prof. Dr. Luis Gregório Godoy de  
Vasconcellos Dias da Silva

Dissertação de mestrado apresentada ao Instituto de  
Física da Universidade de São Paulo, como requisito  
parcial para a obtenção do título de Mestre em Ciências.

Banca Examinadora:

Prof. Dr. Luis Gregório Godoy de Vasconcellos Dias da Silva (IFUSP)

Prof. Dr. Eric de Castro e Andrade (IFSC/USP)

Prof. Dr. Antonio Carlos Ferreira Seridonio (UNESP -Ilha Solteira)

São Paulo  
2019



University of São Paulo  
Physics Institute

# Kondo-Majorana coupling in double quantum dots

Jesús David Cifuentes Pardo

Supervisor: Prof. Dr. Luis Gregório Godoy de Vasconcellos Dias da Silva

Dissertation submitted to the Physics Institute of the University of São Paulo in partial fulfillment of the requirements for the degree of Master of Science.

Examining Committee:

Prof. Dr. Luis Gregório Godoy de Vasconcellos Dias da Silva (IFUSP)  
Prof. Dr. Eric de Castro e Andrade (IFSC/USP)  
Prof. Dr. Antonio Carlos Ferreira Seridonio (UNESP -Ilha Solteira)

São Paulo  
2019



*To my family*

*To that little cheek*

# Acknowledgments

This piece summarizes the main aspects of my work during these two lovely years in Brazil. This has been a wonderful that will always remain in my mind. While working in this thesis, I not only learned about physics, condensed matter and quantum computing, but I also acquired experience in independent research , which allowed me to improve my coding, writing, oral and drawing skills. This manuscript, our upcoming paper on "Majorana manipulation in double quantum dots" and the thesis presentation performed on May 8,2019, are the final outcomes of this enriching experience.

Furthermore, I was gladly surprised by the large condensed matter community in Brazil, which allowed me to participate in several conferences, present my advances, and increase my background. In my way, I met up to three Nobel prizes, including Duncan Haldane, whose work inspired me to study topological materials. Apart from the academic life, I also learned Portuguese and got the chance to get immersed in depth into the amazing Brazilian culture, I loved their music (specially forró) and meet amazing people. Today, I look backwards and I get marveled by all the amazing things that happened in these years and those who allowed everything to be possible.

Therefore, in this brief statement, I want to thank all the wonderful people that received me here in Brazil: My thesis advisor Prof. Luis Gregorio, who suggested to me this project, helped me enter into the USP and guided me ever since with patience and care. I thank my previous advisors, Andres and Cesar, and all my former professors in Brazil and Colombia who prepared for this life of research. I thank to my friends and my room partners, all of them have contributed to make this stage of my life an amazing adventure. And finally, but also very important, I would like to thank my family and that little cheek. Even from far away I have always felt their support, which endeavored me to complete this goal.

I also acknowledge the financial support by: 1) Grants No. 307107/2013 – 2 and 449148/2014 – 9, Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq). 2) Grant No. 2016/18495 – 4, Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP). The Institute of Physics (IFUSP). These grants allowed us to develop and present partial and complete results of this project in workshops and conferences. Without this support nothing of this would have ever been possible. Thank you all.

*"I believe that beginning research is quite similar to that magical situation when babies learn how to talk. Initially, they are just there, listening to conversations. They seem disoriented with the vocabulary, but they are paying attention and they are catching sounds. Soon enough, before you even notice, their own words start to flow."*

Memories from my first research meeting with Professors Andrés Reyes and A.P. Balachandran. Hear me talk.

# Abstract

Majorana zero modes (MZMs) emerging at the edges of topological superconducting wires are a promising platform for fault-tolerant quantum computation. Novel proposals use quantum dots (QDs) coupled to the end of these wires to detect Majorana signatures. This detection method provides the following advantages: 1) This device allows to study the prospective coexistence of Kondo-Majorana signatures, which have been recently reported in experiments. 2) Today's precise experimental control over QDs offers the unique possibility of manipulating MZMs inside multi-dot systems. This innovative idea has enlightened the design of scalable quantum architectures, bringing us closer to the implementation of a topological quantum computer.

The simplest case where Majorana manipulation is possible is in a double quantum dot (DQD). This system offers several possibilities for manipulation of MZMs, including different geometric configurations of the dots, from symmetric and linear couplings to T-dot junctions. In this project, we perform a theoretical study of the transitions of the Majorana signature in these geometries in non-interacting and interacting regimes. By tuning the dot's gate voltages, we will show that it is possible to control the localization of the MZM inside both dots. We will also explore the interplay of these signatures with the Kondo effect, which emerges in non-interacting dots in superposition with the MZM.

We adopt two methods in this project: 1) The Green equations of motion (EOM) allow us to obtain exact expressions for the density of states in coulomb-non-interacting systems. We present the Graph -Gauss-Jordan elimination process as a simple-graphical method to solve the emergent linear systems in the EOM. 2) We use Wilson's numerical renormalization group (NRG) in interacting systems, to study the combined Kondo-Majorana physics. We will test these methods, first in a double quantum dot (DQD) (chapter 3) and then in a QD-Majorana model (chapter 4), where we confirm the results of previous papers [1–3]. Finally, we include the main contribution of this thesis, the study of a DQD coupled to a Majorana chain (chapter 5).

**Keywords:** Double quantum dot; Majorana; Manipulation; Kondo; Spectral density.

# Resumo

O uso das quasi-partículas de Majorana que emergem nas bordas de um supercondutor topológico é uma plataforma promissora para computação quântica. Novas propostas usam quantum dots (QDs) para detectar sinais de Majorana. Este método tem duas vantagens: 1) Os QDs são os melhores dispositivos para estudar a co-existência de Kondo e Majorana, a qual têm sido reportada recentemente em experimentos. 2) O controle experimental preciso sobre os quantum dots que temos hoje em dia oferece a oportunidade única para manipular quasi-partículas de Majoranas dentro de sistemas com vários dots. Esta ideia abriu novos caminhos para o desenho de arquiteturas quânticas, nos aproximando do objetivo de implementar um computador quântico topológico.

O caso mais simples em que se é possível manipular tais quasi-partículas é num quantum dot duplo (DQD). Este modelo oferece várias possibilidades para mover os Majoranas, incluindo múltiplas configurações geométricas dos dots como acoplamentos simétricos, lineares e em junções T. Neste trabalho vamos apresentar uma análise teórica das transições dos sinais de Majorana dentro do DQD em sistemas interagentes e não interagentes. Vamos ver que é possível controlar a localização dos modos zero de Majorana mediante o incremento nas voltagens de gate dos QDs. Também vamos explorar como esses sinais interagem com o efeito Kondo que emerge em superposição com o modo zero de Majorana.

Principalmente, vamos a usar dois métodos neste projecto: 1) Usamos as equações de movimento no formalismo de funções de Green para obter expressões exatas para a densidade de estados em sistemas não interagentes. Vamos apresentar o método the eliminação de Gauss-Jordan com grafos, o qual permite resolver rapidamente o sistema linear emergente nas equações de movimento. 2) Em sistemas Coulomb interagentes usamos NRG, no qual podemos observar a interação entre o Majorana e o efeito Kondo. Vamos testar ambos os métodos nos modelos de um double quantum dot e um QD acoplado com uma cadeia de Majorana, com o qual vamos reproduzir os resultados presentes na literatura. Finalmente, incluímos a maior contribuição deste trabalho, o estudo de um DQD acoplado a uma cadeia de Majorana.

**Palavras chaves:** Ponto quântico duplo; Majorana; Manipulação; Kondo; Densidade espectral.

# List of Figures

1.1	Up: Failed topological evolution from a cup of coffee to a sphere. Down: Smooth evolution of a coffee cup into a donut. Taken from <a href="https://bit.ly/2ETicqM">https://bit.ly/2ETicqM</a>	1
1.2	Illustration of the Kitaev chain for open boundary conditions in the Majorana representation. Dirac fermions are depicted by the big globes. Each one of them contains two Majoranas (small blue and red circles) a)Represents the trivial case where the hopping and the superconducting term approaches to 0. b) The non-trivial topological phase. The coupling is produced between Majoranas in different Dirac fermions By the author	4
1.3	Spectrum of Hamiltonian Equation 1.5 with 30 sites and $t =  \Delta $ s. Method: Numerical diagonalization. By the author	5
1.4	Path of $\hat{H}_k$ for the interval $[-\pi, \pi]$ . a) Trivial phase: The resulting path can be homotopically deformed to a point. b) Non-trivial phase: The path is a non-contractible loop around the unitary circle. By the author	7
1.5	Representation of non-abelian braiding	9
1.6	(a) Setup of a Majorana wire with high spin-orbit coupling, over a superconducting plane and an induced Zeeman field. (b) Splitting of the energy bands with different spins in the non-superconducting regime (red and yellow bands). With superconducting proximity , a gap is opened and the red and blue energy bands are projected forming the black bands. Adapted from [4]	11
1.7	(a) Experimental setup of InSb wire exhibiting a quantized Majorana signature. (b) Up: Observed magnetic field dependence of the conductivity. Down: Zero-bias conductivity. The ZBCP rises up to the expected value $\frac{2e^2}{h}$ forming a quantized plateau at magnetic fields around $0,5T$ Adapted from [5]	12
1.8	a) Braiding proposal b) Basic architecture with four Majorana Zero Modes in a scalable quantum computer.Adapted from (a) [6] (b) [7]	13
2.1	a) Vertical quantum dot. b)Atomic force microscopy picture of two coupled lateral QDs (bright central circles). Gates 1 and 2 act as drain and source voltage. A negative voltage is applied at gates A,B to allow the formation of the droplets inside the free space in the 2D electron gas. Adapted from [8]	16

**LIST OF FIGURES**

---

2.2	Representation of the Density of States of a QD. The gate potential $V_G$ can be tuned to change the Fermi energy of the dot. . . . .	17
2.3	(a) Representation of transport through QD. The red curve represents the hybridized energy level. The gate voltage tunes this level. In the case represented, the energy level is in the middle of the drain and source voltages allowing transport between the leads. (b) Charging diagram of a quantum dot. Differential conductance dependence over the gate voltage $V_G$ and the source-drain voltage ( $V_{SD} = V_S - V_D$ ). Coulomb blockade occurs at the diamond-shaped regions with zero conductance. At these regions the number of electrons is constant and increases 1 by 1 when the gate voltage is scaled up. (b) Adapted from [9] . . . . .	18
2.4	(a) Resistivity minimum near the Kondo temperature $T_k$ . (b) Kondo cloud formed by a singlet grouped around the impurity divided in spin- $\uparrow$ ,spin- $\downarrow$ regions. Adapted from a) [10] , b) [11] . . . . .	21
2.5	Observation of the Kondo effect in a single electron transistor. Color scale temperatures from 15mk (Black) to 800mk(Red) a) Dependence of the zero bias conductance over the gate voltage. A plateau in the conductance peak appears in the odd particle regimes. b) Dependence of the conductance over the gate source drain voltage inside an odd electron regime. A zero bias conductance peak (ZBCP) of height $\frac{2e^2}{h}$ is observed. This is the Kondo signature. Adapted from [12] . . . . .	23
3.1	Graph representation of Gauss-Jordan elimination a) Graph $\mathcal{G}_{d_1d_2}$ b) After the elimination of vertex $c_k$ , the energies of dots $d_1$ and $d_2$ , and the coupling parameter are changed. c) After Gaussian elimination of dot 2 the energy of the remaining dot $\varepsilon_D QD^+$ represents the transport information through $d_1$ of the entire DQD. . . . .	28
3.2	Evolution of the density of states at each QD (Left: dot 1, Right: dot 2) at three distinct arrangements of DQD-lead coupling. The inset at the first column depicts the type of coupling. The purple line represents the tuning variable. The energy unit is $\Gamma_1$ . $e_1 = e_2 = 0$ in all arrangements. (a),(b) The lead is connected to both QDs. Tuning variable: $\Gamma_2$ . (c)(d) Indirect coupling of the second dot through dot one. Tuning variable: $t_{dots}$ . (e)(f) Triangular coupling. Tuning variable: $t_{dots}$ . . . . .	32
3.3	NRG algorithm: The logarithmic discretization of the conduction band according to the discretization parameter $\Lambda$ . The system with the lead attached to the QD-impurity is mapped to the Wilson's chain, such that each step of the chain contains the contributions at different energy scales $\Lambda^{-N/2}$ . . . . .	33
3.4	Schematic representation of the spectrum in the iterative diagonalization process. a)→b): Rescaling. b)→c): Including next site and diagonalizing. c)→d): Truncation of high energy states . Adapted from [13] . . . . .	35

*LIST OF FIGURES*

---

3.5 Evolution of the QD-spectrum vs number of iterations of the code for $U' = 0.5$ , $\epsilon' = -0.25$ , $\Gamma = 2.82 \times 10^{-2}$ . . . . .	39
3.6 Wilson's chain depicted as an open quantum system. Adapted from [14]. . . . .	41
3.7 Diagram of the NRG code . . . . .	42
3.8 Density of states for the DQD attached to the lead at different configurations. Left insets: Setup that is being used. Right insets: Low energy DOS. . . . .	44
4.1 Model for the QD-Majorana system. Solid lines: Hopping interactions: $V_1$ couplings of QD1. Dashed lines: Majorana spin- $\downarrow$ effective couplings (Equation 4.3) $t_1$ . The atomic energy levels appear inside each QD $\epsilon_1$ are tuned by the gate voltages. The coulomb interaction is represented by $U_1$ separates two energy levels. The red dashed horizontal lines represent the Fermi level. . . . .	47
4.2 Graph $\mathcal{G}_M$ representing the transport equations. . . . .	49
4.3 Density of states for a Majorana coupled to a QD under the tuning of different parameter. The tuning parameter is drawn in purple line in the inset model. . . . .	50
4.4 DOS at $t_1 = \Gamma_1$ at the PHS-point. Insets: Left: QD-Majorana model. Right: Zoom to low energy DOS. . . . .	51
4.5 (a)&(b): Dependence of the DOS over the gate voltage $\Delta\epsilon_1$ at $t_1 = \Gamma_1$ . (a)Spin- $\uparrow$ (b) Spin- $\downarrow$ . (c) DOS at the red-dashed horizontal cut in (a)&(b). Insets: Left: QD-Majorana model. Right: Low energy DOS. (d) DOS at the Green-dashed vertical cut in (a)&(b). . . . .	52
4.6 Dependence of the zero-bias conductance over tuning voltage Adapted from [3]. . . . .	52
5.1 Model for the DQD-Majorana system. Solid lines: Hopping interactions ( $t_{dots}$ : inter-dot coupling , $V_1, V_2$ couplings of QD1 and QD2 with the lead. ). Dashed lines: Majorana spin- $\downarrow$ effective couplings (Equation 4.3) $t_1, t_2$ . The atomic energy levels appear inside each QD $\epsilon_1, \epsilon_2$ are tuned by the gate voltages. The coulomb interaction is represented by $U_1, U_2$ . The red dashed horizontal lines represent the Fermi level. . . . .	54
5.2 Graph method applied to a DQD coupled to a Majorana zero mode. a) Initial stage. b) Eliminated vertexes $c_k^\dagger, c_k, d_{2,\downarrow}, d_{2,\downarrow}^\dagger$ in that order. c) Eliminated vertexes $d_{1,\downarrow}^\dagger$ and $f_\downarrow$ , the final energy is $\omega - G_{d_1, d_1^\dagger}(\omega)$ . . . . .	56
5.3 . . . . .	58
5.4 Non-interacting DOS in the symmetric coupling(Figure 5.3(a)) at each QD. First column: Dot 1. Second column: Dot 2. The gate voltages vary at each row. First row: Zero-bias in both dots $\epsilon_1 = \epsilon_2 = 0$ . Second row: $\epsilon_1 = 5\Gamma_1$ , $\epsilon_2 = 0$ . Third row: $\epsilon_1 = 0$ , $\epsilon_2 = -5\Gamma_1$ . Bold blue lines: Spin- $\uparrow$ DOS. Thin red lines: Spin- $\downarrow$ DOS. The insets at the right show which dot carries a Majorana signature, represented by a red dashed circle. Upper: First dot. Lower: Second dot. . . . .	59

## LIST OF FIGURES

---

5.5 Non-interacting DOS of the T-dot coupling Figure 5.3(b). (b). First line (a),(b): $\varepsilon_1 = \varepsilon_2 = 0$ . Second line (c),(d): $\varepsilon_1 = 5\Gamma_1$ , $\varepsilon_2 = 0$ . Third line (e),(f): $\varepsilon_2 = -5\Gamma_1$ , $\varepsilon_1 = 0$ . Blue bold lines: Spin- $\uparrow$ DOS. Red thin lines: Spin- $\downarrow$ DOS. The inset at the upper-right corner of each line indicates which dots exhibit Majorana signature, which is represented by a red dashed circle inside the dot. . . . .	60
5.6 Non-interacting DOS of the set up in Figure 5.3(c). First line (a),(b): $\varepsilon_1 = \varepsilon_2 = 0$ . Second line (c),(d): $\varepsilon_1 = 5\Gamma_1$ , $\varepsilon_2 = 0$ . Third line (e),(f): $\varepsilon_2 = -5\Gamma_1$ , $\varepsilon_1 = 0$ . Blue bold lines: Spin- $\uparrow$ DOS. Red thin lines: Spin- $\downarrow$ DOS. The inset at the upper-right corner of each line indicates which dots exhibit Majorana signature, which is represented by a red dashed circle inside the dot. . . . .	61
5.7 Density of states of both dots in the symmetric coupling without gate voltages between the Majorana and the interacting DQD. Bold blue lines: Spin- $\uparrow$ DOS. Thin red lines: Spin- $\downarrow$ DOS. Inset: Low-energy DOS. . . . .	62
5.8 (a)-(d) Dependence of the density of states of setup in Figure 5.3(a) over $\omega$ and the gate voltage $\Delta\varepsilon_2$ . $\Delta\varepsilon_1 = 0$ Up: Dot 1. Down: Dot 2. Left: Spin- $\uparrow$ . Right: Spin- $\downarrow$ . (e) Evolution of the relation $\frac{\rho_{\uparrow}(0)}{\rho_{\downarrow}(0)}$ for both QDs. While QD2 losses rapidly the Majorana signature, QD1 maintains it till $\Delta\varepsilon_2 \sim 5$ . . . . .	63
5.9 The same as in Figure 5.4 for the interacting DOS in the symmetric coupling (Figure 5.3. . . . .	64
5.10 The same as in Figure 5.4 for the interacting DOS of the T-dot junction Figure 5.3(b). Inset in b): Zoom to low-energy DOS. . . . .	65
5.11 Logarithmic dependence of the density of states for interacting dots coupled in series Figure 5.3(c) over $\omega$ and the gate voltage $\Delta\varepsilon_1$ . $\Delta\varepsilon_2 = 0$ . The dependence over $\Delta\varepsilon_2$ produces similar results. . . . .	65
5.12 The same as in Figure 5.4 for the interacting DOS for interacting dots coupled in series (Figure 5.3(c)). Inset in b): Zoom to low-energy DOS. . . . .	66
5.13 Dependence of the DOS in the symmetric model Figure 5.3(a) over $t_1 = t_2$ and $\omega$ . Up: High energy . Down: Zoom to low energy states. . . . .	67
5.14 Dependence of the DOS in the symmetric model Figure 5.3(a) over $t_1 = t_2$ and $\omega$ . Up: High energy . Down: Zoom to low energy states. . . . .	68
5.15 DOS at both dots for the model in the left inset. The right inset zooms the low-energy DOS in the second dot. . . . .	69
5.16 Logarithmic dependence of the DOS in the second dot for setup in Figure 5.3(b) over the second gate voltage. Cut at $5\Gamma_1$ corresponds to Figure 5.10(f). . . . .	69
6.1 Table of Majorana signatures in the studied cases in interacting dots. . . . .	71

# List of Abbreviations

MZM	Majorana zero mode
MBS	Majorana bound state
QD	Quantum dot
DQD	Double quantum dot
DOS	Density of states
ZBCP	Zero-bias conductance peak
NRG	Numerical renormalization group
EOM	Equations of motion
TS	Topological Superconductor

# Table of Contents

<b>List of Figures</b> . . . . .	iv
<b>List of Abbreviations</b> . . . . .	viii
<b>1 Introduction and Motivation:</b>	
<b>The Pursuit of Majorana Fermions in superconducting wires.</b> . . . . .	1
1.1 The Kitaev Chain . . . . .	3
1.1.1 Topological phase transition . . . . .	6
1.1.2 Non-abelian statistics . . . . .	8
1.1.3 Real implementations of the Kitaev Chain . . . . .	10
1.2 Detecting Majorana zero modes using quantum dots . . . . .	12
1.2.1 Why double quantum dots? . . . . .	14
1.3 Overview . . . . .	14
<b>2 Preliminaries</b> . . . . .	16
2.1 Transport process in quantum dots (QDs) . . . . .	16
2.2 The Anderson Model . . . . .	18
2.3 The Kondo Effect . . . . .	20
2.3.1 Kondo Effect in QDs . . . . .	22
<b>3 Theory and Methods</b> . . . . .	24
3.1 Double Green function formalism and the equations of motion . . . . .	24
3.1.1 Green function of fermion operators and the equations of motion . . . . .	25
3.1.2 Presenting the Graph-Gauss-Jordan elimination process to solve the EOM . . . . .	27
3.1.3 Graph Algorithm . . . . .	30
3.1.4 Non-interacting DOS in a double quantum dot coupled to a lead . . . . .	30
3.2 The Numerical Renormalization Group (NRG) . . . . .	32
3.2.1 From the Renormalization Group to Wilson's Chain . . . . .	33
3.2.2 Iterative Diagonalization . . . . .	36
3.2.3 Symmetries . . . . .	36
3.2.4 Iterative diagonalization in a single QD attached to a metallic lead . . . . .	37
3.2.5 The Density Matrix Renormalization Group (DM-NRG) . . . . .	39

---

*Table of Contents*

---

3.2.6	Specifications of the NRG Code . . . . .	42
3.2.7	NRG results in a double quantum dot coupled to a metallic lead . . . . .	43
<b>4</b>	<b>Coupling a QD with a Majorana zero mode</b> . . . . .	46
4.1	Model . . . . .	46
4.2	Non-interacting QD coupled to Majorana chain . . . . .	48
4.3	Kondo-Majorana physics . . . . .	51
<b>5</b>	<b>Coupling the Majorana Zero Mode to a Double Quantum Dot</b> . . . . .	54
5.1	Applying our methods to the DQD-Majorana system . . . . .	55
5.1.1	Non-interacting Green function: . . . . .	55
5.1.2	NRG for the interacting system . . . . .	57
5.2	Manipulation of Majorana zero modes . . . . .	57
5.2.1	Non-interacting manipulation . . . . .	59
5.2.2	Interacting manipulation . . . . .	62
5.3	Additional results . . . . .	67
5.3.1	Indirect exchange through the Majorana mode . . . . .	67
5.3.2	Indirect Majorana coupling through the lead . . . . .	68
5.3.3	Critical behavior in zero-bias DOS . . . . .	69
<b>6</b>	<b>Conclusions</b> . . . . .	70
	<b>References</b> . . . . .	71
	<b>Bibliography</b> . . . . .	72

## Appendices

<b>A</b>	<b>Appendix</b> . . . . .	80
A.1	From the logarithmic discretization to the Wilson's chain. . . . .	80
A.2	Initial DQD-Majorana Hamiltonian. . . . .	84

# Chapter 1

## Introduction and Motivation: The Pursuit of Majorana Fermions in superconducting wires.

*"It started out with a toy model demonstration, and then I realized it was very good model. You don't understand the full implications until other people start thinking it is true and they observe the big picture [...] Now, that toy model is like Hydrogen atom for topological materials- it turned out to be the first example of topological quantum matter."*

– F. Duncan M. Haldane

In 2001, Alexei Yu Kitaev presented a toy model that could lead to the implementation of topological qubits [15], an innovative idea that promised to sort out the problem of high decoherence in quantum computation. Kitaev used the properties of an exotic quasi-particle bound at the edges of a superconducting wire. This "bound state" was cataloged as a "Majorana fermion", the iconic particle predicted by Ettore Majorana in the 1930s. Majoranas were initially proposed as a real field solution of the Dirac equation, which describes fermions that are their own antiparticle[16]. So far, no fundamental particle with these characteristics has been observed. However, novel exciting proposals are oriented to search these particles at the boundaries of topological superconducting wires.

These "topological superconductors" belong to an emergent group of materials that experience phase transitions without passing through a symmetry breaking, meaning that they cannot



Figure 1.1: Up: Failed topological evolution from a cup of coffee to a sphere. Down: Smooth evolution of a coffee cup into a donut. Taken from <https://bit.ly/2ETicqM>

be characterized by Landau theory. Instead, these phases of matter are described by a new type of order determined by topology. In mathematics, the word topology is used to describe non-local features of surfaces (or manifolds) that are preserved under smooth deformations. The cliché, but always educative, joke to explain this concept says that "Topologists cannot tell the difference between a donut and a coffee cup, since one of them can always be continuously deformed into the other through a sequence of smooth and small alterations" (Figure 1.1).<sup>1</sup> However it wouldn't be possible to topologically transform a soccer ball into a donut since no there is no way of putting a hole into the ball by performing smooth deformations. We then say that coffee and donuts are topologically equivalent to each other, but not to the ball.

The insight of topology into the field of condensed matter physics is the following. Those materials who are attributed a topological characterization are endowed with a characteristic stability under smooth deformations (or adiabatic evolutions). Moreover, they usually present exotic excitations at its boundaries . The most famous example of this behavior is the integer quantum hall effect (IQHE), which exhibits an insulating bulk with electron currents passing through the edges of the material. More importantly, the magnetic field defines conductivity plateaus representing different topological phases. This effect is so robust that it allowed to define with high precision a resistivity standard unit  $R_K = \frac{h}{e^2} = 25812.807557(18)\Omega$ , which had major impact in science and technology.

Kitaev's toy model describes a one dimensional topological chain. Similar to the IQHE, this chain exhibits bizarre edge states, which are documented as bounded Majorana quasi-particles. Besides the topological protection, these Majoranas have another important characteristic. Displayed in multi-state system Majoranas can show anionic non-abelian statistics [17]. This means that if we have three Majorana fermions  $\gamma_1, \gamma_2$  and  $\gamma_3$  and we, somehow, get to exchange the first two Majoranas and then the second and the third, it produces a different state than exchanging the last two Majoranas first and then the first two. This is a very interesting property for quantum computing since it allows to encode information in many body systems. Kitaev pointed out that the combination of these two properties, topological stability and non-abelian statistics, could lead Majoranas to be the base of promising implementation of fault-tolerant quantum computers [? ]. This marked the beginning of a new area called topological quantum computing [18], which combines areas as different as topology, knot theory [19] and solid state physics.

These ideas motivated the pursuit of Majorana fermions in condensed matter systems [4, 20–22]. Indeed, the last decade has been full of excitement, as new experiments have turned some of the theoretical predictions of the 1990s and 2000s into reality. Recently, the first evidence of Majorana bound states in topological superconductors (TS) has been found in multiple experiments [5, 23–26] following the prescription by Oreg et al. [27] and Lutchyn et al. [28]. These experiments have been based on tunneling spectroscopy in junctions between TS and non metallic (NM) leads, where resonances have been observed at zero energy, which is consistent with the presence of Majorana zero-energy modes. All of these facts, have contributed to the

---

<sup>1</sup>For decades, this has been the main reason for the absence of donuts at topology workshops.

“boom” of Majorana wires, which has become one of the hottest research topics in Physics during the last decade.

In this section, we will present a review of the main ideas behind the Kitaev chain (section 1.1) taking special attention on its topological characterization and non-abelian statistics. Finally, we will tell how that model inspired real implementations of Majorana wires in subsection 1.1.3. This discussion will eventually bring us closer to the main motivation of this thesis in section 1.2, the detection Majorana zero modes using quantum dots.

## 1.1 The Kitaev Chain

Kitaev’s tight binding toy model represents an spinless superconducting chain with the following Hamiltonian

$$H = \sum_{i=1}^N \left[ -t(a_i^\dagger a_{i+1} + a_{i+1}^\dagger a_i) - \mu a_i^\dagger a_i + \Delta a_i a_{i+1} + \Delta^* a_{i+1}^\dagger a_i^\dagger \right]. \quad (1.1)$$

Where  $\mu$  is the chemical potential, so that  $\mu a_i^\dagger a_i$  is the energy associated to each step in the chain, and  $t(a_i^\dagger a_{i+1} + a_{i+1}^\dagger a_i)$  represents the interaction between neighboring sites which is determined by the hopping term  $t$ . The remaining terms describe the superconducting properties of the system as it is established by the BCS theory of superconductivity. Here,  $\Delta$  is a complex superconducting parameter with the form  $\Delta = e^{i\theta} |\Delta|$ . The associated terms represent the Cooper pairs which can be created or annihilated at neighboring sites of the system hence breaking particle number. However, the system still preserves fermion parity, a property that will be very important during the rest of the project.

There are different ways to diagonalize this Hamiltonians. The usual method is to perform a Bogoliubov-de Gennes transformation which makes the problem equivalent to diagonalize a  $2N \times 2N$ -matrix. However, Kitaev’s brilliant solution passes through another transform. Lets define the operators  $\gamma_{A,j}$  and  $\gamma_{B,j}$  such that

$$\gamma_{A,j} = e^{i\theta/2} a_j + e^{-i\theta/2} a_j^\dagger, \quad \gamma_{B,j} = -i(e^{i\theta/2} a_j - e^{-i\theta/2} a_j^\dagger). \quad (1.2)$$

It is simple to check that these operators are self-adjoint ( $\gamma_{A,j}^\dagger = \gamma_{A,j}$ ,  $\gamma_{B,j}^\dagger = \gamma_{B,j}$ ) , which implies that they are their own antiparticle. Moreover, they satisfy the fermion anti-commutation relations

$$\begin{aligned} \{\gamma_{A,i}, \gamma_{A,j}\} &= \{\gamma_{B,i}, \gamma_{B,j}\} = 2\delta_{ij}, \\ \{\gamma_{A,i}, \gamma_{B,j}\} &= 0. \end{aligned} \quad (1.3)$$

These relations are characteristic from a Clifford algebra, which basically describes a fermion which is its own antiparticle. A Majorana fermion. Thus, we can think  $\gamma_{A,i}, \gamma_{B,i}$  as Majorana operators.

### 1.1. The Kitaev Chain

---




Figure 1.2: Illustration of the Kitaev chain for open boundary conditions in the Majorana representation. Dirac fermions are depicted by the big globes. Each one of them contains two Majoranas (small blue and red circles) a)Represents the trivial case where the hopping and the superconducting term approaches to 0. b) The non-trivial topological phase. The coupling is produced between Majoranas in different Dirac fermions By the author

Taking the inverse of (1.2) we obtain that each (Dirac) fermion in Hamiltonian (1.1) is composed by two Majorana fermions such that

$$a_j = \frac{e^{-i\theta/2}}{2}(\gamma_{A,j} + i\gamma_{B,j}), \quad a_j^\dagger = \frac{e^{i\theta/2}}{2}(\gamma_{A,j} - i\gamma_{B,j}). \quad (1.4)$$

Hence, we could adventure to say that these Majorana operators are actually splitting the Dirac fermions into real( $\gamma_A$ ) and imaginary ( $\gamma_B$ ) parts ,the same way as complex numbers are a composite of two real numbers. This is just a possible interpretation to elucidate that each Dirac fermion is composed by two types of Majorana quasi-particles just as in Figure 1.2.

The new Kitaev Hamiltonian in the "Majorana representation" looks like

$$H = \frac{i}{2} \sum_{j=1}^N [-\mu \gamma_{A,j} \gamma_{B,j} + (t + |\Delta|) \gamma_{B,j} \gamma_{A,j+1} + (t - |\Delta|) \gamma_{A,j} \gamma_{B,j+1}] + Const, \quad (1.5)$$

Depending on the values of parameters  $\mu, t$  and  $|\Delta|$  we can identify two regimes represented by the following situations:

1. If  $|\Delta| = t = 0$  and  $\mu < 0$ , Hamiltonian (1.5) becomes  $\frac{-i\mu}{2} \sum_j \gamma_{A,j} \gamma_{B,j}$  which represents the coupling of the Majoranas in the same Dirac fermion. (See Figure 1.2 (a))
2. If  $|\Delta| = t > 0$  and  $\mu = 0$ , the situation is much more interesting. The Hamiltonian (1.5) takes the form  $H = 2ti \sum_j \gamma_{B,j} \gamma_{A,j+1}$ . This implies that the coupling is performed between Majoranas of different Dirac fermions. Notably, this configuration leaves the edge Majorana operators ( $\gamma_{A,1}$  and  $\gamma_{B,N}$ ) uncoupled to the system (See Figure 1.2b)). Note that these uncoupled Majorana fermions can be at any state without any repercussion in the energy of the system. This explains the emergence of a ground state localized at the edges of the chain.




Figure 1.3: Spectrum of Hamiltonian 1.5 with 30 sites and  $t = |\Delta|$  s. Method: Numerical diagonalization. By the author

These two situations are representatives of two different topological phases. The trivial phase occurs for  $\frac{\mu}{2t} > 1$  and the non-trivial phase appears when  $\frac{\mu}{2t} < 1$  (See Figure 1.3). The main characteristic of the non-trivial phase is the creation of a robust zero-mode generated by the uncoupled Majorana fermions at the edges of the Kitaev chain. Note that if

$$H = 2ti \sum_j \gamma_{B,j} \gamma_{A,j+1}, \quad (1.6)$$

it is possible to define new Dirac fermion operators as

$$c_j = \frac{1}{\sqrt{2}} (\gamma_{B,j} + i\gamma_{A,j+1}), \quad c_j^\dagger = \frac{1}{\sqrt{2}} (\gamma_{B,j} - i\gamma_{A,j+1}).$$

Then (1.6) becomes

$$H = it \sum_{j=1}^{N-1} (2c_j^\dagger c_j - 1). \quad (1.7)$$

Therefore, a ground state  $|\Omega\rangle$  of this Hamiltonian is an state vacuum at all sites  $j$  from 1 to  $N-1$  ( $c_j|\Omega\rangle = 0$ ). This condition allows some degeneracy since the sites at the boundary are not coupled to the Hamiltonian  $\gamma_{A,1}$  and  $\gamma_{B,N}$ . The Dirac operators formed by these Majoranas

$$c_N = \frac{1}{\sqrt{2}} (\gamma_{B,N} + i\gamma_{A,1}), \quad c_N^\dagger = \frac{1}{\sqrt{2}} (\gamma_{B,N} - i\gamma_{A,1}),$$

### 1.1. The Kitaev Chain

---

can be either occupied ( $c_N^\dagger c_N |\Omega\rangle = 1$ ) or empty ( $c_N^\dagger c_N |\Omega\rangle = 0$ ). Each of these results will have a different fermion parity, that is a symmetry of our Hamiltonian. Indeed, we can define a global parity operator as

$$\mathcal{P} = \prod_{i=1}^N (2c_j^\dagger c_j - 1) = \prod_{i=1}^N -i\gamma_{B,j}\gamma_{A,j+1} = \pm 1. \quad (1.8)$$

In the ground state  $|\Omega\rangle$ , this parity will be defined by the result of  $\gamma_{B,N}\gamma_{A,1}$  because the other states are fixed by ( $c_j|\Omega\rangle = 0$ ). This is an important point, since symmetry protection is actually correlating the two opposite sites of the Kitaev chain i.e. Any attempt to disturb one site of the chain would perturb the other site to guarantee the preservation of fermion parity. This is a great deal, it shows how coherent are these Majorana edge states. A possible explanation for this is the topology of the Kitaev chain, which is the objective of the next subsection.

#### 1.1.1 Topological phase transition

Both regimes described previously can be characterized with a topological parameter. A didactic method to explain is the one used by Alicea[4]. First of all, suppose that we have an infinite Kitaev chain chain ( $N = \infty$ ) in (1.5). This system is translation invariant, hence we can make a Fourier transformation to the momentum space as

$$H = \sum_{k \in BZ} (b'_k \ c'_k) H_k \begin{pmatrix} b'_{-k} \\ c'_{-k} \end{pmatrix}, \quad (1.9)$$

with the Bloch Hamiltonian equal to

$$H_k = \begin{pmatrix} 0 & \frac{-i\mu}{2} + it \cos k + |\Delta| \sin k \\ \frac{i\mu}{2} - it \cos k + |\Delta| \sin k & 0 \end{pmatrix} = (|\Delta| \sin k) \sigma_x + \left(\frac{\mu}{2} - t \cos k\right) \sigma_y. \quad (1.10)$$

Here,  $\sigma_x$  and  $\sigma_y$  are the corresponding Pauli matrices. The Brilloin zone ( $BZ$ ) is the periodic space  $[-\pi, \pi]$  which can be mapped to the unitary circle. Equation (1.10) determines the coordinates of the Bloch Hamiltonian in the base  $\{\sigma_x, \sigma_y\}$ .

We can map these coordinates to the unitary circle by taking the norm of this vector giving

$$\hat{H}_k = \frac{1}{\sqrt{|\Delta|^2 \sin^2 k + (\frac{\mu}{2} - t \cos k)^2}} \begin{pmatrix} |\Delta| \sin k \\ \frac{\mu}{2} - t \cos k \end{pmatrix}. \quad (1.11)$$

Note that  $|\Delta|^2 \sin^2 k + (\frac{\mu}{2} - t \cos k)^2 \neq 0$  for all the values of  $k$  as long as  $\frac{\mu}{2t} \neq 1$ . When  $\frac{\mu}{2t} = 1$ ,  $H_{k=0} = 0$ , so it cannot be normalized. This is the same point were the phase transition occurs!. At any other value of  $\frac{\mu}{2t}$  it is possible to normalize  $H_k$  for all values of  $k \in BZ$ . The result of mapping  $\hat{H}_k$  for all  $k$  is a path onto the unitary circle.




Figure 1.4: Path of  $\hat{H}_k$  for the interval  $[-\pi, \pi]$ . a) Trivial phase: The resulting path can be homotopically deformed to a point. b) Non-trivial phase: The path is a non-contractible loop around the unitary circle. By the author

This path can take two forms as we can observe in Figure 1.4. If  $\frac{\mu}{2t} > 1$  the path is an arc in the upper part of the circle. In the non-trivial phase  $\frac{\mu}{2t} < 1$  the path completes a wind around the circle. This is a topological difference between both phases. While the path described by the trivial phase can be contracted to a single dot, the path described by the non-trivial circle cannot be contracted.

To determine whether the path of a given phase is type a) or type b) we only need to check if  $\hat{H}_{k=0}$  and  $\hat{H}_{k=\pi}$  are the same point  $(0, 1)$  or the opposite points  $(0, 1)$  and  $(0, -1)$ . This can be expressed into a simple equation

$$\hat{H}_{k=0,y}\hat{H}_{k=\pi,y} = \begin{cases} 1 & \text{trivial phase} \\ -1 & \text{non-trivial phase} \end{cases}, \quad (1.12)$$

where  $\hat{H}_{k=0,y}$  is the  $y$ -th component of  $\hat{H}_k$ . The term  $\hat{H}_{k,y}$  is a particular case of the Pfaffian  $\mathcal{P}(k)$ , which widely used to characterize topological phase transitions involving Majorana fermions.

In a more general perspective, the main idea behind this topological characterization relies in the adiabatic theorem. In simple words, the adiabatic theorem says that a slow evolution of a gaped Hamiltonian will produce a smooth evolution of its eigenstates, so that the order of energies remains unchanged.

A keyword in the previous definition is “gaped”. As we can observe in Figure 1.3 the phase transition occurs at  $\frac{\mu}{2t} = 1$ . This point is where the gap of the Hamiltonian closes. In periodic boundary conditions no Majorana zero modes will emerge since there are no edges in the system. Therefore, the edge states with zero energy for  $\frac{\mu}{2t} < 1$  will not appear. We obtain that the gapless point  $\frac{\mu}{2t} = 1$  divides two gapped regimes. According to the adiabatic theorem,

these two regions must be separated, hence meaning that no adiabatic evolution could lead from one region to the other since that would involve crossing through a “gapless” Hamiltonian.

In conclusion, these gapless Hamiltonians are forbidden in an adiabatic evolution. Hence, the banned points can be thought as “holes” in the space of Hamiltonians, which generates spaces with non-trivial topologies. Since adiabatic evolutions can be understood as smooth deformations of the Hamiltonian, the relation with topology becomes clear. Thus, characterizing the phase transitions in the Kitaev chain, as in similar robust materials, is mainly a topological problem, whose solution involves computing topological quantities such as Pfaffians, Chern numbers or Winding numbers, which are always integer values.

All of these ideas lead us to an interesting question. If we have two connected topological materials, one characterized by the number 0 and the other by the number 1, then what should happen at the boundary between these two discrete parameters?. Indeed some excitations are visible at the boundaries of topological materials. In the IQHE, these excitations are quantized currents passing through the edges of the placket. In the Kitaev chain these excitations are the Majorana quasi-particles!.

Finally, note that in a system that preserves certain symmetries, the space of Hamiltonians has more forbidden points. Therefore, these systems have different topological characterizations according to the protected symmetries [29]. This is the case of the Kitaev chain where the topological phase protects the parity of the symmetry under perturbations involving the two opposed Majoranas at the edges. This endowed topological stability combined with Majorana’s non-abelian statistics (next subsection) makes the Kitaev chain a promising platform for quantum computation.

### 1.1.2 Non-abelian statistics

Imagine that we want to exchange two Majorana fermions  $\gamma_1$  and  $\gamma_2$ <sup>2</sup>. This procedure can be performed with an adiabatic evolution of the Hamiltonian  $H(t)$  that exchanges both operators while leaving the system invariant. Therefore, after a period  $T$  we require that

$$\begin{aligned}\gamma_1(T) &\rightarrow \gamma_2(0) \\ \gamma_2(T) &\rightarrow \gamma_1(0)\end{aligned}\tag{1.13}$$

while  $H(0) = H(T)$ .

The adiabatic evolution is then represented by a unitary operator  $U(t) = e^{-\frac{i}{\hbar} \int H(t)}$  and is applied according to Heisenberg’s picture as

$$\gamma_i(T) = U^\dagger(t) \gamma_i U(t).$$

Since Majoranas preserve fermion parity,  $H$  must commute with the parity operator  $P = -i\gamma_1\gamma_2$ . In a Clifford algebra generated by the operators  $\gamma_1$  and  $\gamma_2$  (See algebraic relations (1.3)),

$$[H, \gamma_1\gamma_2] = 0$$

---

<sup>2</sup>This section is inspired on the webpage topocondmat [https://topocondmat.org/w2\\_majorana/braiding.html](https://topocondmat.org/w2_majorana/braiding.html), which contains an amazing tutorial about Majorana fermions and topological condensed matter.




Figure 1.5: Representation of non-abelian braiding .

implies that  $H(t) \propto \gamma_1 \gamma_2$  or  $H(t)$  is a constant. Taking the non-trivial answer we obtain that the evolution operator has the form  $U(t) = e^{\alpha(t)\gamma_1 \gamma_2}$ , where  $\alpha(t)$  is a complex function over  $t$ . We can simplify this exponential noting that  $(\gamma_1 \gamma_2)^2 = -1$  which after Taylor expansion reduces to

$$U(t) = \cos(\alpha(t)) - \gamma_1 \gamma_2 \sin(\alpha(t)). \quad (1.14)$$

Replacing this solution in (1.13) we obtain

$$\begin{aligned} \gamma_1(T) &= \gamma_1 \cos(2\alpha(T)) - \gamma_2 \sin(2\alpha(T)) \rightarrow \gamma_2 \\ \gamma_2(T) &= \gamma_2 \cos(2\alpha(T)) + \gamma_1 \sin(2\alpha(T)) \rightarrow \gamma_1, \end{aligned} \quad (1.15)$$

which can only happen if  $\alpha(T) = \pm \frac{\pi}{4}$ . Hence, we conclude that the exchange operator between both Majoranas is

$$U_{12} = e^{\pm \frac{\pi}{4} \gamma_1 \gamma_2} = \frac{1}{\sqrt{2}} (1 \pm \gamma_1 \gamma_2). \quad (1.16)$$

Note that this exchange does not depend on the evolution, nor the period  $T$ .

Now imagine that we have three Majoranas  $\gamma_1, \gamma_2$  and  $\gamma_3$  and we want to perform the following processes. On the first one, we exchange Majoranas 1 and 2 and then the Majorana in 2 (which was initially at 1) is exchanged with Majorana 3 (Figure 1.5[Left]). On the second process, we invert the order, hence exchanging first exchange Majoranas 2 and 3 and then Majoranas 1 and 2 (Figure 1.5[Right]). These two cases are represented by the following operators respectively

$$\begin{aligned} U_{23}U_{12} &= \frac{1}{2} (1 + \gamma_2 \gamma_3) (1 + \gamma_1 \gamma_2) = \frac{1}{2} (1 + \gamma_2 \gamma_3 + \gamma_1 \gamma_2 + \gamma_3 \gamma_1) \\ U_{12}U_{23} &= \frac{1}{2} (1 + \gamma_1 \gamma_2) (1 + \gamma_2 \gamma_3) = \frac{1}{2} (1 + \gamma_1 \gamma_2 + \gamma_2 \gamma_3 + \gamma_1 \gamma_3). \end{aligned} \quad (1.17)$$

Since  $\gamma_3 \gamma_1 = -\gamma_1 \gamma_3$ , the outcome of both processes is essentially different, which means that it actually matters the order in which the Majoranas are exchanged .

The particles that satisfy this strange property receive the name of non-abelian anyons. While the word "anyon" usually integrates several types of particles including bosons and

fermions, the word non-abelian makes emphasis on those anyons with non-commutative exchange statistics.

Non-abelian statistics is what make anyons a fantastic candidate to implement quantum algorithms. The idea of exchanging anyons can be thought as a braiding code like in Figure 1.5. Since the order of braiding matters, different braiding orders can be associated to distinct algorithms. This generates another form of codifying information which has been extendedly studied in knot theory [19]. And if these anyons were topological, they will be protected from quantum decoherence [30]. To the date, the closest candidates to satisfy both properties (non-abelian statistics and topological characterization) are the Majorana fermions. Nonetheless, the basic braiding protocol that would unleash the keys to topological quantum computation [18] has not been measured yet. Many theoretical proposals have been set up in this direction, but there is still a long experimental road to it.

### 1.1.3 Real implementations of the Kitaev Chain

One of the main problems to implement real devices capable to exhibit Majorana quasi-particles at the boundaries, is that Majorana's are spin-less. Since all materials have fermion doubling, it was necessary to endow the system with a physical property that could separate the spin energy bands. To bypass this problem, Lutchyn et al. proposed using a material with strong spin-orbit Rashba interaction [31] and applying a magnetic field. This would split the energy band by spin, hence destroying fermion doubling.

This idea allowed scientists to designed the first Majorana wires. The recipe consists in growing a semi-conducting wire with high spin-orbit coupling, over an s'wave superconductor and inducing a Zeeman magnetic field (Figure 1.6(a)). Such model is described by a Hamiltonian of the form [4](65)

$$H = H_{\text{wire}} + H_{\Delta} \quad (1.18)$$

with

$$H_{\text{wire}} = \sum_{\sigma \in \{\uparrow, \downarrow\}} \int dx \psi_{\sigma}^{\dagger}(x) \left( \frac{-1}{2m} \partial_x^2 - \mu - i\alpha \sigma_y \partial_x + h \sigma_x \right) \psi_{\sigma}(x), \quad (1.19)$$

$$H_{\Delta} = \int dx \Delta \psi_{\downarrow} \psi_{\uparrow} + \Delta^* \psi_{\uparrow} \psi_{\downarrow}. \quad (1.20)$$

Where  $\psi_{\sigma}^{\dagger}(x)$  creates a particle at  $x$  with spin  $\sigma$ ,  $\mu$  is the chemical potential,  $h$  is the Zeeman splitting energy and  $\alpha > 0$  is the Rashba spin-coupling parameter, favoring spin-align. The induced superconducting proximity is depicted by  $H_{\Delta}$  where  $\Delta$  is the superconducting gap. Note that this system (1.18) has all the ingredients of the Kitaev chain.

If  $\Delta = 0$ , the Bloch Hamiltonian is given by

$$\mathcal{H}_k = \frac{k^2}{2m} - \mu + k\alpha \sigma_y + h \sigma_x. \quad (1.21)$$




Figure 1.6: (a) Setup of a Majorana wire with high spin-orbit coupling, over a superconducting plane and an induced Zeeman field. (b) Splitting of the energy bands with different spins in the non-superconducting regime (red and yellow bands). With superconducting proximity , a gap is opened and the red and blue energy bands are projected forming the black bands. Adapted from [4]

And diagonalizing this matrix we obtain that the band splits as

$$\varepsilon_{\pm}(k) = \frac{k^2}{2m} - \mu \pm \sqrt{(\alpha k)^2 + h^2} \quad (1.22)$$

with opposed spins as observed in the blue and red lines of Figure 1.6(b).

The superconducting proximity effect ( $\Delta > 0$ ) opens a gap that projects the upper and lower bands forming the black bands of Figure 1.6(b) . This separation of both energy channels allows us to think the conduction band as an spin-less . This system enters into the topological phase when

$$h > \sqrt{\Delta^2 + \mu^2}. \quad (1.23)$$

At these points, Majorana bound states emerge at the edges of the wire.

This theoretical proposal led in 2012 to the first observation of Majorana signatures in InSb nanowires <sup>3</sup>, by Mourik et al. from the Kavli Institute at Delft. This was a huge boost to the field which immediately attracted abundant experimental and theoretical work.

In just 6 years, more than 5 groups have documented the observation of Majorana signatures [5, 24–26, 32]. This signature is characterized by the emergence of a robust zero bias conductance peak ZBCP of height  $\frac{2e^2}{h}$  produced by the Majorana zero mode MZM localized at the edges of the wire. Though the first experiments didn't observed such an stable signature, the last year Zhang et al. published a paper documenting the observation of this robust peak with the expected theoretical magnitude in and InSb wire (See Figure 1.7(a)) . As can be observed in Figure 1.7(b) the ZBCP increases up to  $\frac{2e^2}{h}$  for a strong magnetic field, where the system enters

<sup>3</sup>A material with strong spin-orbit coupling and large  $g$  factor.

## 1.2. Detecting Majorana zero modes using quantum dots




Figure 1.7: (a) Experimental setup of InSb wire exhibiting a quantized Majorana signature. (b) Up: Observed magnetic field dependence of the conductivity. Down: Zero-bias conductivity. The ZBCP rises up to the expected value  $\frac{2e^2}{h}$  forming a quantized plateau at magnetic fields around  $0,5T$  Adapted from [5]

the topological phase according to equation (1.23). This zero-bias peak maintains a stable magnitude for a small range of parameters generating a small plateau, which became the first observation of a quantized Majorana signature.

Despite the successful experimental results, there is still certain skepticism about the existence of Majorana fermions, mainly because Majorana zero-modes (MZM) have been found in superposition with similar types of phenomena that produce zero-modes. Some examples of these are the Andreev reflection [33] or even the Kondo peak [34]. New experimental proposals focus on distinguishing MZMs from these effects and implementing braiding protocols [35–37]. One promising idea that could lead to important results in both research lines is coupling Majorana wires with QDs. This is the objective of the following section.

## 1.2 Detecting Majorana zero modes using quantum dots

Liu and Baranger were the first to propose in 2011 the possibility of using QDs to detect Majorana zero modes [2]. When a QD is attached to the end of a Majorana chain in the topological phase, the Majorana Zero Mode at the end of the chain leaks inside the QD [38] producing a zero-bias conductance peak of half a quanta  $\frac{e^2}{2h}$  through the dot. This method of detecting Majorana signatures presents important advantages:

- 1. It does not destroy the entire qubit information:** Other detection methods such as tunneling spectroscopy have the downside that it probes not only the end of the Topological

## 1.2. Detecting Majorana zero modes using quantum dots

Superconductor(TS), but its bulk as well. This completely destroys the qubit information. When attaching the Majorana chain to a QD, the conductivity is measured through the dot which does not disturb the bulk of the chain.

2. **Kondo-Majorana physics:** The observation of Kondo signatures in superconducting devices [39] and the similarities between Kondo and Majorana signatures [38] motivated studies of prospective Kondo-Majorana physics [34, 40]. These two effects could co-exist at temperatures of a few mili-kelvins in quantum impurity systems.

Quantum dots (QD) are artificial and adaptable quantum impurities, which makes them the best device to perform this kind of study. In particular, a QD-Majorana system was already studied by my advisor in a previous paper [3], where he proved that it is possible separate Kondo and Majorana physics by tuning dot's gate voltage or by applying a strong magnetic field.

3. **Manipulation of Majorana zero modes:** Today's precise experimental control over QDs makes us dream with the possibility of implementing scalable braiding proposals Figure 1.8(a) and quantum architectures for topological quantum computation Figure 1.8(b).

These architectures perform adiabatic evolutions similar to the ones described in subsection 1.1.2 to braid Majorana fermions. This operation strongly relies on the possibility of manipulating the Majorana zero modes inside the dots. The mean idea of MZM manipulation is to tune the gate voltage of one dot to induce the Majoranas to "move" into the other dots. In a prospective braiding protocol, as the one described in [6] (Figure 1.8(a)), this manipulation process would have to be performed several times. However, till this moment MZM manipulation hasn't been achieved experimentally.




Figure 1.8: a) Braiding proposal b) Basic architecture with four Majorana Zero Modes in a scalable quantum computer. Adapted from (a) [6] (b) [7].

Nonetheless, the future for this area is still very promising. Recent experiments have documented the observation of Majorana signatures in Majorana-QD devices [26, 41] and Andreev molecules in topological superconductors attached to double quantum dots [42]. The next steps are clearly directed to achieve Majorana manipulation.

### 1.2.1 Why double quantum dots?

The simplest case where Majorana manipulation is possible is in a double quantum dot (DQD). Tunneling Majorana modes in these basic structures have inspired theoretical studies [43–45] and experimental setups confirming the observations of Andreev molecules [42]. Even though quantum tunneling of a MZM into a double dot shows several possibilities for manipulation of MZM, there is still no complete analysis of the transitions of the Majorana signatures between the QDs in this model.

In this thesis we fill this gap by performing a study of the spectral density in a DQD attached to a MZM and a metallic lead. The simplicity of this model allows us to explore analytically different configurations of QD's from symmetric and linear couplings to T-junctions . We considered both non-interacting and interacting regimes, observing major agreement between both approaches about the location of the Majorana signature. While the non-interacting regime is suitable to obtain exact expressions for the Green function, the interacting case shows how the Majorana signature co-exists with strongly correlated phenomena such as the Kondo effect and RKKY interactions [46–48].

## 1.3 Overview

This thesis is integrated by 4-major chapters:

- In chapter 2, we will take a review to the basis of quantum transport in single electron transistors, the Anderson model and the emergence of the Kondo effect in quantum dots.
- In chapter 3 we provide a description of the methods that we will use to study the Double Quantum Dot-Majorana system. To study non-interacting systems we will use the Zubarev's ballistic transport[49] which is optimized through the Graph-Gauss-Jordan elimination algorithm define in subsection 3.1.2. For non-interacting systems we use Wilson's Numerical Renormalization Group (NRG) technique [50]. We will test our methods in a double quantum dot attached to a magnetic field. Hence, background information about double quantum dots systems will be presented in this chapter.
- In chapter 4 We will test our methods in the QD-Majorana system, confirming the results of previous papers [1, 2] .
- The chapter 5 contains our major contributions to this area. Using the methods from chapter 3 and the previous acquired experience with the double quantum dot and the

### *1.3. Overview*

---

QD-Majorana model , we study in a double quantum dot attached to a Majorana zero mode and a metallic lead. We will characterize in this simple model the transitions of the Majorana signature for distinct configurations of the dots. The final results are now part of a paper that we hope to submit for publication in the following months.

# Chapter 2

## Preliminaries

In this preliminary chapter we will give a brief description about transport processes in quantum dots (QDs). We will discuss basic aspects about single electron transistors, the application of the Anderson impurity model in QDs and the subsequent emergence of the Kondo effect, one of the main condensed matter problems of the 20<sup>th</sup>-century.

### 2.1 Transport process in quantum dots (QDs)




Figure 2.1: a) Vertical quantum dot. b) Atomic force microscopy picture of two coupled lateral QDs (bright central circles). Gates 1 and 2 act as drain and source voltage. A negative voltage is applied at gates A,B to allow the formation of the droplets inside the free space in the 2D electron gas. Adapted from [8]

Quantum mechanical effects are visible when the system size is of the order of the de Broglie wavelength [51, (1.1)]

$$\lambda_f = \frac{h}{\sqrt{3m_{\text{eff}}k_B T}}$$

where  $m_{\text{eff}}$  is the electron effective mass in the crystal. Since  $m_{\text{eff}}$  can be much smaller than the free electron mass in some semiconducting materials, size quantization effects can be observed




Figure 2.2: Representation of the Density of States of a QD. The gate potential  $V_G$  can be tuned to change the Fermi energy of the dot.

at system of sizes  $\sim 100\text{nm}$  [9]. A 0D quantum system is a device confined in the three spacial dimensions up to this length-scale. This type of device receive the name of quantum dot (QD).

Nowadays, QDs can be manufactured in different substracts, geometries and orientations [51]. They can be integrated in structures like double quantum dots (Figure 2.1(a)) or can even be grown vertical to the based 2D-electron gas (Figure 2.1(b)). The precise experimental control over these devices allows to design atom-like structures with tunable energy levels. This has important applications on laser physics and in the implementation of single electron transistors.

Ideally, the energy spectrum of a QD is a discrete set of energy levels resembling the spectrum of an atom. This atom-like structure is usually connected to three main gates (See Figure 2.3(a)). Two of them are the source and the drain with voltages  $V_S$  and  $V_D$ , used to control the electric gradient through the QD. When the QD is connected to these leads these energy levels are hybridized with respect to a broadening parameter  $\Gamma$  which increases as the square of the source-drain voltage  $V_{SD}$

$$\Gamma \propto \pi \|V_{SD}\|^2 \quad (2.1)$$

This broadening is depicted in Figure 2.2. Ideally,  $\Gamma \ll \Delta E$  is smaller enough such that the energy levels do not overlap each other.

The third gate is the gate voltage  $V_G$  which allows to tune the energy levels of the dot with high precision. With this three gates it is possible to execute transport measurements through a QD. An electron can pass from the source to the drain if there is an energy level in the middle of the two voltages, just as in Figure 2.3(a). If this condition is not satisfied, the dot enters into a Coulomb blockade region without electron transport between both leads as can be observed in Figure 2.3(b)). Inside the Coulomb blockade regions (black diamonds) the number of electrons is constant. When increasing  $V_G$  a single electron enters into the dot each time the system makes a transition between blockade regions. Since all of these effects can be controlled precisely with the gate voltage, the system described is indeed a single electron transistor (SET).

Note that when  $V_{SD} = 0$ , the condition for conductance can still be satisfied if there is an hybridized state at the level of  $V_S = V_D$ , the Fermi energy. This situation receives the name of zero-bias conductance peak (ZBCP). The effects that can cause robust ZBCPs are quite interesting. In this thesis were will study two of them. The Kondo effect and the Majorana zero




Figure 2.3: (a) Representation of transport through QD. The red curve represents the hybridized energy level. The gate voltage tunes this level. In the case represented, the energy level is in the middle of the drain and source voltages allowing transport between the leads. (b) Charging diagram of a quantum dot. Differential conductance dependence over the gate voltage  $V_G$  and the source-drain voltage ( $V_{SD} = V_S - V_D$ ). Coulomb blockade occurs at the diamond-shaped regions with zero conductance. At these regions the number of electrons is constant and increases 1 by 1 when the gate voltage is scaled up. (b) Adapted from [9]

modes.

## 2.2 The Anderson Model

The Anderson model is used to describe quantum impurity systems [52]. A quantum dot attached to a metallic lead is basically an artificial impurity that can be experimentally designed, modified and manipulated. Hence QDs are the perfect type of structure to test the Anderson mode.

Due to the small confinement space inside these dots, Coulomb repulsion is relevant. However, it is usually impossible to provide a complete analytical description of these kind of systems due to the high correlations generated by this factor. Instead, we can obtain an overall description of the transport through the impurity by neglecting this Coulomb repulsion. This will allow us to obtain some analytic intuition of the models before adventuring with long-lasting numerical simulations of interacting models. During this thesis, we will consider these two regimes as follows

- **Non-interacting systems:** Coulomb repulsion is not relevant . In this case, spin- $\uparrow$  and spin- $\downarrow$  channels are independent. They can be solved analytically through the equations of motion (section 3.1).
- **Interacting systems:** The Coulomb repulsion is relevant. The repulsion factor will be defined by the factor  $U$  which will take a fix value during the entire project. In this case,

spin- $\uparrow$  and spin- $\downarrow$  channels are not independent since the Coulomb repulsion limits the number of particles inside each dot. We will use the Numerical Renormalization Group to treat this case. The intuition acquired from non-interacting systems will help us to select the input parameters of the algorithm.

To begin the description of the Anderson model, first consider that we have a QD (impurity) coupled to the conduction band of a metallic lead. We will define a Coulomb repulsion factor  $U_i$ , which will be set to 0 if the system is non-interacting. Using the Hund rules, we know that the energy levels inside the dot should be filled from lower to higher energies with two electrons with different spin at each state. Each pair of electrons will interact magnetically and electrically. In addition, there is an energy term associated to each electron and a Zeeman splitting factor in case a  $\hat{z}$ -directed magnetic field  $B$  is placed. Considering these interactions, we can obtain a very general expression in second quantization for the QD Hamiltonian of the form [9, (3.2)]

$$H_d = \sum_{i\sigma} \varepsilon_i d_{i\sigma}^\dagger d_{i\sigma} + \sum_i U_i \hat{n}_{i\uparrow} \hat{n}_{i\downarrow} + \sum_{\sigma\sigma', i \neq j} U_{ij} \hat{n}_{i\sigma} \hat{n}_{j\sigma'} - \mu_B g B \sum_i S_i^z + J \sum_{i \neq j} \mathbf{S}_i \cdot \mathbf{S}_j, \quad (2.2)$$

where  $\sigma \in \{\uparrow, \downarrow\}$ ,  $d_{i\sigma}^\dagger$  ( $d_{i\sigma}$ ) is the dot creation(annihilation) operator,  $\hat{n}_{i\sigma} := d_{i\sigma}^\dagger d_{i\sigma}$  is the particle number,  $\mathbf{S}_i$  is the spin-vector,  $\varepsilon_i$  is the energy of the  $i^{\text{th}}$ -level in the dot,  $U_i$  is the Coulomb repulsion between electrons in the same energy level  $i$ ,  $U_{ij}$  is the Coulomb interaction between electrons in different levels (And therefore smaller than  $U_i$ ),  $B$  is an applied magnetic field in the  $\hat{z}$ -direction causing a Zeeman splitting and  $J$  is the term representing the spin coupling between distinct levels

At low temperatures, the quantum interactions occur only with the closest energy level to the Fermi energy. Hence, we can make a single-level approximation, neglecting the other energy levels. Hence, we can set the parameters  $U_i = U_{ij} = 0$  and  $J = 0$ , which significantly reduces the complexity of the dot Hamiltonian

$$H_d = \sum_{\sigma} \varepsilon d_{\sigma}^\dagger d_{\sigma} + U \hat{n}_{\uparrow} \hat{n}_{\downarrow} - \mu_B g B S^z. \quad (2.3)$$

Besides to the dot Hamiltonian, we need to consider the energy of the electrons in the lead  $H_{\text{lead}}$  and the dot-lead interaction  $H_{\text{int}}$ . We can model the conduction band of the lead as an electron gas with the following Bloch Hamiltonian

$$H_{\text{lead}} = \sum_{\mathbf{k}\sigma l} \varepsilon_{\mathbf{k}l} c_{\mathbf{k}\sigma l}^\dagger c_{\mathbf{k}\sigma l}. \quad (2.4)$$

where  $\mathbf{k}$  represents the possible crystal momentums in the leads,  $l \in \{S, D\}$ ,  $c_{\mathbf{k}\sigma l}^\dagger$  ( $c_{\mathbf{k}\sigma l}$ ) creates(annihilates) an electron with momentum  $\mathbf{k}$  and spin  $\sigma$  in the lead  $l$ ,  $\varepsilon_{\mathbf{k}l}$  is the energy of the electron in the leads.

### 2.3. The Kondo Effect

---

On the other hand, the interaction between the dot and the leads is then given by

$$H_{int} = \sum_{\mathbf{k}\sigma l} V_{\mathbf{k}l} c_{\mathbf{k}\sigma l}^\dagger d_\sigma + V_{\mathbf{k}l}^* d_\sigma^\dagger c_{\mathbf{k}\sigma l}, \quad (2.5)$$

where  $V_{\mathbf{k}l}$  is a hopping exchange term between the leads and the QD.

The sum of these three interactions receives the name of Anderson Model.

$$\begin{aligned} H &= H_d + H_{lead} + H_{int} \\ &= \sum_{\sigma} \epsilon d_{\sigma}^\dagger d_{\sigma} + U \hat{n}_{\uparrow} \hat{n}_{\downarrow} - \mu_B g B S^z + \sum_{\mathbf{k}\sigma l} \epsilon_{\mathbf{k}l} c_{\mathbf{k}\sigma l}^\dagger c_{\mathbf{k}\sigma l} + \sum_{\mathbf{k}\sigma l} V_{\mathbf{k}l} c_{\mathbf{k}\sigma l}^\dagger d_{\sigma} + V_{\mathbf{k}l}^* d_{\sigma}^\dagger c_{\mathbf{k}\sigma l}. \end{aligned} \quad (2.6)$$

In this project, we will make two extra modifications to this system. Using the anti-commutation properties of the fermion operators

$$\{d_{\sigma}^\dagger, d_{\sigma'}\} = \delta_{\sigma\sigma'}, \quad \{d_{\sigma}^\dagger, d_{\sigma'}^\dagger\} = \{d_{\sigma}, d_{\sigma'}\} = 0,$$

we get

$$\begin{aligned} (d_{\uparrow}^\dagger d_{\uparrow} + d_{\downarrow}^\dagger d_{\downarrow} - 1)^2 &= \sum_{\sigma} (d_{\sigma}^\dagger d_{\sigma})^2 - 2 \sum_{\sigma} d_{\sigma}^\dagger d_{\sigma} + 2 d_{\uparrow}^\dagger d_{\uparrow} d_{\downarrow}^\dagger d_{\downarrow} - 1 \\ &= 2 d_{\uparrow}^\dagger d_{\uparrow} d_{\downarrow}^\dagger d_{\downarrow} - \sum_{\sigma} d_{\sigma}^\dagger d_{\sigma} - 1. \end{aligned}$$

And replacing this in (2.3) we obtain a nice spin-symmetric form of the dot Hamiltonian

$$\left( \epsilon + \frac{U}{2} \right) d_{\sigma}^\dagger d_{\sigma} + \frac{U}{2} (d_{\sigma}^\dagger d_{\sigma} - 1)^2 - \mu_B g B S^z. \quad (2.7)$$

In addition, it is possible to do a linear transform to the lead operators

$$\frac{1}{\sqrt{V_S^2 + V_R^2}} \begin{bmatrix} V_S & V_R \\ -V_R & V_S \end{bmatrix} \begin{bmatrix} c_{\mathbf{k}\sigma S} \\ c_{\mathbf{k}\sigma D} \end{bmatrix} = \begin{bmatrix} c_{\mathbf{k}\sigma+} \\ c_{\mathbf{k}\sigma-} \end{bmatrix}. \quad (2.8)$$

After this transformation the operator will be decoupled from the dot Hamiltonian  $c_{\mathbf{k}\sigma-}$ . This implies that we can suppose that the dot is coupled with just one lead. In the following chapters we will maintain this convention.

## 2.3 The Kondo Effect

The Kondo effect is still regarded as one of the biggest condensed matter problems in the 20th century. There was an uncountable number of experimental and theoretical physicist that contributed to this problem. The most renown are probably the physicists Jacques Friedel, Jun Kondo and the two Nobel prizes Philip Anderson and Kenneth Wilson [53].




Figure 2.4: (a) Resistivity minimum near the Kondo temperature  $T_K$ . (b) Kondo cloud formed by a singlet grouped around the impurity divided in spin- $\uparrow$ , spin- $\downarrow$  regions. Adapted from a) [10], b) [11]

The history of the Kondo effect began in the early 1930s. By that time it was known that the resistivity of a metal is regulated by different scattering interactions against lattice phonon vibrations  $\rho_{\text{phonon}} \sim T^5$ , other electrons  $\sim T^2$  and static impurities, which is temperature independent. The form of these contribution clearly implies that the resistivity should decay uniformly with a decreasing temperature. Nevertheless, some groundbreaking experiments revealed the observation of a resistance minimum in some metals at temperatures lower than 10K [10] (See Figure 2.4 (a)).

This phenomenon intrigued the scientific community for the following decades until the year 1964 when the physicist Jun Kondo gave the first convincing solution to this puzzle. Kondo attributed the phenomenon to the scattering of the electrons due to the spin-interaction with a small concentration of magnetic impurities in the metal. To describe it he proposed the following interaction Hamiltonian

$$H_K = 2J\hat{\mathbf{S}} \cdot \hat{\mathbf{s}} \quad (2.9)$$

$$= J(2S_zs_z + S_+s_- + S_-s_+), \quad (2.10)$$

with  $S_{\pm} = S_x \pm iS_y$ . The Hamiltonian  $H_K$  which is better known as the Kondo s-d model describes the spin interaction between the spin of the impurity  $\hat{\mathbf{S}}$  and the spin of the particles in the metal  $\hat{\mathbf{s}}$ .

Kondo took  $H_K$  as a perturbation of the electron gas in the metallic lead, with  $J$  the perturbation parameter. While the first order led to no important contribution, Kondo was able to obtain on second order perturbation theory a logarithmic correction on the temperature in the resistivity of the form

### 2.3. The Kondo Effect

---

$$\rho_{imp} \propto \ln \frac{T_K}{T}, \quad (2.11)$$

where  $T_K$  received the name of Kondo temperature. Summing up this term to the other resistivity contributions we obtain the full expression

$$\rho_{metal}(T) = \rho_{imp} + a_e T^2 + b_{Phonon} T^5 + c_m \ln \frac{T_K}{T}. \quad (2.12)$$

When  $T < T_K$  the term  $\ln \frac{T_K}{T}$  increases. Eventually, it compensates the decaying resistivity which finally explained the resistance minimum.

Although Kondo's explanation was initially very successful it also presented a troublesome outcome. The logarithmic term introduced by Kondo diverges when the temperature approximates to 0, hence proving to be inefficient at temperatures well below  $T_K$ . Going to the following orders in perturbation theory also led to divergent contributions to the resistivity. This led physicists to explore non-perturbative approaches to solve the Kondo effect.

By that time, Anderson had already created his famous impurity model. One of his main contributions was the inclusion of the Hubbard term to represent the Coulomb interaction inside the dot, which proofed to be fundamental to understand the Kondo effect. Nonetheless, the Anderson model was a huge numerical challenge. The problem was finally solved by Kenneth Wilson, who was able to effectively diagonalize the Anderson model using a numerical method that combines ideas from scalability and the renormalization group.

Wilson's explanation solved the Kondo problem almost completely. It turned out that below the Kondo temperature [9], which is around

$$k_B T_K = \frac{1}{2} \sqrt{U\Gamma} e^{\epsilon(\epsilon+U)/\Gamma U}, \quad (2.13)$$

the impurity entangles with the low-energy electrons forming a strongly correlated many-body singlet. This singlet surrounds the impurity in a structure formed by alternating regions of spin- $\uparrow$  and spin- $\downarrow$  particles called the Kondo cloud (See Figure 2.4(b)). The Kondo cloud was the first evidence of an entangled many body state, and is predicted to have an astonishing correlation length between  $0.1\mu m$  to  $10\mu m$ . This is a huge number if we think that the impurity(or QD) can have a radius below  $1nm$ . Finally, the minimum in the resistivity is produced when the conduction electrons scatter with the Kondo cloud.

#### 2.3.1 Kondo Effect in QDs

The problem of magnetic impurities in metals can be treated using the Anderson model in a similar form as the transport in quantum dots. Hence, it is not a surprise that the Kondo Effect could also occur these systems. In 1998 the technological advances allowed the observation of the Kondo effect for the first time in a single electro transistor [54]. When an odd number of electrons is in the QD the last level below the Fermi energy is half-occupied, so that the dot can

### 2.3. The Kondo Effect




Figure 2.5: Observation of the Kondo effect in a single electron transistor. Color scale temperatures from 15mK (Black) to 800mK(Red) a) Dependence of the zero bias conductance over the gate voltage. A plateau in the conductance peak appears in the odd particle regimes. b) Dependence of the conductance over the gate source drain voltage inside an odd electron regime. A zero bias conductance peak (ZBCP) of height  $\frac{2e^2}{h}$  is observed. This is the Kondo signature. Adapted from [12]

be considered as a magnetic impurity. The unlocalized electrons in the reservoirs then interact with this localized electron. Spin-exchange can occur as it happened with magnetic impurities in metals. At low temperatures, this magnetic interaction gives rise to strong quantum correlations that favor the formation of a singlet state between the localized electron and the electrons in the leads. As a result, the zero-bias density of states is increased producing a zero-bias conductance peak Figure 2.5(b).

Note that the physical implications of the Kondo effect are different in magnetic impurities in metals and in QDs. In fact, they are complete opposite. While in magnetic impurities the resistivity increase, the Kondo effect in QD's leads no an unexpected raised in the zero-bias conductivity. The reason for this disagreement between both situations are the system dimensions. The scattering against a Kondo singlet in a 3D or 2D systems against magnetic impurities is an obstacle to the conducting electrons. Instead, scattering in 1D-devices enhances the conductivity of the QD because there is only one scattering direction.

As we previously discussed in section 2.1, zero-bias transport in quantum dots can only occur if there is a state at the Fermi energy. The Kondo effect creates a zero bias peak that is present whenever the dot has odd electrons. This explains the zero-bias plateaus observed in Figure 2.5(a). We may think that the new singlet at the Fermi energy is creating a "channel" that allows quantum transport between both sides of the dot. This explains the Kondo peak.

In the following chapters we will give more details about this zero-energy-mode and theory that explained it: The numerical renormalization group.

# Chapter 3

## Theory and Methods

The density of states contains important physical information about the presence of zero-modes which can be caused by the Kondo effect or by a Majorana quasi-particle. Studying this quantity will allow us to observe both effects separately inside each quantum dots. In this chapter we describe the two methods that we will use during the entire project to compute the DOS:

- The first method uses Zubarev's Green function formalism [49]. This approach requires the solution of the equations of motion, a process that we simplify introducing the graph-Gauss-Jordan elimination method (subsection 3.1.2). In the non-interacting regime, this method allows us to obtain an exact analytical expression of this quantity for density of states of the system.
- Interacting systems are more complex and cannot be solved analytically. Instead, we appeal to the renown numerical renormalization group (NRG) to deal with the strong correlations of the interacting Anderson model section 3.2. The NRG is famous for providing the most complete explanation of the Kondo effect. Now we intend to use it to study the co-existence of Kondo and Majorana physics.

Both methods will be tested on the model of a double quantum dot attached to a metallic lead. We will observe that at very low energies, the physics of interacting systems emulate characteristic features of the non-interacting model. The methods developed in this chapter will be tested again in chapter 4 for a QD coupled to a Majorana chain. Finally we will use them to make our leading contribution to this theory, the analysis of a double quantum dot attached to a Majorana wire (chapter 5).

### 3.1 Double Green function formalism and the equations of motion

The Green function  $G$  of a Hamiltonian  $H$  is the operator that satisfies the homogeneous equation

$$\left( i\hbar \frac{\partial}{\partial t} - H \right) G(t-t') = \delta(t-t'). \quad (3.1)$$

This type of differential equations are usually solved with a Fourier transform

$$G_H(\omega) = \int_{-\infty}^{\infty} d(t-t') G(t-t') e^{-i\omega(t-t')}. \quad (3.2)$$

By convention we took  $\hbar = 1$  so that we can unify units of frequency and energy. This turns equation (3.1) into a function of the energy  $\omega$  such that

$$(\omega + is - H)G_H(\omega) = I. \quad (3.3)$$

The term  $+is$  in the previous Hamiltonian is part of a mathematical trick quite common in this theory. During the whole procedure, the Green function acts on the complex field, with  $+is$  making reference to a small imaginary part that allows to avoid singularities in the real line. But when we need to obtain a physical interpretation, we will take the limit  $s \rightarrow 0$  to obtain the result for real energies.

The next step is to decompose  $G_H(\omega)$  in the eigenbase of the Hamiltonian  $\{|\alpha\rangle\}$  by

$$\langle\alpha|G_H(\omega)|\alpha'\rangle = \frac{\delta_{\alpha\alpha'}}{\omega - is - \varepsilon_\alpha} = \frac{\delta_{\alpha\alpha'}(\omega + is - \varepsilon_\alpha)}{(\omega - \varepsilon_\alpha)^2 + s^2}. \quad (3.4)$$

From the Cauchy relation

$$\lim_{s \rightarrow 0} \frac{s}{(\omega - \varepsilon_\alpha)^2 + s^2} = \pi\delta(\omega - \varepsilon_\alpha) \quad (3.5)$$

we obtain

$$Im[\langle\alpha|G_H(\omega)|\alpha'\rangle] = \pi\delta(\omega - \varepsilon_\alpha)\delta_{\alpha,\alpha'}. \quad (3.6)$$

Note that the sum of  $Im[\langle\alpha|G_H(\omega)|\alpha'\rangle]$  over all the eigenstates of  $H$  is simply  $\pi$  times the density of states

$$\rho(\omega) = -\frac{1}{\pi} \sum_{\alpha} Im[\langle\alpha|G_H(\omega)|\alpha'\rangle] = \sum_{\alpha} \delta(\omega - \varepsilon_{\alpha}). \quad (3.7)$$

In the following section we will show how to extend this idea to fermion algebras in the second quantization.

### 3.1.1 Green function of fermion operators and the equations of motion

In many body physics, fermion operators are defined by quantized fields over time satisfying anti-commutation relations, such that for any two independent operators at a time  $t$  ( $A(t), B(t)$ ) we have

$$\{A(t), B(t)\} = \delta_{B,A^\dagger}. \quad (3.8)$$

Two types of Green functions are usually defined in this case.

$$G_{A,B}^r(t, t') = -i\theta(t - t')\langle\{A(t), B(t')\}\rangle. \quad (3.9)$$

$$G_{A,B}^a(t, t') = -i\theta(t' - t)\langle\{A(t), B(t')\}\rangle. \quad (3.10)$$

Where  $\theta(t - t')$  is the Heaviside step function and the brackets  $\langle \rangle$  refer to the statistical mean over thermal states.  $G_{A,B}^r$  is called the retarded Green function. It is non-zero only if  $t > t'$ , such it allows to compute the response of the system after it has been perturbed. The advanced Green function  $G_{A,B}^a$  is the adjoint of  $G_{A,B}^r$  and represents exactly the opposite regime.

Usually, we can obtain all relevant physical properties just with the retarded Green function, including the density of states. To perform this, we need to take the Fourier transform of the retarded green function to enter into the energy domain

$$G_{A,B}(\omega) = \int_{-\infty}^{\infty} dt (t-t') G_{A,B}^r(t,t') e^{-i\omega(t-t')} . \quad (3.11)$$

In this formalism, the DOS is associated to a fermion operator  $A$ . Similar to (3.7), it can be computed as

$$\rho_{A,A^\dagger} = -\frac{1}{\pi} \text{Im} [G_{A,A^\dagger}(\omega)] . \quad (3.12)$$

The density of states contains important physical information related to operator  $A$ . In our case, operator  $A^\dagger$  will be related to the dot's creation operator  $d^\dagger$ . Therefore, computing (3.12) will allow us to observe the hybridization of the dot's discrete states and the creation of other energy levels due to the interaction with the lead and other impurities. In particular, we are interested in studying the zero-modes of the system which are the main signatures of Kondo and Majorana physics.

But before talking more about the DOS, we still need an efficient method to compute the Green function of the system. We can achieve this by analyzing the evolution of the retarded Green function in the time domain. This is determined by Schroedinger's differential equation in the Heisenberg picture

$$i \frac{dA(t)}{dt} = [A(t), H] , \quad (3.13)$$

which allows us to derive

$$i \frac{d}{dt} G_{A,B}^r(t,t') = -i^2 \langle [A(t), B(t)] \rangle \delta(t-t') - i\theta(t-t') \left\langle \left\{ i \frac{dA(t)}{dt}, B(t') \right\} \right\rangle \quad (3.14)$$

$$= \delta_{A^\dagger,B} \delta(t-t') - i\theta(t-t') \langle \{[A(t), H], B(t')\} \rangle \quad (3.15)$$

$$= \delta_{A^\dagger,B} \delta(t-t') + G_{[A,H'],B}^r(t,t') . \quad (3.16)$$

When taking the Fourier transform of (3.16) we obtain the following equation in the energy domain

$$\omega G_{A,B}(\omega) = \delta_{A^\dagger,B} + G_{[A,H'],B}(\omega) . \quad (3.17)$$

In a set of operators  $\{A_1, A_2, \dots\}$  (3.17) defines a system of transport equations describing the flow of state transitions of the operators in our model. This system receives the name of equations of motion. In this thesis we will identify each set of equations with a flow graph. This will be our leading method to compute the green functions of the system.

### 3.1.2 Presenting the Graph-Gauss-Jordan elimination process to solve the EOM

Solving the transport equations involves dealing with a set of linear equations where all the possible variables including  $\omega$ , and the Hamiltonian parameters are assumed to be constant. This can be done by Gauss-Jordan elimination, noting that after each elimination process we need to solve the system in terms of the initial variables. The solution will be a polynomial fraction. When the number of operators in the Hamiltonian increases the number of terms in the polynomial grows-up rapidly according to the number of initial parameters. This reveals the importance of exploring alternative methods that could simplify the solution of this system, and present a readable factorized expression of the final Green function.

The method presented here uses graph theory algorithms that provide a shortcut to Gauss-Jordan elimination [55]. This idea is not entirely new, as it was previously applied to study infinite tight binding lattices [56]. Nevertheless, the real insight of this approach is the possibility of using the graph structure to tackle problems with non-trivial geometries, just like the double quantum dot-Majorana system.

We will explain this method by solving the EOM for a non-interacting ( $U = 0$ ) DQD connected to one lead. According to the Anderson model, the Hamiltonian for this system looks like

$$H = \sum_{i=1}^2 \varepsilon_i d_i^\dagger d_i + t_{dots} d_1^\dagger d_2 + t_{dots}^* d_2^\dagger d_1 + \sum_k \left( V_i d_i^\dagger c_k + V_i^* c_k^\dagger d_i \right) + \varepsilon_k c_k^\dagger c_k. \quad (3.18)$$

Where the operators  $d_1^\dagger$  and  $d_2^\dagger$  create an electron in dots 1 and 2 respectively <sup>4</sup>. Since the system is non-interacting, we ignore the spin-degeneracy of this Hamiltonian. The only new parameter here is the term  $t_{dots}$ , which represents the tunneling between both quantum dots.

Using equation (3.17) with  $B = d_1^\dagger$  and  $A$  shifting among all the other operators we compute the following EOM

$$(\omega - \varepsilon_1) G_{d_1, d_1^\dagger}(\omega) = 1 + t_{dots} G_{d_2, d_1^\dagger}(\omega) + V_1^* \sum_k G_{c_k, d_1^\dagger}(\omega), \quad (3.19)$$

$$(\omega - \varepsilon_k) G_{c_k, d_1^\dagger}(\omega) = V_1 G_{d_1, d_1^\dagger}(\omega) + V_2 G_{d_2, d_1^\dagger}(\omega), \quad (3.20)$$

$$(\omega - \varepsilon_2) G_{d_2, d_1^\dagger}(\omega) = t_{dots} G_{d_1, d_1^\dagger}(\omega) + V_2^* \sum_k G_{c_k, d_1^\dagger}(\omega). \quad (3.21)$$

This system is already closed, hence, it will have a unique solution. The associated matrix form is

$$\begin{bmatrix} \omega - \varepsilon_2 & -V_2 & -t_{dots} \\ -V_2^* & \omega - \varepsilon_k & -V_1 \\ -t_{dots}^* & -V_1^* & \omega - \varepsilon_1 \end{bmatrix} \begin{bmatrix} G_{c_k, d_1^\dagger}(\omega) \\ G_{d_2, d_1^\dagger}(\omega) \\ G_{d_1, d_1^\dagger}(\omega) \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \quad (3.22)$$

<sup>4</sup>This case is different from the Anderson model defined in section 2.2, where the  $d_i^\dagger$  where creation operators in distinct energy levels, not dots.




Figure 3.1: Graph representation of Gauss-Jordan elimination a) Graph  $\mathcal{G}_{d_1d_2}$  b) After the elimination of vertex  $c_k$ , the energies of dots  $d_1$  and  $d_2$ , and the coupling parameter are changed. c) After Gaussian elimination of dot 2 the energy of the remaining dot  $\epsilon_{DQD}^+$  represents the transport information through  $d_1$  of the entire DQD.

By convenience we changed the order of the rows in the matrix and we removed the sum over  $k$  ( $\sum_k$ ) to simplify the algebraic operations. We will insert these terms back in the equations at the end of the procedure.

Although this matrix is not Laplacian, the procedure in [55] can still be applied with the downside of loosing part of the speed-up of the algorithm. We still preserve some of the advantages of using graphs, such as the schematic representation of the transport process, the possibility of taking minimal cuttings and the relation between Gauss-Jordan elimination and random walks [55]. All of these, simplify the complexity of the solution.

Now, our goal is to compute the green function  $G_{d_{1\downarrow}, d_{1\downarrow}^\dagger}(\omega)$ . To this end, we take the graph  $\mathcal{G}_{d_1d_2}$  associated to the matrix (3.22) (See Figure 3.1(a)). The vertexes of this graph are the operators in the first site of the of the green functions  $(d_{1\downarrow}, d_2, c_k, d_1^\dagger)$ .  $d_1^\dagger$  is not included since it only appears in the second sub-index of the green functions. The edges are given by the non-diagonal sites in the matrix multiplied by  $-1$  and there is a self-energy assigned to each vertex, equal to the difference between  $\omega$  and the corresponding diagonal term. These self-energies can also be understood as additional edges connecting each vertex with itself. The plot of the energy parameters in this algorithm is quite important, hence we prefer to keep this name to differentiate them from the other couplings.

The algorithm consists in the following: Each step of Gauss-Jordan elimination leads to a new graph with different energies and couplings. The elimination of a row and column is equivalent to eliminate the corresponding vertex in the graph. For instance, lets eliminate the first row and column of the matrix in (3.22). For it we just need to subtract the rank-1 matrix with the same first row and first column

$$\begin{bmatrix} \omega - \varepsilon_k & -V_2 & -V_1 \\ -V_2^* & \omega - \varepsilon_2 & -t_{dots} \\ -V_1^* & -t_{dots}^* & \omega - \varepsilon_1 \end{bmatrix} - \begin{bmatrix} \omega - \varepsilon_k & -V_2 & -V_1 \\ -V_2^* & \frac{V_2^* V_2}{\omega - \varepsilon_k} & \frac{V_2^* V_1}{\omega - \varepsilon_k} \\ -V_1^* & \frac{V_2 V_1^*}{\omega - \varepsilon_k} & \frac{V_1 V_1}{\omega - \varepsilon_k} \end{bmatrix} \quad (3.23)$$

$$= \begin{bmatrix} 0 & 0 & 0 \\ 0 & \omega - \varepsilon_2 - \frac{V_2^* V_2}{\omega - \varepsilon_k} & -t_{dots} - \frac{V_2^* V_1}{\omega - \varepsilon_k} \\ 0 & -t_{dots}^* - \frac{V_2 V_1^*}{\omega - \varepsilon_k} & \omega - \varepsilon_1 - \frac{V_1 V_1}{\omega - \varepsilon_k} \end{bmatrix} \quad (3.24)$$

The graph associated to this new matrix can be observed in Figure 3.1(b) where operator  $c_k$  has been removed. It is possible to associate the correction to the the energies and couplings to the possible walks passing through the vertex  $c_k$ . For instance  $d_1$ 's energy  $\varepsilon_1$  receives an extra term  $\frac{V_1^* V_1}{\omega - \varepsilon_k}$  representing an additional walk from  $d_1$  to  $d_1$  passing through  $c_k$ . The same logic can be applied to the other coupling terms. The correction to  $t_{dots}$  is  $\frac{V_1^* V_2}{\omega - \varepsilon_k}$  which corresponds to a path from  $d_1$  to  $d_2$  passing through the eliminated vertex  $c_k$ . Note that this term includes the multiplication of both couplings  $V_1, V_2^*$  divided by the difference of  $\omega$  with the self-energy  $\varepsilon_k$  of the vertex. This correspondence between the energy correction and eliminated paths through the graph makes the "vertex-elimination" process a straightforward task.

We now proceed to eliminate vertex  $d_2$ , which leaves just a single vertex as shown in Figure 3.1(c). The self-energy of it can be readily computed with the previous path elimination idea which gives

$$\varepsilon_{DQD}^+ = \varepsilon_1 + \sum_{\mathbf{k}} \frac{V_1 V_1^*}{\omega - \varepsilon_{\mathbf{k}}} + \frac{\left\| t_{dots} + \sum_{\mathbf{k}} \frac{V_1 V_2^*}{\omega - \varepsilon_{\mathbf{k}}} \right\|^2}{\omega - \varepsilon_2 - \sum_{\mathbf{k}} \frac{V_2 V_2^*}{\omega - \varepsilon_{\mathbf{k}}}}, \quad (3.25)$$

where we selectively included the  $\sum_{\mathbf{k}}$ -terms in the places where  $k$  appeared.

As a result of Gauss-Jordan elimination, the linear equation in (3.22) has evolved into the trivial form

$$\begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & \omega - \varepsilon_{DQD}^+ \end{bmatrix} \begin{bmatrix} G_{c_{\mathbf{k}}, d_1^\dagger}(\omega) \\ G_{d_2, d_1^\dagger}(\omega) \\ G_{d_1, d_1^\dagger}(\omega) \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}. \quad (3.26)$$

The Green function is then

$$G_{d_1, d_1^\dagger}(\omega) = \frac{1}{\omega - \varepsilon_{DQD}^+} = \left[ \left( \omega - \varepsilon_1 - \sum_{\mathbf{k}} \frac{V_1 V_1^*}{\omega - \varepsilon_{\mathbf{k}}} \right) - \frac{\left( t_{dots} + \sum_{\mathbf{k}} \frac{V_1 V_2^*}{\omega - \varepsilon_{\mathbf{k}}} \right) \left( t_{dots} + \sum_{\mathbf{k}} \frac{V_1 V_2^*}{\omega - \varepsilon_{\mathbf{k}}} \right)^*}{\omega - \varepsilon_2 - \sum_{\mathbf{k}} \frac{V_2 V_2^*}{\omega - \varepsilon_{\mathbf{k}}}} \right]^{-1}. \quad (3.27)$$

This Green function will appear again in the last chapter when solving the double dot-Majorana system.

### 3.1.3 Graph Algorithm

In this part we summarize the steps of the Graph-Gauss-Jordan elimination process to compute the Green function  $G_{d,d^\dagger}(\omega)$  of an operator  $d$ :

1. Computing the equations of motion with the second term of the Green function fixed in the creation operator  $d^\dagger$ .
2. Mapping the linear system to the associated directed flow graph. Enumerate the vertexes of the graph as  $v_n$ , setting  $v_1 = d$ . The self-energy  $\varepsilon_n$  of each vertex  $v_n$  is initialized as  $\omega$  minus the corresponding diagonal term  $c_{nn}$ . The coupling terms  $c_{ij}$  connecting vertexes  $v_i$  to  $v_j$  are given by the  $(i, j)$ -off-diagonal terms of the matrix multiplied by  $-1$ .
3. Removing one-by-one the vertexes of the graph, starting by the last vertex  $v_N$ . While removing each vertex  $v_n$  the extra-terms of the energies and couplings are computed as follows:
  - (a) Energies: Let  $c_{in}, c_{ni}$  be the coupling constants associated to the edges from  $v_i$  to  $v_n$  and from  $v_n$  to  $v_i$  respectively. In our case  $c_{ni} = c_{in}^*$ . Then there is an indirect path from  $v_i$  to itself passing through  $v_n$ . When  $v_n$  is eliminated, the extra-term added to  $\varepsilon_i$  is  $\frac{c_{in}c_{in}}{\omega - \varepsilon_n}$ .
  - (b) Couplings: Let  $c_{in}, c_{nj}$  be the coupling constants associated to the edges from  $v_i$  to  $v_n$  and from  $v_n$  to  $v_j$ . Then there is an indirect path from  $v_i$  to  $v_j$  passing through  $v_n$ . When  $v_n$  is eliminated, the extra-term added to  $c_{ij}$  is  $\frac{c_{in}c_{nj}}{\omega - \varepsilon_v}$ .

This process is iterated from  $n = N$  till  $n = 1$ .

4. The self-energy in the remaining vertex  $v_1 = d$  is related with the green-function as  $\varepsilon_d = \omega - \frac{1}{G_{d,d^\dagger}(\omega)}$ .

This algorithm will be our main method to find the Green function and therefore the density of states of any non-interacting system.

### 3.1.4 Non-interacting DOS in a double quantum dot coupled to a lead

In this subsection, we describe the remaining steps to obtain the density of states of the double quantum dot from the Green function (3.27). We will plot the results and observe the evolution of the DOS while tuning the parameters of the model.

First note that equation (3.27) depends on the term  $\sum_k \frac{V_i^* V_j}{\omega - \varepsilon_k}$ , which describes the broadening of the DOS when the QD enters in contact with the lead. This broadening is usually named

$\Gamma_i = V_i^* V_i$  (Or  $\Delta$  depending on the text book). In general,  $V_i$  is a function of  $\mathbf{k}$ . However, in the limit of flat-band we can assume that  $V_i$  is constant. Therefore, it is enough to integrate

$$\sum_k \frac{1}{\omega - \varepsilon_k + i\delta} = \int_{-D}^D \frac{d\varepsilon_k}{\omega - \varepsilon_k + i\delta} = -\ln \left( \frac{D - \varepsilon_k + i\delta}{-D - \varepsilon_k + i\delta} \right) \xrightarrow[D \rightarrow \infty]{} -i, \quad (3.28)$$

where we assumed that there is a maximum energy cutoff  $D$  going to infinity in the wide-band limit. Hence

$$-i\Gamma_i = \sum_k \frac{V_i^* V_i}{\omega - \varepsilon_k}. \quad (3.29)$$

We can replace this in equation (3.27) to obtain the real expression for the green function  $G_{d_1 d_1^\dagger}(\omega)$ . The terms of the form  $V_1 V_2^*$  can be replaced for  $\sqrt{\Gamma_1 \Gamma_2}$ , supposing there is no additional complex phase.

Now, remember from (3.12) that the DOS  $\rho$  depends on the imaginary factor of the Green Function  $G_{d_1 d_1^\dagger}(\omega)$ . This term depends in the broadening  $\Gamma$ . If  $\Gamma = 0$  the density of states will be 0 as well. At any other case, one of the dots should be attached to the lead. Let  $\Gamma_1$  be the broadening of this dot. We will take  $\Gamma_1$  as our unit for the rest of this thesis.

In Figure 3.2 we can observe the evolution of the Density of States under certain processes. Each plot includes an inset showing the model applied to the figure. The coupling in purple indicates the tuning variable. We set  $e_1 = e_2 = 0$  so that both dots satisfy PHS. The primary results are the

1. **Coupling QD2. Figure 3.2(a)(b):** At  $\Gamma_2 = 0$  the second dot is decoupled, hence the first dot's DOS is the same of a single dot case. The maximum height is achieved at  $\rho\pi\Gamma_1 = 1$  and the width of this peak is approximately  $\Gamma_1$ , just as in Figure Figure 2.2. When the second dot is attached  $\Gamma_2 > 0$ , the density of states is divided between both dots. At  $\Gamma_1 = \Gamma_2$  the DOS at the Fermi energy is equal to  $\frac{1}{4\pi\Gamma}$  for both dots. For higher values of  $\Gamma_2$ , the DOS in the second dot is higher than in the first one.
2. **Indirect Coupling of QD2. Figure 3.2(c)(d):** This case is interesting. When the second dot is connected indirectly through the first dot, quantum inference splits the central peak in two new states. We will observe later that in the interacting case this procedure can also destroy the Kondo signature. Note that the higher the coupling  $t_{dots}$  is, the greater is the gap between the states. We will usually take  $t_{dots} = 2\Gamma_1$  to make these gap more visible in the NRG simulations.
3. **Breaking Particle Hole Symmetry. Figure 3.2(e)(f):** Suppose we have  $\Gamma_2 = \Gamma_1$ . The "triangular connections" break Particle Hole Symmetry, with the central peak detuning to the positive part of the spectrum. We will avoid this situation during this project, because because PHS-breaking will prevent the Majorana to tunnel inside the DQD. Hence, this model won't lead to any interesting result on Majorana manipulation.




Figure 3.2: Evolution of the density of states at each QD (Left: dot 1, Right: dot 2) at three distinct arrangements of DQD-lead coupling. The inset at the first column depicts the type of coupling. The purple line represents the tuning variable. The energy unit is  $\Gamma_1$ .  $e_1 = e_2 = 0$  in all arrangements. (a),(b) The lead is connected to both QDs. Tuning variable:  $\Gamma_2$ . (c)(d) Indirect coupling of the second dot through dot one. Tuning variable:  $t_{dots}$ . (e)(f) Triangular coupling. Tuning variable:  $t_{dots}$ .

### 3.2 The Numerical Renormalization Group (NRG)

The numerical renormalization group is renowned for providing the most successful and complete explanation of the Kondo problem, of integrating the strong correlations in Coulomb interacting systems. In this chapter, we will review the main ideas of the NRG algorithm including the logarithmic discretization, the construction of Wilson's chain and the iterative diagonalization in symmetry blocks. At the end, we will give some specifications about our code and we will show its results in the model of a double quantum dot attached to a metallic lead.




Figure 3.3: NRG algorithm: The logarithmic discretization of the conduction band according to the discretization parameter  $\Lambda$ . The system with the lead attached to the QD-impurity is mapped to the Wilson's chain, such that each step of the chain contains the contributions at different energy scales  $\Lambda^{-N/2}$ .

### 3.2.1 From the Renormalization Group to Wilson's Chain

The real meaning of the divergent logarithmic term in the resistivity predicted by Kondo is that important contributions at low-energy scales caused by the strong quantum correlations in the system are being neglected by perturbation theory. This problem can be solved by introducing ideas from renormalization group theory. A renormalization approach is more adequate for this type of problem since it assigns an appropriate effective Hamiltonian to each scale of temperature. This provides a more accurate representation of the increasing density of correlated states appearing close the Fermi energy.

In renormalization group theory, the Hamiltonian transformations are performed by an operator  $\mathcal{T}$  that represents an endomorphism in the space of operators.  $\mathcal{T}$  generates the semigroup  $\{1, \mathcal{T}, \mathcal{T}^2, \dots\}$ , that defines a complete set of transformations

$$\mathcal{T}[H_0] = H_1, \mathcal{T}[H_1] = H_2, \dots, \mathcal{T}[H_N] = H_{N+1}, \dots$$

If  $\mathcal{T}$  is a contracting map<sup>5</sup> then it is known that this set of operations should eventually lead to a fix point  $\mathcal{T}^N[H] \xrightarrow{N \rightarrow \infty} H^*$  such that  $\mathcal{T}[H^*] = H^*$ . In numerical simulations,  $N$  will only

---

<sup>5</sup> Let  $\mathcal{O}$  be a set of operators, then  $T$  is a contracting map if  $\mathcal{T}[\mathcal{O}'] \subset \mathcal{O}'$  for every  $\mathcal{O}' \subset \mathcal{O}$ .

### 3.2. The Numerical Renormalization Group (NRG)

---

increase up to a value where  $H_N$  is close enough to the fix point  $H^*$  so that no new significant contributions to the Hamiltonian are obtained. For the purposes of this project, taking  $N = 51$  will be enough for the NRG code to converge.

In the 1970's G.Wilson used this theory to create the famous Numerical Renormalization Group (NRG) [13, 50, 57]. His main idea was to perform a logarithmic discretization of the conduction band in the lead as shown in Figure 3.3.(Left). Taking into account that the leading contributions to the conductance occur at states close to the Fermi energy  $\omega = 0$ , we can define a cut-off ( $|\omega| < D$ ) so that the rest at higher contributions are not relevant. Then we use  $D$  to rescale the energy interval. As you can observe in the figure, the QD is coupled to all these energy states at the same time. The logarithmic discretization gives more relevance to the low energy scales by assigning a different Hamiltonian coupling to each one of them. The discretized intervals are determined by a variable  $\Lambda > 1$ . This value has certain relevance on the convergence of the code. If it is selected to small ( $\Lambda \sim 1$ ) it might never converge, but if it is too high it could produce misleading results. In this thesis we use a definite value of  $\Lambda = 2.5$ . This value was tested several times, giving positive results.

To complete this idea, the NRG code maps the Hamiltonian of the QD-lead system to the Wilson's chain shown in Figure 3.3(Right), so that each step of the chain contains the contributions of a different energy scale. A detailed description of this map is included in the Appendix section A.1. After these steps we obtain a chain Hamiltonian of the form

$$H = H_d + D \sum_{\sigma} \left[ \sqrt{\frac{2\Gamma}{\pi D}} (d_{\sigma}^{\dagger} f_{0\sigma} + f_{0\sigma}^{\dagger} d_{\sigma}) + \frac{1}{2} (1 + \Lambda^{-1}) \sum_{n=0}^{\infty} \Lambda^{-\frac{n}{2}} \xi_n (f_{n\sigma}^{\dagger} f_{n+1,\sigma} + f_{n+1,\sigma}^{\dagger} f_{n\sigma}) \right]. \quad (3.30)$$

In the flat-band approximation the parameters  $\xi_n$  can be obtained analytically [13]

$$\xi_n = \frac{1 - \Lambda^{-n-1}}{(1 - \Lambda^{-2n-1})^{\frac{1}{2}} (1 - \Lambda^{-2n-3})^{\frac{1}{2}}}.$$

From equation (3.30) we define the following sequence of shell Hamiltonians

$$H_{N+1} = T [H_N] = \Lambda^{\frac{1}{2}} H_N + \xi_N (f_{N+1,\sigma}^{\dagger} f_{N,\sigma} + f_{N,\sigma}^{\dagger} f_{N+1,\sigma}), \quad (3.31)$$

with

$$H_{-1} := \frac{2H_d \Lambda^{-1/2}}{D(1 + \Lambda^{-1})}, \quad \xi_{-1} = \sqrt{\frac{2\Gamma}{\pi D}}. \quad (3.32)$$

Note that at each step  $H_N$  is being rescaled by a factor  $\Lambda^{1/2}$  before including the following contribution. This is performed iteratively to eliminate the high order excitations, which allows to obtain an effective model valid for the lower energies. In consequence, each Hamiltonian  $H_N$  represents only the physics that is relevant at the energy scale  $\Lambda^{-N/2}$ .

On the other hand, the Renormalization Group transformation  $\mathcal{T}$  can be defined as




Figure 3.4: Schematic representation of the spectrum in the iterative diagonalization process. a)→b): Rescaling. b)→c): Including next site and diagonalizing. c)→d): Truncation of high energy states . Adapted from [13]

$$\mathcal{T}^N H_{-1} = \frac{1 + \Lambda^{-1}}{2} \Lambda^{\frac{-N}{2}} H_N$$

In the limit  $\xrightarrow{N \rightarrow \infty}$  we should recover the initial Anderson Hamiltonian. In addition, note that the leading coefficients of the contributions to each Hamiltonian  $H_N$  are given by

$$\Lambda^{\frac{-N}{2}} \xi_N \xrightarrow{N \rightarrow \infty} \frac{\Lambda^{\frac{-N}{2}} (1 - \Lambda^{-N})}{1 - \Lambda^{-2N}} \sim \frac{\Lambda^{\frac{-N}{2}}}{1 + \Lambda^{-N}},$$

which decays exponentially with the length of the chain. Therefore, we may think that at some point these new contributions will be so small that the that the map  $\mathcal{T}$  will eventually converge. Formally the theory for NRG convergence is too complex for this thesis. However the results show that the operator that truly converges is  $\mathcal{T}^2$  and not  $\mathcal{T}$ [57]. This has important consequences, for instance the convergence of the code has to be analyzed on odd and even values of  $N$  separately.

To this point, the equation (3.30) and the derived limit of  $H_N$  to the Anderson Hamiltonian are exact expressions. The first approximations will be performed in the following section which describes the iterative diagonalization of the shell Hamiltonians.

### 3.2.2 Iterative Diagonalization

The diagonalization begins with the dot-impurity Hamiltonian  $H_{-1}$ , which must be written in matrix form according to a defined basis. The other steps can be defined by induction. Suppose that the spectrum of  $H_N$  is diagonal on a given basis. Then the NRG code performs each of the following steps:

1. Rescaling the spectrum of  $H_N$  by  $\Lambda^{\frac{1}{2}}$  as defined in (A.12). Figure 3.4 (a)→(b).
2. Adding the next step of the chain to form  $H_{N+1}$  and diagonalizing the new Hamiltonian such that  $H_{N+1} = U_{N+1}^\dagger D_{N+1} U_{N+1}$ . After this step, each of the eigenstates of  $H_N$  will split in up to 4 new energy states (probably degenerate) determined by the new coupling with the  $N+1$  site basis  $|0\rangle, |\uparrow\rangle, |\downarrow\rangle, |\uparrow\downarrow\rangle$ . Figure 3.4 (b)→(c).
3. Shifting the spectrum by a certain factor such that the zero of energy is always the ground state. Figure 3.4 (c)→(d).
4. Numerical cutting: If the number of states in the system exceeds a definite number (1000 in this thesis) the exceeding higher energy states are neglected<sup>6</sup>. This is in agreement with the previously exposed idea of eliminating high order excitations to obtain a valid effective model at low energies. Figure 3.4 (c)→(d).
5. Rotating operators  $f_{N,\sigma}$  by  $U_N f_{N,\sigma} U_N^\dagger$  to start the next operation.

The final outcome of this operations will be the complete spectrum of the Anderson model at each energy level. However we still need to talk about an important speed-up to the code obtain when considering the symmetries of the system.

### 3.2.3 Symmetries

The symmetries of the initial Hamiltonian take a very important role in this iterative diagonalization. Lets suppose that the initial Hamiltonian  $H_d$  has certain symmetries classified by the quantum number  $S$ . Then  $H_d$  can be written in block Hamiltonians over a basis of the form  $|S, i\rangle$ . A diagonalization process of an square matrix with  $L$  rows usually has square order, proportionate to the number of entries  $\mathcal{O} \sim L^2$ . However, if the matrix is organized in blocks of length  $L_j$  such that  $\sum_j L_j = L$ , then the order of diagonalization will be around  $\sum_j L_j^2$  which is in general much smaller than  $(\sum_j L_j)^2 = L^2$ . Therefore the block diagonalization provides an important numerical speed-up to the algorithm.

To maintain this advantage, we must preserve this symmetric structure for the rest of the NRG code. For it, we first need to verify that the picked symmetry also commutes with the

---

<sup>6</sup> This step must be performed carefully to preserve the symmetries of the system. If two states are entangled and one of them is eliminated and the other is not, the program could lead to misleading results. Further discussions to solve this problem are presented in the symmetry subsection.

hopping terms in the chain Hamiltonian. If so, for each step  $N$  of the NRG algorithm the  $H_N$  Hamiltonian can be written in a block diagonal form with basis  $|S_N, i_N\rangle$ . Then it is necessary to define transition rules from the quantum numbers  $S_N$  to  $S_{N+1}$ . By doing this, we assure that the block architecture is transmitted through the entire algorithm, hence reducing the computational time significantly at each step of the NRG chain. This is a key procedure to optimize the algorithm.

In the following subsection we will give a example of this symmetry propagation in the model a quantum dot attached to a metallic lead.

### 3.2.4 Iterative diagonalization in a single QD attached to a metallic lead

Now that we have an iterative representation of the Anderson Model Hamiltonian (3.30), we will describe how the NRG code works for a QD attached to a lead. We start with the dot Hamiltonian.

$$H_d = \frac{1}{D} \left( \epsilon + \frac{U}{2} \right) d_\sigma^\dagger d_\sigma + \frac{U}{2D} (d_\sigma^\dagger d_\sigma - 1)^2. \quad (3.33)$$

Now observe that Hamiltonian (3.33) is already in the diagonal form for the basis  $\{| \uparrow\downarrow \rangle, | \uparrow \rangle, | \downarrow \rangle, | 0 \rangle\}$

$$H_d = \frac{1}{D} \begin{bmatrix} 2\epsilon + \frac{3U}{2} & 0 & 0 & 0 \\ 0 & \epsilon + \frac{U}{2} & 0 & 0 \\ 0 & 0 & \epsilon + \frac{U}{2} & 0 \\ 0 & 0 & 0 & \frac{U}{2} \end{bmatrix}.$$

Lets define  $H_{-1} := \frac{2H_d\Lambda^{-1/2}}{1+\Lambda^{-1}}$  as in 3.32. It is convenient here to define unit-less variables  $\epsilon' = \frac{2\epsilon\Lambda^{-1/2}}{D(1+\Lambda^{-1})}$ ,  $U' = \frac{2U\Lambda^{-1/2}}{D(1+\Lambda^{-1})}$  and  $\Gamma' := \sqrt{\frac{2\Gamma}{\pi D}}$ . Then adding the first site of the Wilson's chain interaction to  $H_d$  we obtain a new Hamiltonian of the form

$$H_0 = \Lambda^{\frac{1}{2}} H_{-1} + \Gamma' \left( d_\sigma^\dagger f_{0\sigma} + f_{0\sigma}^\dagger d_\sigma \right). \quad (3.34)$$

The Hilbert space for this new Hamiltonian must be extended to include the 4 degrees of freedom of the  $f_{0\sigma}^\dagger$  particles, which are given by  $\{| \uparrow\downarrow \rangle, | \uparrow \rangle, | \downarrow \rangle, | 0 \rangle\}$ . Therefore, the total Hilbert space for  $H_0$  is given by a base of the form

$$|s_1\rangle|s_2\rangle := |s_1\rangle \otimes |s_2\rangle \text{ with } |s_i\rangle \in \{| \uparrow\downarrow \rangle, | \uparrow \rangle, | \downarrow \rangle, | 0 \rangle\}.$$

We obtain an space of dimension  $4 \times 4 = 16$ . Now, before adventuring to write the Hamiltonian for  $H_0$  as a  $16 \times 16$ -matrix note that  $H_{-1}$  preserves the total particle number  $\mathcal{N}$  and the total spin  $S$ . We can associate each state to these symmetries as

$$|\uparrow\downarrow\rangle \rightarrow |\mathcal{N}=2, S=0\rangle, |0\rangle \rightarrow |\mathcal{N}=0, S=0\rangle \quad (3.35)$$

$$|\uparrow\rangle \rightarrow |\mathcal{N}=1, S=\frac{1}{2}\rangle, |\downarrow\rangle \rightarrow |\mathcal{N}=1, S=-\frac{1}{2}\rangle. \quad (3.36)$$

The propagation rule for the symmetry is defined with the following identity

$$|\mathcal{N}_1, S_1\rangle \otimes |\mathcal{N}_2, S_2\rangle \subset |\mathcal{N}_1 + \mathcal{N}_2, S_1 + S_2\rangle \quad (3.37)$$

Then we can use  $\mathcal{N}$  and  $S$  as quantum numbers to generate the block structure of  $H_0$ . We will observe that the terms in the diagonal will correspond to the eigenvalues of  $H_{-1}$  for the first space. The non-diagonal terms are the result of the hopping interactions with the first site.

$$H_{\mathcal{N}=0, S=0} :$$

$$|0\rangle|0\rangle \rightarrow \left[ \begin{array}{c} U' \\ 2 \end{array} \right] \quad |\uparrow\downarrow\rangle|\uparrow\downarrow\rangle \rightarrow \left[ \begin{array}{c} 2\varepsilon' + \frac{3U'}{2} \\ 2 \end{array} \right]$$

$$H_{\mathcal{N}=1, S=\frac{1}{2}} :$$

$$\begin{aligned} |\uparrow\rangle|0\rangle &\rightarrow \left[ \begin{array}{cc} \varepsilon' + \frac{U'}{2} & \Gamma' \\ \Gamma' & \frac{U'}{2} \end{array} \right] \\ |0\rangle|\uparrow\rangle &\rightarrow \left[ \begin{array}{cc} \varepsilon' + \frac{U'}{2} & \Gamma' \\ \Gamma' & \frac{U'}{2} \end{array} \right] \end{aligned}$$

$$H_{\mathcal{N}=2, S=-1} :$$

$$|\downarrow\rangle|\downarrow\rangle \rightarrow \left[ \begin{array}{c} \varepsilon' + \frac{U'}{2} \\ 2 \end{array} \right] \quad |\uparrow\rangle|\uparrow\rangle \rightarrow \left[ \begin{array}{c} \varepsilon' + \frac{U'}{2} \\ 2 \end{array} \right]$$

$$H_{\mathcal{N}=2, S=0} :$$

$$\begin{aligned} |\uparrow\downarrow\rangle|0\rangle &\rightarrow \left[ \begin{array}{cccc} 2\varepsilon' + \frac{3U'}{2} & \Gamma' & -\Gamma' & 0 \\ \Gamma' & \varepsilon' + \frac{U'}{2} & 0 & \Gamma' \\ -\Gamma' & 0 & \varepsilon' + \frac{U'}{2} & -\Gamma' \\ 0 & \Gamma' & -\Gamma' & \frac{U'}{2} \end{array} \right] \\ |0\rangle|\uparrow\downarrow\rangle &\rightarrow \left[ \begin{array}{cccc} 2\varepsilon' + \frac{3U'}{2} & \Gamma' & -\Gamma' & 0 \\ \Gamma' & \varepsilon' + \frac{U'}{2} & 0 & \Gamma' \\ -\Gamma' & 0 & \varepsilon' + \frac{U'}{2} & -\Gamma' \\ 0 & \Gamma' & -\Gamma' & \frac{U'}{2} \end{array} \right] \end{aligned}$$

$$H_{\mathcal{N}=3, S=\frac{1}{2}} :$$

$$\begin{aligned} |\uparrow\downarrow\rangle|\uparrow\rangle &\rightarrow \left[ \begin{array}{cc} \varepsilon' + \frac{U'}{2} & -\Gamma' \\ -\Gamma' & \frac{U'}{2} \end{array} \right] \quad |\uparrow\downarrow\rangle|\downarrow\rangle \rightarrow \left[ \begin{array}{cc} \varepsilon' + \frac{U'}{2} & -\Gamma' \\ -\Gamma' & \frac{U'}{2} \end{array} \right] \\ |\uparrow\rangle|\uparrow\downarrow\rangle &\rightarrow \left[ \begin{array}{cc} \varepsilon' + \frac{U'}{2} & -\Gamma' \\ -\Gamma' & \frac{U'}{2} \end{array} \right] \quad |\downarrow\rangle|\uparrow\downarrow\rangle \rightarrow \left[ \begin{array}{cc} \varepsilon' + \frac{U'}{2} & -\Gamma' \\ -\Gamma' & \frac{U'}{2} \end{array} \right] \end{aligned}$$

The next step would be to diagonalize  $H_0$  by blocks ( $H_{\mathcal{N}, S}$ ) and then including the next place in the chain. The following Hamiltonians are generated in the same way from equation (3.31). The symmetries of the new states can be obtained from the propagation rule (A.15). When the number of states surpasses the 1000 states, the code will automatically cutoff the




Figure 3.5: Evolution of the QD-spectrum vs number of iterations of the code for  $U' = 0.5$ ,  $\epsilon' = -0.25$ ,  $\Gamma = 2.82 \times 10^{-2}$ .

higher energy states. However, it is important that a block is not divided during this cutting protocol since it could break the preserved symmetries.

Finally, the spectrum for  $\Lambda = 2.5$  takes the “spaghetti” form in Figure 3.5. Note that before  $N = 30$ , the low-energy contributions generate significant changes in the energy levels. However, the stable spectrum after the step  $N = 30$  is a signature that the code has converged. As we previously declared, it is not  $\mathcal{T}$  but  $\mathcal{T}^2$  the transformation that has fixed points, which explains why it was necessary to plot the even and the odd spectrum separately.

At the end of the NRG code, we obtain a complete list of the spectrum ( $E_{i,N}$ ) and the eigenstates ( $|i,N\rangle$ ) of the Hamiltonian at each step  $N$  of the chain (See Figure 3.5). It is important to keep all of these states since each one of them represents different thermodynamic regimes of the system. While the site  $N = 1$  represents the physics of the system relevant at temperatures around  $\frac{D\Lambda^{-1}}{k_B}$ , the site  $N = 30$  shows the low energy contributions at  $T \sim \frac{D\Lambda^{-30}}{k_B}$  where the ground state is strongly correlated. Furthermore, the states at low temperatures are entangled with the higher energy states. We need to take this into account to extract dynamical quantities of the system, which is the objective of the following section.

### 3.2.5 The Density Matrix Renormalization Group (DM-NRG)

The NRG codes allows us to compute several thermodynamic quantities such as the entropy  $S$ , the free energy and the partition function  $Z(\beta)$ . In addition, we can compute the spin magnetization or dynamical quantities such the density of states, the magnetic susceptibility and the conductivity. To perform this we can use the spectrum obtained in the NRG code to define the

Boltzman distribution of the system. Then we apply the usual methods of statistical mechanics.

In this thesis we will focus in computing the density of states at the impurity (QD). For this, let  $|j\rangle$  and  $|q\rangle$  label a base of eigenstates of the Hamiltonian  $H$ . Now recall the definition of the retarded green function (3.9)

$$G_{d,d^\dagger}(t) = \theta(t) \langle \{d(t)d^\dagger(0)\} \rangle \quad (3.38)$$

$$= \theta(t) \left\langle \left[ e^{\frac{i}{\hbar}Ht} d e^{-\frac{i}{\hbar}Ht} d^\dagger \right] \right\rangle + \theta(-t) \left\langle \left[ d^\dagger e^{\frac{i}{\hbar}Ht} d e^{-\frac{i}{\hbar}Ht} \right] \right\rangle \quad (3.39)$$

$$= \theta(t) \sum_{|j\rangle, |q\rangle} p_j \langle j | e^{\frac{i}{\hbar}Ht} d | q \rangle \langle q | e^{-\frac{i}{\hbar}Ht} d^\dagger | j \rangle + \theta(-t) \sum_{|j\rangle, |q\rangle} p_q \langle q | d^\dagger e^{\frac{i}{\hbar}Ht} | j \rangle \langle j | d e^{-\frac{i}{\hbar}Ht} | q \rangle \quad (3.40)$$

$$= \theta(t) \sum_{|j\rangle, |q\rangle} p_j e^{\frac{-i}{\hbar}t(E_j - E_q)} \|\langle j | d | q \rangle\|^2 + \theta(-t) \sum_{|j\rangle, |q\rangle} p_q e^{\frac{i}{\hbar}t(E_j - E_q)} \|\langle j | d | q \rangle\|^2. \quad (3.41)$$

Where  $p_j := \frac{e^{-\beta E_j}}{Z(\beta)}$  defines the Boltzmann probability of the eigenstate  $|j\rangle$  according to Hamiltonian  $H$ .

It is known that the Fourier transform of an expression of the form  $e^{-ax}\theta(x)$  is  $\frac{1}{\omega + is - a}$ . Then the Green function in the energy domain is

$$G_{d,d^\dagger}(\omega) = \frac{1}{Z(\beta)} \sum_{|j\rangle, |q\rangle} \frac{e^{-\beta E_j} + e^{-\beta E_q}}{\omega + is - E_j + E_q} \|\langle j | d | q \rangle\|^2. \quad (3.42)$$

From the imaginary part of (3.42) we obtain a formula for the spectral density in terms of the eigenstates and energies of the Hamiltonian

$$\rho_d = \frac{1}{Z(\beta)} \sum_{|j\rangle, |q\rangle} \left( e^{-\beta E_j} + e^{-\beta E_q} \right) \delta(\omega - E_j + E_q) \|\langle j | d | q \rangle\|^2. \quad (3.43)$$

This new expression for the DOS can be integrated into the NRG code in different ways . A first method created by Costi *et al.* consists in computing (3.43) with the eigenstates of each shell Hamiltonian  $H_N$  [58]. It is necessary to take into account that the operator  $d$  is constantly rotating after each diagonalization procedure, which produces different representations at each shell Hamiltonian basis. Then, an important part of Costi's algorithm is to obtain these new representations of  $d$  at the  $H_N$ -basis ( $\langle j | d | q \rangle|_N$ ) recursively starting from an input representation  $\langle j | d | q \rangle|_{N=0}$ .

Although Costi's method predicts accurately the DOS at low-energies, it fails to fit the high energy levels. The method that corrects this problem receives the name of Density Matrix Numerical Renormalization Group (DMNRG) [14]. The main idea of DMNRG is to include the entanglement corrections with the lower energy-states using the density matrix formalism. For this, Hofstetter defines the density matrix at the last shell Hamiltonian  $\hat{\rho}_{N_{max}}$  as the thermal mixed state

### 3.2. The Numerical Renormalization Group (NRG)




Figure 3.6: Wilson's chain depicted as an open quantum system. Adapted from [14].

$$\hat{\rho}_{N_{\max}} = \sum_j e^{-\beta E_j} |j\rangle_{N_{\max}} \langle j|, \quad (3.44)$$

where the subindex  $N_{\max}$  suggests that  $|j\rangle_{N_{\max}} \langle j|$  is an eigenstate of  $H_{N_{\max}}$ .

With this new density matrix, we could rewrite (3.39) as

$$G_{d,d^\dagger}^{N_{\max}}(t) = \text{Tr}(\hat{\rho}_{N_{\max}} \theta(t) \{d^\dagger(t), d(0)\}). \quad (3.45)$$

From the Green function  $G_{d,d^\dagger}^{N_{\max}}(t)$  we can obtain the density of states associated to the temperature  $T_{N_{\max}}$ . Nevertheless, these results are not relevant at higher temperatures.

To solve this problem we may think Wilson's chain as an open quantum system where the environment are the low-energy sites of the chain and the system contains the high-energy sites including the QD-impurity as observed in Figure 3.6. Using this analogy, we can readily obtain the density matrix  $\rho_S$  by taking the partial trace over the environment

$$\rho_{\text{sys}} = \text{Tr}_{\text{env}}[\rho_{N_{\max}}]. \quad (3.46)$$

The DM-NRG code applies recursively this idea to obtain the density matrix corresponding to each scale of temperature  $T_N$ . It starts from  $\rho_{N_{\max}}$  defined at (3.44) and obtains  $\rho_{N_{\max}-1}$  by taking the partial trace over the vector space corresponding to the last site of the chain

$$\rho_{N_{\max}-1} = \text{Tr}_{N_{\max}}[\rho_{N_{\max}}]. \quad (3.47)$$

The other density matrices are computed by induction as

$$\rho_{N-1} = \text{Tr}_N[\rho_N], \quad (3.48)$$

and the density of states at each temperature regime can be computed from the green function

$$G_{d,d^\dagger}^N(t) = \text{Tr}(\hat{\rho}_N \theta(t) \{d^\dagger(t), d(0)\}). \quad (3.49)$$

The DM-NRG algorithm produces significantly better results at high energies than Costi's initial idea. Indeed, DM-NRG is still one of the best methods to compute dynamic quantities of an impurity system. In the following subsection we will give some details of how DM-NRG is integrated with the NRG code.




Figure 3.7: Diagram of the NRG code

### 3.2.6 Specifications of the NRG Code

The NRG code used in this thesis was implemented by my thesis advisor Luis Gregorio Dias during his posdoc at the University of Ohio. The general scheme is shown in Figure 3.7. It incorporates different stages. Here we give a brief description of them:

- **Model:** The model defines the type of impurity that we are going to study. It could be a single QD or a more complex structure, such as a DQD or a Majorana-QD system. My main contribution to this code was at this stage. I designed the mode DQD-Majorana, which describes a DQD coupled to a Majorana zero mode. These models preserve different symmetries. In the single dot Hamiltonian presented in subsection 3.2.4 we used the symmetry  $\mathcal{N}S_z$ , which is equivalent to charge-spin  $QS_z$ . However we will find that Majorana systems require another symmetry-type that we call as  $N \uparrow P \downarrow$ . The model is defined by an initial Hamiltonian  $H_{-1}$  and the annihilation operators  $d_{i\sigma}$  which must be submitted in the code written in the block symmetry representation described in subsection 3.2.3.
- **Input:** The input is a '.dat'-file that attributes a numerical value to each parameter of the

model. In addition, it allows to set different code specifications as the number of iterations  $N_{max}$ , the scaling parameter  $\Lambda$  and the maximum number of states before the cut-off. It is also possible to include additional implementations to improve the results of the NRG code such as the Z-trick [59] and the Complete Fock State. In this project, we only used the Z-trick, which significantly improves the spectral resolution at high energies.

- **NRG:** This part of the code mainly integrates the ideas of subsection 3.2.1 and implements the iterative diagonalization described in subsection 3.2.2. Each shell Hamiltonian  $H_n$  is diagonalized. The high-energy eigenstates are cut-off if they exceed the maximum limit. Then, the states are rotated and the eigenvalues are rescaled to include the next step of the Wilson's chain. The symmetry block structure is preserved during the entire loop. NRG's output is a detailed evolution of the spectrum which produces the spaghetti form. In addition, it can print the states and operators that are necessary to start other instances of the code like DM-NRG.
- **Site  $N + 1$ :** This is a small class that creates another site of the chain in the base  $\{| \uparrow\downarrow \rangle, | \uparrow \rangle, | \downarrow \rangle, | 0 \rangle\}$ . This base must be rewritten according to the symmetry quantum number. It is coupled with the matrices at the NRG code.
- **DM-NRG:** As described in subsection 3.2.5, this code generates iteratively the density matrix associated to each energy scale . Then it computes the green function and the density of states. The DOS of the single QD model is an example of its outputs. The plot shows the characteristic Kondo peak at the Fermi energy in the middle of the Coulomb peaks describing the energy states.

This NRG code is implemented in C++. It can be cloned from the Github link <https://git.io/fh9cM>. To optimize the performance of NRG, the code uses the packages Boost, LAPACK and Gnu Scientific Library (GSL), which provide a rapid interface for numerical matrix diagonalization.

### 3.2.7 NRG results in a double quantum dot coupled to a metallic lead

We now intend to observe the results of this code applied to the model of a double quantum dot attached to a metallic lead. The Hamiltonian for this system is the Anderson model with the impurity described by the interacting version of Hamiltonian (3.18)

$$H_{DQD} = \sum_{i=1}^2 \sum_{\sigma} \left( \epsilon_i + \frac{U_i}{2} \right) d_{i\sigma}^\dagger d_{i\sigma} + U_i \left( \sum_{i\sigma} d_{i\sigma}^\dagger d_{i\sigma} - 1 \right)^2 + \sum_{\sigma} t_{dots} d_{1\sigma}^\dagger d_{2\sigma} + \text{h.c} \quad (3.50)$$

The Hilbert space of this system has 16-dimensions and the symmetries in this Hamiltonian are exactly the same than the ones in the single QD case  $\mathcal{N}S$ . Nevertheless, we decided to test the DQD-Majorana mode in this system. By setting the Majorana couplings to  $t_1 = t_2 = 0$  we can




Figure 3.8: Density of states for the DQD attached to the lead at different configurations. Left insets: Setup that is being used. Right insets: Low energy DOS.

decouple the Majorana mode, hence obtaining the DQD model (See chapter 5). The results we obtained are in agreement with previous previous papers [1], which was an important test to confirm the veracity of this new mode.

For the entire thesis we will fix the value of the coulomb repulsion parameters to

$$U_1 = U_2 = 17.7305\Gamma_1. \quad (3.51)$$

We picked these parameters considerably higher than the broadening unit  $\Gamma_1$  to guarantee the appearance of Kondo physics at visible temperatures in comparison with the Coulomb peak.

**NRG-code** We used the same configurations from Figure 3.2. The results appear in Figure 3.8:

1. The single QD attached to a metallic lead is a particular case of the double quantum dot model, where the second dot is not attached  $\Gamma_2 = t_{dots} = 0$ . Figure 3.8 shows the NRG

### 3.2. The Numerical Renormalization Group (NRG)

---

results for this case. The three plots show the external Coulomb peaks at  $e_1 = \frac{U}{2} \sim 8.62\Gamma_1$ , which represent the DOS of the energy levels. In addition, Figure 3.8a shows a central peak at the Fermi energy. Note that there shouldn't be a state in this position since it is inside a gap between two energy levels (Coulomb peaks). **This is the Kondo Peak [53].** This peak explains the zero-bias plateaus in Figure 2.5.

2. In Figure 3.8b we observe the DOS when the two QDs are symmetrically attached. At low energies, the inset shows the appearance of two satellite peaks representing the Ruderman-Kittel-Kasuya-Yosida (RKKY) interaction. This is an anti-ferromagnetic coupling that appears in the interacting case due to the strong correlations, which cause an state-exchange interaction between both dots through the lead at energies around  $\pm 3.56\Gamma^2/U$  [60].
3. The most interesting case is in Figure 3.8c. As we already observed in the non-interacting case, the interference with the second dot completely destroys the zero mode which is now formed by the Kondo peak. This effect is observed at low energies closed to  $t_{dots} = 0.689\Gamma$ . The paper describing this result was a central part of one of my advisor's project [1]. This result encouraged me to formulate the following question. What happens if we attached a Majorana mode to one of these dots?. Would it be destroyed by interference or it will survive to it. Solving this question is one of the main objectives of chapter chapter 5.

With these results we finish the first test of our methods. We will probe them again in the next chapter in the QD-Majorana system, and again in chapter 5 to study the manipulation of Majorana zero modes in a double quantum dot.

## Chapter 4

# Coupling a QD with a Majorana zero mode

In the introduction we pointed out three properties that make particularly promising the QD-Majorana systems. These are:

1. It does not destroy the entire qubit information.
2. The possibility of exploring Kondo-Majorana physics.
3. It provides the unique opportunity of manipulating the Majorana zero modes inside the QDs, which has promising applications in the design of quantum architectures.

In this project we will exploit the second and the third properties to manipulate MZMs in double quantum dot systems in the Kondo regime. But before going through that model, we are going to test our methods in the single dot-Majorana system. As we will observe in this chapter, we were able to reproduce effectively the results of previous papers [2, 3].

### 4.1 Model

In this section we will recreate the results of Liu and Baranger using the methods developed in chapter 3. This will also allow us to probe our methods in a system with Majorana zero modes.

The Hamiltonian for Majorana-QD-lead hybrid system (See Figure 4.1) is given by

$$H = H_{QD-Lead} + H_{M-QD} + H_M. \quad (4.1)$$

Where  $H_{QD-Lead}$  is the Hamiltonian for the non-interacting Anderson model (2.6),  $H_M$  is the Hamiltonian of the Majorana chain and  $H_{M-QD}$  represents the coupling between the QD and the Majorana Fermion at the boundary.

Now, the real question is how to define the coupling between the QD and the MBS. In fact, there are many ways to represent this interaction. One alternative is to replace in  $H_M$  with the entire Kitaev chain Hamiltonian (1.1) (or even with the Majorana chain (1.18)) and then pick  $H_{M-QD}$  as the coupling between the QD and the first site of the chain [38]. A simpler approach is to define an effective coupling with the Majorana operator at the edge of the Majorana chain. Since the Kitaev chain is spin-less, we can choose to couple the Majorana to the spin- $\downarrow$  channel of the QD. It has been debated in previous papers that the magnetic fields applied to these type of




Figure 4.1: Model for the QD-Majorana system. Solid lines: Hopping interactions:  $V_1$  couplings of QD1 . Dashed lines: Majorana spin- $\downarrow$  effective couplings (4.3)  $t_1$ . The atomic energy levels appear inside each QD  $\epsilon_1$  are tuned by the gate voltages. The coulomb interaction is represented by  $U_1$  separates two energy levels. The red dashed horizontal lines represent the Fermi level.

systems -Usually surrounding  $500mT$ - are not big enough to disturb our choice of matching the spin- $\downarrow$  coordinate with the channel coupled to the spin-less MZM [3]. Therefore, the Majorana fermion should be the superposition of the creation and annihilation operators of a spin  $\downarrow$  particle  $f_\downarrow$ :

$$\gamma_1 := \frac{1}{\sqrt{2}} (f_\downarrow^\dagger + f_\downarrow), \gamma_2 := \frac{1}{\sqrt{2}} (f_\downarrow^\dagger - f_\downarrow). \quad (4.2)$$

This makes possible to define an effective coupling between the Majorana Mode and the dot by attaching  $\gamma_1$  with the spin- $\downarrow$  channel in the QD

$$H_{M-QD} = t_1 (d_\downarrow^\dagger \gamma_1 + \gamma_1 d_\downarrow). \quad (4.3)$$

In addition, the two Majoranas at the edges of the chain can have an internal coupling

$$H_M = \epsilon_M \gamma_1 \gamma_2. \quad (4.4)$$

According to Kitaev's initial paper [15], this coupling term decreases exponentially with the length of the chain. This explains why this term is not taken into account in many computations.

Replacing the Dirac fermion operators we obtain the following Hamiltonians

$$H_M = \epsilon_m f_\downarrow^\dagger f_\downarrow, \quad (4.5)$$

$$H_{M-QD} = \frac{t_1}{\sqrt{2}} d_{1\downarrow}^\dagger f_\downarrow + \frac{t_1^*}{\sqrt{2}} f_\downarrow^\dagger d_{1\downarrow} + \frac{t_1}{\sqrt{2}} d_{1\downarrow}^\dagger f_\downarrow^\dagger + \frac{t_1^*}{\sqrt{2}} f_\downarrow d_{1\downarrow}. \quad (4.6)$$

Finally we have the Hamiltonian

$$H = \sum_{k,\sigma} \left( \epsilon_1 + \frac{U_1}{2} \right) d_{1\sigma}^\dagger d_{1\sigma} + \frac{U}{2} (d_{1\sigma}^\dagger d_{1\sigma} - 1)^2 + t_1 d_{1\downarrow}^\dagger \gamma_1 + V d_{1\sigma}^\dagger c_{k\sigma} + \epsilon_m f_\downarrow^\dagger f_\downarrow + \text{h.c.} \quad (4.7)$$

The fidelity of this effective model has been discussed by Ruiz-Tijerina et al. [3] concluding that this model reproduces the same results than coupling a Kitaev chain model in the topological phase to a QD. (This statement is true even for more realistic models of the TS including Rashba spin-orbit interactions and a Zeeman field [3] ).

## 4.2 Non-interacting QD coupled to Majorana chain

In the non-interacting case we can use the ballistic transport equations from section 3.1. The green functions are then determined by the following set of linear equations.

$$(\omega - \epsilon_M) G_{f_\downarrow, d_{1\downarrow}^\dagger}(\omega) = (\omega + \epsilon_M) G_{f_\downarrow^\dagger, d_{1\downarrow}^\dagger}(\omega) = \frac{t_1^*}{\sqrt{2}} \left( G_{d_{1\downarrow}, d_{1\downarrow}^\dagger}(\omega) - G_{d_{1\downarrow}^\dagger, d_{1\downarrow}^\dagger}(\omega) \right) \quad (4.8)$$

$$(\omega - \epsilon_1) G_{d_{1\downarrow}, d_{1\downarrow}^\dagger}(\omega) = 1 + \frac{t_1}{\sqrt{2}} t_1 G_{f_\downarrow, d_{1\downarrow}^\dagger}(\omega) + \frac{t_1}{\sqrt{2}} t_1 G_{f_\downarrow^\dagger, d_{1\downarrow}^\dagger}(\omega) + V_1 \sum_{\mathbf{k}} G_{c_{\mathbf{k}\downarrow}, d_{1\downarrow}^\dagger}(\omega) \quad (4.9)$$

$$(\omega - \epsilon_k) G_{c_{\mathbf{k}}, d_{1\downarrow}^\dagger}(\omega) = V_1^* G_{d_{1\downarrow}, d_{1\downarrow}^\dagger}(\omega) \quad (4.10)$$

$$(\omega + \epsilon_1) G_{d_{1\downarrow}^\dagger, d_{1\downarrow}^\dagger}(\omega) = -\frac{t_1}{\sqrt{2}} G_{f_\downarrow, d_{1\downarrow}^\dagger}(\omega) - \frac{t_1}{\sqrt{2}} G_{f_\downarrow^\dagger, d_{1\downarrow}^\dagger}(\omega) - V_1^* \sum_{\mathbf{k}} G_{c_{\mathbf{k}\downarrow}^\dagger, d_{1\downarrow}^\dagger}(\omega) \quad (4.11)$$

$$(\omega + \epsilon_k) G_{c_{\mathbf{k}}^\dagger, d_{1\downarrow}^\dagger}(\omega) = -V_1^* G_{d_{1\downarrow}, d_{1\downarrow}^\dagger}(\omega). \quad (4.12)$$


The graph representing these green functions is in Figure 4.2 (a) (Look subsection 3.1.2 for details). However, using that  $(\omega - \epsilon_M) G_{f_\downarrow, d_{1\downarrow}^\dagger}(\omega) = (\omega + \epsilon_M) G_{f_\downarrow^\dagger, d_{1\downarrow}^\dagger}(\omega)$  we can take  $G_{f_\downarrow^\dagger, d_{1\downarrow}^\dagger}(\omega)$  out of the equations. After eliminating this term from (4.9) it becomes

$$(\omega - \epsilon_1) G_{d_{1\downarrow}, d_{1\downarrow}^\dagger}(\omega) = 1 + \frac{t_1}{\sqrt{2}} \left( 1 + \frac{\omega - \epsilon_M}{\omega + \epsilon_M} \right) G_{f_\downarrow, d_{1\downarrow}^\dagger}(\omega) + V_1 \sum_{\mathbf{k}} G_{c_{\mathbf{k}\downarrow}, d_{1\downarrow}^\dagger}(\omega) \quad (4.13)$$

$$= 1 + \frac{\sqrt{2}t_1}{\omega + \epsilon_M} G_{f_\downarrow, d_{1\downarrow}^\dagger}(\omega) + V_1 \sum_{\mathbf{k}} G_{c_{\mathbf{k}\downarrow}, d_{1\downarrow}^\dagger}(\omega). \quad (4.14)$$

Similarly,

$$(\omega + \epsilon_1) G_{d_{1\downarrow}^\dagger, d_{1\downarrow}^\dagger}(\omega) = -\frac{\sqrt{2}t_1}{\omega + \epsilon_M} G_{f_\downarrow, d_{1\downarrow}^\dagger}(\omega) - V_1^* \sum_{\mathbf{k}} G_{c_{\mathbf{k}\downarrow}^\dagger, d_{1\downarrow}^\dagger}(\omega). \quad (4.15)$$


 Figure 4.2: Graph  $\mathcal{G}_M$  representing the transport equations.

With these new equations we can construct the graph in Figure 4.2 (b). Using the Graph-Gauss-Jordan algorithm from subsection 3.1.3 we proceed to eliminate vertexes  $c_k$ ,  $c_k^\dagger$  and  $d_1^\dagger$  in that order. The result is the graph in figure Figure 4.2(c) with

$$\epsilon_{M,d_1^\dagger,c^\dagger} = \epsilon_M + \frac{\omega}{\omega + \epsilon_M} \frac{\|t\|^2}{\omega + \epsilon_M + \sum_{\mathbf{k}} \frac{V_1 V_1^*}{\omega + \epsilon_{\mathbf{k}}}}. \quad (4.16)$$

We finally eliminate  $f_d$  to obtain

$$G_{d_{1\downarrow},d_{1\downarrow}^\dagger}(\omega) = \left[ \omega - \epsilon_1 - \sum_{\mathbf{k}} \frac{V_1 V_1^*}{\omega - \epsilon_1} - \frac{\omega}{\omega + \epsilon_M} \frac{\|t\|^2}{\omega - \epsilon_{M,d_1^\dagger,c^\dagger}} \right]^{-1}. \quad (4.17)$$

This is the Green function we have been looking for. After a few algebraic operations it is possible to show that this result is equivalent to the first computation done by Liu and Baranger in the paper [2].

To compute the DOS we need to replace  $\sum \frac{V_1 V_1^*}{\omega - \epsilon_k} = -i\Gamma_1$  as we already did in subsection 3.1.2. Note that these computations are only for the spin- $\downarrow$  channel. The spin- $\uparrow$  channel is even simpler since this channel is not coupled to the Majorana mode by convention. Hence it corresponds to the case of a single quantum dot coupled to a Lead. The results for the DOS can be observed in Figure 4.3. Each figure has an inset showing the model in the Majorana representation. The small blue and red balls are Majorana fermions just as the ones in figure

## 4.2. Non-interacting QD coupled to Majorana chain

Figure 1.2. The Majorana at the edge of the chain is represented by the isolated red ball connected to the QD (Figure 4.3a). The isolated blue ball in Figure 4.3c represents the Majorana at the other edge of the chain which is connected to the sphere by the parameter  $\varepsilon$ .

- **Figure 4.3.(a),(b):** The spin- $\uparrow$  DOS shows the result of coupling the QD with the lead and without Majoranas. When the parameter  $t$  is increased, the MZM is coupled to the spin- $\downarrow$  which causes the dispersion of the DOS. The most relevant signature is the robust height of 0.5 in the DOS that is observed in the central peak for all  $t > 0$ . This mid-height DOS is responsible for the decay of half a quanta in the conductivity of the QD [2].




Figure 4.3: Density of states for a Majorana coupled to a QD under the tuning of different parameter. The tuning parameter is drawn in purple line in the inset model.

- **Figure 4.3.(c),(d):** This time a gate voltage is induced in the dot which breaks PHS. However the robust 0.5-height Majorana signature prevails in the dot even at very high gate voltages where the dot is expected to be empty. This result is similar to the one in [38], where the is attached to the last cite of the Kitaev chain.




Figure 4.4: DOS at  $t_1 = \Gamma_1$  at the PHS-point. Insets: Left: QD-Majorana model. Right: Zoom to low energy DOS..

- **Figure 4.3.(e),(f):** The term  $\varepsilon_M$  couples both Majoranas at the edges of the chain. The strength of this parameter decays exponentially with the length of the Majorana chain so that it is often neglected. Here we observe the consequences of including this parameter in the model. The spin- $\downarrow$  DOS for energies  $\omega < \varepsilon_M$  is forced to take the same value of the the spin- $\uparrow$  DOS. This clearly destroys the Majorana zero mode in the dot.

### 4.3 Kondo-Majorana physics

In interacting quantum dots the Kondo effect is visible at low temperatures even when the QD is attached to a Majorana chain, which allows the study Kondo-Majorana physics. To observe this , we used the NRG code with a fixed Coulomb repulsion of  $U = 17.6\Gamma_1$ , just as in section subsection 3.2.7. Then, particle-hole equilibrium is achieved when  $(\varepsilon_1 + \frac{U_1}{2})\hat{n}_1\sigma$ . Any tunning of the dot's gate voltage must be understood as a displacement  $\Delta\varepsilon_1$  from this equilibrium point.

Figure 4.4 shows the PHS case for a Majorana coupling  $t_1 = \Gamma_1$ . The two small wide peaks at the borders of the plot are the coulomb states. In the right inset of the figure, we observe the low-temperature regime inside the gap. There, two zero modes can be appreciated. While the spin- $\uparrow$  DOS is the same Kondo peak from Figure 3.8a, the spin- $\downarrow$  DOS reveals a Majorana zero mode of half the amplitude of the Kondo peak  $(\frac{0.5}{\pi\Gamma_1})$ . This Majorana signature resembles the one in Figure 4.3a.




Figure 4.5: (a)&(b): Dependence of the DOS over the gate voltage  $\Delta\epsilon_1$  at  $t_1 = \Gamma_1$ . (a) Spin- $\uparrow$  (b) Spin- $\downarrow$ . (c) DOS at the red-dashed horizontal cut in (a)&(b). Insets: Left: QD-Majorana model. Right: Low energy DOS. (d) DOS at the Green-dashed vertical cut in (a)&(b). .




Figure 4.6: Dependence of the zero-bias conductance over tuning voltage Adapted from [3].

It is possible to separate Kondo and Majorana physics by inducing a gate voltage in the dot. As observed in Figure 4.5(a), the gate voltage detunes the Kondo peak from the Fermi energy. Instead, the MZM in Figure 4.5(b) remains at the same position. At  $\Delta\epsilon_1 = 5\Gamma_1$  we can already observe a decaying Kondo peak next to the robust Majorana signature of height  $\frac{0.5}{\pi\Gamma}$  (Figure 4.5(c)). This is more clear in Figure 4.5(d) where the spin- $\uparrow$  DOS decays with  $\Delta\epsilon_1$

while the spin- $\downarrow$  DOS is stable, even at  $\Delta\epsilon_1 \sim \frac{U}{2} = 8.6$  where the dot is supposed to be empty.

This interesting result was already pointed out by Ruiz-Tijerina et al. who proved that increasing the gate voltage would produce a visible decay in the zero bias conductance down to  $\frac{0.5e^2}{h}$  (See Figure 4.6). Hence, allowing to measure the Majorana signature without the superposition with the Kondo peak. This result is clear from Figure 4.5. At  $\Delta\epsilon = 0$  the DOS at the Fermi energy is  $\frac{1}{\pi\Gamma_1}$  for spin- $\uparrow$  and  $\frac{0.5}{\pi\Gamma_1}$  for spin- $\downarrow$ . Instead, for bigger values of  $\Delta\epsilon_1$  only the  $\frac{0.5}{\pi\Gamma_1}$ -height spin- $\downarrow$  peak appears. Since the zero bias conductance at zero temperature is essentially the sum of both spectral densities (times unit correction), Figure 4.5 recovers the results in Figure 4.6.

Another possibility to distinguish Kondo and Majorana physics is quenching the Kondo effect with a strong magnetic field. Similar to what was observed in Figure 4.6, the Kondo peak will be destroyed while the Majorana signature remains stable [3].

In the next chapter we will use the ideas from this chapter to study the model of a DQD attached to a Majorana zero mode.

## Chapter 5

# Coupling the Majorana Zero Mode to a Double Quantum Dot




Figure 5.1: Model for the DQD-Majorana system. Solid lines: Hopping interactions ( $t_{dots}$ : inter-dot coupling ,  $V_1, V_2$  couplings of QD1 and QD2 with the lead. ). Dashed lines: Majorana spin- $\downarrow$  effective couplings (4.3)  $t_1, t_2$ . The atomic energy levels appear inside each QD  $\epsilon_1, \epsilon_2$  are tuned by the gate voltages. The coulomb interaction is represented by  $U_1, U_2$ . The red dashed horizontal lines represent the Fermi level.

The DQD-Majorana model is the most fundamental structure where Majorana manipulation is possible. Tunneling Majorana modes in this device have already inspired a few theoretical studies [43, 44] and experimental setups confirming the observations of Andreev molecules [42]. However, there is still no complete analysis of the transitions of the Majorana signatures between the QDs in this model, even though quantum tunneling of a MZM into a double dot offers several possibilities for MZM manipulation.

In this chapter, we will explore different possibilities for Majorana manipulation in a device consisting of a DQD coupled to a MZM and a metallic lead (See Figure 5.1). The simplicity of this model allows us to explore analytically different geometries of QD's from linear and symmetric couplings to T-junctions (Fig. Figure 5.3). As in the previous models , we will consider both non-interacting and interacting regimes.

The model in Figure 5.1 can be described from the combination of the Hamiltonians of a

### 5.1. Applying our methods to the DQD-Majorana system

---

QD-Majorana system (4.7) and a DQD (3.18). Integrating these models we obtain

$$H = \sum_{i=1}^2 \sum_{k,\sigma} \left( \varepsilon_i + \frac{U_i}{2} \right) d_{i\sigma}^\dagger d_{i\sigma} + \frac{U_i}{2} (d_{i\sigma}^\dagger d_{i\sigma} - 1)^2 + t_i \gamma d_{i\downarrow} + V_i d_{i\sigma}^\dagger c_{k\sigma} + t_{dots} d_{1\sigma}^\dagger d_{2\sigma} + \text{h.c.} \quad (5.1)$$

Where  $V_1, V_2$  is the coupling of dots 1,2 to the lead.  $t_1, t_2$  define the Majorana couplings with each dot.  $t_{dots}$  is the inter-dot coupling.  $\varepsilon_1, \varepsilon_2$  are the energy levels of the dot, which are tuned by the the gate voltage and  $U_1, U_2$  are the coulomb repulsion parameters.

## 5.1 Applying our methods to the DQD-Majorana system

### 5.1.1 Non-interacting Green function:

This new model is a combination between the DQD (Figure 3.1) and the Majorana-QD Figure 4.2(b). We can use the trick in section 4.2 to get rid of the Green function  $G_{f_{\downarrow}, d_{1\downarrow}^\dagger}(\omega)$  for the second Majorana operator. This allows us to obtain the following transport equations

$$\begin{bmatrix} \omega - \varepsilon_1 & -V_1^* & -t_{dots} & -T_1 & 0 & 0 & 0 \\ -V_1 & \omega - \varepsilon_k & -V_2 & 0 & 0 & 0 & 0 \\ -t_{dots}^* & -V_2^* & \omega - \varepsilon_2 & -T_2 & 0 & 0 & 0 \\ -T_1^* & 0 & -T_2^* & \omega - \varepsilon_M & T_2^* & 0 & -T_1 \\ 0 & 0 & 0 & T_2 & \omega + \varepsilon_2 & V_2^* & t_{dots}^* \\ 0 & 0 & 0 & 0 & V_2 & \omega + \varepsilon_k & V_1 \\ 0 & 0 & 0 & T_1 & t_{dots} & V_1^* & \omega + \varepsilon_1 \end{bmatrix} \begin{bmatrix} G_{d_{1\downarrow}, d_{1\downarrow}^\dagger}(\omega) \\ G_{c_{k\downarrow}, d_{1\downarrow}^\dagger}(\omega) \\ G_{d_{2\downarrow}, d_{1\downarrow}^\dagger}(\omega) \\ G_{f_{\downarrow}, d_{1\downarrow}^\dagger}(\omega) \\ G_{d_{2\downarrow}^\dagger, d_{1\downarrow}^\dagger}(\omega) \\ G_{c_{k\downarrow}^\dagger, d_{1\downarrow}^\dagger}(\omega) \\ G_{d_{1\downarrow}^\dagger, d_{1\downarrow}^\dagger}(\omega) \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 1 \end{bmatrix}, \quad (5.2)$$

where  $T_i = \frac{t_i}{\sqrt{\omega + \varepsilon_M}}$ .

The graph representing this equation is in Figure 5.2(a). Using the algorithm in subsection 3.1.3 we start eliminating vertexes  $c_k, c_k^\dagger, d_{2,\downarrow}$  and  $d_{2,\downarrow}^\dagger$  in that order. The self-energies associated to  $d_{1,\downarrow}$  and  $d_{1,\downarrow}^\dagger$  will be similar to the energy of the DQD (3.25) giving

$$\varepsilon_{DQD}^\pm = \pm \varepsilon_1 + \sum_{\mathbf{k}} \frac{V_1 V_1^*}{\omega - \varepsilon_{\mathbf{k}}} + \frac{\left| \pm t_{dots} + \sum_{\mathbf{k}} \frac{V_1 V_2^*}{\omega - \varepsilon_{\mathbf{k}}} \right|^2}{\omega \mp \varepsilon_2 - \sum_{\mathbf{k}} \frac{V_2 V_2^*}{\omega - \varepsilon_{\mathbf{k}}}}. \quad (5.3)$$

There is also a correction in the couplings between the Majorana mode and  $d_{1,\downarrow}, d_{1,\downarrow}^\dagger$  given by

$$T_\pm = \pm t_1 \pm t_2 \frac{\left( \pm t_{dots} + \sum_{\mathbf{k}} \frac{V_1 V_2^*}{\omega - \varepsilon_{\mathbf{k}}} \right)}{\omega \mp \varepsilon_2 - \sum_{\mathbf{k}} \frac{V_2 V_2^*}{\omega - \varepsilon_{\mathbf{k}}}}. \quad (5.4)$$




Figure 5.2: Graph method applied to a DQD coupled to a Majorana zero mode. a) Initial stage. b) Eliminated vertexes  $c_k^+$ ,  $c_k$ ,  $d_{2,\downarrow}$ ,  $d_{2,\dagger\downarrow}$  in that order. c) Eliminated vertexes  $d_{1,\dagger\downarrow}$  and  $f_\downarrow$ , the final energy is  $\omega - G_{d_1,d_1^\dagger}(\omega)$ .

In addition, since the Majorana is in contact with dot 2, there is an extra-term appearing in the Majorana self-energy given by

$$\epsilon_{M2} = \omega - \epsilon_M - \frac{\frac{\omega}{\omega + \epsilon_M} \|t_2\|^2}{\omega - \epsilon_2 - \sum_{\mathbf{k}} \frac{V_2 V_2^*}{\omega - \epsilon_{\mathbf{k}}}} - \frac{\frac{\omega}{\omega + \epsilon_M} \|t_2\|^2}{\omega + \epsilon_2 - \sum_{\mathbf{k}} \frac{V_2 V_2^*}{\omega + \epsilon_{\mathbf{k}}}}. \quad (5.5)$$

It only remains to eliminate out vertexes  $d_1^\dagger$  and  $f_\downarrow$  to obtain the green function

$$G_{d_{1\downarrow},d_{1\downarrow}^\dagger}(\omega) = \frac{1}{\omega - \epsilon_{DQD}^+ - \frac{\|T_+\|^2}{\omega - \epsilon_{M2} - \frac{\|T_-\|^2}{\epsilon_{DQD}^-}}}. \quad (5.6)$$

This simple formula summarizes the transport information through the first dot of the non-interacting Majorana-DQD system. To compute the DOS we just need to replace  $\sum \frac{V_i V_i^*}{\omega - \epsilon_{\mathbf{k}}} = -i\Gamma_i$  as performed in subsection 3.1.2. By plotting the final DOS in Mathematica we were able to observe the transitions of the Majorana mode under manipulation of the model parameters.

### 5.1.2 NRG for the interacting system

The Numerical Renormalization Group (NRG) technique described in section 3.2 is the most successful methods to study interacting quantum impurity models. In this model, the impurity is described by the DQD attached to the MZM. In our code, we set a Coulomb repulsion factor of  $U = 17.3\Gamma_1$  in both dots and a cut-off energy of  $D = 2U = 34.6\Gamma_1$ . The spacing with other energy levels is assumed to be higher than  $D$ , such that only the two coulomb states are relevant for the system dynamics. When  $\varepsilon_i = \frac{U}{2}$  in both dots, the system is in the Particle-Hole-Symmetric region. At this point, each dot has an odd number of electrons, hence, at sufficiently low temperature the system will exhibit characteristic Kondo peaks at the Fermi energy [50]. The coexistence of Kondo and Majorana zero modes is still a point of contention in the area that has never been studied in DQDs and one of the objectives of this part of the project.

To improve the efficiency of the code we used the symmetries of the system to maintain a block structure during NRG's iterative diagonalization process. This model preserves the spin- $\uparrow$  particle number  $\hat{N}_\uparrow$  and the spin- $\downarrow$  parity  $\hat{P}_\downarrow = \pm (+\text{even}, -\text{odd})$ . The spin- $\downarrow$  particle number is not preserved due to superconducting-type Majorana coupling (4.3). The initial Hamiltonian is organized in blocks according to these symmetries. This block structure is preserved during the entire iteration process [13]. To compute the spectral functions, we use the density matrix renormalization group (DM-NRG) described in subsection 3.2.5 in combination with the Z-trick method [59], which improves spectral resolution at high energies.

To initialize the model in Figure 3.7 we set  $H_{-1}$  equal to

$$H_{-1} = \sum_{i=1}^2 \sum_{k,\sigma} \left( \varepsilon_i + \frac{U_i}{2} \right) d_{i\sigma}^\dagger d_{i\sigma} + \frac{U_i}{2} (d_{i\sigma}^\dagger d_{i\sigma} - 1)^2 + t_i (\gamma d_{i,\downarrow} + d_{i,\downarrow}^\dagger \gamma) + t_{dots} (d_{1\sigma}^\dagger d_{2\sigma} + d_{2\sigma}^\dagger d_{1\sigma}), \quad (5.7)$$

and wrote the Hamiltonian in the symmetry-block diagonal representation (see section A.2). We also include manually the Hamiltonian  $H_0$  into the code (See Figure 3.7) to guarantee that both quantum dots are coupled to the first site of the chain. After this, the code follows the standard NRG algorithm and prints the density matrices to initialize DM-NRG. The final result is the spectral density which contains sufficient physical information to study the MZM-DQD model. In the following section, we show how the density of states can be used to simulate the Manipulation process of an MZM inside the DQD.

## 5.2 Manipulation of Majorana zero modes

The density of states provides significant information about the presence of a Majorana zero modes in the dot. We characterize the Majorana signature by a robust zero-mode with two possible heights:

- **Type I:** The spin- $\downarrow$  DOS is the half of the spin- $\uparrow$  DOS at the Fermi energy ( $\rho_\downarrow(0) = \rho_\uparrow(0)$ ).




Figure 5.3: .

- **Type II:** A spin- $\downarrow$  zero mode of height  $\rho_{\downarrow}(0) = \frac{0.5}{\pi\Gamma_1}$ .

In our results we observe several times these two types of signatures. Type I often appears when there is a zero-mode in the spin- $\uparrow$  DOS, which is caused by the Kondo effect in the interacting case. Type II emerges in the remaining situations.

We call MZM manipulation to the "movements" attributed to the Majorana signature under the tunning of the dot gate voltages ( $\epsilon_1, \epsilon_2$ ). This manipulation process is performed in three different set ups that are presented in Figure 5.3 with definite values of  $\Gamma_2, t_{dots}, t_1$  and  $t_2$ . In configuration (a), we coupled the QD symmetrically to the lead and the Majorana mode. With this setup we expect to break the localization of the MZM which should split and tunnel into both dots. In setups (b) and (c) we coupled the second dot indirectly through the first dot. Hence, quantum interference should split the zero mode in two states. Our objective is to observe what occurs with the Majorana signature in this situation. There are two options to connect the MZM in this situation. Attached it directly through the first dot (b) or indirectly through the second dot (c). Both alternatives are geometrically distinct since (b) suggests a T-junction coupling while (c) reflects a connection in series of both QD's between the lead and the MZM.




Figure 5.4: Non-interacting DOS in the symmetric coupling(Figure 5.3(a)) at each QD. First column: Dot 1. Second column: Dot 2. The gate voltages vary at each row. First row: Zero-bias in both dots  $\varepsilon_1 = \varepsilon_2 = 0$ . Second row:  $\varepsilon_1 = 5\Gamma_1$ ,  $\varepsilon_2 = 0$ . Third row:  $\varepsilon_1 = 0$ ,  $\varepsilon_2 = -5\Gamma_1$ . Bold blue lines: Spin- $\uparrow$  DOS. Thin red lines: Spin- $\downarrow$  DOS. The insets at the right show which dot carries a Majorana signature, represented by a red dashed circle. Upper: First dot. Lower: Second dot.

### 5.2.1 Non-interacting manipulation

The non-interacting results for setups (a),(b) and (c) of Figure 5.3 are shown at figures Figure 5.4, Figure 5.5 and Figure 5.6 respectively. Each figure depicts the DOS of dot 1(left) and dot 2(right). The gate voltage is initially 0 in both dots at the first row. In the second row, the gate voltage is turned on to  $\varepsilon_1 = 5\Gamma_1$ , while the second dot remains at  $\varepsilon_2 = 0$ . In the third row the first dot's voltage is off  $\varepsilon_1 = 0$  and we switch on the second dot with a negative voltage of  $\varepsilon_2 = -5\Gamma_1$ . The inset figures at the right side of each row show which dots exhibit Majorana signatures, depicted by a red dashed circle inside the dot. These images will continuously change under the tuning of gate voltages which represents the manipulation of the Majorana signature.

In Figure 5.4 we observe the results for the symmetric coupling setup Figure 5.3(a). In the particle hole symmetric case (first row) the DOS is equal in both dots. Note that the spin- $\downarrow$  (Thin red line) DOS is the half of the spin- $\uparrow$  (Bold blue line) DOS at the Fermi energy ( $\rho_{\downarrow}(0) = \frac{1}{2}\rho_{\uparrow}(0)$ ). This type II Majorana signature is similar to the one observed when a single dot is coupled to a Majorana mode. [2] We may conclude that the Majorana in tunneling inside both




Figure 5.5: Non-interacting DOS of the T-dot coupling Figure 5.3(b). (b). First line (a),(b):  $\varepsilon_1 = \varepsilon_2 = 0$ . Second line (c),(d):  $\varepsilon_1 = 5\Gamma_1$ ,  $\varepsilon_2 = 0$ . Third line (e),(f):  $\varepsilon_2 = -5\Gamma_1$ ,  $\varepsilon_1 = 0$ . Blue bold lines: Spin- $\uparrow$  DOS. Red thin lines: Spin- $\downarrow$  DOS. The inset at the upper-right corner of each line indicates which dots exhibit Majorana signature, which is represented by a red dashed circle inside the dot.

dots breaking the localization of the MZM. If a positive or negative gate voltage is induced in one of the dots, as shown in the second and third row of Figure 5.4(c)-(f), the Majorana zero mode vanishes from that dot. Meanwhile the density of states in the other dot increases while preserving the Majorana signature. This means that the MZM is actually being induced to "leave" this dot and leak into the other by the biased voltage. This is the first example of MZM manipulation.

Another example can occur when the second dot is not directly connected to the lead. In this case, the inter-dot tunneling generates quantum interference which finally destroys the central peak as observed in Figure 5.5(a) at the spin- $\uparrow$  DOS. The spin- $\downarrow$  channel at Figure 5.5(a), which is coupled to the MZM, does not exhibit the characteristic Fermi peak either. Instead, the one half Majorana signature at the Fermi energy ( $\rho_{\downarrow}(0) = \frac{1}{2}\rho_{\uparrow}(0)$ ) appears clearly inside the second dot Figure 5.5(b). This situation prevails when the first dot's gate voltage is turned on Figure 5.5(c)&(d). While the first dot does not seem to exhibit any type of Majorana signature, the second dot's spin- $\downarrow$  DOS exhibits a robust zero-mode of height  $\frac{0.5}{\pi\Gamma}$ . The results are more exciting when the second dot's gate voltage is turned on in Figure 5.5(e)&(f). These figures clearly show how the MZM, previously localized at the second dot, is induced to leave this dot and to return into the first dot. Moreover, the DOS of spin- $\uparrow$  and spin- $\downarrow$  channels are very




Figure 5.6: Non-interacting DOS of the set up in Figure 5.3(c). First line (a),(b):  $\varepsilon_1 = \varepsilon_2 = 0$ . Second line (c),(d):  $\varepsilon_1 = 5\Gamma_1$ ,  $\varepsilon_2 = 0$ . Third line (e),(f):  $\varepsilon_2 = -5\Gamma_1$ ,  $\varepsilon_1 = 0$ . Blue bold lines: Spin- $\uparrow$  DOS. Red thin lines: Spin- $\downarrow$  DOS. The inset at the upper-right corner of each line indicates which dots exhibit Majorana signature, which is represented by a red dashed circle inside the dot.

similar to the spectral densities observed at Figure 5.4(d)(e), which means that the previous interference pattern has disappeared due to this gate voltage.

The results of the third configuration Figure 5.3(c) appear in Figure 5.6. Contrary to what was observed in the previous case, this time the Majorana signature is not destroyed by the interference but instead, the  $\frac{0.5}{\pi\Gamma}$ -height MZM emerges indirectly in the first dot. This is a perfect way to separate the Majorana's spin- $\downarrow$  DOS from the central spin- $\uparrow$  zero-mode which is still destroyed by the interference. In addition, the second dot still exhibits a type I Majorana signature as observed in Figure 5.6(b). In the second row we observe that turning on the gate voltage in dot 1 destroys the Majorana signature in both dots Figure 5.6(c)(d). On the other hand, if the second dot's voltage is switched both dots will preserve their Majorana signature (QD1:type I, QD2: type II), while the spin- $\uparrow$  quantum interference vanishes in the first dot.




Figure 5.7: Density of states of both dots in the symmetric coupling without gate voltages between the Majorana and the interacting DQD. Bold blue lines: Spin- $\uparrow$  DOS. Thin red lines: Spin- $\downarrow$  DOS. Inset: Low-energy DOS.

### 5.2.2 Interacting manipulation

Now we consider a Coulomb repulsion energy of  $U = 17\Gamma_1$  in both dots. The factor  $\frac{U_i}{2}(\sum_{\sigma} \hat{n}_{i\sigma} - 1)^2$  in (5.1) favors states with an odd number of electrons (and holes). In addition, particle-hole equilibrium is now achieved when  $\Delta\varepsilon_i := (\varepsilon_i + \frac{U_i}{2})$ . Any induced gate voltage must be considered as a shifting  $\Delta\varepsilon_i$  from this equilibrium point. Figure 5.7 shows the DOS of both QDs for the symmetric coupling configuration Figure 5.3. The two peaks appearing at around  $8.6\Gamma_1 = \frac{U_i}{2}$  represent the two energy levels spaced by the Coulomb repulsion factor  $U$ . The central spin- $\uparrow$  peak is a consequence of the Kondo effect, [50, 53] while the two satellite peaks observed in the inset are the result of the RKKY indirect interaction between both dots. [46–48] Moreover, the system presents a Majorana signature characterized by half spin- $\downarrow$  DOS at the Fermi energy ( $\rho_{\downarrow}(0) = \frac{1}{2}\rho_{\uparrow}(0)$ ). Note, that in this case the Majorana signature coexists with the Kondo effect in the DQD as already predicted by Ruiz-Tijerina *et al.* for a single dot. [3]

## 5.2. Manipulation of Majorana zero modes




Figure 5.8: (a)-(d) Dependence of the density of states of setup in Figure 5.3(a) over  $\omega$  and the gate voltage  $\Delta\epsilon_2$ .  $\Delta\epsilon_1 = 0$  Up: Dot 1. Down: Dot 2. Left: Spin- $\uparrow$ . Right: Spin- $\downarrow$ . (e) Evolution of the relation  $\frac{\rho_{1\uparrow}(0)}{\rho_{1\uparrow}(0)}$  for both QDs. While QD2 losses rapidly the Majorana signature, QD1 maintains it till  $\Delta\epsilon_2 \sim 5$ .

In this part of the project we are interested in the physics at low energy scales  $\omega \sim \Gamma_1$  close to the Kondo and MZM temperature <sup>7</sup>. At this scale we can observe similar Majorana signatures compared with the non-interacting case. For instance, the inset Figure 5.7 shows the NRG results for the symmetric setup in Figure 5.3(a). In agreement with the non-interacting results, both dots have type I Majorana signatures.

When the gate voltage is turned on, we observe the number of holes ( $\omega > 0$ ) increasing in the spin- $\uparrow$ , while the particles ( $\omega < 0$ ) increase in the spin- $\downarrow$  (See Figure 5.8(a)-(d)). At  $\Delta\epsilon_2 \approx 6\Gamma_1$ , the Coulomb peak overlaps with the Fermi energy in the spin- $\downarrow$  channel, destroying the Majorana zero mode. This event is visible in Figure 5.8(e), where the type one Majorana signature is clearly destroyed in Dot 1 after  $\Delta\epsilon_2 > 6\Gamma_1$ . For  $\Delta\epsilon_2 < 6\Gamma_1$  we observe an stable Ma-

<sup>7</sup>The MZM temperature  $T_M$  can be defined by the energy scale  $k_B T_M$  where the Majorana signature is visible. This effect has been plentifully discussed in a previous paper , together with its relationship with the Kondo temperature [3]. In the regime we are presenting in this thesis, the Majorana and Kondo temperature are similar, hence we are at an energy range where we can observe both effects co-existing. We will present further discussion about this idea in our DQD-Majorana model in a future paper.




Figure 5.9: The same as in Figure 5.4 for the interacting DOS in the symmetric coupling (Figure 5.3).

Majorana signature in dot 1 with  $\rho_{\downarrow}(0) \approx \frac{1}{2}\rho_{\uparrow}(0)$ ). In the second dot the zero modes are absorbed by the dot, which slowly destroys the Majorana signature.

The red cuts in Figure 5.8 are plotted in Figure 5.9 where we depict the results of Majorana manipulation. In agreement with the non-interacting case, the particle hole symmetric model receives both Majorana modes. Whenever a gate voltage is switched on, the Majorana is forced to tunnel into the other dot. We observe similar results for positive and negative gate voltages.

In the T-dot coupling Figure 5.3(b), the spin- $\uparrow$  Kondo peak in Figure 5.10 is destroyed by interference just as in the non-interacting case. This phenomenon had already been predicted for a T-junction of a double quantum dot attached to metallic leads [1]. The insight of our model is that an attached MZM should also disappear due to the same interference. Furthermore, a type I Majorana signature can be observed at very low energies in the inset of Figure 5.10(b). However we have to recognize that both zero-modes decay significantly in the second dot. When the first voltage is turned on, the Majorana mode jumps onto the first dot which presents a type I Majorana signature. This is a clear difference with the non-interacting results where the Majorana signature stayed in the second dot. If the second dot is switched on, a type II Majorana signature appears at very low energies in dot 1, which is coherent with the idea that the Majorana interference should disappear in this case. In Figure 5.10(e) we identify emergence of a Fano resonance at the Fermi energy causing the sharp-asymmetric peak at  $\omega = 0$ . Fano resonances have already been documented in similar models in [61]. We are going to talk more

## 5.2. Manipulation of Majorana zero modes




Figure 5.10: The same as in Figure 5.4 for the interacting DOS of the T-dot junction Figure 5.3(b). Inset in b): Zoom to low-energy DOS.

about this resonance in following section.




Figure 5.11: Logarithmic dependence of the density of states for interacting dots coupled in series Figure 5.3(c) over  $\omega$  and the gate voltage  $\Delta\epsilon_1$ .  $\Delta\epsilon_2 = 0$ . The dependence over  $\Delta\epsilon_2$  produces similar results.

Finally, Figure 5.12 shows the NRG results for the last configuration, where the dots are coupled in series Figure 5.3(c). Notably, the indirectly-attached MZM exhibits a robust type




Figure 5.12: The same as in Figure 5.4 for the interacting DOS for interacting dots coupled in series (Figure 5.3(c)). Inset in b): Zoom to low-energy DOS.

II Majorana signature in the first dot over a destroyed Kondo peak. This is observed clearly in Figure 5.11 where  $\rho_{1\downarrow}$  exhibits a constant  $\frac{0.5}{\pi\Gamma_1}$ -height Majorana peak . This signature is stable under the gate voltage tuning in dot 1 and similar results are obtained in dot 2 . In addition, only in the particle hole symmetric case the second dot presents a type II Majorana signature (Inset Figure 5.12(b)).

We could understand this effect by thinking that the dots in model (c) are attached in series. Therefore both QDs can be thought as extensions of the Kitaev chain, were the first dot is the last place in the wire. Hence the Majorana should be localized at this dot despite the application of gate voltages. This situation is similar to the case of a single dot attached to a Majorana chain, where it is known that the MZM appears in the dot even when this is supposed to be empty [38]. It still remains the doubt about why this effect is not observed in the non-interacting case . On the other hand, there is a Fano resonance at the Fermi energy in the spin- $\downarrow$  DOS Figure 5.12(d)(e) . This zero mode was not identified as a potential Majorana signature since it varies with the values of  $\Delta\epsilon_1$  (Figure 5.11) and  $\Delta\epsilon_2$ .

We are now writing a paper summarizing these results. As we observed, we were able to characterize the transitions of the Majorana signature in different geometric arrangements of the dots. In the following section we will present some ideas that are still in development. We hope they could lead us to future publications




Figure 5.13: Dependence of the DOS in the symmetric model Figure 5.3(a) over  $t_1 = t_2$  and  $\omega$ . Up: High energy . Down: Zoom to low energy states.

## 5.3 Additional results

This section contains additional results which we are considering to study in future publications.

### 5.3.1 Indirect exchange through the Majorana mode

In Figure 5.7 we observed the emergence of satellite peaks at low energies product of anti-ferromagnetic exchange interactions. This exchange interaction can occur through the lead and through the Majorana mode. The reason why we are observing just two satellites is because the Majorana couplings ( $t_1 = t_2$ ) and the broadening parameters ( $\Gamma_1 = \Gamma_2$ ) are about the same order. Hence the peaks are a superposition of both exchange interactions.

We can separate both exchange interactions by observing the dependence of the DOS at different orders of  $t_1 = t_2$  in Figure 5.13. We can distinguish two regimes:

1. Low Majorana coupling  $t_1 = t_2 < 0.5$ : Two additional satellite peaks appear in the spin- $\downarrow$  DOS (See inset in Figure 5.14 for better appreciation). These peaks are similar to the Kondo satellites that appear at high energies. However, since they appear only at low-energies, it is clear that they are produced by the MZM. We conclude that these two




Figure 5.14: Dependence of the DOS in the symmetric model Figure 5.3(a) over  $t_1 = t_2$  and  $\omega$ . Up: High energy . Down: Zoom to low energy states.

satellites are produced by the indirect exchange through the attached Majorana quasi-particle.

2. High Majorana coupling  $t_1 = t_2 > 0.5$ : When the Majorana coupling is high enough, the indirect exchange through the MZM occurs in the same energy scale as the Kondo satellites. Notably, the spin- $\downarrow$  satellite peaks in the high energy regime are not affected by this effect. Instead, a visible inflation of the satellites in the spin- $\uparrow$  DOS is observed. Therefore the MZM is actually correlated with the spin- $\uparrow$  DOS through the satellite peaks. This is unexpected since the Majorana is only coupled to the spin- $\downarrow$  channel. The only explanation for this is that these satellites are formed by a strongly correlated state between the MZM, the dot states and the lead.

### 5.3.2 Indirect Majorana coupling through the lead

Imagine a model where we connect both dots symmetrically to the leads but we only connect the MZM to the first dot. In addition, we do not allow inter-dot tunneling. Then the only connection between the MZM and the second dot would be passing through the first dot and the lead. We didn't expect to see any Majorana signature in the second dot in these conditions. However, Figure 5.15 shows a clear type I Majorana signature in the second dot.

Note also that the density of states in the second dot is very small in comparison with the first dot. This is intriguing since this dot is directly connected with the lead and it is still at the Kondo regime. This ambiguity means that the zero-bias DOS is favored by a direct coupling to an MZM.




Figure 5.15: DOS at both dots for the model in the left inset. The right inset zooms the low-energy DOS in the second dot.




Figure 5.16: Logarithmic dependence of the DOS in the second dot for setup in Figure 5.3(b) over the second gate voltage. Cut at  $5\Gamma_1$  corresponds to Figure 5.10(f).

### 5.3.3 Critical behavior in zero-bias DOS

In Figure 5.10(f) we observe a sharp peak at DOS. This peak is actually a big problem for our results since they are supported on the spectral densities at the Fermi energy. In Figure 5.16 we observe a critical behavior in the zero-bias DOS close to  $\Delta\epsilon_2 = 0$ . This is quite intriguing .

# Chapter 6

## Conclusions

In this work, we developed two methods to study the system of a double quantum dot attached to a Majorana chain. We probed successfully these methods in simpler cases such as the double quantum dot (subsection 3.1.4 , subsection 3.2.7) and the QD-Majorana system (chapter 4). The results obtained in these sections are in agreement with previous papers regarding the Kondo interference in double quantum dots [1], MZM leaking into quantum dots [2] and the co-existence of Kondo-Majorana physics [3].

Moreover, we introduced the Graph-Gauss-Jordan algorithm (subsection 3.1.2) as a simple, didactic, analytical and graphical method to solve the equations of motion. This method allowed us to obtain exact expressions for the non-interacting Green functions at several stages of the project. In particular, in the double quantum dot - Majorana model it proved to be extremely useful to simplify the complexity of the solution given by a fractional polynomial of 9<sup>th</sup>-degree in up to 7 variables. Furthermore, with this expression we were able to predict interesting parameters for simulation in the interacting regime. We hope for its extended use in condensed matter physics.

In chapter 5. we used the methods from chapter 3 to study the transitions of the Majorana zero modes inside a double quantum dot. Comparing the exact analytical solution in the non-interacting system and the NRG results for interacting quantum dots, we were able to characterized the displacements of the MZM inside the double quantum dot for the three setups in Figure 5.3, representing a symmetric coupling, a T-dot junction and a linear coupling of the dots. All of these manipulations are summarized in Figure 6.1 . We observe a considerable agreement regarding the location of the Majorana signature between the interacting and non-interacting results with minor differences:

- Symmetric coupling (Figure 6.1 C.I) : The MZM leaks inside both dots. For interacting dots, the Majorana signature will emerge near the Kondo temperature. At this regime the system presents combined Kondo-Majorana physics . Additional satellite peaks produced by the indirect exchange through the lead and the MZM appear at low energies. If the gate voltage of one dot is turned on, the MZM is induced to tunnel only into the other dot, which is the key to MZM manipulation.
- T-dot coupling (Figure 6.1 C.II): The the spin- $\uparrow$  zero mode at QD1 (The Kondo peak if the system is interacting) is destroyed by quantum interference with the second dot. This interference will also destroy the MZM in the first dot but a type I Majorana signature will still appear in the second dot. The Majorana mode can be induced to tunnel back




Figure 6.1: Table of Majorana signatures in the studied cases in interacting dots.

into the first dot if a gate voltage is applied on the second dot. This signature is visible at very low energies (below  $0.1\Gamma_1$ ) in the interacting case.

- In-series coupling (Figure 6.1 C.III) : An indirect type II Majorana signature is observed in the first dot. This signature is robust, specially in the interacting case, where it is present in all configurations, despite gate voltage tuning. Fano resonances emerge in the second dot. One of these signatures seems to have critical behavior.

Besides MZM manipulation we also pointed out other cases that could lead to future projects, such as the separation of the exchange interactions produced through the lead and through the MZM, the emergence of an indirect Majorana signature passing through the leads and the critical behavior in the T-dot junction when a gate voltage is applied on the second dot.

# Bibliography

- [1] Luis G. G. V. Dias da Silva, Nancy Sandler, Kevin Ingersent, and Sergio E. Ulloa. Transmission in double quantum dots in the Kondo regime: Quantum-critical transitions and interference effects. *Physica E: Low-dimensional Systems and Nanostructures*, 40(5):1002–1005, March 2008. ISSN 1386-9477. doi: 10.1016/j.physe.2007.08.098. URL <http://www.sciencedirect.com/science/article/pii/S1386947707003074>. (document), 1.3, 3.2.7, 3, 5.2.2, 6
- [2] Dong E. Liu and Harold U. Baranger. Detecting a majorana-fermion zero mode using a quantum dot. 84(20). ISSN 1098-0121, 1550-235X. doi: 10.1103/PhysRevB.84.201308. URL <http://arxiv.org/abs/1107.4338>. 1.2, 1.3, 4, 4.1, 4.2, 5.2.1, 6
- [3] David A. Ruiz-Tijerina, E. Vernek, Luis G. G. V. Dias da Silva, and J. C. Egues. Interaction effects on a majorana zero mode leaking into a quantum dot. 91(11):115435. doi: 10.1103/PhysRevB.91.115435. URL <https://link.aps.org/doi/10.1103/PhysRevB.91.115435>. (document), 2, 4, 4.1, 4.1, 4.6, 4.3, 5.2.2, 7, 6
- [4] Jason Alicea. New directions in the pursuit of Majorana fermions in solid state systems. *Rep. Prog. Phys.*, 75(7):076501, 2012. ISSN 0034-4885. doi: 10.1088/0034-4885/75/7/076501. URL <http://stacks.iop.org/0034-4885/75/i=7/a=076501>. (document), 1, 1.1.1, 1.1.3, 1.6
- [5] Hao Zhang, Chun-Xiao Liu, Sasa Gazibegovic, Di Xu, John A. Logan, Guanzhong Wang, Nick van Loo, Jouri D. S. Bommer, Michiel W. A. de Moor, Diana Car, Roy L. M. Op het Veld, Petrus J. van Veldhoven, Sebastian Koelling, Marcel A. Verheijen, Mihir Pendharkar, Daniel J. Pennachio, Borzoyeh Shojaei, Joon Sue Lee, Chris J. Palmstrom, Erik P. A. M. Bakkers, S. Das Sarma, and Leo P. Kouwenhoven. Quantized Majorana conductance. *Nature*, 556:74, March 2018. URL <http://dx.doi.org/10.1038/nature26142>. (document), 1, 1.1.3, 1.7
- [6] Cornelius Malciu, Leonardo Mazza, and Christophe Mora. Braiding Majorana zero modes using quantum dots. *Physical Review B*, 98(16):165426, October 2018. doi: 10.1103/PhysRevB.98.165426. URL <https://link.aps.org/doi/10.1103/PhysRevB.98.165426>. (document), 3, 1.8
- [7] Torsten Karzig, Christina Knapp, Roman M. Lutchyn, Parsa Bonderson, Matthew B. Hastings, Chetan Nayak, Jason Alicea, Karsten Flensberg, Stephan Plugge, Yuval Oreg,

## BIBLIOGRAPHY

---

- Charles M. Marcus, and Michael H. Freedman. Scalable designs for quasiparticle-poisoning-protected topological quantum computation with Majorana zero modes. *Physical Review B*, 95(23):235305, June 2017. doi: 10.1103/PhysRevB.95.235305. URL <https://link.aps.org/doi/10.1103/PhysRevB.95.235305>. (document), 1.8
- [8] Alexander W. Holleitner, Robert H. Blick, Andreas K. Huttel, Karl Eberl, and Jorg P. Kotthaus. Probing and Controlling the Bonds of an Artificial Molecule. *Science*, 297(5578):70–72, July 2002. ISSN 0036-8075, 1095-9203. doi: 10.1126/science.1071215. URL <http://science.sciencemag.org/content/297/5578/70>. (document), 2.1
- [9] Michael Sindel. *Numerical Renormalization Group studies of Quantum Impurity Models in the Strong Coupling Limit*. Text.PhDThesis, Ludwig-Maximilians-Universitat Munchen, January 2005. URL <https://edoc.ub.uni-muenchen.de/3115/>. (document), 2.1, 2.3, 2.2, 2.3, A.1
- [10] W.J. de Haas, J. de Boer, and G.J. van dĂ«n Berg. The electrical resistance of gold, copper and lead at low temperatures. *Physica*, 1(7):1115 – 1124, 1934. ISSN 0031-8914. doi: [https://doi.org/10.1016/S0031-8914\(34\)80310-2](https://doi.org/10.1016/S0031-8914(34)80310-2). URL <http://www.sciencedirect.com/science/article/pii/S0031891434803102>. (document), 2.4, 2.3
- [11] Markus Ternes, Andreas J Heinrich, and Wolf-Dieter Schneider. Spectroscopic manifestations of the kondo effect on single adatoms. *Journal of Physics: Condensed Matter*, 21(5):053001, dec 2008. doi: 10.1088/0953-8984/21/5/053001. URL <https://doi.org/10.1088%2F0953-8984%2F21%2F5%2F053001>. (document), 2.4
- [12] W. G. van der Wiel, S. De Franceschi, T. Fujisawa, J. M. Elzerman, S. Tarucha, and L. P. Kouwenhoven. The Kondo Effect in the Unitary Limit. *Science*, 289(5487):2105–2108, September 2000. ISSN 0036-8075, 1095-9203. doi: 10.1126/science.289.5487.2105. URL <http://science-sciencemag-org.ez67.periodicos.capes.gov.br/content/289/5487/2105>. (document), 2.5
- [13] Ralf Bulla, Theo A. Costi, and Thomas Pruschke. Numerical renormalization group method for quantum impurity systems. *Reviews of Modern Physics*, 80(2):395–450, April 2008. doi: 10.1103/RevModPhys.80.395. URL <https://link.aps.org/doi/10.1103/RevModPhys.80.395>. (document), 3.2.1, 3.2.1, 3.4, 5.1.2, A.1
- [14] Walter Hofstetter. Generalized Numerical Renormalization Group for Dynamical Quantities. *Physical Review Letters*, 85(7):1508–1511, August 2000. doi: 10.1103/PhysRevLett.85.1508. URL <https://link.aps.org/doi/10.1103/PhysRevLett.85.1508>. (document), 3.2.5, 3.6
- [15] A. Yu Kitaev. Unpaired Majorana fermions in quantum wires. *Phys.-Usp.*, 44(10S):131, 2001. ISSN 1063-7869. doi: 10.1070/1063-7869/44/10S/S29. URL <http://stacks.iop.org/1063-7869/44/i=10S/a=S29>. 1, 4.1

## BIBLIOGRAPHY

---

- [16] Frank Wilczek. Majorana returns, September 2009. URL <https://www.nature.com/articles/nphys1380>. 1
- [17] A. Yu Kitaev. Fault-tolerant quantum computation by anyons. *Annals of Physics*, 303(1):2–30, January 2003. ISSN 00034916. doi: 10.1016/S0003-4916(02)00018-0. URL <http://arxiv.org/abs/quant-ph/9707021>. arXiv: quant-ph/9707021. 1
- [18] Jiannis K. Pachos. *Introduction to Topological Quantum Computation*. Cambridge University Press, Cambridge ; New York, 1 edition edition, May 2012. ISBN 978-1-107-00504-4. 1, 1.1.2
- [19] Vladimir G. Turaev. *Quantum Invariants of Knots and 3-Manifolds*. De Gruyter, Berlin, Boston, 3rd corr. ed. edition, 2016. ISBN 978-3-11-043522-1. URL <https://www.degruyter.com/view/product/461906>. 1, 1.1.2
- [20] Liang Fu and C. L. Kane. Superconducting Proximity Effect and Majorana Fermions at the Surface of a Topological Insulator. *Phys. Rev. Lett.*, 100(9):096407, March 2008. doi: 10.1103/PhysRevLett.100.096407. URL <https://link.aps.org/doi/10.1103/PhysRevLett.100.096407>. 1
- [21] Masatoshi Sato, Yoshiro Takahashi, and Satoshi Fujimoto. Non-Abelian Topological Order in \$s\$-\$s\$-Wave Superfluids of Ultracold Fermionic Atoms. *Phys. Rev. Lett.*, 103(2):020401, July 2009. doi: 10.1103/PhysRevLett.103.020401. URL <https://link.aps.org/doi/10.1103/PhysRevLett.103.020401>.
- [22] C.w.j. Beenakker. Search for Majorana Fermions in Superconductors. *Annual Review of Condensed Matter Physics*, 4(1):113–136, March 2013. ISSN 1947-5454. doi: 10.1146/annurev-conmatphys-030212-184337. URL <https://www.annualreviews.org/doi/10.1146/annurev-conmatphys-030212-184337>. 1
- [23] V. Mourik, K. Zuo, S. M. Frolov, S. R. Plissard, E. P. a. M. Bakkers, and L. P. Kouwenhoven. Signatures of Majorana Fermions in Hybrid Superconductor-Semiconductor Nanowire Devices. *Science*, 336(6084):1003–1007, May 2012. ISSN 0036-8075, 1095-9203. doi: 10.1126/science.1222360. URL <http://science-sciencemag.org.ez67.periodicos.capes.gov.br/content/336/6084/1003>. 1, 1.1.3
- [24] Anindya Das, Yuval Ronen, Yonatan Most, Yuval Oreg, Moty Heiblum, and Hadas Shtrikman. Zero-bias peaks and splitting in an Al-InAs nanowire topological superconductor as a signature of Majorana fermions. *Nature Physics*, 8(12):887, December 2012. ISSN 1745-2481. doi: 10.1038/nphys2479. URL <https://www.nature.com/articles/nphys2479>. 1.1.3
- [25] M. T. Deng, C. L. Yu, G. Y. Huang, M. Larsson, P. Caroff, and H. Q. Xu. Anomalous Zero-Bias Conductance Peak in a Nb-InSb Nanowire-Nb Hybrid Device. *Nano Letters*,

## BIBLIOGRAPHY

---

- 12(12):6414–6419, December 2012. ISSN 1530-6984. doi: 10.1021/nl303758w. URL <http://dx.doi.org/10.1021/nl303758w>.
- [26] M. T. Deng, S. Vaitiekenas, E. B. Hansen, J. Danon, M. Leijnse, K. Flensberg, J. Nygård, P. Krogstrup, and C. M. Marcus. Majorana bound state in a coupled quantum-dot hybrid-nanowire system. 354(6319):1557–1562. ISSN 0036-8075, 1095-9203. doi: 10.1126/science.aaf3961. URL <http://science.sciencemag.org/content/354/6319/1557>. 1, 1.1.3, 1.2
  - [27] Yuval Oreg, Gil Refael, and Felix von Oppen. Helical Liquids and Majorana Bound States in Quantum Wires. *Physical Review Letters*, 105(17):177002, October 2010. doi: 10.1103/PhysRevLett.105.177002. URL <https://link.aps.org/doi/10.1103/PhysRevLett.105.177002>. 1
  - [28] Roman M. Lutchyn, Jay D. Sau, and S. Das Sarma. Majorana Fermions and a Topological Phase Transition in Semiconductor-Superconductor Heterostructures. *Physical Review Letters*, 105(7):077001, August 2010. doi: 10.1103/PhysRevLett.105.077001. URL <https://link.aps.org/doi/10.1103/PhysRevLett.105.077001>. 1, 1.1.3
  - [29] Shinsei Ryu, Andreas P. Schnyder, Akira Furusaki, and Andreas W. W. Ludwig. Topological insulators and superconductors: tenfold way and dimensional hierarchy. *New Journal of Physics*, 12(6):065010, June 2010. ISSN 1367-2630. doi: 10.1088/1367-2630/12/6/065010. URL <https://doi.org/10.1088/1367-2630/12/6/065010>. 1.1.1
  - [30] Chetan Nayak, Steven H. Simon, Ady Stern, Michael Freedman, and Sankar Das Sarma. Non-Abelian anyons and topological quantum computation. *Reviews of Modern Physics*, 80(3):1083–1159, September 2008. doi: 10.1103/RevModPhys.80.1083. URL <https://link.aps.org/doi/10.1103/RevModPhys.80.1083>. 1.1.2
  - [31] A. Manchon, H. C. Koo, J. Nitta, S. M. Frolov, and R. A. Duine. New perspectives for Rashba spin-orbit coupling. *Nature Materials*, 14(9):871–882, September 2015. ISSN 1476-4660. doi: 10.1038/nmat4360. URL <https://www.nature.com/articles/nmat4360>. 1.1.3
  - [32] Stevan Nadj-Perge, Ilya K. Drozdov, Jian Li, Hua Chen, Sangjun Jeon, Jungpil Seo, Allan H. MacDonald, B. Andrei Bernevig, and Ali Yazdani. Observation of Majorana fermions in ferromagnetic atomic chains on a superconductor. *Science*, 346(6209):602–607, October 2014. ISSN 0036-8075, 1095-9203. doi: 10.1126/science.1259327. URL <http://science.sciencemag.org/content/346/6209/602>. 1.1.3
  - [33] S. Das Sarma, Jay D. Sau, and Tudor D. Stanescu. A Majorana smoking gun for the superconductor-semiconductor hybrid topological system. *Physical Review B*, 86(22), December 2012. ISSN 1098-0121, 1550-235X. doi: 10.1103/PhysRevB.86.220506. URL <http://arxiv.org/abs/1211.0539>. arXiv: 1211.0539. 1.1.3

## BIBLIOGRAPHY

---

- [34] G. Górski, J. Barański, I. Weymann, and T. Domanski. Interplay between correlations and Majorana mode in proximitized quantum dot. *Scientific Reports*, 8(1):15717, October 2018. ISSN 2045-2322. doi: 10.1038/s41598-018-33529-1. URL <https://www.nature.com/articles/s41598-018-33529-1>. 1.1.3, 2
- [35] David Aasen, Michael Hell, Ryan V. Mishmash, Andrew Higginbotham, Jeroen Danon, Martin Leijnse, Thomas S. Jespersen, Joshua A. Folk, Charles M. Marcus, Karsten Flensberg, and Jason Alicea. Milestones Toward Majorana-Based Quantum Computing. *Physical Review X*, 6(3):031016, August 2016. doi: 10.1103/PhysRevX.6.031016. URL <https://link.aps.org/doi/10.1103/PhysRevX.6.031016>. 1.1.3
- [36] Sankar Das Sarma, Michael Freedman, and Chetan Nayak. Majorana zero modes and topological quantum computation. *npj Quantum Information*, 1:15001, October 2015. ISSN 2056-6387. doi: 10.1038/npjqi.2015.1. URL <https://www.nature.com/articles/npjqi20151>.
- [37] B. van Heck, A. R. Akhmerov, F. Hassler, M. Burrello, and C. W. J. Beenakker. Coulomb-assisted braiding of Majorana fermions in a Josephson junction array. *New Journal of Physics*, 14(3):035019, March 2012. ISSN 1367-2630. doi: 10.1088/1367-2630/14/3/035019. URL <https://doi.org/10.1088/1367-2630/14/3/035019>. 1.1.3
- [38] E. Vernek, P. H. Penteado, A. C. Seridonio, and J. C. Egues. Subtle leakage of a majorana mode into a quantum dot. 89(16):165314. doi: 10.1103/PhysRevB.89.165314. URL <https://link.aps.org/doi/10.1103/PhysRevB.89.165314>. 1.2, 2, 4.1, 4.2, 5.2.2
- [39] Eduardo J. H. Lee, Xiaocheng Jiang, Ramón Aguado, Georgios Katsaros, Charles M. Lieber, and Silvano De Franceschi. Zero-Bias Anomaly in a Nanowire Quantum Dot Coupled to Superconductors. *Physical Review Letters*, 109(18):186802, October 2012. doi: 10.1103/PhysRevLett.109.186802. URL <https://link.aps.org/doi/10.1103/PhysRevLett.109.186802>. 2
- [40] Minchul Lee, Jong Soo Lim, and Rosa Lopez. Kondo effect in a quantum dot side-coupled to a topological superconductor. 87(24):241402. doi: 10.1103/PhysRevB.87.241402. URL <https://link.aps.org/doi/10.1103/PhysRevB.87.241402>. 2
- [41] M.-T. Deng, S. Vaitiekenas, E. Prada, P. San-Jose, J. Nygard, P. Krogstrup, R. Aguado, and C. M. Marcus. Nonlocality of Majorana modes in hybrid nanowires. *Physical Review B*, 98(8):085125, August 2018. doi: 10.1103/PhysRevB.98.085125. URL <https://link.aps.org/doi/10.1103/PhysRevB.98.085125>. 1.2
- [42] Zhaoen Su, Alexandre B. Tacla, Moira Hocevar, Diana Car, Sébastien R. Plissard, Erik P. A. M. Bakkers, Andrew J. Daley, David Pekker, and Sergey M. Frolov. Andreev molecules in semiconductor nanowire double quantum dots. *Nature Communications*,

## BIBLIOGRAPHY

---

- 8(1):585, September 2017. ISSN 2041-1723. doi: 10.1038/s41467-017-00665-7. URL <https://www.nature.com/articles/s41467-017-00665-7>. 1.2, 1.2.1, 5
- [43] Joelson F Silva and E Vernek. Andreev and majorana bound states in single and double quantum dot structures. *Journal of Physics: Condensed Matter*, 28(43):435702, sep 2016. doi: 10.1088/0953-8984/28/43/435702. URL <https://doi.org/10.1088/0953-8984/28/43/435702>. 1.2.1, 5
- [44] T. I. Ivanov. Coherent tunneling through a double quantum dot coupled to Majorana bound states. *Physical Review B*, 96(3):035417, July 2017. doi: 10.1103/PhysRevB.96.035417. URL <https://link.aps.org/doi/10.1103/PhysRevB.96.035417>. 5
- [45] L.S. Ricco, Y. Marques, F.A. Dessotti, M. de Souza, and A.C. Seridonio. Effect of inter-dots electronic repulsion in the majorana signature for a double dot interferometer. *Physica E: Low-dimensional Systems and Nanostructures*, 78:25 – 30, 2016. ISSN 1386-9477. doi: <https://doi.org/10.1016/j.physe.2015.11.025>. URL <http://www.sciencedirect.com/science/article/pii/S1386947715302915>. 1.2.1
- [46] M. A. Ruderman and C. Kittel. Indirect Exchange Coupling of Nuclear Magnetic Moments by Conduction Electrons. *Physical Review*, 96(1):99–102, October 1954. doi: 10.1103/PhysRev.96.99. URL <https://link.aps.org/doi/10.1103/PhysRev.96.99>. 1.2.1, 5.2.2
- [47] Tadao Kasuya. A Theory of Metallic Ferro- and Antiferromagnetism on Zener’s Model. *Progress of Theoretical Physics*, 16(1):45–57, July 1956. ISSN 0033-068X. doi: 10.1143/PTP.16.45. URL <https://academic.oup.com/ptp/article/16/1/45/1861363>.
- [48] Kei Yosida. Magnetic Properties of Cu-Mn Alloys. *Physical Review*, 106(5):893–898, June 1957. doi: 10.1103/PhysRev.106.893. URL <https://link.aps.org/doi/10.1103/PhysRev.106.893>. 1.2.1, 5.2.2
- [49] D. N. Zubarev. DOUBLE-TIME GREEN FUNCTIONS IN STATISTICAL PHYSICS. *Soviet Physics Uspekhi*, 3(3):320, 1960. ISSN 0038-5670. doi: 10.1070/PU1960v003n03ABEH003275. URL <http://iopscience.iop.org/article/10.1070/PU1960v003n03ABEH003275/meta>. 1.3, 3
- [50] Kenneth G. Wilson. The renormalization group: Critical phenomena and the Kondo problem. *Reviews of Modern Physics*, 47(4):773–840, October 1975. doi: 10.1103/RevModPhys.47.773. URL <https://link.aps.org/doi/10.1103/RevModPhys.47.773>. 1.3, 3.2.1, 5.1.2, 5.2.2
- [51] Dieter Bimberg, Marius Grundmann, and Nikolai N. Ledentsov. *Quantum Dot Heterostructures*. Wiley, Chichester, Eng.; New York, 1 edition, March 1999. ISBN 978-0-471-97388-1. 2.1, 2.1

## BIBLIOGRAPHY

---

- [52] P. W. Anderson. Localized Magnetic States in Metals. *Physical Review*, 124(1):41–53, October 1961. doi: 10.1103/PhysRev.124.41. URL <https://link.aps.org/doi/10.1103/PhysRev.124.41>. 2.2
- [53] Alexander Cyril Hewson. *The Kondo Problem to Heavy Fermions*. Cambridge University Press, April 1997. ISBN 978-0-521-59947-4. Google-Books-ID: fPzgHneNFDAC. 2.3, 1, 5.2.2
- [54] D. Goldhaber-Gordon, Hadas Shtrikman, D. Mahalu, David Abusch-Magder, U. Meirav, and M. A. Kastner. Kondo effect in a single-electron transistor. *Nature*, 391(6663):156–159, January 1998. ISSN 1476-4687. doi: 10.1038/34373. URL <https://www.nature.com/articles/34373>. 2.3.1
- [55] Daniel A Spielman. Algorithms, graph theory, and linear equations in laplacian matrices. In *Proceedings of the International Congress of Mathematicians*, volume 4, pages 2698–2722, 2010. URL <http://www.mathunion.org/ICM/ICM2010.4/Main/icm2010.4.2698.2722.pdf>. 3.1.2, 3.1.2
- [56] Hideo Aoki. Decimation method of real-space renormalization for electron systems with application to random systems. *Physica A: Statistical Mechanics and its Applications*, 114(1):538 – 542, 1982. ISSN 0378-4371. doi: [https://doi.org/10.1016/0378-4371\(82\)90345-4](https://doi.org/10.1016/0378-4371(82)90345-4). URL <http://www.sciencedirect.com/science/article/pii/0378437182903454>. 3.1.2
- [57] H. R. Krishna-murthy, J. W. Wilkins, and K. G. Wilson. Renormalization-group approach to the Anderson model of dilute magnetic alloys. 1. Static properties for the symmetric case. *Phys.Rev.*, B21:1003–1043, 1980. doi: 10.1103/PhysRevB.21.1003. 3.2.1, 3.2.1, A.1, A.1
- [58] T. A. Costi, A. C. Hewson, and V. Zlatic. Transport coefficients of the Anderson model via the numerical renormalization group. *Journal of Physics: Condensed Matter*, 6(13):2519, 1994. ISSN 0953-8984. doi: 10.1088/0953-8984/6/13/013. URL <http://stacks.iop.org/0953-8984/6/i=13/a=013>. 3.2.5
- [59] Wanda C. Oliveira and Luiz N. Oliveira. Generalized numerical renormalization-group method to calculate the thermodynamical properties of impurities in metals. *Physical Review B*, 49(17):11986–11994, May 1994. doi: 10.1103/PhysRevB.49.11986. URL <https://link.aps.org/doi/10.1103/PhysRevB.49.11986>. 3.2.6, 5.1.2
- [60] Yan-Hua Liao, Jin Huang, and Wei-Zhong Wang. Real two-stage kondo effect in parallel double quantum dot. *Journal of Magnetism and Magnetic Materials*, 377:354 – 361, 2015. ISSN 0304-8853. doi: <https://doi.org/10.1016/j.jmmm.2014.10.138>. URL <http://www.sciencedirect.com/science/article/pii/S0304885314010610>. 2

- [61] Alexander Schuray, Luzie Weithofer, and Patrik Recher. Fano resonances in Majorana bound states–quantum dot hybrid systems. *Physical Review B*, 96(8):085417, August 2017. doi: 10.1103/PhysRevB.96.085417. URL <https://link.aps.org/doi/10.1103/PhysRevB.96.085417>. 5.2.2

# Appendix A

## Appendix

### A.1 From the logarithmic discretization to the Wilson's chain.

#### Logarithmic Discretization:

We start with an Anderson model Hamiltonian such as the one in (2.6) without magnetic field

$$H = \frac{U}{2} + \sum_{\sigma} \left[ \left( \varepsilon + \frac{U}{2} \right) d_{\sigma}^{\dagger} d_{\sigma} + \frac{U}{2} (d_{\sigma}^{\dagger} d_{\sigma} - 1)^2 + \sum_{\mathbf{k}} \varepsilon_{\mathbf{k}} c_{\mathbf{k}\sigma}^{\dagger} c_{\mathbf{k}\sigma} + V_{\mathbf{k}} d_{\sigma}^{\dagger} c_{\mathbf{k}\sigma} + V_{\mathbf{k}}^* c_{\mathbf{k}\sigma}^{\dagger} d_{\sigma} \right]. \quad (\text{A.1})$$

At low-energies we can assume that QD couples only to s-wave states in the leads[57]. This implies that the Fermi surface is contained in a single, isotropic conduction band extending inside some fixed cutoffs  $-D$  and  $D$ . Thus,  $\varepsilon_{\mathbf{k}}$  only depends on  $|\mathbf{k}|$ . This makes possible to transform the sum over  $\mathbf{k}$  in equation A.1 into an integral over  $\varepsilon$  between the energy cutoffs

$$\begin{aligned} H = \sum_{\sigma} & \left[ \left( \varepsilon + \frac{U}{2} \right) d_{\sigma}^{\dagger} d_{\sigma} + \frac{U}{2} (d_{\sigma}^{\dagger} d_{\sigma} - 1)^2 + \int_{-D}^D d\varepsilon \varepsilon c_{\varepsilon\sigma}^{\dagger} c_{\varepsilon\sigma} \right. \\ & \left. + \int_{-D}^D \sqrt{\rho_{\sigma}(\varepsilon)} d\varepsilon V_{\varepsilon} d_{\sigma}^{\dagger} c_{\varepsilon\sigma} + V_{\varepsilon}^* c_{\varepsilon\sigma}^{\dagger} d_{\sigma} \right]. \end{aligned} \quad (\text{A.2})$$

Here  $c_{\varepsilon\sigma}^{\dagger}$  creates an electron with energy  $\varepsilon$  and  $\rho_{\sigma}(\varepsilon)$  is the density of states of the system per spin, which appears in the integral due to the change of variable from  $\mathbf{k}$  to  $\varepsilon \propto |\mathbf{k}|^2$ . Finally, we ignore the energy dependence of  $\rho$  and  $V_d$  and we replace them by their values in the Fermi energy (This approximation has no great relevance which is justified in [57]) and we renormalize the energy band doing the replacements  $k = \frac{\varepsilon}{D}$  and  $c_{k\sigma} := \sqrt{D} c_{\varepsilon\sigma}$  so that (A.2) becomes

$$\begin{aligned} H = D \sum_{\sigma} & \left[ \frac{1}{D} \left( \varepsilon + \frac{U}{2} \right) d_{\sigma}^{\dagger} d_{\sigma} + \frac{U}{2D} (d_{\sigma}^{\dagger} d_{\sigma} - 1)^2 + \int_1^1 dk k c_{k\sigma}^{\dagger} c_{k\sigma} \right. \\ & \left. + \sqrt{\frac{\Gamma}{\pi D}} \int_1^1 dk d_{\sigma}^{\dagger} c_{k\sigma} + c_{k\sigma}^{\dagger} d_{\sigma} \right] \end{aligned} \quad (\text{A.3})$$

$$= H_d + D \sum_{\sigma} \left[ \int_1^1 dk k c_{k\sigma}^{\dagger} c_{k\sigma} + \sqrt{\frac{\Gamma}{\pi D}} \int_1^1 dk d_{\sigma}^{\dagger} c_{k\sigma} + c_{k\sigma}^{\dagger} d_{\sigma} \right], \quad (\text{A.4})$$

---

### A.1. From the logarithmic discretization to the Wilson's chain.

---

where  $\Gamma = \pi\rho V^2$  is associated to the lever-width [9, (3.5)]. At this point we have our model dependent of three unit-less constants  $\frac{\varepsilon}{D}$ ,  $\frac{U}{2D}$  and  $\frac{\Gamma}{\pi D}$ . The logarithmic discretization starts by defining an scaling parameter  $\Lambda \geq 1$  in diving the energy domain  $[-1, 1]$  into an array of intervals of the form  $\{[\pm\Lambda^{-(n+1)}, \pm\Lambda^n]\}_{n \in \mathbb{N}}$ , as we can observe in Figure 3.3. Note that the width of these intervals is decreasing exponentially by

$$d_n = \Lambda^{-n} (1 - \Lambda^1).$$

Then inside of these energy intervals we can define a set of orthonormal Fourier series of the form

$$\phi_{np}^\pm(\varepsilon) = \begin{cases} \frac{1}{\sqrt{d_n}} e^{\pm i\omega_n p \varepsilon} & \varepsilon \in [\pm\Lambda^{-(n+1)}, \pm\Lambda^n] \\ 0 & \text{a.o.c.} \end{cases} \quad (\text{A.5})$$

with  $\omega_n := \frac{2\pi}{d_n}$  so that  $\phi_{np}^\pm(\pm\Lambda^{-(n+1)}) = \phi_{np}^\pm(\pm\Lambda^{-n})$ . Then we can decompose the creation operators  $c_k^\dagger$  into their interval-Fourier contributions as

$$c_{k\sigma}^\dagger = \sum_{np} \phi_{np}^+(k) c_{np\sigma}^{+\dagger} + \phi_{np}^-(k) c_{np\sigma}^{-\dagger} \quad (\text{A.6})$$

with the new creation operators defined as

$$c_{np\sigma}^{\pm\dagger} := (c_{np\sigma}^\pm)^\dagger = \int_1^1 d\varepsilon [\phi_{np}^\pm(\varepsilon)]^* c_{\varepsilon\sigma}^\dagger.$$

This decomposition (A.6) is a simple consequence of the orthonormality of the functions defined in (A.5). In addition we can readily proof that  $c_{np\sigma}^{\pm\dagger}$ -operators satisfy the anti-commutation relations, so that they are rightful fermionic creation operators.

We can now use (A.6) to replace the  $k$ -dependent terms in hamiltonian (A.3). Then we obtain

$$\begin{aligned} \int_1^1 dk c_{k\sigma}^\dagger d\sigma &= \int_1^1 dk \left( \sum_{np} \phi_{np}^+(k) c_{np\sigma}^{+\dagger} + \phi_{np}^-(k) c_{np\sigma}^{-\dagger} \right) d\sigma \\ &= \left( \sum_{np} \left( \int_1^1 dk \phi_{np}^+(k) \right) c_{np\sigma}^{+\dagger} + \left( \int_1^1 dk \phi_{np}^-(k) \right) c_{np\sigma}^{-\dagger} \right) d\sigma \\ &= \left( \sum_{np} \left( \int_{\Lambda^{-(n+1)}}^{\Lambda^{-n}} dk \frac{e^{i\omega_n p k}}{\sqrt{d_n}} \right) c_{np\sigma}^{+\dagger} + \left( \int_{-\Lambda^{-n}}^{-\Lambda^{-(n+1)}} dk \frac{e^{-i\omega_n p k}}{\sqrt{d_n}} \right) c_{np\sigma}^{-\dagger} \right) d\sigma \\ &= \left( \sum_{np} \sqrt{d_n} \delta_p c_{np\sigma}^{+\dagger} + \sqrt{d_n} \delta_p c_{np\sigma}^{-\dagger} \right) d\sigma \\ &= \sqrt{1 - \Lambda^1} \sum_n \Lambda^{-\frac{n}{2}} (c_{np\sigma}^{+\dagger} + c_{np\sigma}^{-\dagger}) d\sigma. \end{aligned} \quad (\text{A.7})$$

---

### A.1. From the logarithmic discretization to the Wilson's chain.

---

And

$$\begin{aligned}
\int_1^1 dk k c_{k\sigma}^\dagger c_{k\sigma} &= \sum_{n,n',p,p'} \sum_{s,s'=\pm} \left( \int_1^1 k dk \phi_{np}^s(k) \left( \phi_{np'}^{s'}(k) \right)^* \right) c_{np\sigma}^{s\dagger} c_{n'p'\sigma}^{s'} \\
&= \sum_{n,n',p,p'} \sum_{s,s'=\pm} \left( \frac{\delta_{nn'} \delta_{ss'}}{d_n} \int_{\Lambda^{-(n+1)}}^{\Lambda^{-n}} k dk e^{is\omega_n k(p-p')} \right) c_{np\sigma}^{s\dagger} c_{n'p'\sigma}^s \\
&= \sum_{npp's=\pm} \left( \frac{s}{2} \Lambda^{-2n} (1-\Lambda^2) \delta_{pp'} + \frac{1-\delta_{pp'}}{is\omega_n(p-p')} \left[ ke^{is\omega_n k(p-p')} \right]_{\Lambda^{-(n+1)}}^{\Lambda^{-n}} \right) \frac{c_{np\sigma}^{s\dagger} c_{n'p'\sigma}^{s'}}{d_n} \\
&= \frac{1}{2} (1+\Lambda^1) \sum_{np} \Lambda^{-n} (c_{np\sigma}^{+\dagger} c_{np\sigma}^+ - c_{np\sigma}^{-\dagger} c_{np\sigma}^-) \\
&\quad + \sum_n \sum_{p \neq p'} \frac{1-\Lambda^1}{2i\pi(p'-p)} (c_{np\sigma}^{+\dagger} c_{n'p'\sigma}^+ - c_{np'\sigma}^{-\dagger} c_{np\sigma}^-) e^{\frac{2i\pi(p-p')}{1-\Lambda^1}}. \tag{A.8}
\end{aligned}$$

Thus, if we replace (A.7) and (A.8) into (A.3) we will obtain a logarithmic discretization of the hamiltonian. The next part will we to map this discretization to an iterative process that is worth for a numerical computations.

### Mapping the Anderson model to a Chain-Hamiltonian

We are looking for a model just like the one we have in the right part of Figure 3.3(a). This is because a Chain-Hamiltonian will give an iterative approximation of the Anderson model with an increasing (but still controllable) number of degrees of freedom. This will provide the rightful structure for a numerical diagonalization of the hamiltonian.

To do this, observe from equations (A.7),(A.8) that the QD ( $d_\sigma$ ) couples directly only to the operators with  $p = 0$  ( $c_{n0\sigma}^{\pm\dagger}$ ). The  $p \neq 0$  terms will appear in the hamiltonian only because they are coupled to  $c_{np\sigma}^{+\dagger}$  in Equation (A.8). Thus, as a first approximation we can neglect all terms in (A.8) with  $p \neq 0$ . This leaves only the first part of (A.8), so that we can define  $c_{n\sigma}^{\pm\dagger} := c_{np\sigma}^{\pm\dagger}$ . Let

$$f_{0\sigma}^\dagger = \sqrt{\frac{1-\Lambda^1}{2}} \sum_n \Lambda^{-\frac{n}{2}} (c_{n\sigma}^{+\dagger} + c_{n\sigma}^{-\dagger}), \text{ so that } \sqrt{2} f_{0\sigma}^\dagger d_\sigma = \int_1^1 dk c_{k\sigma}^\dagger d_\sigma. \tag{A.9}$$

Note  $\{f_{0\sigma}^\dagger, f_{0\sigma}\} = \frac{1-\Lambda^1}{2} \sum_n 2\Lambda^{-n} = 1$ . Replacing this in (A.3)we get

$$H = H_d + D \sum_\sigma \left[ \sqrt{\frac{2\Gamma}{\pi D}} (d_\sigma^\dagger f_{0\sigma} + f_{0\sigma}^\dagger d_\sigma) + \frac{1}{2} (1+\Lambda^1) \sum_n \Lambda^{-n} (c_{n\sigma}^{+\dagger} c_{n\sigma}^+ - c_{n\sigma}^{-\dagger} c_{n\sigma}^-) \right].$$

### A.1. From the logarithmic discretization to the Wilson's chain.

---

$f_0^\dagger$  will represent the first site of the chain-hamiltonian in Figure 3.3(b) since no other term is coupled to the dot hamiltonian. We also have the coupling term  $\xi_0 = \sqrt{\frac{2\Gamma}{\pi D}}$ . It is possible to obtain the following  $f_m^\dagger$ -operators by supposing a solution of the form

$$f_{m\sigma}^\dagger = \sum_n a_{mn}^+ c_{n\sigma}^{+\dagger} + a_{mn}^- c_{n\sigma}^{-\dagger} = \sum_n \sum_{s=\pm} a_{mn}^s c_{n\sigma}^{s\dagger}, \quad (\text{A.10})$$

such that they satisfy the anti-commutation relations

$$\{f_{m\sigma}^\dagger, f_{m'\sigma'}\} = \delta_{mm'} \delta_{\sigma\sigma'}, \quad \{f_{m\sigma}^\dagger, f_{m\sigma}^\dagger\} = \{f_{m\sigma}^\dagger, f_{m\sigma}^\dagger\} = 0$$

and

$$\frac{1}{2} (1 + \Lambda^1) \sum_n \Lambda^{-n} (c_{n\sigma}^{+\dagger} c_{n\sigma}^+ - c_{n\sigma}^{-\dagger} c_{n\sigma}^-) = \sum_{m=0}^{\infty} \Lambda^{-\frac{m}{2}} \xi_m (f_{m\sigma}^\dagger f_{m+1,\sigma} + f_{m+1,\sigma}^\dagger f_{m\sigma}). \quad (\text{A.11})$$

It is possible to find a solution for this system using the formula of the right part of equation A.11. Since the relation is only given between consecutive terms  $m, m+1$  and we already have the coefficients for  $m = 0$  ( $a_{0n}^s = \sqrt{\frac{1-\Lambda^1}{2}} \Lambda^{-\frac{n}{2}}$ ). Then it is possible to determine the upper coefficients in a recursive way starting from  $m = 0$ . Supposing we can obtain the  $m^{\text{th}}$ -coefficients ( $a_{mn}^s$ ) and then finding iteratively the coefficients of  $m+1$  ( $a_{m+1,n}^s$ ) using the relation given by equation (A.11). This provides a numerical way for obtaining the  $f_{m\sigma}^\dagger$  operators. In fact in our case, where we actually did important assumptions, the problem can be solved analytically obtaining that the final Hamiltonian is given by

$$H = H_d + D \sum_{\sigma} \left[ \sqrt{\frac{2\Gamma}{\pi D}} (d_{\sigma}^\dagger f_{0\sigma} + f_{0\sigma}^\dagger d_{\sigma}) + \frac{1}{2} (1 + \Lambda^1) \sum_{n=0}^{\infty} \Lambda^{-\frac{n}{2}} \xi_n (f_{n\sigma}^\dagger f_{n+1,\sigma} + f_{n+1,\sigma}^\dagger f_{n\sigma}) \right]. \quad (\text{A.12})$$

with

$$\xi_n = \frac{1 - \Lambda^{-n-1}}{(1 - \Lambda^{-2n-1})^{\frac{1}{2}} (1 - \Lambda^{-2n-3})^{\frac{1}{2}}}.$$

The formal recursive-solution of this problem can be found in [13]. Note that equation (A.12) describes the chain hamiltonian model that we were looking for in Figure 3.3(b). Note that in the limit when  $n \rightarrow \infty$

$$\Lambda^{-\frac{n}{2}} \xi_n \rightarrow \frac{\Lambda^{-\frac{n}{2}} (1 - \Lambda^{-n})}{1 - \Lambda^{-2n}} \sim \frac{\Lambda^{-\frac{n}{2}}}{1 + \Lambda^{-n}},$$

which implies an exponential decaying of the hopping term in the chain.

## A.2 Initial DQD-Majorana Hamiltonian.

The impurity Hamiltonian of our model is written in block diagonal form with quantum numbers: spin- $\uparrow$  total number ( $\mathcal{N}_\uparrow$ ) and spin- $\downarrow$  parity ( $P_\downarrow$ ). The initial basis is described by

$$|\uparrow\downarrow\rangle \longrightarrow |\mathcal{N}_\uparrow = 1, P_\downarrow = -1\rangle, |0\rangle \longrightarrow |\mathcal{N}_\uparrow = 0, P_\downarrow = 1\rangle \quad (\text{A.13})$$

$$|\uparrow\rangle \longrightarrow |\mathcal{N}_\uparrow = 1, P_\downarrow = 1\rangle, |\downarrow\rangle \longrightarrow |\mathcal{N}_\uparrow = 0, P_\downarrow = -1\rangle. \quad (\text{A.14})$$

And the propagation rule for the symmetry is defined with the following identity

$$|\mathcal{N}_\uparrow 1, P_\downarrow 1\rangle \otimes |\mathcal{N}_\uparrow 2, P_\downarrow 2\rangle \subset |\mathcal{N}_\uparrow 1 + \mathcal{N}_\uparrow 2, P_\downarrow 1 * P_\downarrow 2\rangle \quad (\text{A.15})$$

Let  $\varepsilon^+ = \varepsilon_1 + \varepsilon_2$  and  $U^+ = U_1 + U_2$ . In this base, the impurity Hamiltonian  $H$  (5.7) is described in the following way:

$$H_{\mathcal{N}_\uparrow=0, P_\downarrow=-1} :$$

$$\begin{aligned} |\downarrow, \downarrow, \downarrow\rangle &\rightarrow \begin{bmatrix} \varepsilon^+ + \frac{U^+}{2} & 0 & -t_1 & t_2 \\ 0 & \frac{U^+}{2} & t_2^* & t_1^* \\ 0, \downarrow, 0\rangle &\rightarrow \begin{bmatrix} -t_1^* & t_2 & \varepsilon_{d_2} + \frac{U^+}{2} & t_{dots} \\ t_2^* & t_1 & t_{dots}^* & \varepsilon_{d_1} + \frac{U^+}{2} \end{bmatrix} \\ |\downarrow, 0, 0\rangle &\rightarrow \end{aligned}$$

$$H_{\mathcal{N}_\uparrow=0, P_\downarrow=1} :$$

$$\begin{aligned} |0, 0, 0\rangle &\rightarrow \begin{bmatrix} \frac{U^+}{2} & 0 & t_1 & t_2 \\ 0 & \varepsilon^+ + \frac{U^+}{2} & t_2^* & -t_1^* \\ |\downarrow, \downarrow, 0\rangle &\rightarrow \begin{bmatrix} t_1^* & t_2 & \varepsilon_{d_1} + \frac{U^+}{2} & t_{dots} \\ t_2^* & -t_1 & t_{dots}^* & \varepsilon_{d_2} + \frac{U^+}{2} \end{bmatrix} \\ |0, \downarrow, \downarrow\rangle &\rightarrow \end{aligned}$$

$$H_{\mathcal{N}_\uparrow=2, P_\downarrow=-1} :$$

$$\begin{aligned} |\uparrow\downarrow, \uparrow\downarrow, \downarrow\rangle &\rightarrow \begin{bmatrix} 2\varepsilon^+ + \frac{3U^+}{2} & 0 & t_1 & t_2 \\ |\uparrow, \uparrow, \downarrow\rangle &\rightarrow \begin{bmatrix} 0 & \varepsilon^+ + \frac{U^+}{2} & t_2^* & -t_1^* \\ t_1^* & t_2 & f(d_1, d_2) & -t_{dots} \\ |\uparrow, \uparrow\downarrow, 0\rangle &\rightarrow \begin{bmatrix} t_2^* & -t_1 & -t_{dots}^* & f(d_2, d_1) \end{bmatrix} \\ |\uparrow\downarrow, \uparrow, 0\rangle &\rightarrow \end{aligned}$$

With  $f(d_i, d_j) = \varepsilon_{d_i} + \frac{U_i}{2} + 2\varepsilon_{d_j} + \frac{3U_j}{2}$ .

$$H_{\mathcal{N}_\uparrow=2, P_\downarrow=1} :$$

$$\begin{aligned} |\uparrow, \uparrow, 0\rangle &\rightarrow \begin{bmatrix} \varepsilon^+ + \frac{U^+}{2} & 0 & -t_1 & t_2 \\ |\uparrow\downarrow, \uparrow\downarrow, 0\rangle &\rightarrow \begin{bmatrix} 0 & 2\varepsilon^+ + \frac{3U^+}{2} & t_2^* & t_1^* \\ -t_1^* & t_2 & f(d_2, d_1) & -t_{dots} \\ |\uparrow\downarrow, \uparrow, \downarrow\rangle &\rightarrow \begin{bmatrix} t_2^* & t_1 & -t_{dots}^* & f(d_1, d_2) \end{bmatrix} \\ |\uparrow, \uparrow\downarrow, \downarrow\rangle &\rightarrow \end{aligned}$$

---

### A.2. Initial DQD-Majorana Hamiltonian.

---

Finally,

$$H_{\mathcal{N}\hat{\uparrow}=1, P_{\downarrow}=-1} : \begin{bmatrix} H_{\mathcal{N}\hat{\uparrow}=0, P_{\downarrow}=-1} & -\mathbb{T}_{dots} \\ -\mathbb{T}^*_{dots} & H_{\mathcal{N}\hat{\uparrow}=2, P_{\downarrow}=-1} \end{bmatrix}$$

$$H_{\mathcal{N}\hat{\uparrow}=1, P_{\downarrow}=1} : \begin{bmatrix} H_{\mathcal{N}\hat{\uparrow}=0, P_{\downarrow}=1} & \mathbb{T}_{dots} \\ \mathbb{T}^*_{dots} & H_{\mathcal{N}\hat{\uparrow}=2, P_{\downarrow}=1} \end{bmatrix}$$

with

$$\mathbb{T} = \begin{bmatrix} t_{dots} & 0 & 0 & 0 \\ 0 & -t_{dots} & 0 & 0 \\ 0 & 0 & -t_{dots} & 0 \\ 0 & 0 & 0 & t_{dots} \end{bmatrix}.$$

This completes the input values for the Majorana-DQD mode in the NRG code.