

Evaluation of Supracardiac Venous Angioplasty and Stenting on Orthostatic Intolerance and Orthostatic Hypotension - The STANDUP 3 Study

Background and Significance

Orthostatic hypotension (OH), defined as a systolic blood pressure drop of ≥ 20 mmHg or diastolic of ≥ 10 mmHg within 3 minutes of standing [1], affects 16–30% of adults aged 65+ (~8.3–15.6 million Medicare beneficiaries in 2023)[2]. OH, a form of orthostatic intolerance (OI), causes dizziness, syncope, and fatigue, increasing risks of mortality, heart disease, stroke[3], atrial fibrillation[4], venous thromboembolism[5], and kidney disease[6]. Current treatments (midodrine, fludrocortisone, droxidopa) are limited by side effects like hypertension[7], urological issues[8], hypokalemia[9], and heart failure[10].

Cerebral vessels integrate fluid drainage [11], with jugular venous valves regulating return [12]. Jugular outflow disturbance contributes to OH [13]. Venous sinus stenosis insights highlight venous roles in neurology [14].

2023 Medicare Part D PUF data show OH drug spending of \$86,789,822.21 for ~126,593 patients: Midodrine HCl: 866,382 prescriptions, 106,462 patients, \$69,853,535.66, Fludrocortisone Acetate: 232,372 prescriptions, 9,115 patients, \$5,824,754.89, Droxidopa (Northera): 724 prescriptions, 16 patients, \$11,111,531.66 (incomplete data).

The STANDUP 2 Study hypothesizes that supracardiac venous impairment exacerbates OH/OI by disrupting baroreflex, venous return, and glymphatic clearance[15–17]. The intervention aims to improve regulation and reduce costs.

Specific Aims

1. Assess efficacy in reducing ME and blood pressure drop.
2. Evaluate OHQ score improvements.

Methods

Single-arm registry trial recruiting 200 patients (age ≥ 18) with OH/OI at St. Francis Hospital, Roslyn, NY.

- Intervention: Venous angioplasty/stenting with RHC; de-escalate medications (droxidopa \rightarrow pyridostigmine \rightarrow fludrocortisone \rightarrow midodrine).
- Endpoints: Primary (ME/blood pressure); secondary (OHQ).
- Safety: Post-procedure monitoring; DSMB review.
- Analysis: Paired t-tests/ANOVA; power 97.8% (n=200, p<0.05).

Potential Benefits and Cost Savings

Intervention may reduce symptoms and medication use. Potential savings: ~\$38,950,298.45/year (50% generics, 10% droxidopa reduction). OH-related costs ~\$1.0–\$2.1B (0.1–0.2% of \$1,029.8B Medicare spending).

Limitations

Suppression underestimates patients (96.7% midodrine, 98.9% fludrocortisone, 4.8% droxidopa rows). Droxidopa data incomplete (975 rows). Costs Part D only. Savings hypothetical. Single-center.

Scientific and Public Health Impact

Targets venous outflow for OH/OI, potentially reducing \$86.8M Part D burden and \$1.0–\$2.1B total costs for ~12 million beneficiaries.

2023 OH Drug Costs Table

Drug Name	Prescriptions	Patients	Cost (\$)
Midodrine HCl	866,382	106,462	69,853,535.66
Fludrocortisone Acetate	232,372	9,115	5,824,754.89
Droxidopa (Northera)	724	16	11,111,531.66
Total	1,099,478	124,593	86,789,822.21

References

1. Kulkarni S, Jenkins D, Dhar A, Mir F. Treating Lows: Management of Orthostatic Hypotension. *J cardiovasc pharmacol* [Internet]. 2024 Sep 3 [cited 2025 Jan 1];84(3):303–15. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC11368167/>
2. Rutan GH, Hermanson B, Bild DE, Kittner SJ, LaBaw F, Tell GS. Orthostatic hypotension in older adults. *The Cardiovascular Health Study. CHS Collaborative Research Group. Hypertension* [Internet]. 1992 Jun;19:508–19. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/1592445>
3. Ricci F, Fedorowski A, Radico F, Romanello M, Tatasciore A, Di Nicola M, et al. Cardiovascular morbidity and mortality related to orthostatic hypotension: A meta-analysis of prospective observational studies. *European heart journal* [Internet]. 2015 Jul 1 [cited 2025 Jan 1];36(25):1609–17. Available from: <https://doi.org/10.1093/eurheartj/ehv093>
4. Ko D, Preis SR, Lubitz SA, McManus DD, Vasan RS, Hamburg NM, et al. Relation of Orthostatic Hypotension With New-Onset Atrial Fibrillation (From the Framingham Heart Study). *The american journal of cardiology* [Internet]. 2018 Mar 1 [cited 2023 Jul 18];121(5):596–601. Available from: <https://www.sciencedirect.com/science/article/pii/S0002914917318611>
5. Bell EJ, Agarwal SK, Cushman M, Heckbert SR, Lutsey PL, Folsom AR. Orthostatic Hypotension and Risk of Venous Thromboembolism in 2 Cohort Studies. *Am j hypertens* [Internet]. 2016 May [cited 2023 Jul 11];29(5):634–40. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5014082/>
6. Franceschini N, Rose K, Astor BC, Couper D, Vupputuri S. Orthostatic hypotension is associated with incident chronic kidney disease: The Atherosclerosis Risk In Communities Study. *Hypertension* [Internet]. 2010 Dec [cited 2023 Jul 26];56(6):1054–9. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3147024/>
7. Sra J, Maglio C, Biehl M, Dhala A, Blanck Z, Deshpande S, et al. Efficacy of Midodrine Hydrochloride in Neurocardiogenic Syncope Refractory to Standard Therapy. *Journal of cardiovascular electrophysiology* [Internet]. 1997 [cited 2023 Jul 18];8(1):42–6. Available from: <https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1540-8167.1997.tb00607.x>
8. Vaidyanathan S, Soni BM, Hughes PL. Midodrine: Insidious development of urologic adverse effects in patients with spinal cord injury: A report of 2 cases. *Adv ther* [Internet]. 2007;24(4):712–20. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/17901021>
9. Burns A, Brown TM, Semple P. Extreme metabolic alkalosis with fludrocortisone therapy. *Postgrad med j* [Internet]. 1983 Aug [cited 2025 Jan 6];59(694):506–7. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2417584/>

10. Willis FR, Byrne GC, Jones TW. Fludrocortisone induced heart failure in Addison's disease. *J paediatr child health* [Internet]. 1994 Jun;30(3):280–1. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/8074919>
11. Agarwal N, Carare RO. Cerebral Vessels: An Overview of Anatomy, Physiology, and Role in the Drainage of Fluids and Solutes. *Front neurol* [Internet]. 2021 Jan 13 [cited 2025 Jun 13];11:611485. Available from: <https://www.frontiersin.org/articles/10.3389/fneur.2020.611485/full>
12. Anderhuber F. [Venous valves in the large branches of superior vena cava]. *Acta Anat (Basel)* [Internet]. 1984;119(3):184–92. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/6380191>
13. Zhou D, Ding JY, Ya JY, Pan LQ, Yan F, Yang Q, et al. Understanding jugular venous outflow disturbance. *Cns neurosci ther* [Internet]. 2018 Jun;24(6):473–82. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29687619>
14. Zhao K, Gu W, Liu C, Kong D, Zheng C, Chen W, et al. Advances in the Understanding of the Complex Role of Venous Sinus Stenosis in Idiopathic Intracranial Hypertension. *J magn reson imaging* [Internet]. 2022 Sep [cited 2025 Sep 24];56(3):645–54. Available from: <https://pmc.ncbi.nlm.nih.gov/articles/PMC9541264/>
15. Shapiro E. Adolf Fick–forgotten genius of cardiology. *Am j cardiol* [Internet]. 1972 Nov 8;30(6):662–5. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/4563215>
16. Normahani P, Shalhoub J, Narayanan S. Repurposing the systemic venous return model for conceptualisation of chronic venous insufficiency and its management. *Phlebology* [Internet]. 2020 Dec [cited 2025 Jan 1];35(10):749–51. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7675772/>
17. Ketch T, Biaggioni I, Robertson R, Robertson D. Four faces of baroreflex failure: Hypertensive crisis, volatile hypertension, orthostatic tachycardia, and malignant vagotonia. *Circulation* [Internet]. 2002 May 28;105(21):2518–23. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/12034659>