

FOUNDATIONS OF REPRESENTATION THEORY

9. EXERCISE SHEET

Jendrik Stelzner

December 6, 2013

Exercise 33:

Assume that ${}_A A \cong {}_A A \oplus {}_A A$ and let $\phi : {}_A A \rightarrow {}_A A \oplus {}_A A$ be an algebra homomorphism. We set

$$(b_0, b_1) := \phi(1)$$

and notice that for all $a \in A$

$$\phi(a) = \phi(a \cdot 1) = a\phi(1) = a(b_0, b_1) = (ab_0, ab_1).$$

Because ϕ is surjective we find $a_0, a_1 \in A$ with

$$\begin{aligned} (1, 0) &= \phi(a_0) = (a_0 b_0, a_0 b_1) \text{ and} \\ (0, 1) &= \phi(a_1) = (a_1 b_0, a_1 b_1). \end{aligned}$$

In particular we have $a_0 b_0 = a_1 b_1 = 1$ and $a_0 b_1 = a_1 b_0 = 0$. Because

$$\phi(b_0 a_0 + b_1 a_1) = (b_0 a_0 b_0 + b_1 a_1 b_0, b_0 a_0 b_1 + b_1 a_1 b_1) = (b_0, b_1) = \phi(1)$$

it follows from the injectivity of ϕ that $b_0 a_0 + b_1 a_1 = 1$.

Now assume that there exist elements $a_0, a_1, b_0, b_1 \in A$ with $a_0 b_0 = a_1 b_1 = 1$, $a_0 b_1 = a_1 b_0 = 0$ and $b_0 a_0 + b_1 a_1 = 1$. We define

$$\psi : {}_A A \rightarrow {}_A A \oplus {}_A A, a \mapsto a(b_0, b_1) = (ab_0, ab_1).$$

It is clear that ψ is an A -module homomorphism. For all $(c_0, c_1) \in {}_A A \oplus {}_A A$ we have

$$\psi(c_0 a_0 + c_1 a_1) = (c_0 a_0 b_0 + c_1 a_1 b_0, c_0 a_0 b_1 + c_1 a_1 b_1) = (c_0, c_1),$$

so ψ is surjective. For $x \in A$ with $\psi(x) = 0$ we have $(xb_0, xb_1) = (0, 0)$, so $(xb_0 a_0, xb_1 a_1) = (0, 0)$ and thus

$$0 = xb_0 a_0 + xb_1 a_1 = x(b_0 a_0 + b_1 a_1) = x \cdot 1 = x.$$

So ψ is injective. This shows that ψ is an A -module isomorphism and therefore ${}_A A \cong {}_A A \oplus {}_A A$.

One trivial example of such an algebra is $A = 0$.