

Mean Field Theory

- a.k.a. Hartree - Fock Methods

Reduce the many-body problem to the problem of a single particle moving in an effective mean field generated by all of the other particles.

Fundamental approximation in condensed matter and nuclear physics.

We will consider nonrelativistic mean field methods.


- Particles interact through an instantaneous potential:

$$V(\vec{x} - \vec{x}') = \sum_{\vec{q}} e^{i\vec{q} \cdot (\vec{x} - \vec{x}')} \tilde{V}(\vec{q})$$

In the language of Second Quantization,

$$\hat{H} = \hat{H}_0 + \hat{V}$$

$$= \sum_{\vec{q}} \frac{\hbar^2}{2m} \hat{a}_{\vec{q}}^\dagger \hat{a}_{\vec{q}} + \frac{1}{2} \sum_{\vec{k}, \vec{q}, \vec{q}'} \tilde{V}(\vec{q}) \hat{a}_{\vec{q}-\vec{q}}^\dagger \hat{a}_{\vec{k}+\vec{q}}^\dagger \hat{a}_{\vec{k}} \hat{a}_{\vec{q}}$$



In the MEAN FIELD APPROXIMATION, the correction to the ground state energy is found by taking the vacuum (i.e., ground state) expectation value of \hat{V} :

$$E_0 \rightarrow E_0 + \langle 0 | \hat{V} | 0 \rangle$$

How do we calculate $\langle 0 | \hat{V} | 0 \rangle$?

Wick's Theorem tells us how:

$$\begin{aligned} & \langle 0 | \hat{a}_{\vec{s}-\vec{q}}^{\dagger} \hat{a}_{\vec{k}+\vec{q}}^{\dagger} \hat{a}_{\vec{k}} \hat{a}_{\vec{s}} | 0 \rangle \\ &= \langle 0 | : \hat{a}_{\vec{s}-\vec{q}}^{\dagger} \hat{a}_{\vec{k}+\vec{q}}^{\dagger} \hat{a}_{\vec{k}} \hat{a}_{\vec{s}} : \rangle + \langle 0 | : \hat{a}_{\vec{s}-\vec{q}}^{\dagger} \hat{a}_{\vec{k}}^{\dagger} : \hat{a}_{\vec{k}+\vec{q}} \hat{a}_{\vec{s}} : \rangle \\ &+ \langle 0 | : \hat{a}_{\vec{s}-\vec{q}}^{\dagger} \hat{a}_{\vec{s}}^{\dagger} : \hat{a}_{\vec{k}+\vec{q}} \hat{a}_{\vec{k}} : \rangle - \langle 0 | : \hat{a}_{\vec{s}-\vec{q}}^{\dagger} \hat{a}_{\vec{k}}^{\dagger} : \hat{a}_{\vec{s}} \hat{a}_{\vec{k}+\vec{q}} : \rangle \\ &+ \langle 0 | : \hat{a}_{\vec{s}-\vec{q}}^{\dagger} \hat{a}_{\vec{s}}^{\dagger} : \hat{a}_{\vec{k}} \hat{a}_{\vec{k}+\vec{q}} : \rangle + \langle 0 | : \hat{a}_{\vec{k}+\vec{q}}^{\dagger} \hat{a}_{\vec{k}}^{\dagger} : \hat{a}_{\vec{s}-\vec{q}} \hat{a}_{\vec{s}} : \rangle \\ &- \langle 0 | : \hat{a}_{\vec{k}+\vec{q}}^{\dagger} \hat{a}_{\vec{s}}^{\dagger} : \hat{a}_{\vec{s}-\vec{q}} \hat{a}_{\vec{k}} : \rangle + \langle 0 | : \hat{a}_{\vec{s}-\vec{q}}^{\dagger} \hat{a}_{\vec{k}}^{\dagger} : \hat{a}_{\vec{s}} \hat{a}_{\vec{k}} : \rangle \\ &- \langle 0 | : \hat{a}_{\vec{s}-\vec{q}}^{\dagger} \hat{a}_{\vec{k}}^{\dagger} \hat{a}_{\vec{s}} \hat{a}_{\vec{k}+\vec{q}} : \rangle + \langle 0 | : \hat{a}_{\vec{s}-\vec{q}}^{\dagger} \hat{a}_{\vec{s}}^{\dagger} \hat{a}_{\vec{k}+\vec{q}} \hat{a}_{\vec{k}} : \rangle \end{aligned}$$

All of the terms left with a normal ordering of operators will not contribute to the VEV.

So, we are left with

$$\langle 0 | \hat{V} | 0 \rangle = \frac{1}{2} \sum_{\vec{k}, \vec{q}} \tilde{V}(\vec{q}) \{ \langle \hat{a}_{\vec{s}-\vec{q}}^{\dagger} \hat{a}_{\vec{k}+\vec{q}}^{\dagger} \rangle \langle \hat{a}_{\vec{k}} \hat{a}_{\vec{s}} \rangle$$

(3)

$$\begin{aligned}
 & + \langle \hat{a}_{\vec{r}-\vec{g}}^\dagger \hat{a}_{\vec{r}} \rangle \langle \hat{a}_{\vec{k}+\vec{g}}^\dagger \hat{a}_{\vec{k}} \rangle \\
 & - \langle \hat{a}_{\vec{r}-\vec{g}}^\dagger \hat{a}_{\vec{k}} \rangle \langle \hat{a}_{\vec{k}+\vec{g}}^\dagger \hat{a}_{\vec{r}} \rangle \}
 \end{aligned}$$

$$\textcircled{1} = \text{Coulomb Term} \equiv C$$

$$\textcircled{2} = \text{Hartree Term} \equiv H$$

$$\textcircled{3} = \text{Fock Term} \equiv F$$

Let's look at

$$H = \frac{1}{2} \sum_{\vec{k}, \vec{r}, \vec{g}} \tilde{V}(\vec{g}) \langle \hat{a}_{\vec{r}-\vec{g}}^\dagger \hat{a}_{\vec{r}} \rangle \langle \hat{a}_{\vec{k}+\vec{g}}^\dagger \hat{a}_{\vec{k}} \rangle$$

$$= \frac{1}{2} \sum_{\vec{k}, \vec{g}} \tilde{V}(\vec{g}) \langle \hat{a}_{\vec{r}}^\dagger \hat{a}_{\vec{r}} \rangle \langle \hat{a}_{\vec{k}}^\dagger \hat{a}_{\vec{k}} \rangle$$

$\vec{g}=0$ corresponds to infinite wavelength - i.e., a constant

IN POSITION
SPACE

$$\begin{aligned}
 & = \frac{1}{2} \tilde{V}(0) \sum_{\vec{k}, \vec{g}} \langle \hat{\psi}^\dagger(\vec{x}) \hat{\psi}(\vec{x}') \rangle \langle \hat{\psi}^\dagger(\vec{y}) \hat{\psi}(\vec{y}') \rangle \\
 & \times \frac{1}{V^2} \int d^3x d^3x' d^3y d^3y' \langle \hat{\psi}^\dagger(\vec{x}) \hat{\psi}(\vec{x}') \rangle \langle \hat{\psi}^\dagger(\vec{y}) \hat{\psi}(\vec{y}') \rangle
 \end{aligned}$$

$$\sum_{\vec{x}} e^{i\vec{g} \cdot (\vec{x} - \vec{x}')}$$

$$= \sum_{\vec{x}} e^{i\vec{g} \cdot (\vec{x} - \vec{x}')} = \dots$$

$$\times e^{i(\vec{r}-\vec{r}') \cdot \vec{x}} e^{-i\vec{r} \cdot \vec{x}'} e^{i(\vec{k}+\vec{g}) \cdot \vec{y}} e^{-i\vec{k} \cdot \vec{y}'}$$

$$= \frac{1}{2} \tilde{V}(0) \sum_{\vec{g}} \int d^3x d^3y \langle \hat{\psi}^\dagger(\vec{x}) \hat{\psi}(\vec{x}) \rangle \langle \hat{\psi}^\dagger(\vec{y}) \hat{\psi}(\vec{y}) \rangle e^{-i\vec{g} \cdot (\vec{x} - \vec{y})}$$

(4)

$$= \frac{1}{2} \int d^3x d^3y v(\vec{x}-\vec{y}) \langle \hat{\psi}^\dagger(\vec{x}) \hat{\psi}(\vec{x}) \rangle \langle \hat{\psi}^\dagger(\vec{y}) \hat{\psi}(\vec{y}) \rangle$$

Now consider

$$F = -\frac{1}{2} \sum_{\vec{k}, \vec{q}, \vec{p}} \tilde{V}(\vec{q}) \langle \hat{a}_{\vec{k}-\vec{q}}^\dagger \hat{a}_{\vec{k}} \rangle \langle \hat{a}_{\vec{k}+\vec{q}}^\dagger \hat{a}_{\vec{k}} \rangle$$

$$= -\frac{1}{2} \sum_{\vec{k}, \vec{q}, \vec{p}} \tilde{V}(\vec{q})$$

$$\times \frac{1}{V^2} \int d^3x d^3x' d^3y d^3y' \langle \hat{\psi}^\dagger(\vec{x}) \hat{\psi}(\vec{x}') \rangle \langle \hat{\psi}^\dagger(\vec{y}) \hat{\psi}(\vec{y}') \rangle$$

$$\times e^{i(\vec{x}-\vec{q}) \cdot \vec{x}} e^{-i\vec{k} \cdot \vec{x}'} e^{i(\vec{k}+\vec{q}) \cdot \vec{y}} e^{-i\vec{k} \cdot \vec{y}'}$$

$$= -\frac{1}{2} \sum_{\vec{q}} \tilde{V}(\vec{q})$$


$$\times \int d^3x d^3y \langle \hat{\psi}^\dagger(\vec{x}) \hat{\psi}(\vec{y}) \rangle \langle \hat{\psi}^\dagger(\vec{y}) \hat{\psi}(\vec{x}) \rangle e^{-i\vec{q} \cdot (\vec{x}-\vec{y})}$$

$$= -\frac{1}{2} \int d^3x d^3y \langle \hat{\psi}^\dagger(\vec{x}) \hat{\psi}(\vec{y}) \rangle \langle \hat{\psi}^\dagger(\vec{y}) \hat{\psi}(\vec{x}) \rangle v(\vec{x}-\vec{y})$$

Diagrammatic Analysis

Remember our original diagram for


$$\hat{V} = \frac{1}{2} \sum_{\vec{k}, \vec{q}, \vec{p}} \tilde{V}(\vec{q}) \hat{a}_{\vec{q}-\vec{p}}^{\dagger} \hat{a}_{\vec{k}+\vec{q}}^{\dagger} \hat{a}_{\vec{k}} \hat{a}_{\vec{p}}$$



The diagrams that correspond to VEV have no external lines - there are no particles in the initial and final state.

For the Hartree term, we contract $\langle \hat{a}_{\vec{q}-\vec{p}}^{\dagger} \hat{a}_{\vec{q}} \rangle$ and $\langle \hat{a}_{\vec{k}+\vec{q}}^{\dagger} \hat{a}_{\vec{k}} \rangle$.


Diagrammatically, this corresponds to the graph



"TADPOLE" DIAGRAM

For the Fock term, we contract $\langle \hat{a}_{\vec{q}-\vec{p}}^{\dagger} \hat{a}_{\vec{k}} \rangle$ and $\langle \hat{a}_{\vec{k}+\vec{q}}^{\dagger} \hat{a}_{\vec{q}} \rangle$.

Diagrammatically,



"OYSTER" DIAGRAM

①

The Hartree-Fock Ground State Energy of a Metal

$$\langle \bar{\Psi}_0 | \hat{H} | \bar{\Psi}_0 \rangle = \langle \bar{\Psi}_0 | \hat{H}_0 | \bar{\Psi}_0 \rangle + \langle \bar{\Psi}_0 | \hat{V} | \bar{\Psi}_0 \rangle$$

$$\langle \bar{\Psi}_0 | \hat{H}_0 | \bar{\Psi}_0 \rangle = \langle \bar{\Psi}_0 | \sum_{\vec{k}\sigma} \epsilon_{\vec{k}} \hat{a}_{\vec{k}\sigma}^\dagger \hat{a}_{\vec{k}\sigma} | \bar{\Psi}_0 \rangle$$

The sum over \vec{k} will go up to \vec{k}_F , the Fermi momentum:

$$N = 2 V \frac{1}{(2\pi)^3} 4\pi \int_0^{k_F} k^2 dk = \frac{V}{\pi^2} \frac{1}{3} k_F^3$$

$$\epsilon_F = \frac{\hbar^2 k_F^2}{2m_e}$$

$$E = 2 V \frac{1}{(2\pi)^3} 4\pi \int_0^{k_F} \frac{\hbar^2 k^2}{2m_e} k^2 dk = \frac{V}{\pi^2} \frac{1}{5} \frac{\hbar^2 k_F^5}{2m_e}$$

$$\epsilon = E/N = \frac{3}{5} \frac{\hbar^2 k_F^2}{2m_e} = \frac{3}{5} \epsilon_F$$

Then

$$\begin{aligned} \langle \bar{\Psi}_0 | \hat{H}_0 | \bar{\Psi}_0 \rangle &= \langle \bar{\Psi}_0 | \sum_{\vec{k}\sigma} \epsilon_{\vec{k}} \hat{a}_{\vec{k}\sigma}^\dagger \hat{a}_{\vec{k}\sigma} | \bar{\Psi}_0 \rangle = \epsilon \langle \bar{\Psi}_0 | \sum_{\vec{k}\sigma} \hat{a}_{\vec{k}\sigma}^\dagger \hat{a}_{\vec{k}\sigma} | \bar{\Psi}_0 \rangle \\ &= \epsilon \langle \bar{\Psi}_0 | \hat{N} | \bar{\Psi}_0 \rangle = \epsilon N \end{aligned}$$

$$|\bar{\Psi}_0\rangle = \prod_{\sigma} \prod_{\vec{k}} \hat{a}_{\vec{k}\sigma}^\dagger |0\rangle = | \begin{array}{ccccccc} \uparrow & \downarrow & \uparrow & \downarrow & \dots & \uparrow & \downarrow \\ \vec{k}_1 & \vec{k}_1 & \vec{k}_2 & \vec{k}_2 & & \vec{k}_N & \vec{k}_N \end{array} \rangle$$

$|\vec{k}| \leq k_F$

$\uparrow \quad \downarrow$
 $\vec{k}_F \quad \vec{k}_F$

$$E_0 = \epsilon N = \frac{3}{5} N \epsilon_F = N \frac{Z_1 Z_2}{Z_3} E_{\text{Rydberg}}$$

$$n_s = \left(\frac{V}{4\pi/3} N \right)^{1/3} / a_B$$

Now for

$$\Delta E^{(F)} = \langle \bar{\Psi}_0 | \hat{V} | \bar{\Psi}_0 \rangle$$

We're going to adopt the JELLUM MODEL where the ions are assumed to provide a uniform positive background, which cancels with the uniform electron distribution in the Hartree term, which is therefore 0.

$$= -\frac{1}{2} \sum_{\vec{k}, \vec{g}, \vec{g}'} \tilde{V}\left(\frac{\vec{g}}{g}\right) \langle \hat{a}_{\vec{g}-\vec{g}}^\dagger \hat{a}_{\vec{k}} \rangle \langle \hat{a}_{\vec{k}+\vec{g}}^\dagger \hat{a}_{\vec{g}'} \rangle$$

$$= -\frac{1}{2} \sum_{\vec{k}, \vec{g}} \tilde{V}(\vec{g}-\vec{k}) \langle \hat{a}_{\vec{k}}^\dagger \hat{a}_{\vec{k}} \rangle \langle \hat{a}_{\vec{g}}^\dagger \hat{a}_{\vec{g}} \rangle$$

$$= 2 \sum_{|\vec{g}| < g_F} \frac{1}{2} \left\{ - \sum_{|\vec{k}| < g_F} \tilde{V}(\vec{g}-\vec{k}) \right\}$$

for
again

$$= 2 \sum_{|\vec{g}| < g_F} \frac{1}{2} \left\{ - \sum_{|\vec{k}| < g_F} \frac{e^2}{\sqrt{\epsilon_0} |\vec{g}-\vec{k}|^2} \right\}$$

Let's look at

$$\sum_{|\vec{k}| < g_F} \frac{-e^2}{\sqrt{\epsilon_0} |\vec{g}-\vec{k}|^2} = \frac{-e^2}{\sqrt{\epsilon_0}} \int \frac{d^3k}{(2\pi)^3} \frac{1}{|\vec{g}-\vec{k}|^2} = \frac{-g_F}{\pi} \left(\frac{e^2}{4\pi\epsilon_0} \right) F\left(\frac{g_F}{g}\right)$$

Then =

$$\begin{aligned}\Delta E^{(F)} &= -2 \sum_{|\vec{r}| < r_F} \frac{1}{2} \left\{ -\frac{r_F}{\pi} \left(\frac{e^2}{4\pi\epsilon_0} \right) F\left(\frac{r_F}{r_F}\right) \right\} \\ &= -V \frac{r_F}{\pi} \left(\frac{e^2}{4\pi\epsilon_0} \right) \int_{|\vec{r}| < r_F} \frac{d^3 r}{(2\pi)^3} F\left(\frac{r}{r_F}\right) \\ &= -\frac{3N}{2\pi} r_F \left(\frac{e^2}{4\pi\epsilon_0} \right) \underbrace{\int_0^1 dx x^2 F(x)}_{1/2} \\ &= -\frac{3N}{4\pi} r_F \left(\frac{e^2}{4\pi\epsilon_0} \right)\end{aligned}$$


Then

$$\frac{\Delta E}{N} = -\frac{3}{4\pi} r_F \left(\frac{e^2}{4\pi\epsilon_0} \right) = -\frac{0.916}{n_s} \text{ Rydbergs electron}$$

Finally

$$\frac{E}{N} = \left(\frac{2.2}{n_s^2} - \frac{0.916}{n_s} \right) \text{ Rydbergs electron}$$

There exists a stable minimum!



	Li	Na	K
n_s	3.22	3.96	4.86