



Rhythm generation

csd2a - sessie 1


Probability based on position in measure




Probability based on position in measure




Probability based on position in measure




Probability based on position in measure




Probability based on position in measure




Probability based on position in measure




Probability based on position in measure



Probability based on position in measure




Probability based on position in measure



```
# generate note for 'Low'  
if (prevMid(cur_index)) {  
    probabilityLow *= 0.1  
}
```


Probability based on position in measure



```
# generate note for 'Low'  
if (prevMid(cur_index)) {  
    probabilityLow *= 0.1  
}
```

Looking back even further?

Probability based on position in measure



1. Generate list per layer.

2. Remove / add notes
reviewing the other layers and
predefined dependencies

Probability based on position in measure



Pros

- offers a lot of control
- ...

Cons

- requires quite some configuration
- requires complex configurations in case of dependencies between layers
- needs a configuration per time signatures

1. Generate list

2. Remove / add n
reviewing the other layers and
predefined dependencies

Rhythm generation strategies

- Probability based on position in measure
- ...

lemand ideeën?

Rhythm generation strategies

- Probability based on position in measure
- 1st order markov chain
- Euclidean rhythms
- Sets of 2s and 3s as building blocks
- Sequence variations on given configuration sets
- Totally different strategy, none of the above, is that possible?

Rhythm generation strategies

- Probability based on position in measure
- 1st order markov chain
- Euclidean rhythms
- Sets of 2s and 3s as building blocks
- Sequence v
- Totally diff

COMBINATIONS ARE POSSIBLE! For example:

1. divide a measure in sets of 2s and 3s
2. apply probability method to define content of these sets

Rhythm generation strategies

- Probability based on position in measure
- 1st order markov chain
- Euclidean rhythms
- Sets of 2s and 3s as building blocks
- Sequence variations on given configuration sets
- Totally different strategy, none of the above, is that possible?

Rhythm generation strategies

- Probability based
- 1st order markov
- Euclidean rhythm
- Sets of 2s and 3s
- Sequence variations on given configuration sets
- Totally different strategy, none of the above, is that possible?

For example, utilizing:

- image
- weather

Q: ... how to create a relevant / interesting mapping?