

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/311125520>

Childhood Trauma: A Comprehensive Review of Effects, Assessments, and Treatments

Thesis · November 2016

CITATIONS

0

READS

6,182

1 author:



[Shannon Eaton Allen](#)

Arizona State University

4 PUBLICATIONS 0 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Childhood Trauma and Toxic Stress - Effects, Resilience, and Recovery [View project](#)

Childhood Trauma

A Comprehensive Review of Effects, Assessments, and Treatments

Shannon Allen

Arizona State University

Childhood Trauma: A Comprehensive Review of Effects, Assessments, and Treatments

Between 1995 and 1997 researchers from Kaiser Permanente and the Centers for Disease Control and Prevention (CDC) conducted a study on a toxin so pervasive that repeated exposure had the potential to modify the brain and its development, DNA, immune and hormonal systems, and behavior (Felitti, et al., 1998). Researchers found that persistent exposure to chronic stress and trauma during childhood¹ increased the lifetime risk of disease and illness, mental illness, emotional and social problems, high-risk health behaviors, and victimization and perpetration (Felitti, et al., 1998). Among children, traumatic experiences are relatively common with more than 65% experiencing at least one traumatic incident during childhood, with half of these experiencing multiple traumatic events before adulthood (Copeland, Keeler, Angold, & Costello, 2007).

Children may experience trauma in many ways: accidents, serious illnesses or injuries, grief, and natural disasters to name a few. Secondhand knowledge of volatile situations can also have a traumatic impact on a child's emotional well-being (Nelson & Wampler, 2000). These traumatic experiences generally produce tolerable levels of stress (Shonkoff, 2010). However, persistently stressful situations from sources such as physical, sexual, and emotional abuse, neglect, family violence and discourse, community violence, caregiver substance abuse, mental illness, and imprisonment have the most pervasive and far-reaching effects (Cook, Blaustein, Spinazzola, & van der Kolk, 2003). These situations rarely occur in isolation, but rather occur in groups (Felitti, et al., 1998). Complex trauma is the repeated exposure to one or more of the aforementioned incidents (Cook, Blaustein, Spinazzola, & van der Kolk, 2003). Toxic stress is a result of complex trauma that occurs when there is frequent activation of the body's stress management systems such as the hormonal and neurochemical systems (Shonkoff, 2010).

¹ "Childhood", "child", and "children" will collectively refer to individuals from ages 0 to 18 unless otherwise specified.

The significant and startling findings of the Kaiser-CDC study has resulted in a call-to-action among medical and mental health professionals to address these findings by establishing trauma-centered assessments and treatments (Steele, 2009). In addition, research on the unique needs of special populations such as military, homeless, disabled, refugee, and lesbian/gay/bisexual/transgender (LGBT) youth need to be considered in the context of trauma (Rapporteur, et al., 2016). Lastly, experts on childhood trauma are calling on child welfare and juvenile justice systems, and schools to establish training, protocols, and rules for dealing with traumatized children (Ko, et al., 2008).

The Adverse Childhood Experiences Study (ACES)

The Adverse Childhood Experiences Study was conducted by Kaiser Permanente and the Centers for Disease Control and Prevention (CDC) between 1995 and 1997 as a response to unexpected findings from a previous CDC study (Felitti, 1993; Felitti, et al., 1998). In 1993, CDC researcher Dr. Vincent Felitti, found that the majority of participants in his obesity study had a history of sexual abuse. He theorized that eating and subsequent weight gain served as a coping mechanism for fear, anxiety, and depression (Felitti, 1993). Felitti's (1993) findings prompted the largest investigation of immediate and long-term effects of childhood trauma.

Methodology

Beginning in 1997 the CDC and Kaiser Permanente collected data in two waves from over 17,000 Health Maintenance Organization members (Felitti, et al., 1998). Female and male participants were equally represented (Felitti, et al., 1998). Racial and ethnic groups consisted of 74.8% white, 11.2% Hispanic, 7.2% Asian and Pacific Islander, and 4.5% African American (Felitti, et al., 1998). The ages of the participants ranged from 19 to 60 and over (Felitti, et al.,

1998). Most participants were college graduates with “some college”, “high school graduate”, and “not high school graduate” represented as well (Felitti, et al., 1998).

In addition to receiving physical exams and comprehensive reviews of health related histories, participants completed a survey consisting of childhood experiences, behaviors, and health statuses (Felitti, et al., 1998). ACE survey questions pertained to the participants’ first 18 years of life (Felitti, et al., 1998). Questions were categorized within three major domains: abuse, household challenges, and neglect (Felitti, et al., 1998). Within the “abuse” category, questions were asked about emotional abuse, physical abuse, and sexual abuse (Felitti, et al., 1998). The “household challenges” category consisted of questions about violence toward their mother, household substance abuse, household mental illness, caregiver divorce or separation, and imprisoned household members (Felitti, et al., 1998). Lastly, the “neglect” category included emotional and physical neglect (Felitti, et al., 1998).

Findings

The prevalence of adverse childhood experiences of the ACE study sample (n=17,337) were as follows: 10.6% of participants had experienced emotional abuse, 28.3% had experienced physical abuse, and 20.7% had experienced sexual abuse (Felitti, et al., 1998). On measures of “household challenges”, 12.7% had a mother who was treated violently, 26.9% reported experiencing household substance abuse, 19.4% reported mental illness within the household, 23.3% reported caregiver separation and/or divorce, and 4.7% had an incarcerated family member (Felitti, et al., 1998). On measures of, “neglect” 14.8% reported emotional neglect and 9.9% reported physical neglect (Felitti, et al., 1998).

Adverse childhood experiences had a dose-response relationship; as the number of traumatic childhood experiences increased, so did lifetime negative physical and mental health

outcomes (Felitti, et al., 1998). In fact, researchers found that ACEs were major contributors to the leading causes of death and disability (Felitti, et al., 1998). ACE score calculations were the sum of “yes” responses to the aforementioned categories (Felitti, et al., 1998). The numbers of adverse childhood experiences (or ACE scores) were reported as follows: 36.1% of participants reported no experiences, 26% reported one experience, 15.9% reported two experiences, 9.5% reported three experiences, and 12.5% reported four or more experiences (Felitti, et al., 1998).

A Continuance of ACES

Beginning in 1984, the CDC began collecting data throughout the United States through their Behavioral Risk Factor Surveillance System (BRFSS) (Centers for Disease Control and Prevention, n.d.). BRFSS is a telephone interview system that collects data on health-risk behaviors, medical conditions, and preventative measures (Centers for Disease Control and Prevention, n.d.). Data is collected in all 50 states from 400,000 adult participants each year (Centers for Disease Control and Prevention, n.d.). The CDC uses this data to implement health promotion programs on local, state, and federal levels (Centers for Disease Control and Prevention, n.d.).

Effects of Complex Trauma

To date over 40 negative physical and mental lifetime health outcomes have been found to have a dose-response relationship to adverse childhood experiences (ACEs) (Centers for Disease Control and Prevention, 2016). Researchers for the Centers for Disease Control and Prevention (CDC) and Kaiser Permanente explained the *lifetime risks* association by illustrating how complex trauma disrupted neurodevelopment which caused social, emotional, and cognitive impairment (see Figure 1) (Felitti, et al., 1998). As a result of these impairments, high-risk behaviors were adopted leading to disability, disease, social and emotional problems, and

ultimately early death (Felitti, et al., 1998). However, neuroscientists have taken closer looks at the brain-body connection and have found how biochemicals are able to alter the body, brain, DNA, and cells in children exposed to toxic stress (Filiano, et al., 2016; Romens, McDonald, Svaren, & Pollak, 2014; Sapolsky, Romero, & Munck, 2000; Shalev, et al., 2013; Weder, et al., 2014). Without effective interventions, these alterations can potentially last throughout the lifespan.

Toxic Stress and the Body

The stress hormone cortisol plays an important role in stress reactions (Sapolsky, Romero, & Munck, 2000). Under normal circumstances high levels of cortisol are released to prepare the body for stressful events by activating immune, memory, and energy responses (Sapolsky, Romero, & Munck, 2000). This typical response is essential to healthy development (Sapolsky, Romero, & Munck, 2000). In the case of toxic stress, where stress-response systems are persistently activated, there is an overdose of cortisol (Sapolsky, Romero, & Munck, 2000). Cortisol overdose is caused by a reaction called gene methylation (Romens, McDonald, Svaren, & Pollak, 2014; Weder, et al., 2014).

During gene methylation, chemical markers adhere to genes that regulate stress responses and stop them from properly doing their jobs (Romens, McDonald, Svaren, & Pollak, 2014; Weder, et al., 2014). The alteration of these genes causes the body to be in a constant state of alert, perpetually releasing stress hormones (Romens, McDonald, Svaren, & Pollak, 2014; Weder, et al., 2014). Researchers found that children exposed to toxic stress showed alternations throughout their entire genome (Weder, et al., 2014). These alterations have been linked to a number of diseases (Romens, McDonald, Svaren, & Pollak, 2014; Weder, et al., 2014).

In addition to genetic alterations, toxic stress can also have adverse cellular effects. Telomeres are protective structures at the end of chromosomes (Shalev, et al., 2013). The destruction of telomeres has been observed in children exposed to toxic stress (Shalev, et al., 2013). As a result, cells prematurely age causing disease and frequent illness (Shalev, et al., 2013).

Until recently it was believed that the immune system had no connection to the brain, but research conducted by the University of Virginia found otherwise (Filiano, et al., 2016). The immune system inundates the body with chemicals that attack and eliminate toxins (Filiano, et al., 2016). The lymphatic system provides the chemical disbursement pathway (Filiano, et al., 2016). It is now known that this pathway also runs throughout the brain via lymphatic vessels (Filiano, et al., 2016). When a child is exposed to toxic stress the immune system is activated and inflammatory lymphatic chemicals flood the brain (Filiano, et al., 2016). The implications of these findings are still being investigated, but it is believed that the lymphatic brain-body pathway, as it pertains to the immune system, is responsible for a range of neurological diseases (Filiano, et al., 2016).

Disease, illness, and injury. Gene methylation, destruction of telomeres, and lymphatic brain-body pathways can lead to the long-term suppression of the immune system, can cause metabolic syndrome, bone mineral loss, high cholesterol, osteoporosis, arthritis, gastrointestinal disease, Cushing's syndrome, and hyperthyroidism (Felitti, et al., 1998; Felitti, 2009; Lupien, et al., 1998; Marie-Mitchell & O'Connor, 2013; Park, et al., 2016; Teicher, Anderson, & Polcari, 2012). In addition, gene methylation can exhaust the adrenal glands (Dube, et al., 2009). As a result, the adrenal gland can no longer produce enough cortisol to meet the body's demand (Dube, et al., 2009). This can cause autoimmune disorders such as lupus, muscular sclerosis,

rheumatoid arthritis, and fibromyalgia (Dube, et al., 2009). Toxic stress is also a suspected cause of Alzheimer's disease in some cases (Filiano, et al., 2016).

Other diseases often suffered by adults with high ACE scores include heart disease, cancer, stroke, liver disease, diabetes, chronic obstructive pulmonary disease (COPD), poor overall quality of life, and shortened lifespan (Anda, et al., 2008; Barile, Edwards, Dhingra, & Thompson, 2015; Brown, et al., 2009; Brown, et al., 2010; Dong, Dube, Felitti, Giles, & Anda, 2003; Dong, et al., 2004; Felitti, et al., 1998; Huang, et al., 2015). More specifically, ACE scores of four or more increase the risk for COPD and hepatitis two and a half times (Anda, et al., 2008; Felitti, et al., 1998). Individuals with ACE scores of seven or higher have three times the risk of contracting lung cancer and three and a half times the risk of developing heart disease (Felitti, et al., 1998). For each adverse childhood experience, a women's risk of contracting an autoimmune disease increases by 20% (Dube, et al., 2009). ACE scores of six or greater decrease the lifespan by nearly 20 years (Brown, et al., 2009). Other medical issues associated with ACEs are frequent headaches, high rates of fetal death, more prescription medications, and susceptibility to accidents (Anda, Brown, Felitti, Dube, & Giles, 2008; Anda, Tietjen, Schulman, Felitti, & Croft, 2010; Hillis, et al., 2004; Isohookana, Riala, Hakko, & Räsänen, 2013). Individuals with ACE scores of five or more had 70% more antidepressant prescription medications and were 6% more likely to experience fetal death than those with ACE scores of zero (Anda, Brown, Felitti, Dube, & Giles, 2008; Felitti, et al., 1998; Hillis, et al., 2004).

Toxic Stress and the Mind

Poor stress response regulation can occur when neural circuits, hormone and neurochemical systems are exposed to frequent activation through toxic stress (Loman & Gunnar, 2010; McEwen, 2008). These systems can become overly reactive or slow to return to

homeostasis during childhood and throughout the lifespan resulting in improper responses to neutral events (Loman & Gunnar; McEwen, 2008). In addition, toxic stress can lead to brain shrinkage, neuroinflammation, and defective neurocircuitry (Bluhm, Williamson, Osuch, Frewen, & Stevens, 2009; Daniels, Frewen, McKinnon, & Lanius, 2011; Ganguly & Brenhouse, 2015; Herringa, et al., 2013; Opel, et al., 2014; Sheridan, Fox, Zeanah, McLaughlin, & Nelson III, 2012; Vythilingam, et al., 2002)

When the developing brain is exposed to toxic stress, a hormone is released that has been found to shrink the size of the hippocampus (Opel, et al., 2014; Vythilingam, et al., 2002). This is known as hippocampal atrophy (Opel, et al., 2014; Vythilingam, et al., 2002). Hippocampal atrophy can negatively affect emotion control and spatial navigation² (Opel, et al., 2014). Magnetic resonance imaging studies have also found less grey matter in the prefrontal cortex and amygdala in children exposed to toxic stress (Sheridan, Fox, Zeanah, McLaughlin, & Nelson III, 2012). In addition, weak neural connections between the prefrontal cortex, amygdala, and hippocampus have been found in children exposed to toxic stress (Herringa, et al., 2013). These collective issues can significantly disrupt a child's ability to make decisions, process emotions such as fear, self-regulate³, learn, and remember throughout their lifespan (Lupien, et al., 1998; Lupien, McEwen, Gunnar, & Heim, 2009).

A process called neural pruning occurs in the developing brain. During this process non-neural brain cells called microglia, which are part of the immune system, perform necessary housekeeping by devouring and digesting unused neurons and synaptic connections (Cunningham, Martínez-Cerdeño, & Noctor, 2013). In instances where children are exposed to

² Long-term and short-term memories are functions of spatial navigation (Basso, 2008). Spatial memory records information about an individual's environment and helps them orientate themselves in order to navigate the world they live in (Basso, 2008).

³ "Self-regulate" refers to an individual's ability to initiate, inhibit, or modulate emotional, cognitive, physiological, and behavioral responses to external stimuli (Baumeister & Vohs, 2004).

toxic stress, the microglia overreact and overdose the brain with destructive neurochemicals which leads to neuroinflammation (Ganguly & Brenhouse, 2015). Neuroinflammation can cause poor mood regulation, deficient decision-making skills, and faulty executive functioning⁴ in both childhood and across a lifetime (Ganguly & Brenhouse, 2015).

Within the brain exists a neurocircuitry network that is always on standby to help with problem solving functions by connecting memory to thought integration (Bluhm, Williamson, Osuch, Frewen, & Stevens, 2009). This “default setting” helps individuals properly respond to the world around them and decide what is important and what is not (Bluhm, Williamson, Osuch, Frewen, & Stevens, 2009). The brains of children who have been exposed to toxic stress lose their connection to the default state (Bluhm, Williamson, Osuch, Frewen, & Stevens, 2009; Daniels, Frewen, McKinnon, & Lanius, 2011). With the brain being unable to return to a healthy “idle” mode, children (and adults) have issues properly interacting with their environment (Daniels, Frewen, McKinnon, & Lanius, 2011).

Mental health. Adverse mental and behavioral health lifetime outcomes associated with ACEs include depression, suicide, memory disturbances, hallucinations, poor academic and work performance, and financial stress (Anda, et al., 2004; Blodgett, 2012; Brown, et al., 2007; Chapman, et al., 2004; Dube, et al., 2001; Whitfield, Dube, Felitti, & Anda, 2005). Compared to ACE scores of zero, scores of four or more increase the risk for depression four and a half times, and increase the risk for suicide 12 times (Felitti, et al., 1998). Individuals with ACE scores of four or more are 8% more likely to miss two or more days of work each month and are 15% more likely to have serious job issues and financial problems than those with ACE scores of zero (Anda, et al., 2004). Mediators between ACEs and poor work performance include marital,

⁴ “Executive functioning” specifically refers to reasoning, problem solving, and planning (Ganguly & Brenhouse, 2015).

family, and sexual problems, depressed mood, anger problems, panic reactions, and somatic symptoms like back pain, joint problems, and headaches (Anda, et al., 2004). Students with three or more ACEs are more likely to score lower on standardized tests, be suspended or expelled, and have substantial absentee problems than children with no ACEs (Blodgett, et al., 2012).

Stress responses can be profoundly altered as well. Hypervigilant responses may include persistently feeling threatened, chronically anxious, defensive or hostile when there is no legitimate threat, and over-reactive or impulsive responses to nonthreatening situations (Loman & Gunnar, 2010). Hyporesponsiveness to hunger, sleepiness, pain, illness, and legitimate threats may also occur (McEwen, 2008).

Some neurobiological issues can rise to the level of mental disorders. One study found that individuals with ACE scores of four or more were diagnosed with nearly eight different DSM disorders throughout their lifetime (Putnam, Harris, & Putnam, 2013). Disorders reported in children exposed to high levels of toxic stress are panic disorder, mood disorder, posttraumatic stress disorder, generalized anxiety disorder, bipolar disorder, major depressive disorder, oppositional defiant disorder, attention-deficit/hyperactivity disorder, autism, and obsessive compulsive disorder (Briggs & Price, 2009; Brown, et al., 2016; Kerns & Lee, 2015; Pantell, Jutte, & Adler, 2014; Post, et al., 2015; Pynoos, Steinberg, & Piacentini, 1999; Trickey, Siddaway, Meiser-Stedman, Serpell, & Field, 2012).

Children exposed to toxic stress are especially vulnerable to social, emotional, and developmental issues. Children can exhibit behaviors such as impulsivity, self-destruction, aggression, poor body regulation (e.g. eating, sleeping, and elimination), and pathological self-soothing (e.g. self-harm) (Cook, Blaustein, Spinazzola, & van der Kolk, 2003). Their self-

concept can also be negatively modified which can lead to low self-esteem, shame, guilt, distorted body image, and an unpredictable or inconsistent sense of self⁵ (Cook, Blaustein, Spinazzola, & van der Kolk, 2003). Cognitive issues such as problems with task completion, attention, planning, anticipation, and understanding responsibility have also been noted (Cook, Blaustein, Spinazzola, & van der Kolk, 2003).

Dissociation and problems with future orientation, attachment and relationships, affect regulation, and developmental delays are the most common effects of toxic stress in children. Dissociation presents as depersonalization, altered states of consciousness, impaired memory, and amnesia (Liotti, 2004; van Dijke, Ford, & van der Hart, 2015). Issues with future orientation include feelings of loss of control, hopelessness, negative expectations, decreased social competency, feelings of powerlessness, distrust in the world (in that it is not safe), fixed mindset⁶, psychological myopia⁷, learned helplessness, risk-taking, and a sense of a truncated lifespan (Wilson & Keane, 2004). Problems with attachment and relationships include apathy, interpersonal issues, isolation, and distrust (Liotti, 2004). Deficiencies in affect regulation cause frequent misunderstandings of internal states⁸, difficulty expressing feelings, and issues with emotional self-regulation (Ford, 2005).

One study found that childhood risk factors had a dose-response relationship with developmental delays (Barth, et al., 2008). Risk factors examined were caregiver mental health problems and substance abuse, household violence, poverty, chronic illness (child), and child maltreatment (Barth, et al., 2008). Children exposed to six risk factors had nearly a 100% chance of being developmentally delayed, five risk factors had an 85% chance, four risk factors

⁵ “Sense of self” refers to how individuals think about, evaluate or perceive themselves (Prebble, Addis, & Tippet, 2013).

⁶ Fixed mindset is the belief that talent and intelligence are inherent and cannot be improved (Hochanadel & Finamore, 2015).

⁷ Psychological myopia is cognitive nearsightedness when an individual has a tendency to make decisions based solely on their personal judgement rather than more relevant external information (Cutter, 1967).

⁸ “Internal states” refers to needs, wants, interests, or desires (Di Cesare, Di Dio, Marchi, & Rizzolatti, 2015).

had a 55% chance, three risk factors had a 30% chance, and one to two risk factors had a 10% chance of being developmentally delayed (Barth, et al., 2008). Developmental delays may occur in children exposed to toxic stress in areas such as object constancy⁹, language, executive functioning, information processing, and spatial awareness¹⁰ (Cook, Blaustein, Spinazzola, & van der Kolk, 2003; Ford, 2005).

Health-risk behaviors. High ACEs increase health-risk behaviors such as smoking, drug and alcohol abuse, obesity, promiscuity and other sexual risk behaviors, unintended pregnancies, sexually transmitted diseases (STD), and teen pregnancy (Anda, et al., 1999; Dube, Anda, Felitti, Edwards, & Croft, 2002; Dube, Felitti, Dong, Chapman, & Anda, 2003; Hillis, Anda, Felitti, & Marchbanks, 2001; Williamson, Thompson, Anda, Dietz, & Felitti, 2002). Compared to individuals with no ACEs, ACE scores of four or more double their risk of becoming a smoker, increase their risk of becoming an alcoholic by seven times, are 12 times more likely to marry an alcoholic, are 10 times more likely to have injected street drugs, are 30 times more likely to have an unintended pregnancy, are three times more likely to have 50 or more sexual partners, and are 14 times more likely to contract a STD (Anda, et al., 1999; Dube, Anda, Felitti, Edwards, & Croft, 2002; Dube, Felitti, Dong, Chapman, & Anda, 2003; Hillis, Anda, Felitti, & Marchbanks, 2001). Data on children and high-risk behaviors related to ACEs is just as bleak. Compared to children with ACE scores of zero, those with ACE scores of four or higher are 20 times more likely to have intercourse by the age of 14 and 20 times more likely to get pregnant (as a teen) (Hillis, et al., 2004). Children with ACE scores of five or more are approximately 30% more

⁹ “Object constancy” is a skill acquired during early development (Lax, Bach, & Burland, 1986). It is an understanding that an absent object or person still exists (Lax, Bach, & Burland, 1986).

¹⁰ “Spatial awareness” refers to an individual’s knowledge of objects in relation to themselves (Spencer, Blades, & Morsley, 1989).

likely to use alcohol and illegal drugs than those with ACE scores of zero (Dube, Felitti, Dong, Chapman, & Anda, 2003; Dube, et al., 2006).

Victimization and perpetration. Increased risks of victimization and perpetration, especially within the context of intimate partner violence and sexual violence, are common in those with high ACE scores. Overall, ACE scores of four or more increase the risk of violence perpetration, violent victimization, and incarceration by more than five times (Bellis, Hughes, Leckenby, Perkins, & Lowey, 2014). Compared to women with ACE scores of zero, those with ACE scores of five or greater are 14% more likely to become a victim of domestic violence and are 30% more likely to be sexually assaulted as an adult (Whitfield, Anda, Dube, & Felitti, 2003). Convicted felons with ACE scores of five or more are 11 times more likely to reoffend within the first year of probation and are 15 times more likely to reoffend within the second year of probation than those with ACE scores of zero (Anda, 2011).

Within the context of ACEs, children are not immune to violent perpetration. A 2010 study found that for each adverse experience¹¹ a child is exposed to, the likelihood of violent perpetration increased from 35% to 144% (Duke, Pettingell, McMorris, & Borowsky, 2010). Interpersonal violent perpetrations measured in the study were delinquency, bullying, physical altercations, dating violence, and bringing a gun to school (Duke, Pettingell, McMorris, & Borowsky, 2010). Self-directed violence included suicide attempt or ideation, and self-mutilation (Duke, Pettingell, McMorris, & Borowsky, 2010). Juvenile justice involved youth have roughly three times more ACEs than youth in the general population (Baglivio, et al., 2014).

Poverty

¹¹ ACEs included in the study were physical abuse, sexual abuse, witnessing abuse, and caregiver drug or alcohol abuse (Duke, Pettingell, McMorris, & Borowsky, 2010).

As Figure 2 shows, the cumulative physical and mental health effects of childhood trauma, if left untreated, have the potential to create a pathway to poverty in adulthood (Frederick & Goddard, 2007). Researchers are able to predict specific patterns of behavior, traits, and brain development in children affected by adverse childhood experiences (Cook, Blaustein, Spinazzola, & van der Kolk, 2003; Blodgett, et al., 2012; Brown, et al., 2016; Felitti, et al., 1998; Teicher, et al., 2003). These behaviors, traits, and brain development modifications include aggressive behavior, diminished understanding of social cues which can lead to conflicts among peers and family members, social isolation, attention problems including diagnoses of attention deficit/hyperactivity disorder and attention deficit disorder, low IQ, poor decision making skills, delays in language and reading, and altered brain lateralization¹² (Blodgett, et al., 2012; Brown, et al., 2016; Cook, Blaustein, Spinazzola, & van der Kolk, 2003; Felitti, et al., 1998; Teicher, et al., 2003).

Children who adopt these behaviors, traits, and brain modifications are at early risk for tobacco, alcohol, and drug use (Romer, 2010). Tobacco, alcohol, and drug use can further complicate adverse behaviors, traits, and brain development associated with trauma which can eventually lead to academic problems like being placed in special education classes, poor grades, delinquency, dropping out, suspension, and expulsion (Cox, Zhang, Johnson, & Bender, 2007; Miller & Plant, 1999). Research has found that poor academic performance increases the likelihood of unemployment, low-wage jobs, the need for public assistance, and incarceration (Reardon, 2011).

There are also indications that poverty *causes* adverse childhood experiences (Child Trends, 2013). Child Trends (2013) analyzed data from the National Survey of Children's

¹² Altered brain lateralization can impact language development, sensation, movement, attachment, social interactions, and the interpretation of nonverbal communications (De Pisapia, et al., 2014; Decety & Lamm, 2007; Hecht, 2014).

Health. Controlling for chronic economic hardship as an adverse childhood experience, children living on or near the poverty line were more than two times more likely to have had three or more adverse experiences than children living above the poverty line (Child Trends, 2013).

However, Dr. Felitti and Dr. Anda (2010), co-authors of the ACE study, contended that poverty, described as lack of money, did not cause poverty. Examining trauma from an epidemiological and intergenerational perspective, Felitti and Anda (2010) explained how the middle class had the highest amount of ACEs. The effect of childhood trauma on caregivers often affects their children (Felitti & Anda, 2010). For example, Felitti and Anda (2010) found that two out of four children with caregiver's who had ACE scores of three or more would end up in poverty because of the effects their own adverse childhood experiences would have in adulthood. So while it may be true that poverty causes ACEs the mitigating factor for children is the effect that ACEs had on their caregivers.

Sources of Trauma

It is estimated that 65% of children have experienced some type of trauma, with half of those experiencing multiple traumatic episodes before adulthood, but not all traumatic experiences have pervasive lifelong effects (Copeland, Keeler, Angold, & Costello, 2007; Finkelhor, Ormrod, & Turner, 2009). Positive stress responses are an important part of development (Gunnar & Quevedo, 2007). For example, common situations such as the first day of school or a routine doctor's visit may cause a temporary increase in heart rate and moderate stress hormone elevations, but generally do not produce lifelong adverse health consequences (Sapolsky, Romero, & Munck, 2000). Stress responses to secondary trauma or events such as medical procedures or serious injuries, loss of a loved one, and more seriously, crime victimization and natural disasters generally produce tolerable levels of stress *if* buffered by

sensitive and responsive caregivers (National Scientific Council on the Developing Child, 2014). However, toxic stress, a component of complex trauma, has the most adverse and significant effects to a child's well-being that lasts throughout their lifetime (Cook, Blaustein, Spinazzola, & van der Kolk, 2003). Because complex trauma has many aggravating factors, exposure rarely occurs in isolation, but rather occurs in groups (Felitti, et al., 1998).

Secondary Trauma

Secondary trauma refers to secondhand knowledge about trauma another has suffered either by hearing them talk about it or by witnessing the aftermath. One significant source of secondary trauma is exposure through events seen on television or the Internet (Villani, 2001). Children exposed to secondary trauma may exhibit the same symptoms as children exposed to more pervasive trauma: trouble concentrating, fearful worldview, sleep problems, regression, and withdrawal; although most of these symptoms are short-lived (Pine, Costello, & Masten, 2005). Secondary trauma also plays a role in traumatic events such as school shootings, natural disasters, and terrorism where the child is not directly involved, but is indirectly exposed (Galea, Nandi, & Vlahov, 2005; Pine, Costello, & Masten, 2005).

A child's response to secondary trauma is largely dependent on protective factors (Masten & Narayan, 2012). Generally, with the support of sensitive and responsive caregivers, children can recover within several weeks (National Scientific Council on the Developing Child, 2014). Younger children exposed to media coverage often cannot reconcile that repeated images are from the same event, not different events (Villani, 2001). It is recommended that young children avoid any media exposure to tragic events (Peek, 2008).

However, school age children are likely to hear about such events outside of the home. For older children, caregivers are encouraged to be aware of their child's potential exposure –

inquire about what they already know, correct misconceptions or misinformation, watch the events on television with them, answer their question, inquire about their feelings regarding the event, show them positive images of people helping each other, and be reassuring about their own safety (Masten & Narayan, 2012; Peek, 2008). Children often formulate their own reactions based on the responses from those around them; therefore, caregivers should model healthy coping mechanisms (Pine, Costello, & Masten, 2005). Professional assistance is recommended if the child's response to the event does not decrease within several weeks.

Traumatic Events

Traumatic events set themselves apart from secondary exposure because they involve the child's direct experiences. They can also produce many of the same symptoms as complex trauma; however, like secondary trauma responses, they are generally temporary (Pine, Costello, & Masten, 2005). Immediate symptoms of trauma related to illness or injury, death of a loved one, crime victimization, and natural disasters include anxiety, trouble concentrating, trouble sleeping, withdrawal, outbursts, increased sensory sensitivity, regression, risky behaviors, illnesses such as headaches and stomachaches, helplessness, aggression, clinginess, and poor academic performance (Green, et al., 1991; Pfefferbaum, et al., 2003; Shannon, Lonigan, Finch, Jr., & Taylor, 1994).

Also like secondary trauma, the effects of traumatic events can be mitigated through protective factors (Masten & Narayan, 2012). Caregivers should encourage children to talk about the event and share their feelings about it and answer their questions (Masten & Narayan, 2012). Caregivers should also look for signs of internalized fears and worries (Masten & Narayan, 2012). Providing a consistent environment and encouraging children to participate in normal activities will help strengthen their resilience (Masten & Narayan, 2012). Lastly, if a

child's reactions to the event do not subside or become increasingly worrisome, professional help is recommended.

Illness and injury. Children can experience psychological and physiological trauma related to pain, injury, illness, and medical procedures (Kazak, et al., 2006). Medical trauma is more often related to the child's perception of the experience rather than the actual severity of the illness or injury (Kazak, et al., 2006). Childhood illness and injury is not uncommon. Five out of 100 children are hospitalized each year for injuries and major acute or chronic illnesses (Fuhrmann, 2010). Each year 20 million children are seen by doctors for accidental injuries (Borse, et al., 2008). There are over 15,000 children diagnosed with cancer each year with 380,000 child cancer survivors (Ward, DeSantis, Robbins, Kohler, & Jemal, 2014). More than 1,000 organ transplant surgeries are performed on children each year and nearly 2,000 more are still waiting (U.S. Department of Health and Human Services, Current U.S. waiting list, 2016; U.S. Department of Health and Human Services, Transplants in the U.S. by recipient age, 2016)

Although most children are resilient and adaptive proceeding an illness or injury, some may experience increased arousal, re-experiencing the event, and avoidance of any associations to the event (Saxe, Vanderbilt, & Zuckerman, 2013). These responses are often born of the subjective belief that the world is no longer a safe place, that they are more vulnerable and helpless, and feelings of uncertainty and fear about medical procedures, outcomes, and their overall safety (Bronfman, Biron Campis, & Koocher, 1998; Kassam-Adams & Winston, 2004; Saxe, Vanderbilt, & Zuckerman, 2013). Children are more vulnerable to medical trauma because they may be unfamiliar with or fearful of the sights and sounds of a hospital setting, they are unable to reconcile the seriousness of their illness or injury (e.g. a broken leg may be perceived as life-threatening to a child), they are separated from their caregivers during medical

procedures, and they are often isolated from social support (e.g. their peers) (Rennick, Johnston, Dougherty, Platt, & Ritchie, 2002).

Death of a loved one. Children who have experience the death of a loved one are faced with three primary challenges: reconciling the event that caused the death, understanding the finality of death, and dealing with loss (Perry & Rubenstein, 1999). Most children are able to recover from loss, but may exhibit grief responses that are alarming or confusing to caregivers (Cohen, Mannarino, & Deblinger, 2006). It is not uncommon for children to experience abnormal sensory episodes such as hearing the loved one's voice or catching glimpses of them (Perry & Rubenstein, 1999). Children may try to avoid the subject or reminders of the person and how they died altogether; they may seem withdrawn or pretend that nothing happened (Cohen, Mannarino, & Knudsen, 2004). Reversely, they may re-experience the loss over and over by asking the same questions more than once, drawing pictures related to the death, or by talking to complete strangers about it (Cohen, Mannarino, & Knudsen, 2004). They may experience intrusive memories through nightmares and recurrent and disturbing thoughts about the death (Perry & Rubenstein, 1999). These are all common coping mechanisms that children may employ (Cohen, Mannarino, & Knudsen, 2004; Perry & Rubenstein, 1999). Generally, as time passes and children are able to make more sense of the loss, their grief responses will eventually diminish (Cohen, Mannarino, & Deblinger, 2006).

However, when children fail to return to normal functioning, their responses escalate, or become more worrisome they may be experiencing traumatic grief (Cohen, Mannarino, Greenberg, Padlo, & Shipley, 2002). Traumatic grief is more common in incidents of unexpected deaths such as accidents, suicides, and homicides, especially if the death is that of a caregiver (Cohen, Mannarino, & Knudsen, 2004). It is much more difficult for children to cope

with traumatic grief and it often times escalate to posttraumatic stress disorder (PTSD) (Cohen, Mannarino, & Knudsen, 2004). Symptoms specific to traumatic grief are enduring and include sleep difficulties, anger outbursts, trouble concentrating, hypervigilance, poor school performance, withdrawal, regression, fatigue, stomachaches, and headaches (Jacobs, Mazure, & Prigerson, 2000). These symptoms can largely be overcome with professional treatment (Cohen, Mannarino, & Knudsen, 2004).

Crime victimization. A child's response to being a victim of a crime is largely dependent on the type of crime (Luthra, et al., 2009). Violent crimes elicit a stronger response and are often associated with complex trauma (Luthra, et al., 2009). But, children can also be traumatized by isolated incidents such as personal robbery, peer assault, or familial property victimization such as burglary related theft and vandalism (Young, 2001). In fact, children between the ages of 12 and 18 are more likely to be victims of personal robbery and peer assault than adults (Child Trends, 2014).

What sets crime victimization apart from other traumatic events is the threat of re-victimization through the criminal justice system (Crenshaw, Stella, O'Neill-Stephens, & Walsen, 2016). Trauma memories are often fragmented, consisting of motor-sensory-visceral-somatic sensations that elicit terror and fear (Crenshaw, Stella, O'Neill-Stephens, & Walsen, 2016). Over time children can reconcile their experiences, make sense of them, put them into context, and can better verbalize their experience (Crenshaw, Stella, O'Neill-Stephens, & Walsen, 2016). However, interactions with the criminal justice system generally do not allow time for recovery (Crenshaw, Stella, O'Neill-Stephens, & Walsen, 2016). Child victim/witnesses are immediately thrust into situations where aggressive tactics and age-inappropriate language is likely to be used (Crenshaw, Stella, O'Neill-Stephens, & Walsen,

2016). Child psychiatrist Judith Herman (2003) stated, “Indeed, if one set out to intentionally design a system for provoking symptoms of posttraumatic stress disorder, it might look very much like a court of law” (p. 159). This stresses the importance of trauma-informed justice systems.

Natural disasters. Children’s responses to natural disasters are partially dependent on their caregiver’s response, how destructive or frightening the disaster was, and how quickly the family is able to recover or return to normal routines (Green, et al., 1991; Shannon, Lonigan, Finch, Jr., & Taylor, 1994). The best predictor of trauma symptoms in children following a natural disaster is, specifically, their mother’s response (McFarlane, 1987). This relationship occurs because a mother’s reaction to trauma, often times, alters her parenting style where she suddenly becomes overprotective (McFarlane, 1987). Children translate this overprotectiveness as maternal vulnerability which makes it more difficult for them to reconcile their own increased sense of vulnerability that often occurs after a natural disaster (McFarlane, 1987).

As with most traumatic events, children will exhibit symptoms such as regression, avoidance or reliving the event, hypervigilance, decreased concentration, sleep problems, and anger outbursts (Lonigan, Shannon, Finch, Jr., Daugherty, & Taylor, 1991). However, most children will recover in a supportive environment (Lonigan, Shannon, Finch, Jr., Daugherty, & Taylor, 1991). Caregivers can nurture resilience by limiting media exposure, monitoring adult conversations, returning to normal activities as soon as possible, being reassuring and hopeful, maintaining household rules and expectations, and being mindful that separation anxiety is common after traumatic events (Lonigan, Shannon, Finch, Jr., Daugherty, & Taylor, 1991; Pfefferbaum, Shaw, & American Academy of Child and Adolescent Psychiatry Committee on Quality Issues, 2013).

Complex Trauma

Complex trauma refers to simultaneous or sequential exposure to multiple traumatic events or persistent chronic situations (Cook, Blaustein, Spinazzola, & van der Kolk, 2003). Toxic stress, a response to complex trauma, causes the frequent activation of the body's stress management systems (Sapolsky, Romero, & Munck, 2000). It is the body's response to toxic stress that is primarily responsible for poor mental and physical health outcomes (Filiano, et al., 2016; Romens, McDonald, Svaren, & Pollak, 2014; Sapolsky, Romero, & Munck, 2000; Shalev, et al., 2013; Weder, et al., 2014). Younger children (ages 0-6) are especially vulnerable to toxic stress because of the plasticity of the developing brain (Gunnar, 2003). In addition, maternal prenatal toxic stress can have adverse effects on the fetus (Glover, 2011; Kinsella & Monk, 2009). Complex trauma generally includes exposure to physical, emotional, and sexual abuse, neglect, caregiver substance abuse, mental illness, and incarceration, family violence, household dysfunction, and community violence (Felitti, et al., 1998). According to the Adverse Childhood Experiences (ACE) Study, nearly 65% of those surveyed had experienced complex trauma (Felitti, et al., 1998).

Early childhood trauma. From conception to young adulthood the body's biology is developing. The development of these systems can be considerably influenced by their environment (Gunnar, 2003). There is a common misperception that children between the ages of 0-6 are incapable of understanding traumatic experiences (Gunnar, 2003). Although young children may not be able to verbalize their feelings as well as older children, early childhood toxic stress can have profound effects on the brain which is very vulnerable during the early stages of development (Gunnar, 2003). The malleability of neural circuits during early brain development makes them very susceptible to modification (Loman & Gunnar, 2010).

Maternal toxic stress during pregnancy can also have adverse effects on fetal development and postnatal vulnerabilities (Glover, 2011; Kinsella & Monk, 2009). The “fetal origins hypothesis” contends that prenatal environments can significantly impact in utero physiology and development which can have lifelong implications (Kinsella & Monk, 2009). Maternal antenatal toxic stress can profoundly modify the fetus’s neurodevelopment which increases the risk for autism, cognitive disorders and disabilities, and affective disorders (Glover, 2011). Abnormal fetal heart rate and activity and adverse effects to fetal autonomic and central nervous systems have also been observed (Kinsella & Monk, 2009). Lifetime outcomes include cardiovascular disease, metabolic syndrome, and double the risk of emotional or behavioral problems (Glover, 2011; Kinsella & Monk, 2009). Like the ACE study, outcomes represent a dose-response effect (Glover, 2011; Kinsella & Monk, 2009).

Sensitive, nurturing, and responsive caregivers can potentially compensate for negative behaviors associated with toxic stress in young children, even being able to prevent elevations in cortisol levels (Nachmias, Gunnar, Mangelsdorf, Parritz, & Buss, 1996). Experts recommend providing children with an environment abundant in social play and exploration (Francis, Diorio, Plotsky, & Meaney, 2002). It is especially important, for children who spend time in childcare services, that caregivers evaluate the environment and service provider’s interactions with the children (Gunnar, Kryzer, Van Ryzin, & Phillips, 2010).

Abuse and neglect. In 2014 a reported 702,000 children were victims of neglect and abuse, 1,580 subsequently died with 80% of those being killed by a caregiver (U.S. Department of Health and Human Services, Administration for Children and Families, Administration on Children, Youth and Families, Children’s Bureau; 2016). Of those who suffered maltreatment, 9% suffered sexual abuse, 18% suffered physical abuse, and 80% suffered neglect (U.S.

Department of Health and Human Services, Administration for Children and Families, Administration on Children, Youth and Families, Children's Bureau; 2016). Children age one or younger had the highest rates of victimization (U.S. Department of Health and Human Services, Administration for Children and Families, Administration on Children, Youth and Families, Children's Bureau; 2016). Studies have indicated that families with lower socioeconomic statuses had higher rates of abuse (Bradley & Corwyn, 2002).

Caregiver attachment in early childhood plays a vital role in how individuals develop. Healthy attachments meet the child's basic needs for safety, love, and belonging (Ainsworth, 1979). It is through attachment that children learn how to trust, interact with others, regulate emotions, build self-esteem, and develop a worldview (Ainsworth, 1979). Children abused and neglected by primary caregivers have a difficult time building healthy attachments (Fergusson & Lynskey, 1997). Children who do not develop healthy attachments to primary caregivers often experience difficulties in maintaining emotional balance, developing trust, connecting with others, developing relationships, and recovering from stress, disappointment, or discouragement (Fergusson & Lynskey, 1997; George, 1996). These effects can last throughout their lifetime (George, 1996; Styron & Janoff-Bulman, 1997).

Physical and emotional abuse. Physical abuse is defined as a deliberate or negligible physical act or attempted physical act by a caregiver or someone that the child depends on for protection that causes pain or injury (Gross & Keller, 1992). Indications of physical abuse include frequent injuries that defy reasonable explanations, conflicting explanations between the caregiver and child, habitually high absentee rates at school, and difficulty walking or moving (due to pain) (Dubowitz & Bennett, 2007; McIntyre, 1987; Reiniger, Robison, & McHugh, 1995). Physical abuse during infancy can cause malformation and dysfunction in certain parts of

the brain (Dallam, 2001; Perry, 2001; U.S. Department of Health and Human Services & Administration on Children, Youth, and Families, 2001). This can lead to delays in language development, cognition (memory, attention, and learning), and socioemotional development in addition to sleep disturbances, hyperactivity, conduct disorders, and posttraumatic stress disorder (Dallam, 2001; Perry, 2001; U.S. Department of Health and Human Services & Administration on Children, Youth, and Families, 2001). Childhood abuse can also result in many lifelong chronic health issues including lung disease, heart disease, high cholesterol, liver disease, obesity, and high blood pressure (Danese, et al., 2009; Felitti, et al., 1998; Gilbert, et al., 2015). One study found that 80% of young adults who had been abused during childhood met the diagnostic criteria for at least one mental disorder; more specifically, depressive disorders, eating disorders, and anxiety disorders (Silverman, Reinherz, & Giaconia, 1996).

Children who experience physical abuse are 1.5 times more likely use illicit drugs in middle adulthood (Widom, Marmorstein, & White, 2006). Abused children are 25% more likely to experience teen pregnancy, delinquency, and academic failure (Kelley, Thornberry, & Smith, 1997). An abused child's likelihood of being arrested as a juvenile is 59% higher than children who are not abused. Childhood abuse also increases the likelihood of criminal behavior as an adult by 28% and perpetrating violent crime by 30% (Wisdom & Maxfield, 2001).

Emotional abuse occurs when a caregiver or person of trust constantly criticizes, rejects, belittles, dominates, and ignores their victim (Garbarino & Garbarino, 1994). Some examples of emotional abuse are withholding affection and comfort, threats of excessive corporal punishment, vocalizing a dislike or hatred for the child, using demeaning or negative terms to describe the child, unreasonable demands and expectations, constant unwarranted blame, and public humiliation (Kent & Wallar, 1998; Sanders & Becker-Lausen, 1995). Behavioral

presentations of emotional abuse include aggressiveness or excessive submissiveness, low self-esteem, antisocial behavior, withdraw, loss of bladder and bowel control, unreasonable fears of events (such as going home) or objects, flat affect, immature or overly mature behavior, attention and affection seeking, uncooperativeness, distrust of authority, and problems connecting with peers (Garbarino & Garbarino, 1994; Jantz & McMurray, 1995; Kent & Wallar, 1998; Sanders & Becker-Lausen, 1995).

Emotional abuse and physical abuse rarely occur in isolation, but are co-occurring patterns (Korfmacher, 1998). Children often have no expectation that it will get better and are sometimes afraid to admit or report abuse due to a fear of being punished, fear of being removed from their home, guilt and shame, or because they believe that it is normal or deserved (Ney, Fung, & Wickett, 1994; Palmer, Brown, Rae-Grant, & Loughlin, 1999). However, professionals who work with children are encouraged not to automatically assume that a child is being abused, to ask open-ended questions, and listen for inconsistencies before drawing a conclusion (Crosson-Tower, 2003; Radford, Glaser, & MacMillan, 2009).

Sexual abuse. Sexual abuse includes a range of sexual behaviors between a child and adult (or older child) that is intended to sexually stimulate the perpetrator (Finkelhor, 1979). The most common incidents include oral, anal, or vaginal intercourse and sexual fondling, touching, or kissing (Finkelhor, 1979). However, sexual abuse may also include sexual exploitation (e.g. pornography or prostitution), forcing a child to watch or look at pornographic materials, voyeurism, or exhibitionism (Finkelhor, 1979). Abusers may “groom” their victims by manipulating them with activities or gifts intended to make them feel special (Craven, Brown, & Gilchrist, 2006). Once the act has occurred the perpetrator may use threats or deception to coerce the child into silence (Finkelhor, 1987).

Sexually abused children may have difficulty sleeping, nightmares, depression, anxiety, regression (e.g. bedwetting or thumb sucking), poor school performance, poor personal hygiene, and may use sexual words, have sexual knowledge, or engage in sexual play activities that are inappropriate for their age (Browne & Finkelhor, 1986; Caldwell, Hodson, Craig, & Edgar, 2005; Cohen, Deblinger, Mannarino, & Steer, 2004; Mansell, Sobsey, & Moskal, 1998; Paolucci, Genuis, & Violato, 1999). Sexually abused boys are more likely to exhibit externalized behaviors such as anger outbursts, cruelty to others, running away, disassociation, and conduct disorders (Dube, et al., 2005; Paolucci, Genuis, & Violato, 1999). Sexually abused girls are more likely to become withdrawn and depressed (Dube, et al., 2005). Older children may engage in self-harm and suicide attempts (Dube, et al., 2005).

It is estimated that one in five girls and one in 20 boys will be sexually abused before the age of 18 (Finkelhor, 2008). Children between the ages of seven and 13 are the most vulnerable (Finkelhor, 2008). Three out four children who are sexually abused are usually victimized by someone they know (Kilpatrick, Saunders, & Smith, 2003). However, only about 30% of sexual abusers are family members (Snyder, 2000). Most (about 60%) perpetrators are non-relative acquaintances (e.g. family friends, neighbors, and babysitters) (Snyder, 2000).

Of females who experienced childhood sexual abuse by a family member, 63% also reported rape or attempted rape after the age of 14 (Lalor & McElvaney, 2010). Adolescent and teenage females who were sexually abused as a child were two times more likely to have had intercourse by age 15, three times more likely to become pregnant as a teen, two times more likely to not use birth control, and over two times more likely to have had more than one sexual partner compared to those who had not been victimized (Stock, Bell, Boyer, & Connell, 1997). Adolescent and teenage males who were sexually abused during early childhood were three

times more likely to be promiscuous, two times more likely to have had unprotected sex, and five times more likely to have caused teenage pregnancy compared to those with no history of abuse (Homma, Wang, Saewyc, & Kishor, 2012). Children who live with only one parent or live in adversely dysfunctional homes are at higher risk for being sexually victimized (Finkelhor, 2009). Like with physical and emotional abuse, child victims of sexual abuse may be afraid to report it (Palmer, Brown, Rae-Grant, & Loughlin, 1999).

Neglect. Neglect occurs when a caregiver repeatedly withholds care that is appropriate for the child's age (Dubowitz, 2016). Common forms of neglect are physical neglect, medical neglect, emotional neglect, educational neglect, environmental neglect, inadequate supervision, and newborns addicted or exposed to drugs. Physical neglect includes nutritional neglect, clothing neglect, and abandonment (Gaudin, 1995; Harrington, Zuravin, DePanfilis, Ting, & Dubowitz, 2002). Physical neglect also includes "shuttling" which is repeatedly leaving a child in the care of others for long periods of time (e.g. days, weeks, or months) and "expulsion" which is kicking a child out of the home without arrangements to provide them with adequate shelter or care by others (Gaudin, 1995). Other types of physical neglect are careless disregards for the child's safety and wellbeing such reckless driving with the child in the car (to include intoxicated driving), failure to provide age appropriate car restraints, leaving a child in the car unsupervised, and not providing adequate hygiene (Gaudin, 1995).

Medical neglect includes denying or delaying a child's healthcare needs such as not giving a child needed medication, not providing preventative dental or medical care, not getting care for a sick or injured child, and not providing mental health care (Frank, Drotar, Cook, Bleiker, & Kasper, 2001). Emotional neglect occurs when caregivers fail to provide adequate attention and nurturing or denies the child social interactions (Erickson & Egeland, 2002).

Emotional neglect is thought to have more adverse and longer lasting effects than physical neglect (Goldman, Salus, Wolcott, & Kennedy, 2003). Educational neglect involves failing to assist the child with homework and allowing chronic truancy (Mennen, Kim, Sang, & Tricketta, 2010).

Environmental neglect involves exposure to in-home hazards such as illegal and legal drugs, poisons, electrical wires, stairs, weapons, secondhand smoke, and unsanitary household conditions like rodent or bug infestations, exposed excrement, rotting food, and failing to provide basic utilities such as heat, water, and sewer (Dubowitz, 2016; Dubowitz, Pitts, & Black, 2004). Inadequate supervision includes leaving a young child home alone or with younger siblings to care for, allowing the child to partake in dangerous or illegal activities, leaving a child with an inappropriate caregiver, and depending on the age, not properly supervising the child inside and outside of the home (Dubowitz, 2016; Zuravin, 2001). Lastly, children who are prenatally exposed to drugs and alcohol are at higher risk for physical, mental, and developmental disabilities (Zuckerman & Bresnahan, 1991). Prenatal exposure to drugs and alcohol increases a child's risk of being abused or neglected by 20% (Chasnoff & Lowder, 1999).

Neglect has been found to have adverse developmental effects that can last throughout a lifetime: health, physical, intellectual, cognitive, emotional, psychological, social, and behavioral delays have been observed (DePanfilis, 2006). Indicators of child neglect are malnourishment (bloated stomach or unusually thin), frequently mentions that they are left home alone or that they care for younger siblings, dirty or poorly fitted clothing, improperly dressed for weather, untreated illnesses, injuries, or dental issues, poor hygiene, and frequently appears tired (DePanfilis, 2006). It is important to note that isolated incidents of a caregiver not providing for a child are not necessarily neglect (DePanfilis, 2006). It is also important to

consider socioecological factors such as poverty and social isolation when evaluating neglect (Dubowitz, Black, Starr, & Zuravin, 1993; Erickson & Egeland, 2002). In these cases, social support can often alleviate the problem (Dubowitz, Black, Starr, & Zuravin, 1993; Erickson & Egeland, 2002).

Family dysfunction. Family dysfunction encompasses several interrelated and overlapping conditions. Children exposed to caregiver substance abuse and addiction, caregiver mental illness, domestic violence, and incarceration, divorce and separation, and general household instability are more prone to an array of behavioral, psychiatric, and medical issues (Felitti, et al., 1998). Children living in dysfunctional households often feel shameful of their situation and are afraid to bring friends to their home (Anda, et al., 2004). This can lead to social isolation (Dube, Felitti, Dong, Chapman, & Anda, 2003).

Because of the lack of consistency in dysfunctional homes, children become mistrusting and confused (Dong, et al., 2004). Inconsistency can also lead to low self-esteem and decreased internal locus of control¹³ (McEwen, 2013). They may also have overwhelming feelings of guilt, ambivalence, fear, and insecurity (Repetti, Taylor, & Seeman, 2002). The prevalence of depression, anxiety, eating disorders, sociopathy, attention deficit/hyperactivity, and suicide attempts among these children are much higher than that of their peers (Felitti, et al., 1998). Academic issues can also arise from children being tired and unable to concentrate at school, having little support completing homework, and they have more learning disabilities, miss more days of schools, repeat more grades, and have higher levels of expulsion (Blodgett, et al., 2012; Leahy, 2015; Miller & Plant, 1999). Higher incidents of headaches and migraines, gastrointestinal disorders, and asthma have been observed in children exposed to household dysfunction (Felitti, et al., 1998).

¹³ Internal locus of control is the belief that one has control over or can influence outcomes (Lefcourt, 1991).

Caregiver substance abuse and addiction. Substance abuse is often times an underlying factor in incidents of domestic violence, child abuse and neglect, caregiver incarceration, separation, divorce, and dysfunctional household environments (Dong, et al., 2004). One study found that children exposed to caregiver alcohol abuse were 25 times more likely to be sexually abused, 20 times more likely to be physically abused, and 25 times more likely to suffer emotional abuse than children who had never experienced caregiver alcohol abuse (Anda, 2011). Children of alcoholics (COA) and children of substance abusers (COSA) are six times more likely to witness domestic violence (Anda, 2011). COAs and COSAs are three to four times more likely to become alcoholics and are at a substantially higher risk of becoming drug addicts compared to children of non-addicts (Dube, Anda, Felitti, Edwards, & Croft, 2002). Most recent estimates indicate that 8.3 million children in the United States live with a substance addicted caregiver with the majority of those caregivers dependent on alcohol and the remaining addicted to illicit drugs (Substance Abuse and Mental Health Services Administration - Office of Applied Studies, 2008).

The behavioral effects that alcohol and drugs have on caregivers can be very confusing to children; they may not understand that substances are controlling their caregiver's behavior (Famularo, Kinscherff, & Fenton, 1992). A caregiver may be loving in some instances and withdrawn in others, leaving children feeling anxious due to the unpredictability of their caregiver's behavior (Famularo, Kinscherff, & Fenton, 1992; Wisdom, Ireland, & Glynn, 1995). Caregiver alcohol and drug abuse can also create dangerous and traumatic situations for children such as caregiver intoxicated driving, excessive angry outbursts, and criminal activity (Wisdom, Ireland, & Glynn, 1995). In addition, children may be put in positions to care for their siblings and to manage age inappropriate household duties (Magura & Laudet, 1996).

Caregiver mental illness. It is estimated that 3% of children live with a severely mentally ill caregiver (Pretis & Dimova, 2008). According to the World Health Organization (2001), this number is predicted to rise. The longstanding, sudden, or repetitive nature of caregiver mental illness can be very overwhelming for children because, by nature, they have limited coping skills, defense mechanisms, and social support (Udwin, 1993). Due to stigma and having little knowledge about mental illness, children are often unwilling to discuss their situation with outsiders, leaving them isolated and feeling shame or guilt (Battaglino, 1987; Heller, Roccoforte, & Cook, 1997). Children may also experience grief and traumatic loss; feeling that the caregiver they once knew is gone or during incidents of caregiver hospitalization, which, for severely mentally ill patients, can last from one month to a year (Mattejat & Remschmidt, 2008; Mental Health America of Wisconsin, 2011). Children often become secondary to a caregiver's mental illness as family decisions are based on how it will affect the mentally ill caregiver (Leahy, 2015).

Caregiver mental illness can adversely affect parenting. For example, mentally ill caregivers may have less eye contact, physical interactions, and positive facial expressions (Mattejat & Remschmidt, 2008). Mothers specifically may have reduced empathy and a reduced ability to recognize, interpret, and respond to their child's emotional signals (Mattejat & Remschmidt, 2008). Parenting styles may be inconsistent, fluctuating between permissive and controlling. Mentally ill caregivers may also find it difficult to set consistent boundaries, control their child's behavior, have positive verbal expressions and communications, and often overreact to common stressors (Mattejat & Remschmidt, 2008). As children get older, mentally ill parents may have difficulties supporting their child's development, more specifically in areas of

autonomy¹⁴, competence, and independence (Mattejat & Remschmidt, 2008). Older children may also find it difficult to identify with mentally ill parents and are more frequently drawn into parental conflicts (Mattejat & Remschmidt, 2008).

Domestic violence. Domestic violence is defined, within the psychological framework, as patterns of behavior that occurs, generally, between intimate partners (Holden, 2003). Domestic violence is grounded in a need to control and doesn't only apply to physical or sexual violence, but also psychological abuse through threats of violence and economical abuse (Baker, Jaffe, Ashbourne, & Carter, 2002). A 2011 study found that 25.6 percent of the children surveyed (n = 4549) had been exposed to family violence at some point during their lives (Hamby, Finkelhor, Turner, & Ormrod, 2011). Children under the age of five are most likely to witness domestic violence than any other age (Fantuzzo, Boruch, Beriama, Atkins, & Marcus, 1997). Children who witness domestic violence often learn that violence equals power and control and sometimes affection (Holt, Buckley, & Whelan, 2008).

Symptoms of children exposed to domestic violence are largely dependent on their stage of development, the severity of the incident(s), the number of incidents, and their caregiver's response to them after the incident (Fantuzzo, Boruch, Beriama, Atkins, & Marcus, 1997). Common symptoms in children up to age five include regression, separation anxiety, disruptions in eating and sleep habits, impulsivity, increased aggression, over-reactive outbursts, hyperactivity, constant worry about their safety or the safety of their caregiver, lack of responsiveness to rules and boundaries, and may frequently complain of headaches or stomachaches (Baker, Jaffe, Ashbourne, & Carter, 2002; Jackson, Cram, & Seymour, 2000; Rossman, Hughes, & Rosenburg, 2000; Taylor, Zuckerman, Harik, & Groves, 1994). Younger children may also become distressed when exposed noises or visual images associated with the

¹⁴ Autonomy refers to one's ability to make their own choices; free will (Deci & Ryan, 2002).

violence and may be reluctant to engage in exploratory play (Baker, Jaffe, Ashbourne, & Carter, 2002). Symptoms of children age six to 11 include nightmares, difficulty concentrating, numbing¹⁵, difficulty establishing relationships with peers, and avoidance of social situations (Carlson, 1984; Rossman, Hughes, & Rosenberg, 2000). Common symptoms in children age 12 to 18 include antisocial behavior, substance abuse, running away, depression, generalized anxiety, academic failure, and violent or abusive behavior in dating relationships (Edleson, 1999; Jackson, Cram, & Seymour, 2000).

Regardless of frequency, children who are exposed to domestic violence are at an increased risk for abuse (Herrenkohl, Sousa, Tajima, Herrenkohl, & Moylan, 2008). Children who are frequently exposed to domestic violence are 42% more likely to be sexually abused, 60% more likely to be physically abused, and 65% more likely to be emotionally abused than children with no exposure (Anda, 2011). Children who have been exposed to domestic violence one to two times are 7% more likely to be sexually abused, 20% more likely to be physically abused, and 15% more likely to be emotionally abused than children who are not exposed to domestic violence (Anda, 2011).

In addition, children exposed to domestic violence are more likely to experience other adverse childhood experiences. For example, children who have a battered mother are 95% more likely to have one other ACE, 82% more likely to have two other ACEs, 64% more likely to have three other ACEs, 48% more likely to have four other ACEs, and 52% more likely to have five or more ACEs than their peers (Anda, 2011). ACE scores can also predict ones likelihood of becoming a victim of domestic violence in adulthood. Women with ACE scores of five or more are 14% more likely to be a victim of domestic violence (Anda, 2011).

¹⁵ Emotional numbing refers to an individual's difficulty in experiencing positive emotions while at the same time being able to experience negative emotions (Ford & Kidd, 1998).

Incarcerated caregiver(s). It is estimated that 2.3 million children have at least one incarcerated caregiver; however, only half of these children lived with their parent(s) prior to incarceration and the majority that did, lived with their mother (Mumola, 2000). The mean age of children with an incarcerated parent is eight; mothers assume responsibility 90% of the time when a father goes to jail while fathers assume responsibility only 30% of the time (Mumola, 2000). Unfortunately, the incarceration rate among mothers has been consistently rising over the last 10 years (Mumola, 2000). In many cases, grandparents become primary caregivers (Mumola, 2000).

The effect that caregiver incarceration has on a child depends on several factors: the age of the child, familial conditions prior to the incarceration, quality of alternative housing arrangements, and access to the incarcerated caregiver (Baunach, 1985; Bloom & Steinhart, 1993; Johnson, 1991; Johnston, 1995; Mumalo, 2000; Myers et al., 1999; Sack, Seidler, & Thomas, 1976; Sroufe, 1988; Thompson, 1998). Witnessing the arrest can be very traumatic for children and many suffer flashbacks of the event and have nightmares (Jose-Kampfner, 1995). It is estimated that one in five children are present at the time of the arrest with half of those being age six or under (Johnson, 1991). Some argue that concealing a caregiver's incarceration from a child reduces stress; however, others argue that children are better able to cope if the event is explained to them in age appropriate terms (Ayers, Sandler, West, & Roosa, 1996; Becker & Margolin, 1967; Snyder-Joy & Carlo, 1998). One study indicated that children who were uninformed of their caregiver's incarceration were more fearful and anxious than those who were informed (Johnson, 1995).

Caregiver incarceration can have an especially adverse lifelong impact on infants. Mother-infant bonding is critical to a child's development (Myers, Smarch, Amlund-Hagen, &

Kennon, 1999). When caregivers, especially mothers, are incarcerated the infant does not have the opportunity to form a healthy attachment (Myers, Smarch, Amlund-Hagen, & Kennon, 1999). These children often have lifelong emotional and behavioral problems (Myers, Smarch, Amlund-Hagen, & Kennon, 1999). In addition to developing insecure attachments to primary caregivers, toddlers have the capacity to understand changes in their circumstances when a caregiver is incarcerated (Johnson, 1995, Thompson, 1998). Changes such as relocation and divorce can negatively affect a child's cognitive abilities and relationships (Sroufe, 1988).

One study found that 70% of toddlers with incarcerated mothers had psychological and emotional problems (Baunach, 1985). These problems included shame, guilt, anxiety, withdrawal, depression, hypervigilance, eating disorders, anger, and aggression (Bloom & Steinhart, 1992; Fishman, 1983; Gaudin, 1984; Johnston, 1995; Jose-Kampfner, 1995; Sack, Seidler, & Thomas, 1976). Problems associated with children age six and up are generally observed in the context of peer relationships and school-related issues (Sack, Seidler, & Thomas, 1976). One study found that 70% of children with an incarcerated parent exhibited poor academic performance and 5% had behavioral problems in the classroom (Stanton, 1980). Some children are reluctant to go back to school after their parent's arrest due to fear of being ostracized or teased (Jose-Kampfner, 1991). As school-aged children get older, drop-out rates and suspensions are higher for children with an incarcerated parent than their peers (Trice, 1997).

High quality caregiver-child relationships can serve as protective factors to help buffer a child's anxiety and grief when a caregiver is incarcerated, however; most incarcerated caregivers have a history of insufficient parenting skills so these protective factors are often unavailable (Johnson, 1991; Myers, Smarch, Amlund-Hagen, & Kennon, 1999; Thompson, 1998). Most times children are better able to cope when placed in the care of a familiar family member,

however; this too can present challenges (Bloom & Steinhart, 1993; Mumalo, 2000). Many families are not financially prepared to unexpectedly take in a child and in the case of grandparents; there may be physical boundaries and generational complications (Bloom & Steinhart, 1993; Mumalo, 2000). Visiting an incarcerated caregiver in prison can be very intimidating and scary for children; however, research indicates that anxiety will subside if children can visit their caregiver regularly (Bloom & Steinhart, 1993; Simon & Landis, 1991). Frequent visitation can help calm a child's concerns about their caregiver's safety and helps build caregiver/child emotional bonds (Sack, 1977).

Separation and divorce. Regardless of whether or not parents are cooperative, compromising, and mature during separation and divorce it is likely that children will experience some psychological stress (Wallerstein & Lewis, 2004). Factors that contribute to a child's response to separation and divorce are the duration, frequency, and intensity of parental conflict, the quality of the child/parent relationship, and how well parents are able to focus on the child's needs before, during, and after the separation and divorce (Richards & Ely, 1998). Generally if children were not exposed parental conflicts, had sensitive and nurturing relationships with their parents, and parents were responsive to the child's needs, most children eventually adapt to separation and divorce (Wallerstein & Lewis, 2004).

Still, children of separation and divorce may experience a sense of abandonment and loss, anger, denial, and sadness and more behavioral and academic problems than their peers (Franke, 1983). One study indicated that 39% of girls and 29% of boys had high levels of posttraumatic stress symptoms after their parent's divorce (Joseph, Mynard, & Mayall, 2000). However, the lifelong effects on children of separation and divorce seem to be more adverse and indicate a delayed response and externalized reactions (Richards & Ely, 1998).

College aged females whose parents are divorced are less likely to attend college than their peers from stable families (Richards & Ely, 1998). Females are also 45% more likely to marry before the age of 20 compared to 15.6% of females whose parents remained together (Richards, 1996). Both males and females of divorced families are more likely to cohabitate (Richards & Ely, 1998). Research indicates that early marriage and cohabitation are risk factors for separation and divorce, so it is no surprise that separation and divorce rates are higher among adults whose parents were also separated or divorced (Amato, 2000; Graefe & Lichter, 1999; Moore & Waite, 1981). Overall, adult children of divorce have issues achieving intimate relationships and committing to marriage and parenthood (Wallerstein & Lewis, 2004). Lastly, adult children of split families are also more likely to experience financial difficulties and poverty and more psychological illnesses (Richards & Ely, 1998).

General household dysfunction. Household dysfunction is a set of conditions that inhibit healthy family functioning (Dong, et al., 2004; Forward, 1989). It is generally a culmination of separation and divorce, caregiver incarceration, mental illness, and substance abuse, and domestic violence (Dong, et al., 2004). However, household dysfunction can also include constant or extreme conflicts between parents, children, and other members of the household, lack of empathy among family members, marginalization of certain family members, lack of quality family time, disrespect of boundaries, unrealistic expectations, harsh or dogmatic discipline, over-parenting, ruling by fear, failure to set age appropriate rules and boundaries, and social isolation (Dong, et al., 2004; Forward, 1989; Frick, 1994; Higgins, 2003; Kazdin, 1995).

Community violence. Community violence is not only violence committed against an individual outside of the home, but knowledge of, participating in, and witnessing violence in their community. It is estimated that by the age of five, inner-city children have been directly

exposed to community violence, more specifically, gang activity and shootings (Bell & Jenkins, 1991). A 2015 national survey found that 54.5% of the children surveyed had been physically assaulted, 40.2% had been victims of property crimes, 27.5% had witnessed an assault, and 7.9% had been indirectly exposed to community violence (Finkelhor, Turner, Shattuck, Hamby, & Kracke, 2015).

One study found that children exposed to community violence in Chicago had the same trauma symptoms as children in war-torn areas like the West Bank, Mozambique, and Cambodia (Garbarino, Dubrow, Kostelny, & Pardo, 1992). Common complaints from children exposed to community violence are: mistrust, constant fear and focus on survival, doing anything to stay safe such as carrying a weapon, are on-guard all of the time, fear for their family's safety, especially younger siblings, feelings of a truncated lifespan, difficulty concentrating, nightmares, hypervigilance, constant anger, stomachaches and headaches, reluctance to form bonds, and a preoccupation with violence (National Child Traumatic Stress Network, 2000). Like exposure to domestic violence, exposure to high levels of community violence is associated with behavioral, emotional, social, and cognitive problems in older children and can be long-lasting (Cooley-Quille, Turner, & Beidel, 1995; Pynoos & Eth, 1985; Pynoos & Nader, 1988; Terr, 1983).

Posttraumatic symptoms are common in children exposed to community violence. Symptoms include numbing, flat affect, sleep interruptions, avoidance or re-experience, and exaggerated startle responses (Pynoos & Nader, 1988). In addition, many children exposed to community violence experience low self-esteem, memory lapses, fatigue, aggression, impulsivity, social withdrawal, and poor school performance (Pynoos & Nader, 1988). Non-school aged children can also be affected by community violence and often display passive and regressive symptoms (Peterson, Luborsky, & Seligman, 1983).

Trauma Screening and Assessment

Screenings are often referred to as norm-referenced tests in that they are standardized, are evaluated the same way, and are measured against statistical means of normative responses (Eabon & Abrahamson, n.d.; Kerig, Ford, & Olafson, 2014). As illustrated in Table 1, screening tools are generally pre-written questionnaires that may involve self-reports, caregiver report, or provider report (e.g. teacher, caseworker, or guidance counselor) (Kisiel, Conradi, Fehrenbach, Torgerson, & Briggs, 2014). In cases where caregivers are not the source of trauma, it is important to engage the family in the screening process; explain the purpose of the screening, how information obtained in the screening will be used, and refer the family to appropriate resources if necessary (Kisiel, Conradi, Fehrenbach, Torgerson, & Briggs, 2014). Because trauma screening is descriptive rather than diagnostic, it can be universally implemented by non-clinicians upon initial contact with the child (Kerig, Ford, & Olafson, 2014). Due to the cost-effectiveness of screening, it can be used in a variety of settings such as schools, juvenile justice and child welfare systems, and shelters to determine if a child needs to be referred for a more comprehensive assessment (Briere & Spinazzola, 2009; Strand, Sarmiento, & Pasquale, 2005; Kerig, Ford, & Olafson, 2014).

Trauma assessment is more focused and comprehensive than screening and is usually done by a mental health professional (Kerig, Ford, & Olafson, 2014). Assessments implement a variety of components such as observations, standardized measures, formal and informal interviews with the child, teachers, social services workers, caregivers, and parents, and a review of school and medical records (Eabon & Abrahamson, n.d.; Kerig, Ford, & Olafson, 2014). In general, assessments usually results in a diagnosis, case conceptualization, treatment, and progress and outcome evaluations (Kerig, Ford, & Olafson, 2014).

Trauma assessment should evaluate two major elements: exposure and reaction (Briere & Spinazzola, 2009). Exposure assessment involves examining what type of trauma the child was exposed to and how long the exposure lasted (Briere & Spinazzola, 2009). Assessing reaction involves evaluating the child's response to exposure such as behavioral, emotional, and physical presentations (Briere & Spinazzola, 2009). It is important to note that not all children who experience trauma will exhibit adverse reactions (Briere & Spinazzola, 2009). Reactions to look for are hyperarousal when discussing the event(s), intrusive memories about the event, behavioral problems, avoidance of triggers, aggressiveness, anxiety, depression, irritability, and interpersonal problems (Briere & Spinazzola, 2009; Kiesel, Conradi, Fehrenbach, Torgerson, & Briggs, 2014; Strand, Sarmiento, & Pasquale, 2005). During the assessment process it is important to be mindful of the child's stage of development, cognitive skills, race, age, and culture and how those factors may impact their emotional interpretation of traumatic events (Briere & Spinazzola, 2009; Nader, 2007).

Treatment

Brains, bodies, and lives are somewhat resilient. The use of effective treatment methods and the integration of resilience factors can help victims of trauma improve and lead productive lives (Strand, Hansen, & Courtney, 2013). Core components of effective trauma focused treatments should include psychoeducation about loss and trauma reminders, grief reactions, posttraumatic stress symptoms, instruction on emotional self-regulation, relapse prevention, safety, and adaptive routines, motivational interviewing, client advocacy, parent/child relationship, trauma narrative construction, and frequent evaluations of treatment plans (Courtois, 2004; Ford & Courtois, 2013; Strand, Hansen, & Courtney, 2013). Many empirically

supported trauma focused treatment methods have been established specifically for complex trauma¹⁶.

Trauma Affect Regulation Group Education and Therapy (TARGET) theoretical perspective is based on developmental trauma and cognitive behavioral therapy and focuses on trauma regulation, trauma processing, relationship repair, and social engagement using strength-based and resilience-enhancing approaches (Ford & Russo, 2006). TARGET also can incorporate historical trauma, spiritual beliefs, families with low socioeconomic statuses, and community norms (Ford & Russo, 2006). TARGET is ideal for juvenile justice, school, and child welfare settings (Agosti, Conradi, Halladay, & Langan, 2013; Brunzell, Waters, & Stokes, 2015; Ford & Blaustein, 2013).

Ford and Hawke (2012) tested the efficacy of TARGET in a juvenile justice setting. They found that TARGET and behavioral issues had a dose-response relationship; the more TARGET sessions the youth attended the fewer punitive and disciplinary actions were taken against them (Ford & Hawke, 2012). According to Ford and Hawke (2012), this outcome supported their hypothesis that teaching affect regulation positively contributed to the well-being and safety of the youth residing in juvenile detention. In a similar study, Marrow and colleagues found that the implementation of TARGET in juvenile detention facilities reduced depression and increased optimism and hope among the residents (Marrow, Knudsen, Olafson, & Bucher, 2012).

Structured Psychotherapy for Adolescents Responding to Chronic Stress (SPARCS) is a form of cognitive behavioral therapy that addresses issues such as dissociation, affect regulation, impulsivity, self-regulation, somatization, future orientation, and relationships (Brom, Pat-Horenczyk, & Ford, 2009). It is designed to specifically to address trauma related to physical

¹⁶ See Table 2

and sexual abuse and community violence that is ongoing by helping youth to cope through improved self-efficacy and establishing positive relationships (Habib & DeRosa, 2008).

SPARCS is culturally sensitive and incorporates “meaning making” – what trauma means in the context of cultural beliefs (Habib & DeRosa, 2008). SPARCS has successfully been implemented in schools, group homes, residential treatment facilities, outpatient settings, boarding school, outpatient clinics, foster care, homeless shelters, and juvenile justice facilities (Dorsey, Briggs, & Woods, 2011; Ford, Chapman, Connor, & Cruise, 2012; Weiner, Schneider, & Lyons, 2009; Zelechoski, et al., 2013).

The Mental Health Services and Policy Program of the Illinois Department of Children and Family Services conducted a pilot study of SPARCS involving youth ages 13-21 from two different child welfare agencies (Kisiel, Marett, Markiwicz, & Fehrenbach, 2008). Participation criteria included having at least one traumatic experiences categorized as moderate to severe and noticeable difficulties dealing with the trauma (Kisiel, Marett, Markiwicz, & Fehrenbach, 2008). After time-limited exposure to SPARCS, participants showed decreases in traumatic stress symptoms, emotional and behavioral issues, health risk behaviors and increases in positive life functioning and strengths (Kisiel, Marett, Markiwicz, & Fehrenbach, 2008).

Integrative Treatment of Complex Trauma (ITCT) for adolescents and children takes a multidimensional approach, employing several different models: attachment, cognitive behavioral, exposure, affect regulation, mindfulness, triggers, and psychoeducation (Lawson & Quinn, 2013). ITCT addresses physical, sexual, and emotional abuse, traumatic grief, community violence, and medical trauma (Lanktree & Briere, 2013). It also addresses issues related to social marginalization and poverty (Lanktree, et al., 2012). ITCT is appropriate for use in individual, family, or group settings, in hospitals, and inpatient and outpatient clinics and has

been especially effective in homeless children and youth in the juvenile justice system (Lanktree, et al., 2012).

Lanktree and colleagues (2012) examined ITCT effectiveness among youth living in high risk communities. They found that ITCT significantly reduced posttraumatic stress symptoms, anxiety, and depression and greatly reduced sexual issues, anger, and dissociation (Lanktree, et al., 2012). The effects of ITCT were dose-dependent; the more sessions the youth attended the higher the positive outcomes (Lanktree, et al., 2012). ITCT has also been found to be an effective treatment for substance abuse in traumatized adolescents and teens (Briere & Lanktree, 2014).

Attachment, Regulation, and Competency (ARC) therapy is based on attachment theory, child development, resilience factors, and addresses attachment, self-regulation, and social competency¹⁷ (Hodgdon, Kinniburgh, Gabowitz, Blaustein, & Spinazzola, 2013). It primarily targets early childhood trauma and ongoing adverse experiences (Kinniburgh, Blaustein, Spinazzola, & van der Kolk, 2005). ARC is unique in that considers contextual and cultural factors related to the child, the child's caregivers, and the social environment of the child (Kinniburgh, Blaustein, Spinazzola, & van der Kolk, 2005). The goal of ARC is to not only treat the child, but also develop a trauma informed milieu and systems that interact with the child (Kinniburgh, Blaustein, Spinazzola, & van der Kolk, 2005). ARC is appropriate for parent workshops, home based programs, individual, group, and family therapy (Hodgdon, Kinniburgh, Gabowitz, Blaustein, & Spinazzola, 2013).

¹⁷ Social competency is defined as one's ability to effectively communicate with others, to be amiable or congenial in social situations, and to appropriately view themselves in relation to others (Semrud-Clikeman, 2007).

A pilot study of the ARC framework was conducted in a residential facility that treated youth exposed to complex trauma (Hodgdon, Kinniburgh, Gabowitz, Blaustein, & Spinazzola, 2013). Significant reductions in internalizing and externalizing behaviors, posttraumatic stress symptoms, and use of restraints were observed (Hodgdon, Kinniburgh, Gabowitz, Blaustein, & Spinazzola, 2013). Another study of adopted children with a history of complex trauma found significant reductions in depression, anger, anxiety, dissociation, and internalizing and externalizing issues after 16 weeks of individual and group ARC treatments (Hodgdon, Blaustein, Kinniburgh, Peterson, & Spinazzola, 2016). The positive effects of ARC treatment were observed during a 12-month follow up assessment (Hodgdon, Blaustein, Kinniburgh, Peterson, & Spinazzola, 2016).

Trauma Systems Therapy (TST) is partially grounded in Bronfenbrenner's social-ecological model in that a disruption in one sphere can interfere with other spheres (Dorsey, Briggs, & Woods, 2011). Because TST involves exploring multiple spheres in the child's life, a multidisciplinary team is usually engaged in treatment (Saxe, Ellis, & Brown, 2015). The team may include important figures in the child's life such as spiritual leaders, teachers, and caseworkers (Saxe, Ellis, & Brown, 2015). Although usually applied in individual therapy settings, TST can be adapted to apply to group settings such as schools, families, and neighborhoods (Ellis, Fogler, Hansen, Forbes, & Navalta, 2012). TST is executed in three stages: safety focused treatment, regulation focused treatment, and beyond trauma treatment (Saxe, Ellis, & Kaplow, 2006).

Brown and colleagues tested the efficacy of TST in a Boston, MA residential treatment facility (Brown, McCauley, Navalta, & Saxe, 2013). They found that the regular implementation of TST resulted in a decrease in the use of physical restraints among residents (Brown,

McCauley, Navalta, & Saxe, 2013). They also found a significant increase in placement stability with children in foster care that had been exposed to complex trauma (Brown, McCauley, Navalta, & Saxe, 2013). Compared to an average of 3.4 placement moves, after TST, placement moves were reduced to an average of 1.4 (Brown, McCauley, Navalta, & Saxe, 2013).

Trauma Focused Cognitive Behavioral Therapy (TF-CBT) addresses common emotional and social issues related to trauma including posttraumatic stress disorder, attachment and relationship problems, academic problems, cognitive issues, depression, and anxiety (Deblinger, Mannarino, Cohen, Runyon, & Steer, 2011). It also addresses trauma processing, trust, safety, behavior, affect, and thought regulation, family communication, and parenting skills (Cohen & Mannarino, 2008; Deblinger, Mannarino, Cohen, Runyon, & Steer, 2011). TF-CBT is designed to for trauma related to sexual abuse, traumatic grief, domestic violence, terrorism, and natural disasters (Cohen & Mannarino, 2008). TF-CBT has been modified to work in public systems (e.g. child welfare, juvenile justice, and schools) as well as special populations (e.g. military) (Cohen & Mannarino, 2008; Cohen, Mannarino, Berliner, & Deblinger, 2000; Deblinger, Mannarino, Cohen, Runyon, & Steer, 2011).

The use of TF-CBT on young children between the ages of three to six who had been sexually abused yielded positive findings (Scheeringa, Weems, Cohen, Amaya-Jackson, & Guthrie, 2011). Posttraumatic stress symptoms were most significantly reduced followed by depression, anxiety, and oppositional defiant disorder even after a six month follow up (Scheeringa, Weems, Cohen, Amaya-Jackson, & Guthrie, 2011). Research also indicates that TF-CBT is useful in the treatment of posttraumatic stress disorder in children exposed to isolated traumatic events (March, Amaya-Jackson, Murray, & Schulte, 1998).

Real Life Heroes (RLF) is a resilience-based treatment that can be used in a variety of settings (Richardson, Kagan, Henry, Delorenzo, & Brophy, 2001). It specifically addresses abandonment, loss, domestic violence, disasters, medical trauma, neglect, and physical, emotional, and sexual abuse and is most suitable for complex posttraumatic stress (Kagan, 2007; Richardson, Kagan, Henry, Delorenzo, & Brophy, 2001). RLF is generally utilized in children who lack safe and stable home environments and healthy attachments to caregivers (Kagan, Douglas, Hornik, & Kratz, 2008). RLF employs several tools such as yoga, story-telling, mindfulness, multi-sensory and creative arts to restore, improve, or build relationships and affect regulation (Kagan, Douglas, Hornick, & Kratz, 2008; Kagan & Spinazzola, 2013).

A pilot study including 41 children exposed to household dysfunction, their caregivers, and clinicians assessed the effectiveness of Real Life Heroes (Kagan, Douglas, Hornik, & Kratz, 2008). After four months of exposure to the Real Life Heroes framework, children had significant reductions in behavioral issues (Kagan, Douglas, Hornik, & Kratz, 2008). After a year of treatment, children showed caregiver security and attachment improvements (Kagan, Douglas, Hornik, & Kratz, 2008).

Child-Parent Psychotherapy (CPP) integrates attachment theory, theories of development, social learning theory, psychodynamic theory, and cognitive behavioral theory to teach affect regulation, safety, relationships, and trauma responses (Lieberman, Ippen, & Van Horn, 2006; Lieberman, Van Horn, & Ippen, 2005). CPP is especially helpful when trauma has had an adverse effect on the child-parent relationship (Cicchetti, Rogosch, & Toth, 2006). It specifically addresses physical, emotional, and sexual abuse, neglect, and domestic violence (Lieberman, Ippen, & Van Horn, 2006; Lieberman, Van Horn, & Ippen, 2005). CPP has been successful utilized for treatment of refugee trauma (Cohen & Mannarino, 2010).

Many studies on CPP examined the role attachment played in childhood trauma. For example, Cicchetti, Rogosch, and Toth (2006) conducted a study on the effectiveness of CPP on mothers of emotionally neglected infants. In addition to CPP, other treatments were evaluated; psychoeducational parenting intervention and community standards and controls (Cicchetti, Rogosch, & Toth, 2006). After 12 months of treatment, mothers and infants engaged in CPP showed the most substantial increases in secure attachments (Cicchetti, Rogosch, & Toth, 2006). Findings were similar in a study involving toddlers (Cicchetti, Rogosch, & Toth, 2000; Cicchetti, Toth, & Rogosch, 1999).

Strengthening Family Coping Resources (SFCR) is whole family therapy intended to address ongoing stress and trauma such as community violence or poverty (Kiser, Donohue, Hodgkinson, & Medoff, 2010). The theoretical framework includes a multitude of approaches that can be tailored to meet family-specific needs such as family resilience theory, attachment theory, coping theory, family ritual and routine theory, and social support theory (Kiser, Donohue, Hodgkinson, & Medoff, 2010). SFCR nurtures protective factors like safety, social support and resources, crisis management, consistency, togetherness, and positive affect (Kiser, Donohue, Hodgkinson, & Medoff, 2010).

A study on the viability of SFCR among low-income, urban families found that the treatment was successful in a multifamily group format (Kiser, Donohue, Hodgkinson, Medoff, & Black, 2010). Children who were exposed to complex trauma saw reductions in posttraumatic symptoms, emotional and behavioral issues, re-experiencing, arousal symptoms, depression, anxiety, social problems, attention issues, and aggressive behaviors (Kiser, Donohue, Hodgkinson, Medoff, & Black, 2010). In a similar study on urban youth living in poverty, Kiser (2007) noted a key protective factor born of SFCR. She found that strengthening rituals and

routines within the family unit helped nurture structure and stability which reduced the risk of trauma symptoms in children living in violent communities (Kiser, 2007).

Future Considerations

A wealth of research on childhood trauma has been conducted since the ACE study. Trauma-focused assessment and treatments have been established (Steele, 2009). The mental/medical implication of ACEs has helped bridge the gap between mental health care and physical health care. However, professionals are still breaking ground on trauma assessments and treatments in special populations such as military, homeless, disabled, refugee, and lesbian/gay/bisexual/transgender (LGBT) youth (Rapporteur, et al., 2016). In addition, experts on childhood trauma are calling on child welfare and juvenile justice systems, and schools to consider the implications that childhood trauma has on health, behavior, and development by implementing trauma informed practices and protocols (Ko, et al., 2008).

Special Populations

Some of the most vulnerable members of society are often overlooked in trauma research. In addition to potential trauma exposures that all children face, military, homeless, disabled, refugee, and LGBT youth have unique challenges and are at greater risk for complex trauma than the general population (Alfano, Lau, Balderas, Bunnell, & Beidel, 2016; Bostwick, 2007; Buckner, Beardslee, & Bassuk, 2004; Chambon, 1989; Goldson, 2001). Pathways to quality interventions for special populations exposed to trauma include cultural sensitivity, better accessibility to resources, encouraging positive identities, and understanding unique risk and resilience factors (Sue, Arredondo, & McDavis, 1992).

Military. There are a number of important concepts that providers treating children of military members should be mindful of, but none so important as the fact that they are always

in transition (Hall, 2008). The average military child moves six to nine times between age five and 18 (Hall, 2008). Children in military families can experience a great deal of stress related to frequent moves, but most notably, children whose caregiver(s) deploy during wartime experience the highest amount of psychological stress (Lincoln, Swift, & Shorteno-Fraser, 2008). During the conflicts in Iraq and Afghanistan alone, two million children had a caregiver that deployed, 19,000 had a caregiver wounded while serving, and over 2,200 children lost a caregiver (Chartrand, Molinda, Frank, White, & Shope, 2008; Sogomonyan & Cooper, 2010).

Compared to civilian children, military children have higher rates of academic struggles, peer and family conflicts, and mental health problems (Alfano, Lau, Balderas, Bunnell, & Beidel, 2016). In addition, research has indicated an increase in child maltreatment rates among military children of a deployed parent (Alfano, Lau, Balderas, Bunnell, & Beidel, 2016). Lastly, a 2014 study found that adverse childhood experiences (ACEs) were more common in military members than their civilian counterparts and as the ACE study illustrated, ACEs are usually intergenerational (Blosnich, Dichter, Cerulli, Batten, & Bossarte, 2014; Felitti, et al., 1998). Research into military children and trauma is just breaking ground. Professionals are calling on research with more rigorous methodologies, more longitudinal studies than can measure the long-term effects of military life on children, research that examines a more extensive range of parental and contextual influences and objective measures of stress, and most importantly, an evaluation of resilience and adaptive factors that can help facilitate trauma-informed support services (Alfano, Lau, Balderas, Bunnell, & Beidel, 2016).

Homeless. There are approximately 2.5 million homeless children in the United States (Bassuk, DeCandia, Beach, & Berman, 2014). In addition to the stressors of homelessness such as poverty and shelter life, homeless children are often deprived of the most basic needs:

nutrition, adequate clothing, health and dental care, social interactions, and sometimes adequate education (Armstrong, 2009; Samuels, Shinn, & Buckner, 2010; Wadsworth et al., 2008). They are twice as likely to suffer physical illnesses, have a learning disability, and repeat a grade as other children (Buckner, Beardslee, & Bassuk, 2004).

By the time a homeless child turns eight, they have a one and three chance of developing a mental disorder (Goodman, Saxe, & Harvey, 1991). The high rates of mental disorders in homeless children are compounded by the fact that opportunities for mental health interventions are rare (Samuels, Shinn, & Buckner, 2010). In fact, more than one-fifth of preschool aged homeless children have emotional, behavioral, and developmental issues pervasive enough to require professional treatment, but less than one-third receive intervention (Burt, et al., 1999).

Shelters have generally been recognized as having volatile environments where children are often re-traumatized so it is important for shelter workers to be able to recognize trauma symptoms in children and be prepared to refer them to appropriate services (Burt, et al., 1999). Schools have been identified as places where homeless children find comfort, stability, and support. Studies indicate that school-based interventions have improved the development of positive relationships, self-regulation, and self-control (Goodwin & Miller, 2013; Swick, 2005). It is recommended that schools who have not adopted trauma-informed interventions for homeless children start by providing environments that promote safety and hope and most importantly provide a continuum of services (Swick, 2005).

Disabled. Children with disabilities are more likely to have adverse experiences in nearly every category of complex trauma than children without disabilities (Goldson, 2001). According to one study, children with disabilities were nearly 5% more likely to suffer physical abuse and nearly 2% more likely to suffer sexual abuse than their peers (Goldson, 2001). The

majority of physical abuse perpetrators are immediate caregivers while the majority of sexual abuse perpetrators are other family members (Sullivan & Knutson, 2000). Incidents of sexual abuse against disabled children are four times more common in institutions than at home (Blatt & Brown, 1986).

Individuals with disabilities are four times more likely to be victims of crime than individuals without disabilities (Petersilia, 2001). Unfortunately, crimes against the disabled are less likely to be investigated due to processing and communication problems (Avrin, Charlton, & Tallant, 1998; Senn, 1988). Exposure to trauma is more likely to cause developmental delays in disabled children than in children without disabilities because disabled children are predisposed to certain conditions like impaired resiliency and emotional problems (Burrows & Kochurka, 1995).

The assessment and delivery of treatment for disabled individuals requires special considerations due to potential communication impairments, impairments of abstract, fluid, and flexible thought, poor generalization skills, and a generally imperviousness to ambiguity (Avrin, Charlton, & Tallant, 1998). Specific communication and behavioral difficulties that may arise in clinical settings include misunderstandings of nonverbal communications and social cues, intrusive behaviors, idiosyncratic language, a tendency to ask many questions, especially rote questions, one-sided conversations, and an inability to measure the effect their behavior has on others (Avrin, Charlton, & Tallant, 1998). Mental health professionals are advised to use slow speech - presenting concepts one at a time, visual interpretations of their spoken messages, frequently ask for feedback to assess whether the client comprehends, be specific, and most importantly, involve caregivers in treatment (Avrin, Charlton, & Tallant, 1998).

Refugees. Refugee children are faced with a different set of challenges than naturalized children, but most times trauma occurs prior to fleeing their home country (Birman & Tran, 2008). Children may have experienced torture, malnutrition, property loss, physical assault, rape, and have likely witnessed a great deal of violence and death (Birman & Tran, 2008; Hollifield, et al., 2005). The flight process can also be very traumatic for children (Hollifield, et al., 2005). Children can also suffer post-migration trauma that experts consider similar or worse than war-related trauma (Heptinstall, Sethna, & Taylor, 2004; Silove, Sinnerbrink, Field, Manicavasagar, & Steel, 1997).

Common stressors for refugee children include issues related to resettlement, acculturation, and isolation (Chambon, 1989). Resettlement stress may include loss of social support, poverty, inadequate housing, and inaccessibility to resources (Chambon, 1989; Lustin, et al., 2004). Acculturation can be especially difficult for refugee children because usually they were forced to leave their home country (Fazel, Reed, Panter-Brick, & Stein, 2012). There may be language barriers that prevent them from forming relationships with peers and they have issues developing a healthy identity that encompasses both their old country and new country. Isolation can have the most adverse effects on refugee children (Fazel, Reed, Panter-Brick, & Stein, 2012). They can experience discrimination, harassment, and loss of social status (Fazel, Reed, Panter-Brick, & Stein, 2012).

Refugee children who experience trauma are likely to present with the same symptoms as other children; however, multicultural competencies are especially important when working with refugee children (Bala, Mooren, & Kramer, 2014). Culture can influence how refugees communicate their symptoms and what symptoms they report (Kleinman, 1988). Culture can also influence stigma, coping skills, social support, the type of help they seek, and even if they

seek help at all (Lustin, et al., 2004; Pumariega, Rothe, & Pumariega, 2005). It is also important to be mindful of culture-bound syndromes which are symptoms that may be more common in some cultures than others (Guarnaccia & Rogler, 1999). Lastly, due to the fact that caregivers also experienced trauma, whole family counseling is important (Chambon, 1989).

LGBT. LGBT youth are the most underrepresented in trauma research (Killen-Harvey, 2006). They are not only vulnerable to trauma that all youth may experience, but also are at increased risk for harassment, abuse, and rejection which can result in higher incidents of mental illness, suicide, and hate crimes (Bostwick, 2007; D'Augelli, Grossman, & Starks, 2006; House, Van Horn, Coppeans, & Stepleman, 2011). It is estimated that LGBT youth are 25% more likely to attempt suicide, are 84% more likely to have their life threatened, and 45% more likely to be assaulted than their peers (Bostwick, 2007; D'Augelli, Grossman, & Starks, 2006). They are also two times more likely to develop a mood or anxiety disorder, posttraumatic stress disorder, and body dysphoria disorder than their heterosexual peers (Bostwick, 2007). LGBT youth also miss considerably more days of school, are more likely to become homeless than their peers (due to being kicked out of the home), and are more likely to participate in health-risk behaviors like self-injury and drug/alcohol abuse (Bostwick, 2007; House, Van Horn, Coppeans, & Stepleman, 2011; Kann, et al., 2011).

Many mental health issues that LGBT youth experience are related to the process of developing and expressing their identity, therefore, it is important that professionals do not pressure them to “come out” (Greene, Bieschke, Perez, & DeBord, 2007). Key tenants for the successful treatment of LGBT youth include understanding the difference between being LGBT and same sex sexual encounters, inclusive terminology, providing a safe-space where the client can be vulnerable, develop competencies on sexual identity development, and connect the client

to resources where they can interact with other LGBT youth (Greene, Bieschke, Perez, & DeBord, 2007; Murphy, Rawlings, & Howe, 2002). It is also important that professionals working with LGBT youth acknowledge their client's struggles, isolation, and resilience (Greene, Bieschke, Perez, & DeBord, 2007).

Trauma-Informed Systems

A study by The National Center for Child Traumatic Stress identified several deficiencies in systems (e.g. child welfare, juvenile justice, and schools) that serve children likely exposed to trauma (Taylor & Siegfried, 2005). Of the agencies studied, most agencies did not conduct posttraumatic stress assessments on children who had experienced maltreatment (Taylor & Siegfried, 2005). Less than half of the agencies trained staff on evidence-based treatments for trauma exposure and over a third did no training on trauma assessment (Taylor & Siegfried, 2005). One third of the agencies studied never made referrals to treatment providers based on trauma assessments (Taylor & Siegfried, 2005). Interagency communication was mostly nonexistent; referring agencies seldom supplied in-depth information related to the child's history on trauma with schools being reported as the least helpful (Taylor & Siegfried, 2005).

The National Center for Child Traumatic Stress has devised a trauma-informed framework for systems working with children to consider (de Arellano, Ko, Danielson, & Sprague, 2008). These guidelines are applicable to child welfare, juvenile justice, and school systems (de Arellano, Ko, Danielson, & Sprague, 2008). Effective trauma-informed systems includes the following: routine trauma screenings, culturally competent evidence-based treatments, psychoeducation, strengthens protective and resilience factors, addresses caregiver trauma and its impact on children, and integration and continuity of care across childcare systems (de Arellano, Ko, Danielson, & Sprague, 2008).

Child Welfare. Many children placed in the child welfare system are there because they had already experienced some type of complex trauma (Pecora, Jensen, Romanelli, Jackson, & Ortiz, 2009). However, an overview of current child welfare systems indicate that few are trauma-informed with regards to training, performance, policy, assessment, and treatments (Agosti, Conradi, Halladay, & Langan, 2013). Effective trauma-informed child welfare practices include trauma knowledge building, trauma-focused assessments and treatments, case planning and management, identification of external trauma-informed services, and partnerships and collaboration with other systems (Agosti, Conradi, Halladay, & Langan, 2013).

Basic tenants of trauma knowledge building include staff training, staff coaching and support, trauma-based psychoeducation for caregivers including foster parents, and addressing secondary trauma that workers may experience when working with trauma victims (Agosti, Conradi, Halladay, & Langan, 2013). General guidelines for trauma-focused assessments and treatments include modifying assessments and treatments to meet the needs of trauma victims and sharing assessments with other providers such as pediatricians, therapists, and school counselors (Agosti, Conradi, Halladay, & Langan, 2013). Important practices involving case planning and management include providing birth parents with information about their child's placement in group homes or foster care, giving foster parents a comprehensive review of the child's trauma history, facilitating healthy relationships between foster parents and biological parents, and conducting team meetings that include all individuals directly involved in the child's life (Agosti, Conradi, Halladay, & Langan, 2013).

Examples of identifying external trauma-informed services include increasing in-house access to trauma-focused treatments and a comprehensive list of trauma-informed services in the event a child needs to be referred to a practice outside of the child welfare system (Agosti,

Conradi, Halladay, & Langan, 2013). Lastly, partnerships with other systems can create more opportunities for training (Agosti, Conradi, Halladay, & Langan, 2013). In addition, partnerships can help systems that generally do not engage in mental health treatments (such as court systems) learn about childhood trauma (Agosti, Conradi, Halladay, & Langan, 2013).

Juvenile Justice. Trauma-informed procedures and policies can make juvenile justice systems more effective by providing physical and psychological safety for all parties involved (Dierkhising, Ko, & Goldman, 2013). Trauma-informed juvenile justice systems support recovery, prevents further traumatization, and recognizes the effects that trauma has on behavior (Dierkhising, Ko, & Goldman, 2013). It is recommended that juvenile justice systems avoid practices that may further aggravate trauma symptoms such as coercion, shackling, and cuffing (Dierkhising, Ko, & Goldman, 2013). They should mandate staff psychoeducation on coping strategies and triggers (Dierkhising, Ko, & Goldman, 2013).

Children in the juvenile justice system should be afforded trauma-informed legal representation (Dierkhising, Ko, & Goldman, 2013). Effective juvenile justice systems should also provide trauma-specific treatment while considering the age, gender identification, sex, socioeconomic status, cognitive and emotional development, and culture (Dierkhising, Ko, & Goldman, 2013). Lastly, they should ensure collaboration with other agencies (Dierkhising, Ko, & Goldman, 2013).

Schools. Trauma can impact a child's ability to learn and to function effectively in a classroom environment (National Child Traumatic Stress Network Schools Committee, 2008). Often times school is a child's only safe environment where they can build relationships with compassionate individuals who can nurture their success (Cole, et al., 2006). The Flexible Framework is an outline by which to assist schools in constructing a trauma-informed and

trauma-sensitive environment (Cole, et al., 2006). The Flexible Framework includes six key elements: school wide infrastructure and culture, teacher and staff training, coordination with mental health professionals, and trauma-sensitive non-academic strategies, protocols, procedures, and policies (Cole, et al., 2006).

Educators can also help children cope with trauma by maintaining consistency, allowing students to make choices when appropriate (sometimes children feel a loss of control after a traumatic event), set limits and assert logical consequences for rule breaking, recognize trauma related behavioral issues, increase encouragement and support, be sensitive to potential triggers including anniversaries of traumatic events, and be mindful of other student's reactions to traumatized classmates (National Child Traumatic Stress Network Schools Committee, 2008). Educators can also provide safe spaces for children to discuss what has happened to them (National Child Traumatic Stress Network Schools Committee, 2008). Children should be given simple answers and educators should clarify misconceptions (National Child Traumatic Stress Network Schools Committee, 2008). Lastly, children exposed to traumatic events or living in a toxic environment may need modifications in homework such as shortened assignments and after school tutoring (National Child Traumatic Stress Network Schools Committee, 2008).

Conclusion

Since the Adverse Childhood Experiences Study (ACES), researchers have amassed in-depth studies into the consequences of toxic stress. Despite the wealth of knowledge available, high ACE scores continue to be a world-health problem, but there are success stories (World Health Organization, 2009). Arizona has established an ACE Consortium which has embarked on a successful public campaign (Stevens, 2014). Since its inception in 2006 the ACE Consortium has trained over 450 community members who offer community presentations and

workshops on ACEs (Stevens, 2014). Their first presentation reached 13,000 Arizonians (Stevens, 2014). They also offer an easy-to-understand webpage and downloadable presentations (Stevens, 2014). Because of the ACE Consortium, Arizona's largest healthcare provider now includes routine trauma screenings for families and the Arizona Department of Health Services now includes the ACE questionnaire in their Behavioral Risk Factor Surveillance System (Stevens, 2014).

Beginning in 2011, Iowa has conducted two ACE surveys with one underway and another scheduled (Stevens, 2014). The state has also hosted two ACE summits (Stevens, 2014). Walla Walla, Washington has 42 community networks working to educate the community on the consequences of ACEs (Stevens, 2014). They also launched the Children's Resilience Initiative in 2010, which raises awareness on brain development and ACEs (Stevens, 2014). Lincoln High School in Walla Walla has gained national attention for their implementation of trauma-informed training for educators (Stevens, 2014).

Inspired by the ACE study, Father Jeff Putthoff, and Dr. Jeffrey Brenner of Camden, NJ found ways to help their community utilizing their unique skill sets (Hochman, 2014). Father Putthoff founded Hopework N' Camden which is a trauma-informed GED provider and offers skills training to help youth form pathways to college and/or employment (Hochman, 2014). In 2013, Father Putthoff also hosted a trauma summit which was attended by community activists, law enforcement, clergy, and educators (Hochman, 2014). He is also working on a collaborative trauma-focused effort called Healing 10 (Hochman, 2014).

Dr. Brenner, a Camden physician, changed his entire practice to reflect biomedical information learned from the ACE study and other trauma related research (Hochman, 2014). As executive director of the Camden Coalition of Healthcare Providers (CCHP), he has

implemented routine trauma screenings in medical offices around Camden. In 2014 Father Putthoff joined forces with Dr. Brenner to train CCHP AmeriCorps volunteers (Hochman, 2014).

The implications of the ACE study continue to infiltrate communities all over the United States from trauma education for professionals in and around Philadelphia to trauma-sensitive judicial systems in Florida to the first trauma-informed city of Tarpon Springs, FL to a trauma-informed NFL (Finkel, 2014; Hochman, 2014; Stevens, 2014). Policy makers are already incorporating findings of the ACE study into legislation. Vermont has proposed a bill that will require health providers to perform ACE screenings (Till, 2014). Washington State passed legislation which developed private partnerships to further trauma-informed and resilience-building practices (ACEs Public-Private Initiative, n.d.). In 2014, California proposed a bill aimed at lowering adverse childhood experiences (Bocanegra, Bonta, Bradford, Buchanan, & Calderon, 2014). Though progress may be slow, these success stories and legislative efforts are proof that moving towards a trauma-sensitive society is critical to public health.

References

- Abidin, R. R. (1995). *Parenting Stress Index, third edition: Professional manual*. Odessa, FL: Psychological Assessment Resources.
- ACEs Public-Private Initiative. (n.d.). *The Washington State ACEs*. Seattle, WA: ACEs Public-Private Initiative. Retrieved from <http://www.appi-wa.org/>
- Achenbach, T. M. (2011). Child Behavior Checklist. *Encyclopedia of Clinical Neuropsychology*, 1, 546-552. doi:10.1007/978-0-387-79948-3_1529
- Agosti, J., Conradi, L., Halladay, G. J., & Langan, H. (2013). *Using trauma-informed child welfare practice to improve placement stability breakthrough series collaborative: Promising practices and lessons learned*. Los Angeles, CA and Durham, NC: National Center for Child Traumatic Stress. Retrieved from http://www.nctsn.org/sites/default/files/assets/pdfs/using_ticw_bsc_final.pdf
- Agosti, J., Conradi, L., Halladay, G. J., & Langan, H. (2013). *Using trauma-informed child welfare practice to improve placement stability breakthrough series collaborative: Promising practices and lessons learned*. Los Angeles, CA & Durham, NC: National Center for Child Traumatic Stress. Retrieved from http://www.nctsn.org/sites/default/files/assets/pdfs/using_ticw_bsc_final.pdf
- Ainsworth, M. D. (1979). Attachment as related to mother-infant interaction. *Advances in the Study of Behavior*, 9, 1-51. doi:10.1016/S0065-3454(08)60032-7
- Alfano, C. A., Lau, S., Balderas, J., Bunnell, B. E., & Beidel, D. C. (2016). The impact of military deployment on children: Placing developmental risk in context. *Clinical Psychology Review*, 43, 17-29. doi:10.1016/j.cpr.2015.11.003

- Amato, P. R. (2000). The consequences of divorce for adults and children. *Journal of Marriage and Family*, 62(4), 1269–1287. doi:10.1111/j.1741-3737.2000.01269.x
- Anda, R. (2011, January 12). *Adverse childhood experiences: Connecting a developmental lens to the health of society* [PowerPoint slides]. Retrieved from https://www.in.gov/dcs/files/2Dr_Anda5steps.pdf
- Anda, R. F., Brown, D. W., Dube, S. R., Bremner, J. D., Felitti, V. J., & Giles, W. H. (2008). Adverse childhood experiences and chronic obstructive pulmonary disease in adults. *American Journal of Preventive Medicine*, 34(5), 396-403. doi:10.1016/j.amepre.2008.02.002
- Anda, R. F., Brown, D. W., Felitti, V. J., Dube, S. R., & Giles, W. H. (2008). Adverse childhood experiences and prescription drug use in a cohort study of adult HMO patients. *BMC Public Health*, 8(1), 198. doi:10.1186/1471-2458-8-198
- Anda, R. F., Croft, J. B., Felitti, V. J., Nordenberg, D., Giles, W. H., Williamson, D. F., & Giovino, G. A. (1999). Adverse childhood experiences and smoking during adolescence and adulthood. *Journal of the American Medical Association*, 282(17), 1652-1658. doi:10.1001/jama.282.17.1652
- Anda, R. F., Fleisher, V. I., Felitti, V. J., Edwards, V. J., Whitfield, C. L., Dube, S. R., & Williamson, D. F. (2004). Childhood abuse, household dysfunction, and indicators of impaired worker performance. *The Permanente Journal*, 8(1), 30-38. Retrieved from <http://www.thepermanentejournal.org/files/Winter2004/childhood.pdf>
- Anda, R., Tietjen, G., Schulman, E., Felitti, V., & Croft, J. (2010). Adverse childhood experiences and frequent headaches in adults. *Headache*, 50(9), 1473-1481. doi:10.1111/j.1526-4610.2010.01756.x

- Armsden, G. C., & Greenberg, M. T. (1987). The Inventory of Parent and Peer Attachment: Relationships to well-being in adolescence. *Journal of Youth and Adolescence*, 16(5), 427-454. doi:10.1007/BF02202939
- Armstrong, A. (2009). *Myths of poverty: Realities for students*. Springfield, IL: The Illinois School Board Journal. doi:https://www.iasb.com/journal/j111209_02.cfm
- Armstrong, J. G., Putnam, F. W., Carlson, E. B., Libero, D. Z., & Smith, S. R. (1997). Development and validation of a measure of adolescent dissociation: The Adolescent Dissociative Experiences Scale. *Journal of Nervous and Mental Disorders*, 185(8), 491-497. doi:10.1097/00005053-199708000-00003
- Avrin, S., Charlton, M., & Tallant, B. (1998). *Diagnosis and treatment of clients with developmental disabilities*. Aurora, CO: Aurora Mental Health Center.
- Ayers, T. S., Sandler, I. N., West, S. G., & Roosa, M. W. (1996). A dispositional and situational assessment of children's coping: Testing alternative models of coping. *Journal of Personality*, 64(4), 923-958. doi:10.1111/j.1467-6494.1996.tb00949.x
- Baglivio, M. T., Epps, N., Swartz, K., Huq, M. S., Sheer, A., & Hardt, N. S. (2014). The prevalence of adverse childhood experiences (ACE) in the lives of juvenile offenders. *Journal of Juvenile Justice*, 3(2), 1-14. Retrieved from <http://www.journalofjuvjustice.org/JOJJ0302/JOJJ0302.pdf>
- Baker, L. L., Jaffe, P. G., Ashbourne, L., & Carter, J. (2002). *Children exposed to domestic violence: An early childhood educator's handbook to increase understanding and improve community responses*. London, Ontario: Centre for Children and Families in the Justice System. Retrieved from <http://www.lfcc.on.ca/ece-us.PDF>

- Bala, J., Mooren, T., & Kramer, S. A. (2014). Cultural competence in the treatment of political refugees based on system approaches. *Clinical Neuropsychiatry*, 11(1), 32-39. Retrieved from <http://www.clinicalneuropsychiatry.org/pdf/balaweb.pdf>
- Barile, J. P., Edwards, V. J., Dhingra, S. S., & Thompson, W. W. (2015). Associations among county-level social determinants of health, child maltreatment, and emotional support on health-related quality of life in adulthood. *Psychology of Violence*, 5(2), 183-191. doi:10.1037/a0038202
- Barth, R. P., Scarborough, A. A., Lloyd, E. C., Losby, J. L., Casanueva, C., & Mann, T. (2008). *Developmental status and early intervention service needs of maltreated children*. Washington, DC: Institute for Social and Economic Development. Retrieved from <http://files.eric.ed.gov/fulltext/ED501753.pdf>
- Basso, D. (2008). Spatial navigation. *Cognitive Processing*, 9(4), 227–228. doi:10.1007/s10339-008-0224-0
- Bassuk, E. L., DeCandia, C. J., Beach, C. A., & Berman, F. (2014). *America's youngest outcasts: A report card on child homelessness*. Waltham, MA: American Institutes for Research and the National Center on Family Homelessness. Retrieved from <http://www.air.org/sites/default/files/downloads/report/Americas-Youngest-Outcasts-Child-Homelessness-Nov2014.pdf>
- Battaglini, L. (1987). Family empowerment through self-help groups. In A. B. Hatfield (Ed.), *Families of the mentally ill: Meeting the challenges* (pp. 43-52). San Francisco: Josey-Bass.
- Baumeister, R. F., & Vohs, K. D. (2004). *Handbook of self-regulation: Research, theory, and applications*. New York, NY: Guilford Press.

- Baunach, P. J. (1985). *Mothers in prison*. Brunswick, NJ: Transaction Books.
- Becker, D., & Margolin, F. (1967). How surviving parents handled their young children's adaptation to the crisis of loss. *American Journal of Orthopsychiatry*, 37(4), 753–757. doi:10.1111/j.1939-0025.1967.tb00517.x
- Bell, C., & Jenkins, E. (1991). Traumatic stress and children. *Journal of Health Care for the Poor and Underserved*, 2(1), 175-185. Retrieved from 10.1353/hpu.2010.0089
- Bellis, M. A., Hughes, K., Leckenby, N., Perkins, C., & Lowey, H. (2014). National household survey of adverse childhood experiences and their relationship with resilience to health-harming behaviors in England. *BMC Medicine*, 12(72), 1-10. doi:10.1186/1741-7015-12-72
- Birman, D., & Tran, N. (2008). Psychological distress and adjustment of Vietnamese refugees in the United States: Association with pre- and postmigration factors. *The American Journal of Orthopsychiatry*, 78(1), 109-120. doi:10.1037/0002-9432.78.1.109
- Blatt, E. R., & Brown, S. W. (1986). Environmental influences on incidents of alleged child abuse and neglect in New York state psychiatric facilities: Toward an etiology of institutional child maltreatment. *Child Abuse and Neglect*, 10(2), 171-180. doi:10.1016/0145-2134(86)90078-5
- Blodgett, C. (2012). *Adopting ACEs screening and assessment in in child serving systems*. Pullman, WA: Washington State University. Retrieved from <http://ext100.wsu.edu/cafru/wp-content/uploads/sites/65/2015/03/Complex-Trauma-Research-ACE-Screening-and-Assessment-in-Child-Serving-Systems-7-12-final.pdf>
- Blodgett, C., Harrington, R., Lohan, J., Short, R., Turner, N., & Winikoff, J. (2012). *Adverse childhood experience and developmental risk in elementary schoolchildren*. Pullman,

- WA: Washington State University. Retrieved from <http://ext100.wsu.edu/cafru/wp-content/uploads/sites/65/2015/02/Adverse-Childhood-Experience-and-Developmental-Risk-in-Elementary-Schoolchildren-Research-Briefx.pdf>
- Bloom, B., & Steinhart, D. (1992). *Why punish the children? A reappraisal of the children of incarcerated mothers in America*. New York, NY: National Council on Crime and Delinquency. Retrieved from http://www.nccdglobal.org/sites/default/files/publication_pdf/why-punish-the-children.pdf
- Blosnich, J. R., Dichter, M. E., Cerulli, C., Batten, S. V., & Bossarte, R. M. (2014). Disparities in adverse childhood experiences among individuals with a history of military service. *JAMA Psychiatry*, 71(9), 1041-1048. doi:10.1001/jamapsychiatry.2014.724
- Bluhm, R. L., Williamson, P. C., Osuch, E. A., Frewen, P. A., & Stevens, T. K. (2009). Alterations in default network connectivity in posttraumatic stress disorder related to early-life trauma. *Journal of Psychiatry and Neuroscience*, 34(3), 187-194. Retrieved from <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2674971/?tool=pmcentrez>
- Bocanegra, R., Bonta, R., Bradford, S., Buchanan, J. T., & Calderon, I. (2014). *Assembly concurrent resolution no. 155—Relative to childhood brain development*. Sacramento, CA: California Legislature. Retrieved from http://www.leginfo.ca.gov/pub/13-14/bill/asm/ab_0151-0200/acr_155_bill_20140528_introduced.htm
- Borse, N. N., Gilchrist, J., Dellinger, A. M., Rudd, R. A., Ballesteros, M. F., & Sleet, D. A. (2008). *CDC childhood injury report*. Washington, DC: Centers for Disease Control and Prevention. Retrieved from <https://www.cdc.gov/safechild/images/CDC-childhoodinjury.pdf>

- Bostwick, W. B. (2007). *Mental health issues among gay, lesbian, bisexual, and transgender (GLBT) people*. Arlington, VA: National Alliance on Mental Illness. Retrieved from [http://www.namainys.org/images/uploads/pdfs/Mental%20Health%20Issues%20among%20Gay,%20Lesbian,%20Bisexual,%20and%20Transgender%20\(GLBT\)%20People.pdf](http://www.namainys.org/images/uploads/pdfs/Mental%20Health%20Issues%20among%20Gay,%20Lesbian,%20Bisexual,%20and%20Transgender%20(GLBT)%20People.pdf)
- Bradley, R. H., & Corwyn, R. F. (2002). Socioeconomic status and child development. *Annual Review of Psychology*, 53, 371-399. doi:10.1146/annurev.psych.53.100901.135233
- Briere, J. (1996). *Trauma Symptom Checklist for Children (TSCC), Professional manual*. Odessa, FL: Psychological Assessment Resources.
- Briere, J., & Lanktree, C. B. (2014). *Treating substance use issues in traumatized adolescents and young adults: Key principles and components*. Los Angeles, CA: USC Adolescent Trauma Training Center. Retrieved from <http://keck.usc.edu/adolescent-trauma-training-center/wp-content/uploads/sites/169/2015/04/Treating-Substance-Use-Issues-20131106.pdf>
- Briere, J., & Spinazzola, J. (2009). Assessment of the sequelae of complex trauma. In J. Courtois, & J. Ford (Eds.), *Treating complex traumatic stress disorders* (pp. 104-123). New York, NY: Guilford Press.
- Briere, J., Johnson, K., Bissada, A., Damon, L., Crouch, J., Gil, E., . . . Ernst, V. (2001). The Trauma Symptom Checklist for Young Children (TSCYC): Reliability and association with abuse exposure in a multi-site study. *Child Abuse and Neglect*, 25(8), 1001–1014. doi:10.1016/S0145-2134(01)00253-8
- Briggs, E. S., & Price, I. R. (2009). The relationship between adverse childhood experiences and obsessive-compulsive symptoms and beliefs: The role of anxiety, depression, and

- experiential avoidance. *Journal of Anxiety Disorders*, 23(8), 1037–1046.
doi:10.1016/j.janxdis.2009.07.004
- Brom, D., Pat-Horenczyk, R., & Ford, J. D. (2009). *Treating traumatized children: Risk, resilience and recovery*. New York, NY: Routledge.
- Bronfman, E. T., Biron Campis, L., & Koocher, G. P. (1998). Helping children to cope: Clinical issues for acutely injured and medically traumatized children. *Professional Psychology, Research and Practice*, 29(6), 574-581. doi:10.1037/0735-7028.29.6.574
- Brown, A. D., McCauley, K., Navalta, C. P., & Saxe, G. N. (2013). Trauma Systems Therapy for children in residential care: Improving emotion regulation and the social environment of traumatized children and youth in congregate care. *Journal of Family Violence*, 28(7), 693-703. doi:10.1007/s10896-013-9542-9
- Brown, D. W., Anda, R. F., Edwards, V. J., Felitti, V. J., Dube, S. R., & Giles, W. H. (2007). Adverse childhood experiences and childhood autobiographical memory disturbance. *Child Abuse and Neglect*, 9, 961-969. doi:10.1016/j.chiabu.2007.02.011
- Brown, D. W., Anda, R. F., Felitti, V. J., Edward, V. J., Malarcher, A. M., Croft, J. B., & Giles, W. H. (2010). Adverse childhood experiences and the risk of lung cancer. *BMC Public Health*, 10(20). doi:10.1186/1471-2458-10-20
- Brown, D. W., Anda, R. F., Tiemeier, H., Felitti, V. J., Edwards, V. J., Croft, J. B., & Giles, W. H. (2009). Adverse childhood experiences and the risk of premature mortality. *American Journal of Preventive Medicine*, 37(5), 389–396. doi:10.1016/j.amepre.2009.06.021
- Brown, N. M., Brown, S. N., Briggs, R. D., German, M., Belamarich, P. F., & Oyeku, S. O. (2016). Associations between adverse childhood experiences and ADHD diagnosis and severity. *Academic Pediatrics*, 1-7. Retrieved from

http://www.academicpedsjnl.net/pb/assets/raw/Health%20Advance/journals/acap/ACAP_900_approved.pdf

Browne, A., & Finkelhor, D. (1986). Impact of child sexual abuse: A review of the research.

Psychological Bulletin, 99(1), 66-77. doi:10.1037/0033-2909.99.1.66

Brunzell, T., Waters, L., & Stokes, H. (2015). Teaching with strengths in trauma-affected

students: A new approach to healing and growth in the classroom. *American Journal of Orthopsychiatry*, 85(1), 3-9. doi:10.1037/ort0000048

Buckner, J. C., Beardslee, W. R., & Bassuk, E. L. (2004). Exposure to violence and low-income

children's mental health: Directed, moderated, and mediated relations. *American Journal of Orthopsychiatry*, 74(4), 413-423. doi:10.1037/0002-9432.74.4.413

Burrows, H. C., & Kochurka, K. A. (1995). The assessment of children with disabilities who

report sexual abuse: A special look at those most vulnerable. In T. Ney (Ed.), *True and false allegations of child sexual abuse: Assessment and case management* (pp. 275-289).

New York, NY: Brunner/Mazel.

Burt, M. A., Aron, L. Y., Douglas, T., Valente, J., Lee, E., & Iwen, B. (1999). *Homelessness:*

Programs and the people they serve - Findings of the national survey of homeless assistance providers and clients. Washington, DC: Interagency Council on the Homeless.

Caldwell, P. H., Hodson, E., Craig, J. C., & Edgar, D. (2005). Bedwetting and toileting problems

in children. *The Medical Journal of Australia*, 182(4), 190-195. Retrieved from

<https://www.mja.com.au/journal/2005/182/4/4-bedwetting-and-toileting-problems-children>

Carlson, B. E. (1984). Children's observations of interpersonal violence. In A. R. Edwards (Ed.),

Battered women and their families (pp. 147-167). New York, NY: Springer.

Centers for Disease Control and Prevention. (2016). *About the CDC-Kaiser ACE study: Major findings*. Retrieved October 22, 2016, from

<https://www.cdc.gov/violenceprevention/acestudy/about.html>

Centers for Disease Control and Prevention. (n.d.). *Behavioral Risk Factor Surveillance System: About BRFSS*. Retrieved November 23, 2016, from

<http://www.cdc.gov/brfss/about/index.htm>

Centers for Disease Control and Prevention. (n.d.). *The ACE Pyramid*. Retrieved November 23, 2016, from <https://www.cdc.gov/violenceprevention/acestudy/about.html>

Chambon, A. (1989). Refugee families' experiences: Three family themes - Family disruption, violent trauma and acculturation. *Journal of Systematic Therapies*, 8.

doi:10.1521/jsst.1989.8.bonus.3

Chapman, D. P., Anda, R. F., Felitti, V. J., Dube, S. R., Edwards, V. J., & Whitfield, C. L.

(2004). Adverse childhood experiences and the risk of depressive disorders in adulthood.

Journal of Affective Disorders, 82(2), 217–225. doi:10.1016/j.jad.2003.12.013

Chartrand, M. M., Molinda, M., Frank, D. A., White, L. F., & Shope, T. R. (2008). Effect of parents' wartime deployment on the behavior of young children in military families.

Archives of Pediatric and Adolescent Medicine, 162(11), 1009-1014.

doi:10.1001/archpedi.162.11.1009

Chasnoff, I. J., & Lowder, L. A. (1999). Prenatal AOD use and risk for maltreatment: A timely

approach for intervention. In H. Dubowitz (Ed.), *Neglected children: Research, practice, and policy* (pp. 132–155). Thousand Oaks, CA: Sage.

Child Trends. (2013). *Adverse experiences: Indicators on children and youth*. Bethesda, MD:

Child Trends Databank. Retrieved from http://www.childtrends.org/wp-content/uploads/2013/07/124_Adverse_Experiences.pdf

Child Trends. (2014). *Violent crime victimization*. Bethesda, MD: Child Trends Databank.

Retrieved from http://www.childtrends.org/indicators/violent-crime-victimization/#_edn2

Cicchetti, D., Toth, S. L., & Rogosch, F. A. (1999). The efficacy of toddler-parent psychotherapy to increase attachment security in offspring of depressed mothers.

Attachment and Human Development, 1(1), 34-66. doi:10.1080/14616739900134021

Cicchetti, D., Rogosch, F. A., & Toth, S. L. (2000). The efficacy of toddler-parent psychotherapy for fostering cognitive development in offspring of depressed mothers. *Journal of*

Abnormal Child Psychology, 28(2), 135-148. doi:10.1023/A:1005118713814

Cicchetti, D., Rogosch, F. A., & Toth, S. L. (2006). Fostering secure attachment in infants in maltreating families through preventive interventions. *Development and*

Psychopathology, 18(3), 623-649. doi:10.1017/S0954579406060329

Cloitre, M., Stolbach, B. C., Herman, J. L., van der Kolk, B., Pynoos, R., Wang, J., & Petkova, E. (2009). A developmental approach to complex PTSD: Childhood and adult cumulative trauma as predictors of symptom complexity. *Journal of Traumatic Stress*, 22(5), 1-10.

doi:10.1002/jts.20444

Cohen, J. A., & Mannarino, A. P. (1996). Factors that mediate treatment outcome of sexually abused preschool children. *Journal of the American Academy of Child and Adolescent*

Psychiatry, 35(10), 1402-1410. doi:10.1097/00004583-199610000-00028

Cohen, J. A., & Mannarino, A. P. (2008). Trauma-focused Cognitive Behavioural therapy for children and parents. *Child and Adolescent Mental Health*, 13(4), 158–162.

doi:10.1111/j.1475-3588.2008.00502.x

Cohen, J. A., & Mannarino, A. P. (2010). Psychotherapeutic options for traumatized children.

Current Opinion in Pediatrics, 22(5), 605–609. doi:10.1097/MOP.0b013e32833e14a2

Cohen, J. A., Deblinger, E., Mannarino, A. P., & Steer, R. A. (2004). A multisite, randomized controlled trial for children with sexual abuse – Related PTSD symptoms. *Journal of the American Academy of Child and Adolescent Psychiatry*, 43(4), 393–402.

doi:10.1097/00004583-200404000-00005

Cohen, J. A., Mannarino, A. P., & Deblinger, E. (2006). *Treating trauma and traumatic grief in children and adolescents*. New York, NY: Guilford Press.

Cohen, J. A., Mannarino, A. P., & Knudsen, K. (2004). Treating childhood traumatic grief: A pilot study. *Child and Adolescent Psychiatry*, 43(10), 1225–1233.

doi:10.1097/01.chi.0000135620.15522.38

Cohen, J. A., Mannarino, A. P., Berliner, L., & Deblinger, E. (2000). Trauma-focused Cognitive Behavioral Therapy for children and adolescents: An empirical update. *Journal of Interpersonal Violence*, 15(11), 1202–1223. doi:10.1177/088626000015011007

Cohen, J. A., Mannarino, A. P., Greenberg, T., Padlo, S., & Shipley, C. (2002). Childhood traumatic grief: Concepts and controversies. *Trauma, Violence, & Abuse*, 3(4), 307–327.

doi:10.1177/1524838002237332

Cole, S. F., O'Brien, J. G., Gadd, M. G., Ristuccia, J., Wallace, D. L., & Gregory, M. (2006).

Helping traumatized children learn. Boston, MA: Massachusetts Advocates for Children.

Retrieved from <https://traumasensitiveschools.org/wp-content/uploads/2013/06/Helping-Traumatized-Children-Learn.pdf>

Cook, A., Blaustein, M., Spinazzola, J., & van der Kolk, B. (2003). *Complex trauma in children and adolescents*. Los Angeles, CA and Durham, NC: The National Child Traumatic Stress Network . Retrieved from

http://www.nctsn.org/nctsn_assets/pdfs/edu_materials/ComplexTrauma_All.pdf

Cooley-Quille, M. R., Turner, S. M., & Beidel, D. C. (1995). Emotional impact of children's exposure to community violence: A preliminary study. *Journal of the American Academy of Child and Adolescent Psychiatry*, 34(10), 1362-1368. doi:10.1097/00004583-199510000-00022

Copeland, W. E., Keeler, G., Angold, A., & Costello, E. J. (2007). Traumatic events and posttraumatic stress in childhood. *Archives of General Psychiatry*, 64(5), 577-584. doi:10.1001/archpsyc.64.5.577

Courtois, C. A. (2004). Complex trauma, complex reactions: Assessment and treatment. *Psychotherapy: Theory, Research, Practice, Training*, 41(4), 412-425. doi:10.1037/0033-3204.41.4.412

Cox, R. G., Zhang, L., Johnson, W. D., & Bender, D. R. (2007). Academic performance and substance use: Findings from a state survey of public high school students. *Journal of School Health*, 77(3), 109–115. doi:10.1111/j.1746-1561.2007.00179.x

Craven, S., Brown, S., & Gilchrist, E. (2006). Sexual grooming of children: Review of literature and theoretical considerations. *Journal of Sexual Aggression*, 12(3), 287-299. doi:10.1080/13552600601069414

- Crenshaw, D. A., Stella, L., O'Neill-Stephens, E., & Walsen, C. (2016). Developmentally and trauma-sensitive courtrooms. *Journal of Humanistic Psychology*.
doi:10.1177/0022167816641854
- Crosson-Tower, C. (2003). *The role of educators in preventing and responding to child abuse and neglect*. Washington, DC: U.S. Department of Health and Human Services.
Retrieved from <https://secure.ce-credit.com/articles/101012/101012.pdf>
- Cunningham, C. L., Martínez-Cerdeño, V., & Noctor, S. C. (2013). Microglia regulate the number of neural precursor cells in the developing cerebral cortex. *Journal of Neuroscience*, 33(10), 4216-4233. doi:10.1523/JNEUROSCI.3441-12
- Cutter, F. (1967). Psychological myopia. *Journal of Clinical Psychology*, 23(1), 118-119.
doi:10.1002/1097
- Dallam, S. J. (2001). The long-term medical consequences of childhood maltreatment. In K. Franey, R. Geffner, & R. Falconer (Eds.), *The cost of child maltreatment: Who pays? We all do* (pp. 1-14). San Diego, CA: Family Violence and Sexual Assault Institute.
- Danese, A., Moffitt, T. E., Harrington, H., Milne, B. J., Polanczyk, G., Pariante, C. M., . . . Caspi, A. (2009). Adverse childhood experiences and adult risk factors for age-related disease. *Archives of Pediatrics and Adolescent Medicine*, 163(12), 1135-1143.
doi:10.1001/archpediatrics.2009.214
- Daniels, J. K., Frewen, P., McKinnon, M. C., & Lanius, R. A. (2011). Default mode alterations in posttraumatic stress disorder related to early-life trauma: A developmental perspective. *Journal of Psychiatry and Neuroscience*, 36(1), 56-59. doi:10.1503/jpn.100050

- D'Augelli, A. R., Grossman, A. R., & Starks, M. T. (2006). Childhood gender atypicality, victimization, and PTSD among lesbian, gay, and bisexual youth. *Journal of Interpersonal Violence, 21*(11), 1462-1482. doi:10.1177/0886260506293482
- de Arellano, M. A., Ko, S. J., Danielson, C. K., & Sprague, C. M. (2008). *Trauma-informed interventions: Clinical and research evidence and culture-specific information project*. Los Angeles, CA and Durham, NC: National Center for Child Traumatic Stress. Retrieved from http://www.nctsn.org/sites/default/files/assets/pdfs/CCG_Book.pdf
- De Pisapia, N., Serra, M., Rigo, P., Jager, J., Papinutto, N., Esposito, G., . . . Bornstein, M. H. (2014). Interpersonal competence in young adulthood and right laterality in white matter. *Journal of Cognitive Neuroscience, 26*(6), 1257-1265. doi:10.1162/jocn_a_00534
- Deblinger, E., Mannarino, A. P., Cohen, J. A., Runyon, M. K., & Steer, R. A. (2011). Trauma-focused Cognitive Behavioral Therapy for children: Impact of the trauma narrative and treatment length. *Depression and Anxiety, 28*(1), 67–75. doi:10.1002/da.20744
- Decety, J., & Lamm, C. (2007). The role of the right temporoparietal junction in social interaction: How low-level computational processes contribute to meta-cognition. *Neuroscientist, 13*(6), 580-593. doi:10.1177/1073858407304654
- Deci, E. L., & Ryan, R. M. (2002). *Handbook of self-determination research*. Rochester, NY: University of Rochester Press.
- DePanfilis, D. (2006). *Child neglect: A guide for prevention, assessment, and intervention*. Washington, DC: U.S. Department of Health and Human Services, Administration for Children and Families, Administration on Children, Youth and Families, Children's Bureau, Office on Child Abuse and Neglect.

- Derogatis, L. R., & Melisaratos, N. (1983). The Brief Symptom Inventory: An introductory report. *Psychological Medicine*, 13(3), 595-605. doi:10.1017/S0033291700048017
- Di Cesare, G., Di Dio, C., Marchi, M., & Rizzolatti, G. (2015). Expressing our internal states and understanding those of others. *Proceedings of the National Academy of Sciences of the United States of America*, 112(33), 10331–10335. doi:10.1073/pnas.1512133112
- Dierkhising, C. B., Ko, S., & Goldman, J. H. (2013). *Trauma-informed juvenile justice roundtable: Current issues and new directions in creating trauma-informed juvenile justice systems*. Los Angeles, CA and Durham, NC: National Center for Child Traumatic Stress. Retrieved from http://www.nctsn.org/sites/default/files/assets/pdfs/jj_trauma_brief_introduction_final.pdf
- Dong, M., Anda, R. F., Felitti, V. J., Dube, S. R., Williamson, D. F., Thompson, T. J., . . . Giles, W. H. (2004). The interrelatedness of multiple forms of childhood abuse, neglect, and household dysfunction. *Child Abuse and Neglect*, 28(7), 771–784. doi:10.1016/j.chiabu.2004.01.008
- Dong, M., Dube, S. R., Felitti, V. J., Giles, W. H., & Anda, R. F. (2003). Adverse childhood experiences and self-reported liver disease: New insights into the causal pathway. *Archives of Internal Medicine Journal*, 163(16), 1949-1956. doi:10.1001/archinte.163.16.1949
- Dong, M., Giles, W. H., Felitti, V. J., Dube, S. R., Williams, J. E., Chapman, D. P., & Anda, R. F. (2004). Insights into causal pathways for ischemic heart disease: Adverse Childhood Experiences Study. *Circulation*, 110(13), 1761-1766. doi:10.1161/01.CIR.0000143074.54995.7F

- Dorsey, S., Briggs, E. C., & Woods, B. A. (2011). Cognitive-behavioral treatment for posttraumatic stress disorder in children and adolescents. *Child and Adolescent Psychiatric Clinics of North America*, 20(2), 255–269. doi:10.1016/j.chc.2011.01.006
- Dube, S. R., Anda, R. F., Felitti, V. J., Chapman, D. P., Williamson, D. F., & Giles, W. H. (2001). Childhood abuse, household dysfunction and the risk of attempted suicide throughout the life span: Findings from the Adverse Childhood Experiences Study. *Journal of the American Medical Association*, 286(24), 3089-3096. doi:10.1001/jama.286.24.3089
- Dube, S. R., Anda, R. F., Felitti, V. J., Edwards, V. J., & Croft, J. B. (2002). Adverse childhood experiences and personal alcohol abuse as an adult. *Addictive Behaviors*, 27(5), 713-725. doi:10.1016/S0306-4603(01)00204-0
- Dube, S. R., Anda, R. F., Whitfield, C. L., Brown, D. W., Felitti, V. J., Doug, M., & Giles, W. H. (2005). Long-term consequences of childhood sexual abuse by gender of victim. *American Journal of Preventive Medicine*, 28(5), 430-438. doi:10.1016/j.amepre.2005.01.015
- Dube, S. R., Fairweather, D., Pearson, W. S., Felitti, V. J., Anda, R. F., & Croft, J. B. (2009). Cumulative childhood stress and autoimmune disease. *Psychosomatic Medicine*(71), 243-250. doi:10.1097/PSY.0b013e3181907888
- Dube, S. R., Felitti, V. J., Dong, M., Chapman, D. P., & Anda, R. F. (2003). Childhood abuse, neglect, and household dysfunction and the risk of illicit drug use: The Adverse Childhood Experiences Study. *Pediatrics*, 111(3), 564-572. doi:10.1542/peds.111.3.564
- Dube, S. R., Miller, J. W., Brown, D. W., Giles, W. H., Felitti, V. J., Dong, M., & Anda, R. F. (2006). Adverse childhood experiences and the association with ever using alcohol and

- initiating alcohol use during adolescence. *Journal of Adolescent Health*, 38(4), 444.e1–444.e10. doi:10.1016/j.jadohealth.2005.06.006
- Dubowitz, D. H. (2016). Child neglect: Confronting the challenges. *30th Annual San Diego Conference on Child and Family Maltreatment*. San Diego, CA: Maryland Department of Human Resources, U.S. Department of Health and Human Services - Office on Child Abuse and Neglect, Centers for Disease Control and Prevention, Doris Duke Charitable Foundation.
- Dubowitz, H., & Bennett, S. (2007). Physical abuse and neglect of children. *The Lancet*, 369(9576), 1891–1899. doi:10.1016/S0140-6736(07)60856-3
- Dubowitz, H., Black, M., Starr, R. H., & Zuravin, S. (1993). A conceptual definition of child neglect. *Criminal Justice and Behavior*, 20(1), 8-26. doi:10.1177/0093854893020001003
- Dubowitz, H., Pitts, S. C., & Black, M. M. (2004). Measurement of three major subtypes of child neglect. *Child Maltreatment*, 9(4), 344–356. doi:10.1177/1077559504269191
- Duke, N. N., Pettingell, S. L., McMorris, B. J., & Borowsky, I. W. (2010). Adolescent violence perpetration: Associations with multiple types of adverse childhood experiences. *Pediatrics*, 125(4), 778-786. doi:10.1542/peds.2009-0597
- Eabon, M. F., & Abrahamson, D. (n.d.). *Understanding psychological testing and assessment*. Retrieved November 29, 2016, from American Psychological Association: <http://www.apa.org/helpcenter/assessment.aspx>
- Edleson, J. L. (1999). Children's witnessing of adult domestic violence. *Journal of Interpersonal Violence*, 14(8), 839-870. doi:10.1177/088626099014008004
- Ellis, H. B., Fogler, J., Hansen, S., Forbes, P., & Navalta, C. P. (2012). Trauma Systems Therapy: 15-month outcomes and the importance of effecting environmental change.

- Psychological Trauma: Theory, Research, Practice, and Policy*, 4(6), 624-630.
doi:10.1037/a0025192
- Erickson, M. F., & Egeland, B. (2002). Child neglect. In J. E. Myers, L. Berliner, J. Briere, C. T. Hendrix, C. Jenny, & T. A. Reid (Eds.), *The APSAC handbook on child maltreatment* (pp. 3-20). Thousand Oaks, CA: Sage.
- Evers-Szostak, M., & Sanders, S. (1992). The Children's Perceptual Alteration Scale (CPAS): A measure of children's dissociation. *Dissociation*, 5(2), 91-97. Retrieved from https://scholarsbank.uoregon.edu/xmlui/bitstream/handle/1794/1642/Diss_5_2_5_OCR_rev.pdf?sequence=4&isAllowed=y
- Famularo, R., Kinscherff, R., & Fenton, T. (1992). Parental substance abuse and the nature of child maltreatment. *Child Abuse and Neglect*, 16(4), 475-483. doi:10.1016/0145-2134(92)90064-X
- Fantuzzo, J., Boruch, R., Beriama, A., Atkins, M., & Marcus, S. (1997). Domestic violence and children: Prevalence and risk in five major U.S. cities. *Journal of the American Academy of Child and Adolescent Psychiatry*, 36(1), 116-122. doi:10.1097/00004583-199701000-00025
- Fazel, M., Reed, R. V., Panter-Brick, C., & Stein, A. (2012). Mental health of displaced and refugee children resettled in high-income countries: Risk and protective factors. *The Lancet*, 379(9812), 266-282. doi:10.1016/S0140-6736(11)60051-2
- Felitti, V. J. (1993). Childhood sexual abuse, depression, and family dysfunction in adult obese patients: A case control study. *Southern Medical Journal*, 86(7), 732-736.
doi:10.1097/00007611-199307000-00002

- Felitti, V. J. (2009). Adverse childhood experiences and adult health. *Academic Pediatrics*, 9(3), 131-132. doi:10.1016/j.acap.2009.03.001
- Felitti, V. J., & Anda, R. F. (2010). The relationship of adverse childhood experiences to adult medical disease, psychiatric disorders, and sexual behavior: Implications for healthcare. In R. A. Lanius, E. Vermetten, & C. Pain (Eds.), *The impact of early life trauma on health and disease: The hidden epidemic* (pp. 77-87). New York, NY: Cambridge University Press.
- Felitti, V. J., Anda, R. F., Nordenberg, D., Williamson, D. F., Spitz, A. M., Edwards, V., . . . Marks, J. S. (1998). Relationship of childhood abuse and household dysfunction to many of the leading causes of death in adults: The Adverse Childhood Experiences (ACE) Study. *American Journal of Preventative Medicine*, 14(4), 245-258. doi:10.1016/S0749-3797(98)00017-8
- Fergusson, D. M., & Lynskey, M. T. (1997). Physical punishment/maltreatment during childhood and adjustment in young adulthood. *Child Abuse and Neglect*, 21(7), 617-630. doi:10.1016/S0145-2134(97)00021-5
- Filiano, A. J., Xu, Y., Tustison, N. J., March, R. L., Baker, W., Smirnov, I., . . . Kipnis, J. (2016). Unexpected role of interferon- γ in regulating neuronal connectivity and social behaviour. *Nature*, 535(7612), 425-429. doi:10.1038/nature18626
- Finkel, E. (2014, September 24). *Trauma-informed judges take gentler approach, administer problem-solving justice to stop cycle of ACEs*. Retrieved November 22, 2016, from ACEs too high: <https://acestoohigh.com/2014/09/24/trauma-informed-judges-take-gentler-approach-administer-problem-solving-justice-to-stop-cycle-of-aces/>
- Finkelhor, D. (1979). *Sexually victimized children*. New York, NY: The Free Press.

- Finkelhor, D. (1987). The trauma of child sexual abuse: Two models. *Journal of Interpersonal Violence*, 2(4), 348-366. doi:10.1177/088626058700200402
- Finkelhor, D. (2008). *Childhood victimization: Violence, crime and abuse in the lives of young people*. New York, NY: Oxford University Press.
- Finkelhor, D. (2009). The prevention of childhood sexual abuse. In S. McLanahan, R. Haskins, C. Paxson, & I. Sawhill (Eds.), *Preventing child maltreatment* (pp. 169-187). Princeton, NJ: The Future of Children.
- Finkelhor, D., Ormrod, R. K., & Turner, H. A. (2009). Lifetime assessment of poly-victimization in a national sample of children and youth. *Child Abuse and Neglect*, 33(7), 403-411. doi:10.1016/j.chiabu.2008.09.012
- Finkelhor, D., Turner, H., Shattuck, A., Hamby, S., & Kracke, K. (2015). *Children's exposure to violence, crime, and abuse: An update*. Office of Juvenile Justice and Delinquency Prevention. Washington, DC: U.S. Department of Justice. Retrieved from <http://www.ojjdp.gov/pubs/248547.pdf>
- Fishman, S. H. (1983). The impact of incarceration on children of offenders. In M. Frank (Ed.), *Children of exceptional parents* (pp. 169-192). New York, NY: Hayworth Press.
- Fletcher, K. E. (1994). *Dimensions of Stressful Events Rating Scale*. Worcester, MA: University of Massachusetts.
- Ford, J. D. (2005). Treatment implications of altered affect regulation and information processing following child maltreatment. *Psychiatric Annals*, 35(5), 410-419.
- Ford, J. D., & Blaustein, M. E. (2013). Systemic self-regulation: A framework for trauma-informed services in residential juvenile justice programs. *Journal of Family Violence*, 28(7), 665-677. doi:10.1007/s10896-013-9538-5

Ford, J. D., & Courtois, C. A. (2013). *Treating complex traumatic stress disorders in children and adolescents*. New York, NY: Guilford.

Ford, J. D., & Hawke, J. (2012). Trauma affect regulation psychoeducation group attendance is associated with reduced disciplinary incidents and sanctions in juvenile detention facilities. *Journal of Aggression, Maltreatment, and Trauma*, 21(4), 365–384.
doi:10.1080/10926771.2012.673538

Ford, J. D., & Kidd, P. (1998). Early childhood trauma and disorders of extreme stress as predictors of treatment outcome with chronic posttraumatic stress disorder. *Journal of Traumatic Stress*, 11(4), 743–761. doi:10.1023/A:1024497400891

Ford, J. D., & Russo, E. (2006). Trauma-focused, present-centered, emotional self-regulation approach to integrated treatment for posttraumatic stress and addiction: Trauma adaptive recovery group education and therapy (TARGET). *American Journal of Psychotherapy*, 60(4), 335-355. Retrieved from
http://www.cttntraumatraining.org/uploads/4/6/2/3/46231093/target-am_j_psychotherapy.pdf

Ford, J. D., & Russo, E. (2006). Trauma-focused, present-centered, emotional self-regulation approach to integrated treatment for posttraumatic stress and addiction: Trauma Adaptive Recovery Group Education and Therapy (TARGET). *American Journal of Psychotherapy*, 60(4), 335-355. Retrieved from
http://www.cttntraumatraining.org/uploads/4/6/2/3/46231093/target-am_j_psychotherapy.pdf

- Ford, J. D., Chapman, J., Connor, D. F., & Cruise, K. R. (2012). Complex trauma and aggression in secure juvenile justice settings. *Criminal Justice and Behavior*, 39(6), 694-724.
doi:10.1177/0093854812436957
- Forward, S. (1989). *Toxic parents: Overcoming their hurtful legacy and reclaiming your life*. New York, NY: Bantam Books.
- Francis, D., Diorio, J., Plotsky, P. M., & Meaney, M. J. (2002). Environmental enrichment reverses the effects of maternal separation on stress reactivity. *Journal of Neuroscience*, 22(18), 7840-7843. Retrieved from
<http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.334.2094&rep=rep1&type=pdf>
- Frank, D. A., Drotar, D., Cook, J., Bleiker, J., & Kasper, D. (2001). Failure to thrive. In R. M. Reece, & S. Ludwig (Eds.), *Child abuse: Medical diagnosis and management* (pp. 307-337). Philadelphia, PA: Lippincott, Williams and Wilkins.
- Franke, L. (1983). *Growing up divorced*. New York, NY: Linden Press.
- Frederick, J., & Goddard, C. (2007). Exploring the relationship between poverty, childhood adversity and child abuse from the perspective of adulthood. *Child Abuse Review*, 16(5), 323-341. doi:10.1002/car.971
- Frick, P. J. (1994). Family dysfunction and the disruptive behavior disorders: A review of recent empirical findings. *Advances in Clinical Child Psychology*, 16, 203-226.
doi:10.1007/978-1-4757-9041-2_7
- Friedrich, W. N., Grambsch, P., Damon, L., Hewitt, S. K., Koverola, C., Lang, R. A., . . . Broughton, D. (1992). Child Sexual Behavior Inventory: Normative and clinical comparisons. *Psychological Assessment*, 4(3), 303-311. doi:10.1037/1040-3590.4.3.303

- Fuhrmann, K. (2010). *Treating pediatric traumatic medical stress*. Clovis, CA: Association for Play Therapy. Retrieved from <http://www.a4pt.org/download.cfm?ID=28862>
- Galea, S., Nandi, A., & Vlahov, D. (2005). The epidemiology of post-traumatic stress disorder after disasters. *Epidemiologic Reviews*, 27(1), 78-91. doi:10.1093/epirev/mxi003
- Ganguly, P., & Brenhouse, H. C. (2015). Broken or maladaptive? Altered trajectories in neuroinflammation and behavior after early life adversity. *Developmental Cognitive Neuroscience*, 11, 18-30. doi:10.1016/j.dcn.2014.07.001
- Garbarino, J., & Garbarino, A. (1994). *Emotional maltreatment of children*. Chicago, IL: National Committee to Prevent Child Abuse.
- Garbarino, J., Dubrow, N., Kostelny, K., & Pardo, C. (1992). *Children in danger: Coping with the consequences of community violence*. San Francisco, CA: Jossey-Bass.
- Gaudin, Jr., J. M. (1984). Social work roles and tasks with incarcerated mothers. *Social Casework*, 65, 279-286.
- Gaudin, Jr., J. M. (1995). *Child Neglect: A Guide for Intervention*. Washington, DC: National Center on Child Abuse and Neglect.
- George, C. (1996). A representational perspective of child abuse and prevention: Internal working models of attachment and caregiving. *Child Abuse and Neglect*, 20(5), 411-424. doi:10.1016/0145-2134(96)00016-6
- Gerard, A. B. (1994). *Parent-Child Relationship Inventory (PCRI) manual*. Los Angeles, CA: WPS.
- Gilbert, L. K., Breiding, M. J., Merrick, M. T., Thompson, W. W., Ford, D. C., Dhingra, S. S., & Parks, S. E. (2015). Childhood adversity and adult chronic disease: an update from ten

- states and the District of Columbia, 2010. *American Journal of Preventive Medicine*, 48(3), 345-349. doi:10.1016/j.amepre.2014.09.006
- Gilbert, L. K., Breiding, M. J., Merrick, M. T., Thompson, W. W., Ford, D. C., Dhingra, S. S., & Parks, S. E. (2015). Childhood adversity and adult chronic disease: An update from ten states and the District of Columbia, 2010. *American Journal of Preventive Medicine*, 48(3), 345-349. doi:10.1016/j.amepre.2014.09.006
- Gioia, G. A., Isquith, P. K., Guy, S. C., & Kenworthy, L. (2000). Test review: Behavior Rating Inventory of Executive Function. *Child Neuropsychology*, 6(3), 235-238. doi:10.1076/chin.6.3.235.3152
- Glover, V. (2011). *The effects of prenatal stress on child behavioural and cognitive outcomes start at the beginning*. Montreal, Quebec: Encyclopedia on Early Childhood Development. Retrieved from <http://www.child-encyclopedia.com/sites/default/files/textes-experts/en/872/the-effects-of-prenatal-stress-on-child-behavioural-and-cognitive-outcomes-start-at-the-beginning.pdf>
- Goldman, J., Salus, M. K., Wolcott, D., & Kennedy, K. Y. (2003). *A coordinated response to child abuse and neglect: The foundation for practice*. Washington, DC: U.S. Department of Health and Human Services, Administration for Children and Families, Administration on Children - Youth and Families, Children's Bureau, Office on Child Abuse and Neglect. Retrieved from <http://centerforchildwelfare.fmhi.usf.edu/kb/trprev/a%20coordinated%20response%20to%20child%20abuse%20and%20neglect-the%20foundation%20for%20practice.pdf>
- Goldson, E. (2001). Maltreatment among children with disabilities. *Infants and Young Children*, 13(4), 44-54. doi:10.1097/00001163-200113040-00010

- Goldson, E. (2001). Maltreatment among children with disabilities. *Infants and Young Children*, 13(4).
- Goodman, L., Saxe, L., & Harvey, M. (1991). Homelessness as psychological trauma: Broadening perspectives. *American Psychologist*, 46(11), 1219-1225. doi:10.1037/0003-066X.46.11.1219
- Goodman, R. (1997). The Strengths and Difficulties Questionnaire: A research note. *Journal of Child Psychology and Psychiatry*, 38(5), 581–586. doi:10.1111/j.1469-7610.1997.tb01545.x
- Goodwin, B., & Miller, K. (2013). Teaching self-regulation has long-term benefits. *Educational Leadership*, 70(8), 80-81.
- Graefe, D. R., & Lichter, D. T. (1999). Life course transitions of American children: Parental cohabitation, marriage, and single motherhood. *Demography*, 36(2), 205–217. doi:10.2307/2648109
- Gratz, K. L., & Roemer, L. (2004). Multidimensional assessment of emotion regulation and dysregulation: Development, factor structure, and initial validation of the difficulties in Emotion Regulation Scale. *Journal of Psychopathology and Behavioral Assessment*, 26(1), 41-54. doi:10.1023/B:JOBA.00000007455.08539.94
- Green, B. L., Korol, M., Grace, M. C., Vary, M. G., Leonard, A. C., Gleser, G. C., & Smitson-Cohen, S. (1991). Children and disaster: Age, gender, and parental effects on PTSD symptoms. *Journal of the American Academy of Child and Adolescent Psychiatry*, 30(6), 945-951. doi:10.1097/00004583-199111000-00012
- Greene, B., Bieschke, K. L., Perez, R. M., & DeBord, K. A. (2007). Delivering ethical psychological services to lesbian, gay, and bisexual clients. In K. J. Bieschke, R. M.

- Perez, & K. A. DeBord (Eds.), *Handbook of counseling and psychotherapy with lesbian, gay, bisexual, and transgender clients* (pp. 181-199). Washington, DC: American Psychological Association. doi:10.1037/11482-007
- Gross, A. B., & Keller, H. R. (1992). Long-term consequences of childhood physical and psychological maltreatment. *Aggressive Behavior, 18*(3), 171–185. doi:10.1002/1098-2337(1992)18:3<171::AID-AB2480180302>3.0.CO;2-I
- Guarnaccia, P. J., & Rogler, L. H. (1999). Research on culture-bound syndromes: New directions. *The American Journal of Psychiatry, 156*(9), 1322-1327. doi:10.1176/ajp.156.9.1322
- Gunnar, M. R. (2003). Integrating neuroscience and psychosocial approaches in the study of early experiences. In J. A. King, C. F. Ferris, & I. I. Lederhendler (Eds.), *Roots of mental illness in children* (pp. 238-247). New York, NY: The New York Academy of Sciences.
- Gunnar, M. R., Kryzer, E., Van Ryzin, M. J., & Phillips, D. A. (2010). The rise in cortisol in family day care: Associations with aspects of care quality, child behavior, and child sex. *Child Development, 81*(3), 851-869. doi:10.1111/j.1467-8624.2010.01438.x.
- Gunnar, M., & Quevedo, K. (2007). The neurobiology of stress and development. *Annual Review of Psychology, 58*, 145-173. doi:10.1146/annurev.psych.58.110405.085605
- Habib, M., & DeRosa, R. (2008). *Coping and meaning making: Essential components for complex trauma treatment with adolescents*. Chicago, IL: International Society for Traumatic Stress Studies.
- Habib, M., & Labruna, V. (2011). Clinical considerations in assessing trauma and PTSD in adolescents. *Journal of Child and Adolescent Trauma, 4*(3), 198-216. doi:10.1080/19361521.2011.597684

- Hall, L. K. (2008). *Counseling military families: What mental health professionals need to know*. New York, NY: Routledge.
- Hamby, S. L., Finkelhor, D., Ormond, R., & Turner, H. (2004). *The Juvenile Victimization Questionnaire (JVQ): Administration and scoring manual*. Durham, NH: Crimes Against Children Research Center.
- Hamby, S., Finkelhor, D., Turner, H., & Ormrod, R. (2011). *Children's exposure to intimate partner violence and other family violence*. Office of Juvenile Justice and Delinquency Prevention. Washington, DC: U.S. Department of Justice. Retrieved from <http://www.unh.edu/ccrc/pdf/jvq/NatSCEV-Children's%20Exposure-Family%20Violence%20final.pdf>
- Harrington, D., Zuravin, S., DePanfilis, D., Ting, L., & Dubowitz, H. (2002). The neglect scale: Confirmatory factor analyses in a low-income sample. *Child Maltreatment*, 7(4), 359-368. doi:10.1177/107755902237266
- Harter, S. (1982). The Perceived Competence Scale for Children. *Child Development*, 53(1), 87-97. doi:10.1111/j.1467-8624.1982.tb01295.x
- Hecht, D. (2014). Cerebral lateralization of pro- and anti-social tendencies. *Experimental Neurobiology*, 23(1), 1-27. doi:10.5607/en.2014.23.1.1
- Heller, T., Roccoforte, J. A., & Cook, J. A. (1997). Predictors of support group participation among families of persons with mental illness. *Family Relations*, 46(4), 437-442. doi:10.2307/585103
- Heptinstall, E., Sethna, V., & Taylor, E. (2004). PTSD and depression in refugee children: Associations with pre-migration trauma and post-migration stress. *European Child and Adolescent Psychiatry*, 13(6), 373-380. doi:10.1007/s00787-004-0422-y

- Herman, J. L. (2003). The mental health of crime victims: Impact of legal intervention. *Journal of Traumatic Stress, 16*(2), 159-166. doi:10.1023/A:1022847223135
- Herrenkohl, T. I., Sousa, C., Tajima, E. A., Herrenkohl, R. C., & Moylan, C. A. (2008). Intersection of child abuse and children's exposure to domestic violence. *Trauma, Violence, and Abuse, 9*(2), 84-99. doi:10.1177/1524838008314797
- Herringa, R. J., Birn, R. M., Ruttle, P. L., Burghy, C. A., Stodola, D. E., Davidson, R. J., & Essex, M. J. (2013). Childhood maltreatment is associated with altered fear circuitry and increased internalizing symptoms by late adolescence. *Proceedings of the National Academy of Sciences of the United States of America, 110*, 19119–19124. doi:10.1073/pnas.1310766110
- Higgins, D. J. (2003). Maltreatment and family dysfunction in childhood and the subsequent adjustment of children and adults. *Journal of Family Violence, 18*(2), 107–120. doi:10.1023/A:1022841215113
- Hillis, S. D., Anda, R. F., Dube, S. R., Felitti, V. J., Marchbanks, P. A., & Marks, J. S. (2004). The association between adverse childhood experiences and adolescent pregnancy, long-term psychosocial consequences, and fetal death. *Pediatrics, 113*(2), 320-327. doi:10.1452/peds.113.2.320
- Hillis, S. D., Anda, R. F., Felitti, V. J., & Marchbanks, P. A. (2001). Adverse childhood experiences and sexual risk behaviors in women: A retrospective cohort study. *Family Planning Perspectives, 33*(5), 206-211. doi:10.2307/2673783
- Hochanadel, A., & Finamore, D. (2015). Fixed and growth mindset in education and how grit helps students persist in the face of adversity. *Journal of International Education Research, 11*(1), 48-50. doi:10.19030/jier.v11i1.9099

Hochman, A. (2014, September 30). *The Camden story: A physician and a priest plant seeds of repair*. Retrieved November 22, 2016, from ACEs too high:

<https://acestoohigh.com/2014/09/30/the-camden-story-a-physician-and-a-priest-plant-seeds-of-repair-2/>

Hochman, A. (2014, September 25). *The Philadelphia story: Education and activism converge in "ACEs epicenter"*. Retrieved November 22, 2016, from ACEs too high:

<https://acestoohigh.com/2014/09/25/the-philadelphia-story-education-and-activism-converge-in-aces-epicenter/>

Hodgdon, H. B., Blaustein, M., Kinniburgh, K., Peterson, M. L., & Spinazzola, J. (2016).

Application of the ARC model with adopted children: Supporting resiliency and family well being. *Journal of Child and Adolescent Trauma*, 9(1), 43–53. doi:10.1007/s40653-015-0050-3

Hodgdon, H. B., Kinniburgh, K., Gabowitz, D., Blaustein, M. E., & Spinazzola, J. (2013).

Development and implementation of trauma-informed programming in youth residential treatment centers using the ARC framework. *Journal of Family Violence*, 28(7), 679–692. doi:10.1007/s10896-013-9531-z

Holden, G. W. (2003). Children exposed to domestic violence and child abuse: Terminology and taxonomy. *Clinical Child and Family Psychology Review*, 6(3), 151-160.

doi:10.1023/A:1024906315255

Hollifield, M., Eckert, V., Warner, T. D., Jenkins, J., Krakow, B., Ruiz, J., & Westermeyer, J.

(2005). Development of an inventory for measuring war-related events in refugees.

Comprehensive Psychiatry, 46(1), 67-80. doi:10.1016/j.comppsy.2004.07.003

- Holt, S., Buckley, H., & Whelan, S. (2008). The impact of exposure to domestic violence on children and young people: A review of the literature. *Child Abuse and Neglect*, 32(8), 797–810. doi:10.1016/j.chiabu.2008.02.004
- Homma, Y., Wang, N., Saewyc, E., & Kishor, N. (2012). The relationship between sexual abuse and risky sexual behavior among adolescent boys: A meta-analysis. *Journal of Adolescent Health*, 51(1), 18-24. doi:10.1016/j.jadohealth.2011.12.032
- House, A. S., Van Horn, E., Coppeans, C., & Stepleman, L. M. (2011). Interpersonal trauma and discriminatory events as predictors of suicidal and nonsuicidal self-injury in gay, lesbian, bisexual, and transgender persons. *Journal of Traumatology*, 17(2), 75-85. doi:10.1177/1534765610395621
- Huang, H., Yan, P., Shan, Z., Chen, S., Li, M., Luo, C., . . . Liu, L. (2015). Adverse childhood experiences and risk of type 2 diabetes: A systematic review and meta-analysis. *Metabolism*, 64(11), 1408-1418. doi:10.1016/j.metabol.2015.08.019
- Ippen, C. G., Ford, J., Racusin, R., Acker, M., Bosquet, M., Rogers, K., . . . Edwards, J. (2002). *Traumatic Events Screening Inventory - Parent report revised*. San Francisco, CA: The Child Trauma Research Project of the Early Trauma Network, The National Center for PTSD Dartmouth Child Trauma Research Group.
- Isohookana, R., Riala, K., Hakko, H., & Räsänen, P. (2013). Adverse childhood experiences and suicidal behavior of adolescent psychiatric inpatients. *European Child and Adolescent Psychiatry*, 22(1), 13-22. doi:10.1007/s00787-012-0311-8
- Jackson, M., Cram, F., & Seymour, F. W. (2000). Violence and sexual coercion in high school students' dating relationships. *Journal of Family Violence*, 15(1), 23-36. doi:10.1023/A:1007545302987

- Jacobs, S., Mazure, C., & Prigerson, H. (2000). Diagnostic criteria for traumatic grief. *Death Studies*, 24(3), 185-199. doi:10.1080/074811800200531
- Janoff-Bulman, R. (1989). Assumptive worlds and the stress of traumatic events: Applications of the schema construct. *Social Cognition*, 7(2), 113-136. doi:10.1521/soco.1989.7.2.113
- Jantz, G. L., & McMurray, A. (1995). *Healing the scars of emotional abuse*. Grand Rapids, MI: Revell.
- Johnson, D. (1991). *Jailed mothers*. Pasadena, CA: Pacific Oaks Center for Children of Incarcerated Parents.
- Johnson, D. (1995). Effects of parental incarceration. In K. Gabel, & D. Johnson (Eds.), *Children of incarcerated parents* (pp. 59-88). New York, NY: Lexington Books.
- Jones, R. E., Leen-Feldner, E. W., Olatunji, B. O., & Hawks, E. (2009). Psychometric properties of the affect intensity and reactivity measure adapted for youth (AIR-Y). *Psychological Assessment*, 21(2), 162-175. doi:10.1037/a0015358
- Jose-Kampfner, C. (1995). Post-traumatic stress reactions in children of imprisoned mothers. In K. Gabel, & D. Johnston (Eds.), *Children of incarcerated parents* (pp. 89-100). New York, NY: Lexington Books.
- Joseph, S., Mynard, H., & Mayall, M. (2000). Life-events and post-traumatic stress in a sample of English adolescents. *Journal of Community and Applied Social Psychology*, 10(6), 475-482. doi:10.1002/1099-1298(200011/12)10:6<475::AID-CASP578>3.0.CO;2-F
- Kagan, R. (2007). *Real Life Heroes practitioners manual*. New York, NY: Routledge.
- Kagan, R., & Spinazzola, J. (2013). Real Life Heroes in residential treatment: Implementation of an integrated model of trauma and resiliency-focused treatment for children and

- adolescents with complex PTSD. *Journal of Family Violence*, 28(7), 705-715.
doi:10.1007/s10896-013-9537-6
- Kagan, R., Douglas, A. N., Hornick, J., & Kratz, S. L. (2008). Real Life Heroes pilot study: Evaluation of a treatment model for children with traumatic stress. *Journal of Child and Adolescent Trauma*, 1(1), 5-22. doi:10.1080/19361520801929845
- Kamphaus, R. W., Petoskey, M. D., Cody, A. H., Rowe, E. W., Huberty, C. J., & Reynolds, C. R. (1999). A typology of parent rated child behavior for a national U.S. sample. *Journal of Child Psychology and Psychiatry*, 40(4), 607-616. doi:10.1111/1469-7610.00477
- Kann, L., Olsen, E. O., McManus, T., Kinchen, S., Chyen, D., Harris, W. A., & Wechsler, H. (2011). *Sexual identity, sex of sexual contacts, and health-risk behaviors among students in grades 9-12*. Washington, DC: Centers for Disease Control and Prevention. Retrieved from <https://www.cdc.gov/mmwr/preview/mmwrhtml/ss6007a1.htm>
- Kassam-Adams, N., & Winston, F. K. (2004). Predicting posttraumatic stress after hospitalization for pediatric injury. *Journal of the American Academy of Child & Adolescent Psychiatry*, 43(4), 403-411. doi:10.1097/00004583-200404000-00006
- Kazak, A. E., Kassam-Adams, N., Schneider, S., Zelikovsky, N., Alderfer, M. A., & Rourke, M. (2006). An integrative model of pediatric medical traumatic stress. *Journal of Pediatric Psychology*, 31(4), 343-355. doi:10.1093/jpepsy/jsj054
- Kazdin, A. E. (1995). Child, parent and family dysfunction as predictors of outcome in cognitive-behavioral treatment of antisocial children. *Behaviour Research and Therapy*, 33(3), 271-281. doi:10.1016/0005-7967(94)00053-M

- Kelley, B. T., Thornberry, T. P., & Smith, C. A. (1997). *In the wake of childhood maltreatment*. Washintgon, DC: U.S. Department of Justice. Retrieved from <https://www.ncjrs.gov/pdffiles1/165257.pdf>
- Kent, A., & Wallar, G. (1998). The impact of childhood emotional abuse: An extension of the child abuse and trauma scale. *Child Abuse and Neglect*, 22(5), 393-399. doi:10.1016/S0145-2134(98)00007-6
- Kerig, P. K., Ford, J. D., & Olafson, E. (2014). *Assessing exposure to psychological trauma and posttraumatic stress symptoms in the juvenile justice population*. Los Angeles, CA and Durham, NC: National Center for Child Traumatic Stress. Retrieved from http://www.nctsn.org/sites/default/files/assets/pdfs/assessing_trauma_in_jj_2014.pdf
- Kerns, C., & Lee, B. (2015). The relationship between adverse childhood experiences and autism spectrum disorder in an epidemiological sample from the United States. *European Psychiatry*, 30, 737. doi:10.1016/S0924-9338(15)30585-X
- Killen-Harvey, A. (2006). *Culture and trauma brief: Trauma among lesbian, gay, bisexual, transgender, or questioning youth*. Los Angeles, CA, and Durham, NC: National Child Traumatic Stress Network.
- Kilpatrick, D. J., Saunders, B. E., & Smith, D. W. (2003). *Youth victimization: Prevalence and implications*. Washington, DC: U.S. Department of Justice.
- Kinniburgh, K. J., Blaustein, M., Spinazzola, J., & van der Kolk, B. A. (2005). Attachment, self-regulation, and competency. *Psychiatric Annals*, 35(5), 424-430. Retrieved from <http://psychrights.org/research/Digest/CriticalThinkRxCites/kinniburgh.pdf>

- Kinsella, M. T., & Monk, C. (2009). Impact of maternal stress, depression and anxiety on fetal neurobehavioral development. *Clinical Obstetrics and Gynecology*, 52(3), 425-440.
doi:10.1097/GRF.0b013e3181b52df1
- Kiser, L. J. (2007). Protecting children from the dangers of urban poverty. *Clinical Psychology Review*, 27(2), 211–225. doi:10.1016/j.cpr.2006.07.004
- Kiser, L. J., Donohue, A., Hodgkinson, S., & Medoff, D. (2010). Strengthening Family Coping Resources: The feasibility of a multifamily group intervention for families exposed to trauma. *Journal of Traumatic Stress*, 23(6), 802–806. doi:10.1002/jts.20587
- Kisiel, C., Conradi, L., Fehrenbach, T., Torgerson, E., & Briggs, E. C. (2014). Assessing the effects of trauma in children and adolescents in practice settings. *Journal of Child and Adolescent Psychiatric Clinics of North America*, 23(2), 223-242.
doi:10.1016/j.chc.2013.12.007
- Kisiel, C., Fehrenbach, T., Small, L., & Lyons, J. S. (2009). Assessment of complex trauma exposure, responses, and service needs among children and adolescents in child welfare. *Journal of Child and Adolescent Trauma*, 2(3), 143-160.
doi:10.1080/19361520903120467
- Kisiel, C., Marett, J., Markiwicz, J., & Fehrenbach, T. (2008). *Evidence based interventions for traumatized youth in Illinois child welfare*. Evanston, IL: Northwestern University Feinberg School Mental Health Services and Policy Program. Retrieved from <http://www.tapartnership.org/docs/presentations/socMeetingSummer2011/day1/Workshop%2045-%20Evidence-Based%20Interventions%20for%20Traumatized%20Youth/Evidence-Based%20Interventions.pdf>

- Kleinman, A. (1988). *Rethinking psychiatry: From cultural category to personal experience*. New York, NY: The Free Press.
- Ko, S. J., Ford, J. D., Kassam-Adams, N., Berkowitz, S. J., Wilson, C., Wong, M., . . . Layne, C. M. (2008). Creating trauma-informed systems: Child welfare, education, first responders, health care, juvenile justice. *Professional Psychology: Research and Practice*, 39(4), 396-404. doi:10.1037/0735-7028.39.4.396
- Korfmacher, J. (1998). *Emotional neglect: Being hurt by what is not there*. Chicago, IL: National Committee to Prevent Child Abuse.
- Kroes, G., Veerman, J. W., & De Bruyn, E. J. (2005). The impact of the big five personality traits on reports of child behavior problems by different informants. *Journal of Abnormal Child Psychology*, 33(2), 231–240. doi:10.1007/s10802-005-1830-2
- Lalor, K., & McElvaney, R. (2010). Child sexual abuse, links to later sexual exploitation/high risk sexual behavior and prevention/treatment programmes. *Trauma, Violence and Abuse*, 11(4), 159-177. doi:10.1177/1524838010378299
- Lanktree, C. B., & Briere, J. (2013). Integrative treatment of complex trauma (ITCT) for children and adolescents. In J. D. Ford, & C. A. Courtois (Eds.), *Treating complex trauma traumatic stress disorders with children and adolescents: An evidence-based guide* (pp. 143-161). New York, NY: Guilford.
- Lanktree, C. B., Briere, J., Godbout, N., Hodges, M., Chen, K., Trimm, L., . . . Freed, W. (2012). Treating multi-traumatized, socially-marginalized children: Results of a naturalistic treatment outcome study. *Journal of Aggression, Maltreatment and Trauma*, 21(8), 813-828. doi:10.1080/10926771.2012.722588

- Lawson, D. M., & Quinn, J. (2013). Complex trauma in children and adolescents: Evidence-based practice in clinical settings. *Journal of Clinical Psychology, 69*(5), 497–509. doi:10.1002/jclp.21990
- Lax, R. F., Bach, S., & Burland, J. A. (Eds.). (1986). *Self and object constancy: Clinical and theoretical perspectives*. New York, NY: Guilford Press.
- Leahy, M. A. (2015). Children of mentally ill parents: Understanding the effects of childhood trauma as it pertains to the school setting. *International Journal of Educational Research, 71*, 100–107. doi:10.1016/j.ijer.2015.04.005
- Lefcourt, H. M. (1991). Locus of control. In J. P. Robinson, P. R. Shaver, & L. S. Wrightsman (Eds.), *Measures of personality and social psychological attitudes* (pp. 413-499). San Diego, CA: Academic Press.
- Lieberman, A. F., Ippen, C. G., & Van Horn, P. (2006). Child-Parent Psychotherapy: 6-Month follow-up of a randomized controlled trial. *The American Academy of Child and Adolescent Psychiatry, 45*(8), 913–918. doi:10.1097/01.chi.0000222784.03735.92
- Lieberman, A. F., Van Horn, P., & Ippen, C. G. (2005). Toward evidence-based treatment: Child-Parent Psychotherapy with preschoolers exposed to marital violence. *The American Academy of Child and Adolescent Psychiatry, 44*(12), 1241–1248. doi:10.1097/01.chi.0000181047.59702.58
- Lincoln, A., Swift, E., & Shorteno-Fraser, M. (2008). Psychological adjustment and treatment of children and families with parents deployed in military combat. *Journal of Child Psychology, 64*(8), 984-992. doi:10.1002/jclp.20520

- Liotti, G. (2004). Trauma, dissociation, and disorganized attachment: Three strands of a single braid. *Psychotherapy: Theory, Research, Practice, Training*, 41(4), 472-486.
doi:10.1037/0033-3204.41.4.472
- Loman, M., & Gunnar, M. R. (2010). Early experience and the development of stress reactivity and regulation in children. *Neuroscience and Biobehavioral Reviews*, 34(6), 867-876.
doi:10.1016/j.neubiorev.2009.05.007
- Lonigan, C. J., Shannon, M. P., Finch, Jr., A. J., Daugherty, T. K., & Taylor, C. M. (1991). Children's reactions to a natural disaster: Symptom severity and degree of exposure. *Advances in Behaviour Research and Therapy*, 13(3), 135-154. doi:10.1016/0146-6402(91)90002-R
- Lupien, S. J., de Leon, M., de Santi, S., Convit, A., Tarshish, C., Nair, N. P., . . . Meaney, M. J. (1998). Cortisol levels during human aging predict hippocampal atrophy and memory deficits. *Nature Neuroscience*, 1(1), 69-73. doi:10.1038/271
- Lupien, S. J., McEwen, B. S., Gunnar, M. R., & Heim, C. (2009). Effects of stress throughout the lifespan on the brain, behaviour and cognition. *Nature Reviews Neuroscience*, 10, 434-445. doi:10.1038/nrn2639
- Lustin, S. L., Kia-Keating, M., Knight, W. G., Geltman, P., Ellis, H., Kinzie, J. D., . . . Saxe, G. N. (2004). Review of child and adolescent refugee mental health. *Journal of the American Academy of Child and Adolescent Psychiatry*, 43(1), 24-36.
doi:10.1097/00004583-200401000-00012
- Luthra, R., Abramovitz, R., Greenberg, R., Schoor, A., Newcorn, J., Schmeidler, J., . . . Chemtob, C. M. (2009). The relationship between type of trauma exposure and

- posttraumatic stress disorder among urban children and adolescents. *Journal of Interpersonal Violence*, 24(11), 1919-1927. doi:10.1177/0886260508325494
- Lyons, J., Kiser, L., Beck, V., Connors, K., Gardner, S., & Strieder, F. (2009). *Family Assessment of Needs and Strengths - Trauma version*. Chicago, IL: Praed Foundation.
- Magura, S., & Laudet, A. B. (1996). Parental substance abuse and child maltreatment: Review and implications for intervention. *Children and Youth Services Review*, 18(3), 193-220. doi:10.1016/0190-7409(96)00001-1
- Mannarino, A. P., Cohen, J. A., & Berman, S. R. (1994). The Children's Attributions and Perceptions Scale: A new measure of sexual abuse-related factors. *Journal of Clinical Child Psychology*, 23(2), 204-211. doi:10.1207/s15374424jccp2302_9
- Mansell, S., Sobsey, D., & Moskal, R. (1998). Clinical findings among sexually abused children with and without developmental disabilities. *Mental Retardation*, 36(1), 12-22. doi:10.1352/0047-6765(1998)036<0012:CFASAC>2.0.CO;2
- March, J. S., Amaya-Jackson, L., Murray, M. C., & Schulte, A. (1998). Cognitive-Behavioral Psychotherapy for children and adolescents with posttraumatic stress disorder after a single-incident stressor. *Journal of the American Academy of Child and Adolescent Psychiatry*, 37(6), 585-593. doi:10.1097/00004583-199806000-00008
- Marie-Mitchell, A., & O'Connor, T. G. (2013). Adverse childhood experiences: Translating knowledge into identification of children at risk for poor outcomes. *Academic Pediatrics*, 13(1), 14-19. doi:10.1016/j.acap.2012.10.006
- Marrow, M., Knudsen, K., Olafson, E., & Bucher, S. (2012). The value of implementing TARGET within a trauma-informed juvenile justice setting. *Journal of Child and Adolescent Trauma*, 5(3), 257-270. doi:10.1080/19361521.2012.697105

- Masten, A. S., & Narayan, A. J. (2012). Child development in the context of disaster, war, and terrorism: Pathways of risk and resilience. *Annual Review of Psychology*, 63, 227-257. doi:10.1146/annurev-psych-120710-100356
- Mattejat, F., & Remschmidt, H. (2008). The children of mentally ill parents. *Deutsches Ärzteblatt International*, 105(23), 413-418. doi:10.3238/arztebl.2008.0413
- McEwen, B. S. (2008). Central effects of stress hormones in health and disease: Understanding the protective and damaging effects of stress and stress mediators. *European Journal of Pharmacology*, 583(2-3), 174-185. doi:10.1016/j.ejphar.2007.11.071
- McEwen, B. S. (2013). The brain on stress: Toward an integrative approach to brain, body, and behavior. *Perspectives on Psychological Science*, 8(6), 673-675. doi:10.1177/1745691613506907
- McFarlane, A. C. (1987). Posttraumatic phenomena in a longitudinal study of children following a natural disaster. *Journal of the American Academy of Child and Adolescent Psychiatry*, 26(5), 764-769. doi:10.1097/00004583-198709000-00025
- McIntyre, T. C. (1987). Teacher awareness of child abuse and neglect. *Child Abuse and Neglect*, 11(1), 133-135. doi:10.1016/0145-2134(87)90042-1
- Mennen, F. E., Kim, K., Sang, J., & Tricketta, P. K. (2010). Child neglect: Definition and identification of youth's experiences in official reports of maltreatment. *Child Abuse and Neglect*, 34(9), 647-658. doi:10.1016/j.chiabu.2010.02.007
- Mental Health America of Wisconsin. (2011). *When a parent has a mental illness: Issues and challenges*. Madison, WI, Milwaukee, WI. Retrieved from [https://www.mhawisconsin.org/Data/Sites/1/media/fact-sheets/sf_issues\[1\].pdf](https://www.mhawisconsin.org/Data/Sites/1/media/fact-sheets/sf_issues[1].pdf)

- Mezzich, A. C., Tarter, R. E., Giancola, P. R., & Kirisci, L. (2001). The Dysregulation Inventory: A new scale to assess the risk for substance use disorder. *Journal of Child and Adolescent Substance Abuse, 10*(4), 35-43. doi:10.1300/J029v10n04_04
- Miller, P., & Plant, M. (1999). Truancy and perceived school performance: An alcohol and drug. *Alcohol and Alcoholism, 34*(6), 886-893. doi:10.1093/alcalc/34.6.886
- Moore, K. A., & Waite, L. J. (1981). Marital dissolution, early motherhood and early marriage. *Social Forces, 60*(1), 20-40. doi:10.1093/sf/60.1.20
- Moos, R. A. (1993). *Coping Responses Inventory (CRI)*. Lutz, FL: PAR.
- Moos, R. H. (1990). Conceptual and empirical approaches to developing family-based assessment procedures: Resolving the case of the Family Environment Scale. *Family Process, 29*(2), 199–208. doi:10.1111/j.1545-5300.1990.00199.x
- Mumola, C. J. (2000). *Incarcerated parents and their children*. Washington, DC: U.S. Department of Justice. Retrieved from <https://www.bjs.gov/content/pub/pdf/iptc.pdf>
- Muris, P., Vlaeyen, J., & Meesters, C. (2001). The relationship between anxiety sensitivity and fear of pain in healthy adolescents. *Behaviour Research and Therapy, 39*(11), 1357–1368. doi:10.1016/S0005-7967(01)00018-3
- Murphy, J. A., Rawlings, E. I., & Howe, S. R. (2002). A survey of clinical psychologists on treating lesbian, gay, and bisexual clients. *Professional Psychology: Research and Practice, 33*(2), 183-189. doi:10.1037/0735-7028.33.2.183
- Myers, B. J., Smarch, T. M., Amlund-Hagen, K., & Kennon, S. (1999). Children of incarcerated mothers. *Journal of Child and Family Studies, 8*(1), 11-25. doi:10.1023/A:1022990410036

- Nachmias, M., Gunnar, M. R., Mangelsdorf, S., Parritz, R., & Buss, K. A. (1996). Behavioral inhibition and stress reactivity: The moderating role of attachment security. *Child Development, 67*(2), 508-522. doi:10.1111/j.1467-8624.1996.tb01748.x
- Nader, K. (2007). Culture and the assessment of trauma in youths. In J. P. Wilson, & C. Tang (Eds.), *Cross-cultural assessment of psychological trauma and PTSD* (pp. 169-196). New York, NY: Springer Science Business Media.
- Nader, K. O., Newman, E., Weathers, F. W., Kaloupek, D. G., Kriegler, J. A., & Blake, D. D. (2004). *National Center for PTSD Clinician-Administered PTSD Scale for Children and Adolescents (CAPS-CA) interview booklet*. Los Angeles, CA: Western Psychological Services.
- National Child Traumatic Stress Network. (2000). *Community violence: Reactions and actions to dangerous times*. Los Angeles, CA and Durham, NC: National Center for Child Traumatic Stress. Retrieved from http://www.nctsn.org/sites/default/files/assets/pdfs/community_violence_reactions_actions.pdf
- National Child Traumatic Stress Network. (n.d.). *National Child Traumatic Stress Network empirically supported treatments and promising practices*. Retrieved November 24, 2016, from National Child Traumatic Stress Network: <http://www.nctsn.org/resources/topics/treatments-that-work/promising-practices>
- National Child Traumatic Stress Network. (n.d.). *Standardized measures to assess complex trauma*. Retrieved November 24, 2016, from National Child Traumatic Stress Network: <http://www.nctsn.org/trauma-types/complex-trauma/standardized-measures-assess-complex-trauma>

- National Child Traumatic Stress Network Schools Committee. (2008). *Trauma Toolkit for Educators*. Los Angeles, CA and Durham, NC: National Center for Child. Retrieved from http://www.nctsn.org/sites/default/files/assets/pdfs/Child_Trauma_Toolkit_Final.pdf
- National Scientific Council on the Developing Child. (2014). *Excessive stress disrupts the architecture of the developing brain*. (Working Paper No. 3). Cambridge, MA: Harvard University - Center on the Developing Child. Retrieved from http://46y5eh11fhgw3ve3ytpwxt9r.wpengine.netdna-cdn.com/wp-content/uploads/2005/05/Stress_Disrupts_Architecture_Developing_Brain-1.pdf
- National Scientific Council on the Developing Child. (2014). *Excessive stress disrupts the architecture of the developing Brain*. (Working Paper No. 3). Cambridge, MA: Harvard University - Center on the Developing Child. Retrieved from http://46y5eh11fhgw3ve3ytpwxt9r.wpengine.netdna-cdn.com/wp-content/uploads/2005/05/Stress_Disrupts_Architecture_Developing_Brain-1.pdf
- Negy, C., Lachar, D., Gruber, C. P., & Garza, N. D. (2001). The Personality Inventory for Youth: Validity and comparability of English and Spanish versions for regular education and juvenile justice samples. *Journal of Personality Assessment*, 76(2), 250-263. doi:10.1207/S15327752JPA7602_08
- Nelson, B. S., & Wampler, K. S. (2000). Systemic effects of trauma in clinic couples: An exploratory study of secondary trauma resulting from childhood abuse. *Journal of Marital and Family Therapy*, 26(2), 171–184. doi:10.1111/j.1752-0606.2000.tb00287.x
- Ney, P. G., Fung, T., & Wickett, A. R. (1994). The worst combinations of child abuse and neglect. *Child Abuse and Neglect*, 18(9), 705-714. doi:10.1016/0145-2134(94)00037-9

- Opel, N., Redlich, R., Zwanzger, P., Grotegerd, D., Arolt, V., Heindel, W., . . . Dannlowski, U. (2014). Hippocampal atrophy in major depression: A function of childhood maltreatment rather than diagnosis? *Neuropsychopharmacology*, 39(12), 2723-2731.
doi:10.1038/npp.2014.145
- Palmer, S. E., Brown, R. A., Rae-Grant, N. I., & Loughlin, M. J. (1999). Responding to children's disclosure of familial abuse: What survivors tell us. *Child Welfare*, 78(2), 259-282.
- Pantell, M. S., Jutte, D., & Adler, N. (2014). Association between adverse childhood experiences and psychiatric conditions in childhood. In *142nd APHA Annual Meeting and Exposition 2014*. Washington, DC: American Public Health Association. Retrieved from <https://apha.confex.com/apha/142am/webprogram/Paper312359.html>
- Paolucci, E. O., Genuis, M. L., & Violato, C. (1999). A meta-analysis of the published research on the effects of child sexual abuse. *The Journal of Psychology*, 135(1), 17-36.
doi:10.1080/00223980109603677
- Park, S. H., Videlock, E. J., Shih, W., Presson, A. P., Mayer, E. A., & Chang, L. (2016). Adverse childhood experiences are associated with irritable bowel syndrome and gastrointestinal symptom severity. *Neurogastroenterology and Motility*, 28(8), 1252-1260.
doi:10.1111/nmo.12826
- Pecora, P. J., Jensen, P. S., Romanelli, L. H., Jackson, L. J., & Ortiz, A. (2009). Mental health services for children placed in foster care: An overview of current challenges. *Child Welfare*, 88(1), 5-26. Retrieved from <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3061347/>

- Peek, L. (2008). Children and disasters: Understanding vulnerability, developing capacities, and promoting resilience - An introduction. *Children, Youth, and Environments*, 18(1), 1-29.
Retrieved from
<http://www.jstor.org.ezproxy1.lib.asu.edu/stable/10.7721/chilyoutenvi.18.1.0001>
- Pelcovitz, D., van der Kolk, B., Roth, S., Mandel, F., Kaplan, S., & Resick, P. (1997).
Development of a criteria set and a Structured Interview for Disorders of Extreme Stress
(SIDES). *Journal of Traumatic Stress*, 10(1), 3-16. doi:10.1002/jts.2490100103
- Perry, B. D. (2001). The neurodevelopmental impact of violence in childhood. In D. Schetky, &
E. Benedek (Eds.), *Textbook of child and adolescent forensic psychiatry* (pp. 221–238).
Washington, DC: American Psychiatric Press.
- Perry, B. D., & Rubenstein, J. (1999). The child's loss: Death, grief and mourning: How
caregivers can help children exposed to traumatic death. *Parent and Caregiver Education
Series*, 3, 1. Child Trauma Academy.
- Petersilia, J. R. (2001). Crime victims with developmental disabilities. *Criminal Justice and
Behavior*, 28(6), 655-694. doi:10.1177/009385480102800601
- Peterson, C., Luborsky, L., & Seligman, M. E. (1983). Attributions and depressive mood shifts:
A case study using the symptom-context method. *Journal of Abnormal Psychology*,
92(1), 96-103. doi:10.1037//0021-843X.92.1.96
- Pfefferbaum, B., Shaw, J. A., & American Academy of Child and Adolescent Psychiatry
Committee on Quality Issues. (2013). Practice parameter on disaster preparedness.
Journal of the American Academy of Child and Adolescent Psychiatry, 11, 1224-1238.
doi:10.1016/j.jaac.2013.08.014

- Pfefferbaum, P., Seale, T. W., Brandt, E. N., Pfefferbaum, D. E., Rainwater, D., & Rainwater, S. M. (2003). Media exposure in children one hundred miles from a terrorist bombing. *Annals of Clinical Psychiatry, 15*(1), 1-8. doi:10.3109/10401230309085664
- Piers, E. V., & Harris, D. B. (2002). *Piers-Harris Children's Self-Concept Scale – Second edition*. North Tonawanda, NY: Multi-Health Systems.
- Pine, D. S., Costello, J., & Masten, A. (2005). Trauma, proximity, and developmental psychopathology: The effects of war and terrorism on children. *Neuropsychopharmacology, 30*(10), 1781–1792. doi:10.1038/sj.npp.1300814
- Post, R. M., Altshuler, L. L., Kupka, R., McElroy, S. L., Frye, M. A., Rowe, M., . . . Nolen, W. A. (2015). Verbal abuse, like physical and sexual abuse, in childhood is associated with an earlier onset and more difficult course of bipolar disorder. *Bipolar Disorders, 17*(3), 323-330. doi:10.1111/bdi.12268
- Prebble, S. C., Addis, D. R., & Tippet, L. J. (2013). Autobiographical memory and sense of self. *Psychological Bulletin, 139*(4), 815-840. doi:10.1037/a0030146
- Pretis, M., & Dimova, A. (2008). Vulnerable children of mentally ill parents: Towards evidence-based support for improving resilience. *Support for Learning, 23*(3), 152–159. doi:10.1111/j.1467-9604.2008.00386.x
- Pumariega, A. J., Rothe, E., & Pumariega, J. B. (2005). Mental health of immigrants and refugees. *Community Mental Health Journal, 41*(5), 581–597. doi:10.1007/s10597-005-6363-1
- Putnam, F. W., & Helmers, K. (1993). Development, reliability, and validity of a child dissociation scale. *Child Abuse and Neglect, 17*(6), 731-741. doi:10.1016/S0145-2134(08)80004-X

- Putnam, K. T., Harris, W. W., & Putnam, F. W. (2013). Synergistic childhood adversities and complex adult psychopathology. *Journal of Traumatic Stress, 26*(4), 435-442.
doi:10.1002/jts.21833
- Pynoos, R. S., & Eth, S. (1985). Children traumatized by witnessing acts of personal violence: Homicide, rape, or suicide behavior. In S. Eth, & R. S. Pynoos (Eds.), *Post-traumatic stress disorder in children* (pp. 17-43). Washington, DC: American Psychiatric Press.
- Pynoos, R. S., Steinberg, A. M., & Piacentini, J. C. (1999). A developmental psychopathology model of childhood traumatic stress and intersection with anxiety disorders. *Biological Psychiatry, 46*(11), 1542–1554. doi:10.1016/S0006-3223(99)00262-0
- Pynoos, R., & Nader, K. (1988). Psychological first aid and treatment approaches to children exposed to community violence: Research implications. *Journal of Traumatic Stress, 1*, 445-473. doi:10.1002/jts.2490010406
- Pynoos, S., Rodriguez, N., Steinberg, A., Stuber, M., & Frederick, C. (1998). *UCLA PTSD index for DSM-IV*. Los Angeles, CA: University of California, Los Angeles.
- Radford, L., Glaser, D., & MacMillan, H. L. (2009). Recognising and responding to child maltreatment. *The Lancet, 373*(9658), 167–180. doi:10.1016/S0140-6736(08)61707-9
- Rapporteur, C. M., Committee on National Statistics, Board on Behavioral, Cognitive, and Sensory Sciences, Division of Behavioral and Social Sciences and Education, Board on Health Sciences Policy, Health and Medicine Division, & National Academies of Sciences, Engineering, and Medicine. (2016). *Measuring trauma: Workshop summary*. Washington, DC: National Academies Press. doi:10.17226/23526
- Reardon, S. F. (2011). Widening academic achievement gap between the rich and the poor: New evidence and possible explanations. In G. J. Duncan, & R. J. Murnane (Eds.), *Whither*

- opportunity?: Rising Inequality, schools, and children's life chances* (pp. 91-116). New York, NY: Russell Sage Foundation.
- Reich, W. (2000). Diagnostic Interview for Children and Adolescents (DICA). *Journal of the American Academy of Child and Adolescent Psychiatry*, 39(1), 59–66.
doi:10.1097/00004583-200001000-00017
- Reiniger, R., Robison, E., & McHugh, M. (1995). Mandated training of professionals: A means for improving reporting of suspected child abuse. *Child Abuse and Neglect*, 19(1), 63-69.
doi:10.1016/0145-2134(94)00105-4
- Rennick, J. E., Johnston, C. C., Dougherty, G., Platt, R., & Ritchie, J. A. (2002). Children's psychological responses after critical illness and exposure to invasive technology. *Journal of Developmental and Behavioral Pediatrics*, 23(3), 133-144.
doi:10.1097/00004703-200206000-00002
- Repetti, R. L., Taylor, S. E., & Seeman, T. E. (2002). Risky families: Family social environments and the mental and physical health of offspring. *Psychological Bulletin*, 128(2), 330-366. doi:10.1037/0033-2909.128.2.330
- Richards, M. P. (1996). The long term implications for children of disruptions in parental relations. In J. Brannen, & R. Edwards (Eds.), *Perspectives on parenting and childhood: Looking back and moving forward* (pp. 36-48). Swindon, England: Economic and Social Research Council.
- Richards, M. P., & Ely, M. (1998). Children's well-being, parental divorce and family life: What seems to make the difference in the longer term? In A. Smith (Ed.), *Children's potential: minimising risk and maximising resiliency* (pp. 7-24). Dunedin, New Zealand: University of Otago Press.

- Richardson, M., Kagan, R., Henry, J., Delorenzo, E., & Brophy, M. (2001). *HEROES project six month data analysis*. Albany, NY: Parsons Child and Family Center.
- Romens, S. E., McDonald, J., Svaren, J., & Pollak, S. D. (2014). Associations between early life stress and gene methylation in children. *Child Development, 86*(1), 303–309.
doi:10.1111/cdev.12270
- Romer, D. (2010). Adolescent risk taking, impulsivity, and brain development: Implications for prevention. *Developmental Psychobiology, 53*(2), 263–276. doi:10.1002/dev.20442
- Rossmann, B. B., Hughes, H. M., & Rosenberg, M. S. (2000). *Children and interparental violence: The impact of exposure*. Philadelphia, PA: Brunner/Mazel.
- Sack, W. H. (1977). The children of imprisoned fathers. *Psychiatry, 40*(2), 163–173.
doi:10.1521/00332747.1977.11023929
- Sack, W. H., Seidler, J., & Thomas, S. (1976). The children of imprisoned parents: A psychosocial exploration. *American Journal of Orthopsychiatry, 46*(4), 618–628.
doi:10.1111/j.1939-0025.1976.tb00960.x
- Samuels, J., Shinn, M., & Buckner, J. (2010). *Homeless children: Update on research, policy, programs, and opportunities*. Delmar, NY: Policy Research Associates, Inc.
- Sanders, B., & Becker-Laussen, E. (1995). The measurement of psychological maltreatment: Early data on the child abuse and trauma scale. *Child Abuse and Neglect, 19*(3), 315–323.
doi:10.1016/S0145-2134(94)00131-6
- Sapolsky, R. M., Romero, L. M., & Munck, A. (2000). How do glucocorticoids influence stress responses? Integrating permissive, suppressive, stimulatory, and preparative actions. *Endocrine Reviews, 21*(1), 55–89. doi:10.1210/edrv.21.1.0389

- Saxe, G. N., Ellis, B. H., & Brown, A. B. (2015). *Trauma Systems Therapy for traumatized children and teens*. New York, NY: Guilford Press.
- Saxe, G. N., Ellis, B. H., & Kaplow, J. (2006). *Collaborative care for traumatized children and teens: The Trauma Systems Therapy approach*. New York, NY: Guilford Press.
- Saxe, G., Vanderbilt, D., & Zuckerman, B. (2013). Traumatic stress in ill and injured children. *PTSD Research Quarterly*(2), 14. Washington, DC: The National Center for Post-Traumatic Stress Disorder. Retrieved from www.ptsd.va.gov/professional/newsletters/research-quarterly/V14N2.pdf
- Scheeringa, M. S., Weems, C. F., Cohen, J. A., Amaya-Jackson, L., & Guthrie, D. (2011). Trauma-focused cognitive-behavioral therapy for posttraumatic stress disorder in three-through six year-old children: A randomized clinical trial. *Journal of Child Psychology and Psychiatry*, 52(8), 853–860. doi:10.1111/j.1469-7610.2010.02354.x
- Semrud-Clikeman, M. (2007). *Social competence in children*. New York, NY: Springer Science.
- Senn, C. Y. (1988). *Vulnerable: Sexual abuse and people with an intellectual handicap*. Downsview, Ontario: G. Allan Roehner Institute.
- Shalev, I., Entringer, S., Wadhwa, P. D., Wolkowitz, O. M., Puterman, E., Lin, J., & Epel, E. S. (2013). Stress and telomere biology: A lifespan perspective. *Psychoneuroendocrinology*, 38(9), 1834-1842. doi:10.1016/j.psyneuen.2013.03.010
- Shannon, M. P., Lonigan, C. J., Finch, Jr., A. J., & Taylor, C. M. (1994). Children exposed to disaster: I. Epidemiology of post-traumatic symptoms and symptom profiles. *Journal of the American Academy of Child and Adolescent Psychiatry*, 33(1), 80-93. doi:10.1097/00004583-199401000-00012

Sheridan, M. A., Fox, N. A., Zeanah, C. H., McLaughlin, K. A., & Nelson III, C. A. (2012).

Variation in neural development as a result of exposure to institutionalization early in childhood. *Proceedings of the National Academy of Sciences of the United States*, 109(32), 12927-12932. doi:10.1073/pnas.1200041109

Shonkoff, J. P. (2010). Building a new biodevelopmental framework to guide the future of early childhood policy. *Child Development*, 81(1), 357–367. doi:10.1111/j.1467-8624.2009.01399.x

Silberg, J. L. (1996). *The Dissociative Features Profile*. Lutherville, MD: The Sidran Foundation.

Silove, D., Sinnerbrink, I., Field, A., Manicavasagar, V., & Steel, Z. (1997). Anxiety, depression and PTSD in asylum-seekers: Associations with pre-migration trauma and post-migration stressors. *The British Journal of Psychiatry*, 351-357. doi:10.1192/bjp.170.4.351

Silverman, A. B., Reinherz, H. Z., & Giaconia, R. M. (1996). The long-term sequelae of child and adolescent abuse: A longitudinal community study. *Child Abuse and Neglect*, 20(8), 709–723. doi:10.1016/0145-2134(96)00059-2

Simon, R., & Landis, J. (1991). *The crimes women commit: The punishments they receive*. Lexington, MA: Lexington Books.

Snyder, C. R., Hoza, B., Pelham, W. E., Rapoff, M., Ware, L., Danovsky, M., . . . Stahl, K. J. (1997). The development and validation of the Children's Hope Scale. *Journal of Pediatric Psychology*, 22(3), 399-421. doi:10.1093/jpepsy/22.3.399

Snyder, H. N. (2000). *Sexual assault of young children as reported to law enforcement: Victim, incident, and offender characteristics*. Washington, DC: U.S. Department of Justice. Retrieved from <https://www.bjs.gov/content/pub/pdf/saycrle.pdf>

- Snyder-Joy, Z. K., & Carlo, T. A. (1998). Parenting through prison walls: Incarcerated mothers and children's visitation programs. In S. L. Miller (Ed.), *Crime control and women: Feminist Implications of criminal justice policy* (pp. 130-150). Thousand Oaks, CA: Sage.
- Sogomonyan, F., & Cooper, J. L. (2010). *Trauma faced by children of military families: What every policymaker should know*. New York, NY: National Center for Children of Poverty. Retrieved from http://www.nccp.org/publications/pdf/text_938.pdf
- Spencer, C., Blades, M., & Morsley, K. (1989). *The child in the physical environment: The development of spatial knowledge and cognition*. New York, NY: Wiley.
- Sroufe, L. A. (1988). The role of infant-caregiver attachment in development. In J. Belsky, & T. M. Nezworski (Eds.), *Clinical implications of attachment* (pp. 18-30). Mahwah, NJ: Lawrence Erlbaum Associates.
- Stanton, A. M. (1980). *When mothers go to jail*. Lanham, MD: Lexington Books.
- Steele, W. (2009). *Trauma informed care: A history of helping, a history of excellence*. Grosse Pointe Woods, MI: The National Institute for Trauma and Loss in Children. Retrieved from <http://assets1.simpleclickcms.com/500051/ticsingle.pdf>
- Steven, J. E. (2014, September 17). *Tarpon Springs, FL, first trauma-informed city, embraces messy path toward peace*. Retrieved November 22, 2016, from ACEs too high: <https://acestoohigh.com/2014/09/17/tarponsprings/>
- Stevens, J. E. (2014, October 28). *Arizona ACE Consortium spreads awareness, influences prevention of childhood trauma*. Retrieved November 22, 2016, from ACEs too high: <https://acestoohigh.com/2014/10/28/arizona-ace-consortium-spreads-awareness-influences-prevention-of-childhood-trauma-effects/>

- Stevens, J. E. (2014, October 7). *Children's Resilience Initiative in Walla Walla, WA, draws spotlight to trauma-sensitive school*. Retrieved November 22, 2016, from ACEs too high: <https://acestoohigh.com/2014/10/07/childrens-resilience-initiative-in-walla-walla-wa-draws-spotlight-to-trauma-sensitive-school/>
- Stevens, J. E. (2014, September 25). *How the NFL can stop abuse AND keep its players on the field*. Retrieved November 22, 2016, from ACEs too high: <https://acestoohigh.com/2014/09/25/how-the-nfl-can-stop-abuse-and-keep-its-players-on-the-field/>
- Stevens, J. E. (2014, October 2014). *State data fuels the ACEs conversation in Iowa*. Retrieved November 22, 2016, from ACEs too high: <https://acestoohigh.com/2014/10/16/state-data-fuels-the-aces-conversation-in-iowa/>
- Stock, J. L., Bell, M. A., Boyer, D. K., & Connell, F. A. (1997). Adolescent pregnancy and sexual risk-taking among sexually abused girls. *Family Planning Perspectives, 29*(5), 200-227. doi:10.1363/2920097
- Stokes, J., Pogge, D., Sarnicola, J., & McGrath, R. (2009). Correlates of the MMPI: A Psychopathology Five (PSY-5) facet scales in an adolescent inpatient sample. *Journal of Personality Assessment, 91*(1), 48-57. doi:10.1080/00223890802484167
- Stolbach, B. C. (1997). The Children's Dissociative Experiences Scale and Posttraumatic Symptom Inventory: Rationale, development, and validation of a self-report measure. *Dissertation Abstracts International, 58*(3), 1548B.
- Strand, V. C., Hansen, S., & Courtney, D. (2013). Common elements across evidence-based trauma treatment: Discovery and implications. *Advances in Social Work, 14*(2), 334-354.

Retrieved from

<https://journals.iupui.edu/index.php/advancesinsocialwork/article/view/3052/16378>

Strand, V., Sarmiento, T., & Pasquale, L. (2005). Assessment and screening tools for trauma in children and adolescents: A review. *Trauma, Violence and Abuse*, 6(1), 55-78.

doi:10.1177/1524838004272559

Styron, T., & Janoff-Bulman, R. (1997). Childhood attachment and abuse: Long-term effects on adult attachment, depression, and conflict resolution. *Child Abuse and Neglect*, 21(10), 1015-1023. doi:10.1016/S0145-2134(97)00062-8

Substance Abuse and Mental Health Services Administration. (2008). *Results from the 2007 national survey on drug use and health: National findings*. Rockville, MD: Department of Health and Human Services. Retrieved from

http://www.alcoholandcrime.org/images/uploads/pdf_research/nsduh_2007.pdf

Sue, D. W., Arredondo, P., & McDavis, R. J. (1992). Multicultural counseling competencies and standards: A call to the profession. *Journal of Counseling and Development*, 70(4), 477–486. doi:10.1002/j.1556-6676.1992.tb01642.x

Sullivan, P., & Knutson, J. (2000). Maltreatment and disabilities: A population-based epidemiological study. *Child Abuse and Neglect*, 24(10), 1257–1273. doi:10.1016/S0145-2134(00)00190-3

Swick, K. (2005). Helping homeless families overcome barriers to successful functioning. *Early Childhood Education Journal*, 33(5), 195–200. doi:10.1007/s10643-005-0044-0

Taylor, L., Zuckerman, B., Harik, V., & Groves, B. M. (1994). Witnessing violence by young children and their mothers. *Journal of Developmental and Behavioral Pediatrics*, 15(2), 120-123. doi:10.1097/00004703-199404000-00010

- Taylor, N., & Siegfried, C. B. (2005). *Helping children in the child welfare system heal from trauma: A systems integration approach*. Los Angeles, CA and Durham, NC: National Child Traumatic Stress Network Systems Integration Working Group. Retrieved from http://www.nctsn.org/sites/default/files/assets/pdfs/A_Systems_Integration_Approach.pdf
- Teicher, M. H., Andersen, S. L., Polcari, A., Anderson, C. M., Navalta, C. P., & Kim, D. M. (2003). The neurobiological consequences of early stress and childhood maltreatment. *Neuroscience and Biobehavioral Reviews*, 27(1-2), 33-44. doi:10.1016/S0149-7634(03)00007-1
- Teicher, M. H., Anderson, C. M., & Polcari, A. (2012). Childhood maltreatment is associated with reduced volume in the hippocampal subfields CA3, dentate gyrus, and subiculum. *Proceedings of the National Academy of Sciences of the United States of America*, 109(9), 563–572. doi:10.1073/pnas.1115396109
- Terr, L. (1983). Chowchilla revisited: The effects of psychic trauma four years after a school-bus kidnapping. *American Journal of Psychiatry*, 140(12), 1543-1550. doi:10.1176/ajp.140.12.1543
- The National Child Traumatic Stress Network. (n.d.). *Tips for parents on media coverage*. Retrieved from http://www.nctsn.org/sites/default/files/assets/pdfs/tips_for_parents_media_final.pdf
- Thompson, R. A. (1998). Emotional competence and the development of self. *Psychological Inquiry*, 9(4), 308-309. doi:10.1207/s15327965pli0904_14
- Till, G. (2014). H. 762, *The adverse childhood experience questionnaire*. Montpelier, VT: Vermont Legislature. Retrieved from <http://www.leg.state.vt.us/docs/2014/bills/Intro/H-762.pdf>

- Trice, A. D. (1997). *Risk and protective factors for school and community problems for children of incarcerated women*. Washington, DC: Society for Research in Child Development.
- Trickey, D., Siddaway, A. P., Meiser-Stedman, R., Serpell, L., & Field, A. P. (2012). A meta-analysis of risk factors for post-traumatic stress disorder in children and adolescents. *Clinical Psychology Review, 32*(2), 122–138. doi:10.1016/j.cpr.2011.12.001
- U.S. Department of Health and Human Services. (2016). *Organ procurement and transplantation network: Current U.S. waiting list*. Washington, DC: U.S. Department of Health and Human Services. Retrieved November 3, 2016, from <https://optn.transplant.hrsa.gov/data/view-data-reports/national-data/#>
- U.S. Department of Health and Human Services. (2016). *Organ procurement and transplantation network: Transplants in the U.S. by recipient age*. Washington, DC: U.S. Department of Health and Human Services. Retrieved from <https://optn.transplant.hrsa.gov/data/view-data-reports/national-data/#>
- U.S. Department of Health and Human Services, & Administration on Children, Youth, and Families. (2001). *Understanding the effects of maltreatment on early brain development*. Washington, DC: Government Printing Office. Retrieved from <https://www.childwelfare.gov/pubPDFs/earlybrain.pdf>
- U.S. Department of Health and Human Services, Administration for Children and Families, Administration on Children, Youth and Families, & Children's Bureau. (2016). *Child maltreatment 2014*. Washington, DC. Retrieved from <http://www.acf.hhs.gov/sites/default/files/cb/cm2014.pdf>
- U.S. Department of Health and Human Services, Administration for Children and Families, Administration on Children, Youth and Families, & Children's Bureau. (2016). *Child*

- maltreatment 2014*. Washington, DC. Retrieved from
<http://www.acf.hhs.gov/sites/default/files/cb/cm2014.pdf>
- Udwin, O. (1993). Children's reactions to traumatic events. *Journal of Child Psychology and Psychiatry*, 34(2), 115–127. doi:10.1111/j.1469-7610.1993.tb00974.x
- van Dijke, A., Ford, J. D., & van der Hart, O. (2015). Association of childhood complex trauma and dissociation with complex posttraumatic stress disorder symptoms in adulthood. *Journal of Trauma and Dissociation*, 16(4), 428-441.
doi:10.1080/15299732.2015.1016253
- Villani, S. (2001). Impact of media on children and adolescents: A 10-year review of the research. *Journal of the American Academy of Child and Adolescent Psychiatry*, 40(4), 392–401. doi:10.1097/00004583-200104000-00007
- Vythilingam, M., Heim, C., Newport, J., Miller, A. H., Anderson, E., Bronen, R., . . . Bremner, J. D. (2002). Childhood trauma associated with smaller hippocampal volume in women with major depression. *American Journal of Psychiatry*, 159(12), 2072–2080.
doi:10.1176/appi.ajp.159.12.2072
- Wadsworth, M., Raviv, T., Reinhard, C., Wolff, B., Santiago, C., & Einhorn, L. (2008). An indirect effects model of the association between poverty and child functioning: The role of children's poverty-related stress. *Journal of Loss and Trauma*, 13(2-3), 156-185.
doi:10.1080/15325020701742185
- Wallerstein, J. S., & Lewis, J. M. (2004). The unexpected legacy of divorce: Report of a 25-year study. *Psychoanalytic Psychology*, 21(3), 353–370. doi:10.1037/0736-9735.21.3.353

- Ward, E., DeSantis, C., Robbins, A., Kohler, B., & Jemal, A. (2014). Childhood and adolescent cancer statistics. *CA: A Cancer Journal for Clinicians*, 64(2), 83-103.
doi:10.3322/caac.21219
- Way, I. F., Applegate, B., Cai, X., Frank, L. K., Black-Pond, C., & Yelsma, P. (2010). Children's Alexithymia Measure (CAM): A new instrument for screening difficulties with emotional expression. *Journal of Child and Adolescent Trauma*, 3(4), 303-318.
doi:10.1080/19361521.2010.523778
- Weder, N., Zhang, H., Yang, B. Z., Simen, A., Jackowski, A., Lipschitz, D., . . . Kaufman, J. (2014). Child abuse, depression, and methylation in genes involved with stress, neural plasticity, and brain circuitry. *Journal of the American Academy of Child and Adolescent Psychiatry*, 53(4), 417-424. doi:10.1016/j.jaac.2013.12.025
- Weiner, D., Schneider, A., & Lyons, J. (2009). Evidence-based treatments for trauma among culturally diverse foster care youth: Treatment retention and outcomes. *Children and Youth Services Review*, 31(11), 1199-1205. doi:10.1016/j.childyouth.2009.08.013
- Wells, M. G., Burlingame, G. M., & Lambert, M. J. (1999). The Youth Outcome Questionnaire (Y-OQ). In M. E. Maruish (Ed.), *The use of psychological testing for treatment planning and outcomes assessment: Volume 1: General considerations* (pp. 450-468). Mahwah, NJ: Lawrence Erlbaum Associates.
- Wells, M. G., Burlingame, G. M., Lambert, M. J., Hoag, M. J., & Hope, C. A. (1996). Conceptualization and measurement of patient change during psychotherapy: Development of the Outcome Questionnaire and Youth Outcome Questionnaire. *Psychotherapy: Theory, Research, Practice, Training*, 33(2), 275-283. doi:10.1037/0033-3204.33.2.275

- Whitfield, C. L., Anda, R. F., Dube, S. R., & Felitti, V. J. (2003). Violent childhood experiences and the risk of intimate partner violence in adults. *Journal of Interpersonal Violence*, 18(2), 166-185. doi:10.1177/0886260502238733
- Whitfield, C. L., Dube, S. R., Felitti, V. J., & Anda, R. F. (2005). Adverse childhood experiences and hallucinations. *Child Abuse and Neglect*, 29(7), 797–810. doi:10.1016/j.chiabu.2005.01.004
- Widom, C., Marmorstein, N., & White, H. (2006). Childhood victimization and illicit drug use in middle adulthood. *Psychology of Addictive Behaviors*, 20(4), 394-403. doi:10.1037/0893-164X.20.4.394
- Williamson, D. F., Thompson, T. J., Anda, R. F., Dietz, W. H., & Felitti, V. (2002). Body weight and obesity in adults and self-reported abuse in childhood. *International Journal of Obesity and Related Metabolic Disorders*, 26(8), 1075-1082. doi:10.1038/sj.ijo.0802038
- Wilson, J. P., & Keane, T. M. (2004). *Assessing psychological trauma and PTSD*. New York, NY: The Guilford Press.
- Wisdom, C. S., & Maxfield, M. G. (2001). *An update on the "cycle of violence"*. Washington, DC: U.S. Department of Justice. Retrieved from <https://www.ncjrs.gov/pdffiles1/nij/184894.pdf>
- Wisdom, C. S., Ireland, T., & Glynn, P. J. (1995). Alcohol abuse in abused and neglected children followed-up: Are they at increased risk? *Journal of Studies on Alcohol*, 56(2), 207–217. doi:10.15288/jsa.1995.56.207
- World Health Organization. (2001). *Mental disorders affect one if four people*. Geneva, Switzerland. Retrieved November 16, 2016, from http://www.who.int/whr/2001/media_centre/press_release/en/

- World Health Organization. (2009). *Addressing adverse childhood experiences to improve public health: Expert consultation*. Geneva, Switzerland: World Health Organization. Retrieved from http://www.who.int/violence_injury_prevention/violence/activities/adverse_childhood_experiences/global_research_network_may_2009.pdf
- Young, M. A. (2001). *Victim assistance: Frontiers and fundamentals*. Alexandria, VA: National Organization for Victim Assistance.
- Zelechowski, A. D., Sharma, R., Beserra, K., Miguel, J. L., DeMarco, M., & Spinazzola, J. (2013). Traumatized youth in residential treatment settings: Prevalence, clinical presentation, treatment, and policy implications. *Journal of Family Violence*, 28(7), 639–652. doi:10.1007/s10896-013-9534-9
- Zimet, G. D., Dahlem, N. W., Zimet, S. G., & Farley, G. K. (1988). The Multidimensional Scale of Perceived Social Support. *Journal of Personality Assessment*, 52(1), 30-41. doi:10.1207/s15327752jpa5201_2
- Zuckerman, B., & Bresnahan, K. (1991). Developmental and behavioral consequences of prenatal drug and alcohol exposure. 38(6), 1387-1406. doi:10.1016/S0031-3955(16)38226-8
- Zuravin, S. (2001). Issues pertinent to defining child neglect. In T. D. Morton, & B. Salovitz (Eds.), *The CPS response to child neglect: An administrator's guide to theory, policy, program design and case practice* (pp. 2-1–2-22). Duluth, GA: National Resource on Child Maltreatment.




Figure 1. The ACE Pyramid. Reprinted from “Mechanism by which adverse childhood experiences influence health and well-being throughout the lifespan”, by Centers for Disease Control and Prevention, n.d. Retrieved from <https://www.cdc.gov/violenceprevention/acestudy/about.html>




Figure 2. Pathways to poverty. Adapted from “Adverse childhood experiences: Connecting a developmental lens to the health of society – Pathways to poverty [PowerPoint slides]”, by Anda, 2011. Retrieved from https://www.in.gov/dcs/files/2Dr_Anda5steps.pdf

Table 1.

Standardized Measures

Name of Measure	Domains	Targeted Age	Format	Source
Structured Interview for Disorders of Extreme Stress SIDES	Affect Dysregulation Somatization Attention/Consciousness Self-Perception Relations with Others Systems of Meaning	17-18+	Clinician-Based Interview Self-Report (SIDES-SR)	Pelcovitz et al., 1997
Structured Interview for Disorders of Extreme Stress Adolescent Version SIDES-A	Affect Dysregulation Somatization Attention/Consciousness Self-Perception Relations with Others Systems of Meaning	12-18	Clinician-Based Interview	Habib & Labruna, 2011
Child and Adolescent Needs and Strengths- Trauma Comprehensive CANS Trauma	Attachment Affect Dysregulation Behavioral Control Dissociation Biology Trauma Exposure Caregiver Functioning Strengths	3-18	Clinician Rating/Information Integration Tool	Kisiel, Fehrenbach, Small, & Lyons, 2009
Trauma Symptom Checklist for Children TSCC	Anger Anxiety Depression Dissociation Fantasy Sexual Concerns Somatic/Biology	8-16	Self-Report	Briere, 1996
Trauma Symptom Checklist for Young Children TSCYC	Traumatic Stress Somatic/Biology Anxiety/Mood	3-12	Caregiver Report	Briere et al., 2001
Youth Outcome Questionnaire YOQ – 2.01	Interpersonal Distress Somatic Concerns Interpersonal Relationships Social Problems Behavioral Control	4-17	Caregiver Report	Wells, Burlingame, Lambert, Hoag, & Hope, 1996
Youth Outcome Questionnaire Self-Report YOQ – 2.0 SR	Interpersonal Distress Somatic Concerns Interpersonal Relationships Social Problems Behavioral Dysfunction	12-18	Self-Report	Wells, Burlingame, & Lambert, 1999

Name of Measure	Domains	Targeted Age	Format	Source
Children's Alexithymia Measure CAM	Affect Dysregulation	5-17	Observer Report	Way, et al., 2010
Difficulties in Emotional Regulation Scale DERS	Affect Dysregulation	18-60	Self-Report	Gratz & Roemer, 2004
Five Factor Personality Inventory for Children FFPI-C	Affect Dysregulation Behavioral Control	9-18	Self-Report	Kroes, Veerman, & De Bruyn, 2005
Minnesota Multiphasic Personality Inventory for Adolescents MMPI-A	Affect Dysregulation Somatic/Biology (Hypochondriasis and Hysteria Scales)	14-18	Self-Report	Stokes, Pogge, Sarnicola, & McGrath, 2009
Personality Inventory for Youth/ Personality Inventory for Children, Second Edition PIC-2	Affect Dysregulation Cognition Somatic/Biology	5-19	Parent Report Youth Self-Report Teacher Report	Negy, Lachar, Gruber, & Garza, 2001
Affect Intensity and Reactivity Measure for Youth AIR-Y	Affect Dysregulation	10-17	Self-Report	Jones, Leen-Feldner, Olatunji, & Hawks, 2009
Abbreviated Dysregulation Inventory ADI	Affect Dysregulation Cognitive Behavioral Control	12-18	Self-Report	Mezzich, Tarter, Giancola, & Kirisci, 2001
Attachment Questionnaire for Children AQC	Attachment	9-18	Self-Report	Muris, Vlaeyen, & Meesters, 2001
Inventory of Parent and Peer Attachment IPPA	Attachment	3	5 Story Completion Tasks	Armsden & Greenberg, 1987
Child Dissociative Checklist CDC	Dissociation	5-12	Caregiver Report	Putnam & Helmers, 1993
Adolescent Dissociative Experiences Scale A-DES	Dissociation	11-18	Self-Report	Armstrong, Putnam, Carlson, Libero, & Smith, 1997
Children's Perceptual Alterations Scale CPAS	Dissociation	8-12	Self-Report	Evers-Szostak & Sanders, 1992
Dissociative Features Profile (DFP)	Dissociation	5-17	Clinician report	Silberg, 1996
Children's Dissociative Experiences Scale & Posttraumatic Symptom Inventory CDES & PTSI	Dissociation	6-12	Self-report	Stolbach, 1997 Cloitre, et al., 2009

Name of Measure	Domains	Targeted Age	Format	Source
Children's Attributions and Perceptions Scale CAPS	Cognition	7-17	Clinician Interview	Mannarino, Cohen, & Berman, 1994
Children's Hope Scale	Cognition	8-19	Self-Report	Snyder, et al., 1997
Behavior Rating Inventory of Executive Function BRIEF	Cognition	5-18	Caregiver Report Teacher Report	Gioia, Isquith, Guy, & Kenworthy, 2000
Behavior Assessment System for Children-2 BASC-2	Cognition Behavioral Control Somatic/Biology	2 – 21	Youth Self-Report Caregiver Report Teacher Report	Kamphaus, et al., 1999
Child Behavior Checklist CBCL	Behavioral Control Affect Dysregulation Somatic/Biology Cognition	1.5-18	Caregiver Report	Achenbach, 2011
Child Sexual Behavior Inventory CSBI	Behavioral Control	2-12	Caregiver Report	Friedrich, et al., 1992
Perceived Competence Scales for Children	Behavioral Control	Not Specified	Self-Report	Harter, 1982
Piers-Harris Children's Self-Concept Scale, 2nd ed. Piers-Harris-2	Self-Concept	7-18	Self-report	Piers & Harris, 2002
Traumatic Events Screening Inventory Parent Report Revised TESI-CRF-R	Trauma Exposure/History	4-18	Caregiver Report Youth Report Interview Versions	Ippen, et al., 2002
Juvenile Victimization Questionnaire JVQ	Trauma Exposure/History	2-17	Caregiver Report Youth Report	Hamby S. L., Finkelhor, Ormond, & Turner, 2004
UCLA PTSD Reaction Index	Trauma Exposure/History (Exposure Portion) PTSD Symptoms	7-18	Caregiver Report Youth Report	Pynoos, Rodriguez, Steinberg, Stuber, & Frederick, 1998
Trauma History Checklist and Interview THC	Trauma Exposure/History	13 - adult	Self-Report Interview	Habib & Labruna, 2011
Dimensions of Stressful Events Rating Scale DOSE	Trauma Exposure/History	6-18	Clinician Administered	Fletcher, 1994
Parent-Child Relationship Inventory PCRI	Trauma Exposure/History	3-15	Caregiver Report	Gerard, 1994
Clinician Administered PTSD Scale for Children and Adolescents CAPS-CA	PTSD Symptoms	8-15	Clinician Administered Youth Interview	Nader, et al., 2004

Name of Measure	Domains	Targeted Age	Format	Source
Diagnostic Interview for Children and Adolescents DICA	PTSD Symptoms	6-17	Semi-Structured Interview	Reich, 2000
Family Environment Scale FES	Caregiver/Family Functioning Response to Trauma	5-11: Pictorial Adaptation 12-18+	Caregiver Report Youth Report	Moos, 1990
Family Assessment of Needs and Strengths: Trauma Exposure and Adaptation FANS-TEA	Caregiver/Family Functioning Response to Trauma	Not Specified	Clinician Interview	Lyons, et al., 2009
Parenting Stress Index-Short Form PSI-SF	Caregiver/Family Functioning & Response to Trauma	10-12	Caregiver Report	Abidin, 1995
Parent Emotional Reaction Questionnaire PERQ	Caregiver/Family Functioning Response to Trauma	Not Specified	Caregiver Report	Cohen & Mannarino, 1996
Brief Symptom Inventory BSI	Caregiver/Family Functioning Response to Trauma	13+	Caregiver Report	Derogatis & Melisaratos , 1983
Coping Responses Inventory Youth CRI-Y	Resilience, Coping, & Strengths	12-18	Self-Report	Moos, 1993
Children's Coping Strategies Checklist CCSC	Resilience, Coping, & Strengths	9 – 13	Self-Report	Ayers, Sandler, West, & Roosa, 1996
Multidimensional Scale of Perceived Social Support MSPSS	Resilience, Coping, & Strengths	Children, adolescents, and adults	Self-report	Zimet, Dahlem, Zimet, & Farley, 1988
World Assumptions Scale WAS	Resilience, Coping, & Strengths	13+	Self-report	Janoff-Bulman, 1989
Strengths and Difficulties Questionnaire SDQ	Resilience, Coping, & Strengths	3-18+	Caregiver Report Teacher Report Youth Report	Goodman, 1997

Note. Adapted from “Standardized measures to assess complex trauma,” by National Child Traumatic Stress Network, n.d. Retrieved from <http://www.nctsn.org/trauma-types/complex-trauma/standardized-measures-assess-complex-trauma>

Table 2

Trauma Treatment

Name of Intervention	Targeted Populations	Modality
Adapted Dialectical Behavior Therapy for Special Populations (DBT-SP)	8-21; both males and females; for youth experiencing a wide range of traumas	Individual
Alternatives for Families - A Cognitive Behavioral Therapy (AF-CBT)	School-age children; for youth experiencing a wide range of traumas	Individual and Family
Assessment-Based Treatment for Traumatized Children: Trauma Assessment Pathway (TAP)	0-18; both males and females; for children who have experienced a wide range of traumas	Individual, Family, and Systems
Attachment and Biobehavioral Catch-up (ABC)	Birth – 24 months; both males and females; for low-income families who have experienced neglect, abuse, domestic violence, placement instability	Individual and Family
Attachment, Self-Regulation, and Competence (ARC): A Comprehensive Framework for Intervention with Complexly Traumatized Youth	2-21; both males and females; for children, caregivers, and systems that have experienced a wide range of traumas	Individual, Family, and Systems
Child Adult Relationship Enhancement (CARE)	Children of all ages and their caregivers; both males and females	Family and Systems
Child and Family Traumatic Stress Intervention (CFTSI)	7-18; both males and females; for parents and children who may have complex trauma histories	Individual, Family, and Systems
Child Development-Community Policing Program	0-18+; both males and females; for children and families in the aftermath of crime and violence.	Individual, Family, and Systems
Child-Parent Psychotherapy (CPP)	0-6; both males and females; for youth who have experienced a wide range of traumas and parents with chronic trauma	Individual, Family, and Systems
Cognitive Behavioral Intervention for Trauma in Schools (CBITS)	10-15; both males and females; for children who have experienced a wide range of traumas	Individual, Family, and Systems
Combined Parent Child Cognitive-Behavioral Approach for Children and Families At-Risk for Child Physical Abuse (CPC-CBT)	4-17; both male and female; for families with a history of physical abuse and inappropriate physical discipline/coercive parenting strategies	Individual, Group, and Family
Combined TF-CBT and SSRI Treatment (2007)	10-18; females	Individual and Family
Community Outreach Program (COPE)	4-18; both males and females; for traumatized children who are presenting with behavior or social-emotional problems	Individual and Family
Culturally Modified Trauma-Focused Treatment (CM-TFT)	4-18; both males and females; Latino/Hispanic; for youth who have experienced a wide range of traumas	Individual and Family
Family Advocate Program	18-70; both males and females; for youth who present with anxiety, depression, PTSD symptoms, and/or traumatic loss	Family
Forensically-Sensitive Therapy	4-17; predominantly female; for youth presenting problems ranging from anxiety and depression to risk-taking behaviors and functional impairment. Program is designed for a mental health clinic.	Individual and Family
Group Treatment for Children Affected by Domestic Violence	5-no upper limit; both males and females; for children and their nonoffending parents who have been exposed to DV	Individual, Family, and Systems
Honoring Children, Making Relatives	3-7; both males and females; for American Indian and Alaska Native children	Individual and Family
Honoring Children, Mending the Circle	3-18; both males and females; for American Indian and Alaska Native children	Individual
Honoring Children, Respectful Ways	3-12; both males and females; for American Indian and Alaska Native children	Individual

Name of Intervention	Targeted Populations	Modality
Integrative Treatment of Complex Trauma (ITCT-C, ITCT-A)	2-21; both males and females; for Hispanic-American, African-American, Caucasian, Asian-American; for youth who may have complex trauma histories	Individual, Family, and Systems
International Family Adult and Child Enhancement Services (IFACES)	6-12; both males and females; for refugee and immigrant children who have experienced trauma as a result of war or displacement	Individual
Let's Connect (LC)	3-15; both males and females; for children who have experienced a wide range of traumas	Individual and Family
Parent-Child Interaction Therapy (PCIT)	2-12; both males and females	Individual, Family, and Systems
Problematic Sexual Behavior-Cognitive-Behavioral Therapy for School-Age Children (PSB-CBT-S)	7-12; both males and females; for children with problematic sexual behavior may or may not have a history of trauma	Individual, Family, and Systems
Psychological First Aid (PFA)	0-120; both males and females; for individuals immediately following disasters, terrorism, and other emergencies	Individual
Real Life Heroes (RLH)	6-12, plus adolescents (13-19) with delays in social, emotional or cognitive functioning; both males and females; for children who have experienced a wide range of traumas	Individual, Family, and Systems
Risk Reduction through Family Therapy (RRTF)	13-18, both males and females; for adolescents and family; primary trauma type is childhood sexual abuse/sexual assault	Family
Safe Harbor Program	6-21; both males and females; provided in schools for children and adolescents exposed to trauma and violence who may present with a range of problems and symptoms	Individual, Group, and Family
Safety, Mentoring, Advocacy, Recovery, and Treatment (SMART)	3-11; both males and females; to date the model has been effectively used with primarily African-American children; majority of families are low income	Individual, Family, and Systems
Sanctuary Model	4-no upper limit; both males and females; evidence-supported template for system change based on the active creation and maintenance of a nonviolent, democratic, productive community to help people heal from trauma	Systems
Sanctuary Model Plus (IRIS Project)	Children and adolescents placed in residential treatment centers and their families	Group and Systems
Skills for Psychological Recovery (SPR)	5-120; both males and females	Individual and Family
Skills Training in Affective and Interpersonal Regulation/Narrative Story-Telling (STAIR/NST)	12-21; for females who have experienced sexual/physical abuse and a range of additional traumas, including community violence, domestic violence, and sexual assault	Individual and Group
Southeast Asian Teen Village	adolescents; females, Southeast Asian (mostly Hmong)	Group
Streetwork Project	13-23; both males and females; harm reduction program good with a wide variety of ethnic/racial groups, religious group, and the LGBTQ community	Individual, Group, and Systems
Strengthening Family Coping Resources (SFCR)	0-no upper limit; both males and females; for families experiencing economic hardship	Family
Structured Psychotherapy for Adolescents Responding to Chronic Stress (SPARCS)	12-21; both males and females; for adolescents with Complex Trauma, e.g. adolescents exposed to chronic interpersonal trauma (such as ongoing physical abuse) and/or separate types of trauma (e.g. community violence, sexual assault).	Group
Trauma Adapted Family Connections (TA-FC)	0-18; both males and females; who reside in the household; families experiencing complex development trauma, at risk of neglect	Individual, Group, and Family

Name of Intervention	Targeted Populations	Modality
Trauma Affect Regulation: Guide for Education and Therapy (TARGET)	10-18+; both males and females; for children and caregivers experiencing traumatic stress; very frequently with single parents or with families whose children have limited contact with biological parents (e.g., foster kids, residential placements), and diversity of religious affiliations	Individual, Family, Group, and Systems
Trauma and Grief Component Therapy for Adolescents (TGCT-A)	12-20; both males and females; for trauma-exposed or traumatically bereaved older children and adolescents	Individual, Family, Group, and Systems
Trauma-Focused Cognitive Behavioral Therapy (TF-CBT)	3-21; both males and females; for children with Posttraumatic Stress Disorder (PTSD) or other problems related to traumatic life experiences, and their parents or primary caregivers	Individual and Family
Trauma-Focused Coping in Schools (TFC) (AKA: Multimodality Trauma Treatment Trauma-Focused Coping-MMTT)	6-18; both males and females; for children exposed to single incident trauma and targets posttraumatic stress disorder (PTSD) and collateral symptoms of depression, anxiety, anger, and external locus of control	Individual and Group
Trauma-Informed Organizational Self-Assessment	6-19; both males and females; for children who have experienced a wide range of traumas	Individual, Family, and Systems
Trauma Systems Therapy (TST)	6-19; both males and females; for youth who have experienced a wide range of traumas	Systems
Trauma Systems Therapy for Refugees (TST-R) (2016) (PDF)	10-18; both males and females; newly arriving, recently resettled, and established refugee youth and communities	Systems

Note. Reprinted from “National Child Traumatic Stress Network empirically supported treatments and promising practices”, by National Child Traumatic Stress Network, n.d. Retrieved from <http://www.nctsn.org/resources/topics/treatments-that-work/promising-practices>