

Solving Problems Involving Linear Functions

Jonathan R. Bacolod

Sauyo High School

How to Solve Problems Involving Linear Functions?

1. Understand and analyze the problem.

How to Solve Problems Involving Linear Functions?

1. Understand and analyze the problem.
 - 1.1 Determine the dependent variable $f(x)$ and independent variable x in the problem.

How to Solve Problems Involving Linear Functions?

1. Understand and analyze the problem.
 - 1.1 Determine the dependent variable $f(x)$ and independent variable x in the problem.
 - 1.2 Determine the initial state b of the dependent variable.

How to Solve Problems Involving Linear Functions?

1. Understand and analyze the problem.
 - 1.1 Determine the dependent variable $f(x)$ and independent variable x in the problem.
 - 1.2 Determine the initial state b of the dependent variable.
 - 1.3 Compute for the slope.

How to Solve Problems Involving Linear Functions?

1. Understand and analyze the problem.
 - 1.1 Determine the dependent variable $f(x)$ and independent variable x in the problem.
 - 1.2 Determine the initial state b of the dependent variable.
 - 1.3 Compute for the slope.
2. Use the facts of the problem to form a linear function.

How to Solve Problems Involving Linear Functions?

1. Understand and analyze the problem.
 - 1.1 Determine the dependent variable $f(x)$ and independent variable x in the problem.
 - 1.2 Determine the initial state b of the dependent variable.
 - 1.3 Compute for the slope.
2. Use the facts of the problem to form a linear function.
3. Graph the linear function.

Example 1

It has been observed that a particular plant's growth is directly proportional to time. It measured 2 cm when it arrived at the nursery and 2.5 cm exactly one week later. If the plant continues to grow at this rate, determine the function that represents the plant's growth and graph it.

Example 1

Step 1: Understand and analyze the problem.

It has been observed that a particular plant's growth is directly proportional to time. It measured 2 cm when it arrived at the nursery and 2.5 cm exactly one week later. If the plant continues to grow at this rate, determine the function that represents the plant's growth and graph it.

Example 1

Step 1: Understand and analyze the problem.

It has been observed that a particular plant's growth is directly proportional to time. It measured 2 cm when it arrived at the nursery and 2.5 cm exactly one week later. If the plant continues to grow at this rate, determine the function that represents the plant's growth and graph it.

Let: $f(x) =$

Example 1

Step 1: Understand and analyze the problem.

It has been observed that a particular plant's growth is directly proportional to time. It measured 2 cm when it arrived at the nursery and 2.5 cm exactly one week later. If the plant continues to grow at this rate, determine the function that represents the plant's growth and graph it.

Let: $f(x)$ = plant's height

Example 1

Step 1: Understand and analyze the problem.

It has been observed that a particular plant's growth is directly proportional to time. It measured 2 cm when it arrived at the nursery and 2.5 cm exactly one week later. If the plant continues to grow at this rate, determine the function that represents the plant's growth and graph it.

Let: $f(x)$ = plant's height
 x =

Example 1

Step 1: Understand and analyze the problem.

It has been observed that a particular plant's growth is directly proportional to time. It measured 2 cm when it arrived at the nursery and 2.5 cm exactly one week later. If the plant continues to grow at this rate, determine the function that represents the plant's growth and graph it.

Let:

$f(x)$	=	plant's height
x	=	time measured per week

Example 1

Step 1: Understand and analyze the problem.

It has been observed that a particular plant's growth is directly proportional to time. It measured 2 cm when it arrived at the nursery and 2.5 cm exactly one week later. If the plant continues to grow at this rate, determine the function that represents the plant's growth and graph it.

Let:	$f(x)$	=	plant's height
	x	=	time measured per week
Given:	b	=	

Example 1

Step 1: Understand and analyze the problem.

It has been observed that a particular plant's growth is directly proportional to time. It measured 2 cm when it arrived at the nursery and 2.5 cm exactly one week later. If the plant continues to grow at this rate, determine the function that represents the plant's growth and graph it.

Let:	$f(x)$	=	plant's height
	x	=	time measured per week
Given:	b	=	2 cm (initial height)

Example 1

Step 1: Understand and analyze the problem.

It has been observed that a particular plant's growth is directly proportional to time. It measured 2 cm when it arrived at the nursery and 2.5 cm exactly one week later. If the plant continues to grow at this rate, determine the function that represents the plant's growth and graph it.

Let:	$f(x)$	=	plant's height
	x	=	time measured per week
Given:	b	=	2 cm (initial height)
Find:	m	=	

Example 1

Step 1: Understand and analyze the problem.

It has been observed that a particular plant's growth is directly proportional to time. It measured 2 cm when it arrived at the nursery and 2.5 cm exactly one week later. If the plant continues to grow at this rate, determine the function that represents the plant's growth and graph it.

Let:	$f(x)$	=	plant's height
	x	=	time measured per week
Given:	b	=	2 cm (initial height)
Find:	m	=	$2.5 - 2$

Example 1

Step 2: Use the facts of the problem to form a linear function.

It has been observed that a particular plant's growth is directly proportional to time. It measured 2 cm when it arrived at the nursery and 2.5 cm exactly one week later. If the plant continues to grow at this rate, determine the function that represents the plant's growth and graph it.

Let:	$f(x)$	=	plant's height
	x	=	time measured per week
Given:	b	=	2 cm (initial height)
Find:	m	=	$2.5 - 2 = 0.5$ cm per week

Example 1

Step 2: Use the facts of the problem to form a linear function.

It has been observed that a particular plant's growth is directly proportional to time. It measured 2 cm when it arrived at the nursery and 2.5 cm exactly one week later. If the plant continues to grow at this rate, determine the function that represents the plant's growth and graph it.

Let:	$f(x)$	=	plant's height
	x	=	time measured per week
Given:	b	=	2 cm (initial height)
Find:	m	=	$2.5 - 2 = 0.5$ cm per week
Function:	$f(x)$	=	$mx + b$

Example 1

Step 2: Use the facts of the problem to form a linear function.

It has been observed that a particular plant's growth is directly proportional to time. It measured 2 cm when it arrived at the nursery and 2.5 cm exactly one week later. If the plant continues to grow at this rate, determine the function that represents the plant's growth and graph it.

Let:	$f(x)$	=	plant's height
	x	=	time measured per week
Given:	b	=	2 cm (initial height)
Find:	m	=	$2.5 - 2 = 0.5$ cm per week
Function:	$f(x)$	=	$mx + b$
	$f(x)$	=	

Example 1

Step 2: Use the facts of the problem to form a linear function.

It has been observed that a particular plant's growth is directly proportional to time. It measured 2 cm when it arrived at the nursery and 2.5 cm exactly one week later. If the plant continues to grow at this rate, determine the function that represents the plant's growth and graph it.

Let:	$f(x)$	=	plant's height
	x	=	time measured per week
Given:	b	=	2 cm (initial height)
Find:	m	=	$2.5 - 2 = 0.5$ cm per week
Function:	$f(x)$	=	$mx + b$
	$f(x)$	=	$0.5x + 2$

Example 1

Step 3: Graph the linear function.

$$f(x) = 0.5x + 2$$

Example 1

Step 3: Graph the linear function.

$$f(x) = 0.5x + 2$$
$$m = \frac{\text{rise}}{\text{run}} = 0.5 = \frac{1}{2},$$

Example 1

Step 3: Graph the linear function.

$$f(x) = 0.5x + 2$$

$$m = \frac{\text{rise}}{\text{run}} = 0.5 = \frac{1}{2}, b = 2$$

Example 1

Step 3: Graph the linear function.

$$f(x) = 0.5x + 2$$

$$m = \frac{\text{rise}}{\text{run}} = 0.5 = \frac{1}{2}, b = 2$$




Example 1

Step 3: Graph the linear function.

$$f(x) = 0.5x + 2$$

$$m = \frac{\text{rise}}{\text{run}} = 0.5 = \frac{1}{2}, b = 2$$




Example 1

Step 3: Graph the linear function.

$$f(x) = 0.5x + 2$$

$$m = \frac{\text{rise}}{\text{run}} = 0.5 = \frac{1}{2}, b = 2$$




Example 1

Step 3: Graph the linear function.

$$f(x) = 0.5x + 2$$

$$m = \frac{\text{rise}}{\text{run}} = 0.5 = \frac{1}{2}, b = 2$$




Example 1

Step 3: Graph the linear function.

$$f(x) = 0.5x + 2$$

$$m = \frac{\text{rise}}{\text{run}} = 0.5 = \frac{1}{2}, b = 2$$




Example 1

Step 3: Graph the linear function.

$$f(x) = 0.5x + 2$$

$$m = \frac{\text{rise}}{\text{run}} = 0.5 = \frac{1}{2}, b = 2$$



Example 2

Emman often rides a taxi from one place to another. The standard fare in riding a taxi is Php 40 as a flag-down rate plus Php 5 for every 200 meters or a fraction of it.

Determine the function that represents the fare and graph it.

Example 2

Step 1: Understand and analyze the problem.

Emman often rides a taxi from one place to another. The standard fare in riding a taxi is Php 40 as a flag-down rate plus Php 5 for every 200 meters or a fraction of it. Determine the function that represents the fare and graph it.

Example 2

Step 1: Understand and analyze the problem.

Emman often rides a taxi from one place to another. The standard fare in riding a taxi is Php 40 as a flag-down rate plus Php 5 for every 200 meters or a fraction of it. Determine the function that represents the fare and graph it.

Let: $f(x) =$

Example 2

Step 1: Understand and analyze the problem.

Emman often rides a taxi from one place to another. The standard fare in riding a taxi is Php 40 as a flag-down rate plus Php 5 for every 200 meters or a fraction of it. Determine the function that represents the fare and graph it.

Let: $f(x)$ = standard fare

Example 2

Step 1: Understand and analyze the problem.

Emman often rides a taxi from one place to another. The standard fare in riding a taxi is Php 40 as a flag-down rate plus Php 5 for every 200 meters or a fraction of it. Determine the function that represents the fare and graph it.

Let: $f(x)$ = standard fare
 x =

Example 2

Step 1: Understand and analyze the problem.

Emman often rides a taxi from one place to another. The standard fare in riding a taxi is Php 40 as a flag-down rate plus Php 5 for every 200 meters or a fraction of it. Determine the function that represents the fare and graph it.

Let:

$f(x)$	=	standard fare
x	=	distance measured per 200 m.

Example 2

Step 1: Understand and analyze the problem.

Emman often rides a taxi from one place to another. The standard fare in riding a taxi is Php 40 as a flag-down rate plus Php 5 for every 200 meters or a fraction of it. Determine the function that represents the fare and graph it.

Let: $f(x)$ = standard fare
 x = distance measured per 200 m.
Given: b =

Example 2

Step 1: Understand and analyze the problem.

Emman often rides a taxi from one place to another. The standard fare in riding a taxi is Php 40 as a flag-down rate plus Php 5 for every 200 meters or a fraction of it. Determine the function that represents the fare and graph it.

Let:	$f(x)$	=	standard fare
	x	=	distance measured per 200 m.
Given:	b	=	Php 40 (initial fare)

Example 2

Step 1: Understand and analyze the problem.

Emman often rides a taxi from one place to another. The standard fare in riding a taxi is Php 40 as a flag-down rate plus Php 5 for every 200 meters or a fraction of it. Determine the function that represents the fare and graph it.

Let:	$f(x)$	=	standard fare
	x	=	distance measured per 200 m.
Given:	b	=	Php 40 (initial fare)
Find:	m	=	

Example 2

Step 2: Use the facts of the problem to form a linear function.

Emman often rides a taxi from one place to another. The standard fare in riding a taxi is Php 40 as a flag-down rate plus Php 5 for every 200 meters or a fraction of it. Determine the function that represents the fare and graph it.

Let:	$f(x)$	=	standard fare
	x	=	distance measured per 200 m.
Given:	b	=	Php 40 (initial fare)
Find:	m	=	Php 5 per 200 m.

Example 2

Step 2: Use the facts of the problem to form a linear function.

Emman often rides a taxi from one place to another. The standard fare in riding a taxi is Php 40 as a flag-down rate plus Php 5 for every 200 meters or a fraction of it. Determine the function that represents the fare and graph it.

Let:	$f(x)$	=	standard fare
	x	=	distance measured per 200 m.
Given:	b	=	Php 40 (initial fare)
Find:	m	=	Php 5 per 200 m.
Function:	$f(x)$	=	$mx + b$

Example 2

Step 2: Use the facts of the problem to form a linear function.

Emman often rides a taxi from one place to another. The standard fare in riding a taxi is Php 40 as a flag-down rate plus Php 5 for every 200 meters or a fraction of it. Determine the function that represents the fare and graph it.

Let:	$f(x)$	=	standard fare
	x	=	distance measured per 200 m.
Given:	b	=	Php 40 (initial fare)
Find:	m	=	Php 5 per 200 m.
Function:	$f(x)$	=	$mx + b$
	$f(x)$	=	

Example 2

Step 2: Use the facts of the problem to form a linear function.

Emman often rides a taxi from one place to another. The standard fare in riding a taxi is Php 40 as a flag-down rate plus Php 5 for every 200 meters or a fraction of it. Determine the function that represents the fare and graph it.

Let:	$f(x)$	=	standard fare
	x	=	distance measured per 200 m.
Given:	b	=	Php 40 (initial fare)
Find:	m	=	Php 5 per 200 m.
Function:	$f(x)$	=	$mx + b$
	$f(x)$	=	$5x + 40$

Example 2

Step 3: Graph the linear function.

$$f(x) = 5x + 40$$

Example 2

Step 3: Graph the linear function.

$$f(x) = 5x + 40$$

$$m = \frac{\text{rise}}{\text{run}} = 5 = \frac{5}{1},$$

Example 2

Step 3: Graph the linear function.

$$f(x) = 5x + 40$$


$$m = \frac{\text{rise}}{\text{run}} = 5 = \frac{5}{1}, b = 40$$

Example 2

Step 3: Graph the linear function.

$$f(x) = 5x + 40$$

$$m = \frac{\text{rise}}{\text{run}} = 5 = \frac{5}{1}, b = 40$$




Example 2

Step 3: Graph the linear function.

$$f(x) = 5x + 40$$

$$m = \frac{\text{rise}}{\text{run}} = 5 = \frac{5}{1}, b = 40$$




Example 2

Step 3: Graph the linear function.

$$f(x) = 5x + 40$$

$$m = \frac{\text{rise}}{\text{run}} = 5 = \frac{5}{1}, b = 40$$




Example 2

Step 3: Graph the linear function.

$$f(x) = 5x + 40$$

$$m = \frac{\text{rise}}{\text{run}} = 5 = \frac{5}{1}, b = 40$$




Example 2

Step 3: Graph the linear function.

$$f(x) = 5x + 40$$

$$m = \frac{\text{rise}}{\text{run}} = 5 = \frac{5}{1}, b = 40$$




Example 2

Step 3: Graph the linear function.

$$f(x) = 5x + 40$$

$$m = \frac{\text{rise}}{\text{run}} = 5 = \frac{5}{1}, b = 40$$



Example 3

A pay phone service charges Php 5 for the first three minutes and Php 1 for every minute additional or a fraction thereof. How much will a caller have to pay if his call lasts for 8 minutes? Write a rule that best describes the problem and draw its graph using any method.

Example 3

Step 1: Understand and analyze the problem.

A pay phone service charges Php 5 for the first three minutes and Php 1 for every minute additional or a fraction thereof. How much will a caller have to pay if his call lasts for 8 minutes? Write a rule that best describes the problem and draw its graph using any method.

Example 3

Step 1: Understand and analyze the problem.

A pay phone service charges Php 5 for the first three minutes and Php 1 for every minute additional or a fraction thereof. How much will a caller have to pay if his call lasts for 8 minutes? Write a rule that best describes the problem and draw its graph using any method.

Let: $f(x) =$

Example 3

Step 1: Understand and analyze the problem.

A pay phone service charges Php 5 for the first three minutes and Php 1 for every minute additional or a fraction thereof. How much will a caller have to pay if his call lasts for 8 minutes? Write a rule that best describes the problem and draw its graph using any method.

Let: $f(x) =$ call charge

Example 3

Step 1: Understand and analyze the problem.

A pay phone service charges Php 5 for the first three minutes and Php 1 for every minute additional or a fraction thereof. How much will a caller have to pay if his call lasts for 8 minutes? Write a rule that best describes the problem and draw its graph using any method.

Let: $f(x)$ = call charge
 x =

Example 3

Step 1: Understand and analyze the problem.

A pay phone service charges Php 5 for the first three minutes and Php 1 for every minute additional or a fraction thereof. How much will a caller have to pay if his call lasts for 8 minutes? Write a rule that best describes the problem and draw its graph using any method.

Let:

$f(x)$	=	call charge
x	=	time (in minutes) after 3 min.

Example 3

Step 1: Understand and analyze the problem.

A pay phone service charges Php 5 for the first three minutes and Php 1 for every minute additional or a fraction thereof. How much will a caller have to pay if his call lasts for 8 minutes? Write a rule that best describes the problem and draw its graph using any method.

Let:	$f(x)$	=	call charge
	x	=	time (in minutes) after 3 min.
Given:	b	=	

Example 3

Step 1: Understand and analyze the problem.

A pay phone service charges Php 5 for the first three minutes and Php 1 for every minute additional or a fraction thereof. How much will a caller have to pay if his call lasts for 8 minutes? Write a rule that best describes the problem and draw its graph using any method.

Let:	$f(x)$	=	call charge
	x	=	time (in minutes) after 3 min.
Given:	b	=	Php 5 (initial charge)

Example 3

Step 1: Understand and analyze the problem.

A pay phone service charges Php 5 for the first three minutes and Php 1 for every minute additional or a fraction thereof. How much will a caller have to pay if his call lasts for 8 minutes? Write a rule that best describes the problem and draw its graph using any method.

Let:	$f(x)$	=	call charge
	x	=	time (in minutes) after 3 min.
Given:	b	=	Php 5 (initial charge)
Find:	m	=	

Example 3

Step 2: Use the facts of the problem to form a linear function.

A pay phone service charges Php 5 for the first three minutes and Php 1 for every minute additional or a fraction thereof. How much will a caller have to pay if his call lasts for 8 minutes? Write a rule that best describes the problem and draw its graph using any method.

Let:	$f(x)$	=	call charge
	x	=	time (in minutes) after 3 min.
Given:	b	=	Php 5 (initial charge)
Find:	m	=	Php 1 per minute

Example 3

Step 2: Use the facts of the problem to form a linear function.

A pay phone service charges Php 5 for the first three minutes and Php 1 for every minute additional or a fraction thereof. How much will a caller have to pay if his call lasts for 8 minutes? Write a rule that best describes the problem and draw its graph using any method.

Let:	$f(x)$	=	call charge
	x	=	time (in minutes) after 3 min.
Given:	b	=	Php 5 (initial charge)
Find:	m	=	Php 1 per minute
Function:	$f(x)$	=	$mx + b$

Example 3

Step 2: Use the facts of the problem to form a linear function.

A pay phone service charges Php 5 for the first three minutes and Php 1 for every minute additional or a fraction thereof. How much will a caller have to pay if his call lasts for 8 minutes? Write a rule that best describes the problem and draw its graph using any method.

Let:	$f(x)$	=	call charge
	x	=	time (in minutes) after 3 min.
Given:	b	=	Php 5 (initial charge)
Find:	m	=	Php 1 per minute
Function:	$f(x)$	=	$mx + b$
	$f(x)$	=	

Example 3

Step 2: Use the facts of the problem to form a linear function.

A pay phone service charges Php 5 for the first three minutes and Php 1 for every minute additional or a fraction thereof. How much will a caller have to pay if his call lasts for 8 minutes? Write a rule that best describes the problem and draw its graph using any method.

Let:	$f(x)$	=	call charge
	x	=	time (in minutes) after 3 min.
Given:	b	=	Php 5 (initial charge)
Find:	m	=	Php 1 per minute
Function:	$f(x)$	=	$mx + b$
	$f(x)$	=	$x + 5$

Example 3

Step 3: Graph the linear function.

$$f(x) = x + 5$$

Example 3

Step 3: Graph the linear function.

$$f(x) = x + 5$$

$$m = \frac{\text{rise}}{\text{run}} = 1 = \frac{1}{1},$$

Example 3

Step 3: Graph the linear function.

$$f(x) = x + 5$$


$$m = \frac{\text{rise}}{\text{run}} = 1 = \frac{1}{1}, b = 5$$

Example 3

Step 3: Graph the linear function.

$$f(x) = x + 5$$

$$m = \frac{\text{rise}}{\text{run}} = 1 = \frac{1}{1}, b = 5$$




Example 3

Step 3: Graph the linear function.

$$f(x) = x + 5$$

$$m = \frac{\text{rise}}{\text{run}} = 1 = \frac{1}{1}, b = 5$$




Example 3

Step 3: Graph the linear function.

$$f(x) = x + 5$$

$$m = \frac{\text{rise}}{\text{run}} = 1 = \frac{1}{1}, b = 5$$




Example 3

Step 3: Graph the linear function.

$$f(x) = x + 5$$

$$m = \frac{\text{rise}}{\text{run}} = 1 = \frac{1}{1}, b = 5$$




Example 3

Step 3: Graph the linear function.

$$f(x) = x + 5$$

$$m = \frac{\text{rise}}{\text{run}} = 1 = \frac{1}{1}, b = 5$$




Example 3

Step 3: Graph the linear function.

$$f(x) = x + 5$$

$$m = \frac{\text{rise}}{\text{run}} = 1 = \frac{1}{1}, b = 5$$




Example 3

Step 3: Graph the linear function.

$$f(x) = x + 5$$

$$m = \frac{\text{rise}}{\text{run}} = 1 = \frac{1}{1}, b = 5$$



Example 3

The graph goes through the coordinates $(5, 10)$, therefore a caller has to pay Php 10 if his call lasts for 8 minutes.

Example 4

A motorist drives at a constant rate of 60 kph. If his destination is 240 kilometers away from his starting point, how many hours will it take him to reach his destination? Write a rule that best describes the problem and draw its graph using any method.

Example 4

Step 1: Understand and analyze the problem.

A motorist drives at a constant rate of 60 kph. If his destination is 240 kilometers away from his starting point, how many hours will it take him to reach his destination? Write a rule that best describes the problem and draw its graph using any method.

Example 4

Step 1: Understand and analyze the problem.

A motorist drives at a constant rate of 60 kph. If his destination is 240 kilometers away from his starting point, how many hours will it take him to reach his destination? Write a rule that best describes the problem and draw its graph using any method.

Let: $f(x) =$

Example 4

Step 1: Understand and analyze the problem.

A motorist drives at a constant rate of 60 kph. If his destination is 240 kilometers away from his starting point, how many hours will it take him to reach his destination? Write a rule that best describes the problem and draw its graph using any method.

Let: $f(x)$ = distance travelled (in kilometers)

Example 4

Step 1: Understand and analyze the problem.

A motorist drives at a constant rate of 60 kph. If his destination is 240 kilometers away from his starting point, how many hours will it take him to reach his destination? Write a rule that best describes the problem and draw its graph using any method.

Let: $f(x)$ = distance travelled (in kilometers)
 x =

Example 4

Step 1: Understand and analyze the problem.

A motorist drives at a constant rate of 60 kph. If his destination is 240 kilometers away from his starting point, how many hours will it take him to reach his destination? Write a rule that best describes the problem and draw its graph using any method.

Let:

$f(x)$	=	distance travelled (in kilometers)
x	=	time travelled (in hours)

Example 4

Step 1: Understand and analyze the problem.

A motorist drives at a constant rate of 60 kph. If his destination is 240 kilometers away from his starting point, how many hours will it take him to reach his destination? Write a rule that best describes the problem and draw its graph using any method.

Let:	$f(x)$	=	distance travelled (in kilometers)
	x	=	time travelled (in hours)
Given:	b	=	

Example 4

Step 1: Understand and analyze the problem.

A motorist drives at a constant rate of 60 kph. If his destination is 240 kilometers away from his starting point, how many hours will it take him to reach his destination? Write a rule that best describes the problem and draw its graph using any method.

Let:	$f(x)$	=	distance travelled (in kilometers)
	x	=	time travelled (in hours)
Given:	b	=	0 km (initial distance travelled)

Example 4

Step 1: Understand and analyze the problem.

A motorist drives at a constant rate of 60 kph. If his destination is 240 kilometers away from his starting point, how many hours will it take him to reach his destination? Write a rule that best describes the problem and draw its graph using any method.

Let:	$f(x)$	=	distance travelled (in kilometers)
	x	=	time travelled (in hours)
Given:	b	=	0 km (initial distance travelled)
Find:	m	=	

Example 4

Step 2: Use the facts of the problem to form a linear function.

A motorist drives at a constant rate of 60 kph. If his destination is 240 kilometers away from his starting point, how many hours will it take him to reach his destination? Write a rule that best describes the problem and draw its graph using any method.

Let:	$f(x)$	=	distance travelled (in kilometers)
	x	=	time travelled (in hours)
Given:	b	=	0 km (initial distance travelled)
Find:	m	=	60 kilometers per hour

Example 4

Step 2: Use the facts of the problem to form a linear function.

A motorist drives at a constant rate of 60 kph. If his destination is 240 kilometers away from his starting point, how many hours will it take him to reach his destination? Write a rule that best describes the problem and draw its graph using any method.

Let:	$f(x)$	=	distance travelled (in kilometers)
	x	=	time travelled (in hours)
Given:	b	=	0 km (initial distance travelled)
Find:	m	=	60 kilometers per hour
Function:	$f(x)$	=	$mx + b$

Example 4

Step 2: Use the facts of the problem to form a linear function.

A motorist drives at a constant rate of 60 kph. If his destination is 240 kilometers away from his starting point, how many hours will it take him to reach his destination? Write a rule that best describes the problem and draw its graph using any method.

Let:	$f(x)$	=	distance travelled (in kilometers)
	x	=	time travelled (in hours)
Given:	b	=	0 km (initial distance travelled)
Find:	m	=	60 kilometers per hour
Function:	$f(x)$	=	$mx + b$
	$f(x)$	=	

Example 4

Step 2: Use the facts of the problem to form a linear function.

A motorist drives at a constant rate of 60 kph. If his destination is 240 kilometers away from his starting point, how many hours will it take him to reach his destination? Write a rule that best describes the problem and draw its graph using any method.

Let:	$f(x)$	=	distance travelled (in kilometers)
	x	=	time travelled (in hours)
Given:	b	=	0 km (initial distance travelled)
Find:	m	=	60 kilometers per hour
Function:	$f(x)$	=	$mx + b$
	$f(x)$	=	$60x$

Example 4

Step 3: Graph the linear function.

$$f(x) = 60x$$

Example 4

Step 3: Graph the linear function.

$$f(x) = 60x$$
$$m = \frac{\text{rise}}{\text{run}} = 60 = \frac{60}{1},$$

Example 4

Step 3: Graph the linear function.


$$f(x) = 60x$$
$$m = \frac{\text{rise}}{\text{run}} = 60 = \frac{60}{1}, b = 0$$

Example 4

Step 3: Graph the linear function.

$$f(x) = 60x$$

$$m = \frac{\text{rise}}{\text{run}} = 60 = \frac{60}{1}, b = 0$$




Example 4

Step 3: Graph the linear function.

$$f(x) = 60x$$

$$m = \frac{\text{rise}}{\text{run}} = 60 = \frac{60}{1}, b = 0$$




Example 4

Step 3: Graph the linear function.

$$f(x) = 60x$$

$$m = \frac{\text{rise}}{\text{run}} = 60 = \frac{60}{1}, b = 0$$




Example 4

Step 3: Graph the linear function.

$$f(x) = 60x$$

$$m = \frac{\text{rise}}{\text{run}} = 60 = \frac{60}{1}, b = 0$$




Example 4

Step 3: Graph the linear function.

$$f(x) = 60x$$

$$m = \frac{\text{rise}}{\text{run}} = 60 = \frac{60}{1}, b = 0$$




Example 4

Step 3: Graph the linear function.

$$f(x) = 60x$$

$$m = \frac{\text{rise}}{\text{run}} = 60 = \frac{60}{1}, b = 0$$




Example 4

Step 3: Graph the linear function.

$$f(x) = 60x$$

$$m = \frac{\text{rise}}{\text{run}} = 60 = \frac{60}{1}, b = 0$$



Example 4

The graph goes through the coordinates $(4, 240)$, therefore it will take the motorist 4 hours to reach his destination.

Thank you for watching.