

Estimation of multivariate generalized gamma convolutions through Laguerre expansions

O. Laverny, E. MASIELLO, V. MAUME-DESCHAMPS, D. RULLIÈRE

Université Claude Bernard Lyon 1 & SCOR SE, Université Claude Bernard Lyon 1 & École des Mines de Saint-Étienne

Email : laverny@math.univ-lyon1.fr

Mots Clés : Multivariate Generalized Gamma Convolutions, Estimation, Thorin measure, Laguerre basis, Infinite Divisibility, Analytics combinatorics in several variables.

Biographie – Oskar Laverny is an actuary by formation, fond of statistics and code. Oskar’s PhD is funded through a CIFRE grant between the UCBL and SCOR SE, a reinsurance company. Our research is articulated around the mathematical and statistical study of dependence structures in high-dimensions, and the development of statistical theory and associated computational tools.

Resumé :

The class \mathcal{G}_1 of univariate generalized gamma convolution, defined as the class of weak limits of independent convolutions of gamma distributions, was first introduced by Thorin [7, 8] as a tool to show the infinite divisibility of log-normal and Pareto distributions. Although originating from this very practical question, its study led to many improvements of the theory, well summarized by Bondesson [1] and is still an active field nowadays (see, e.g., [4, 6]). By definition, \mathcal{G}_1 is closed under independent convolutions, but as it appears recently [2], it is also closed by independent products of random variables. Pareto, log-normal, α -stable, Weibull, and many other distributions are in this class, which makes it a nice framework for many applications fields such as climate events modeling, insurance, etc.

An analogue multivariate class \mathcal{G}_d was constructed by Bondesson [2], following an old idea of Cherian [3], by convolving comonotonous multivariate gamma distributions. Finally, we also consider $\mathcal{G}_{d,n}$, the subclass containing convolutions of at most n d -variates gamma. Although not much is known about \mathcal{G}_d , a random vector $\mathbf{X} \in \mathcal{G}_{d,n}$ follows an additive risk-factor structure: there exists gamma random variables $Y_{i,j}$ such that $\forall i \in \{1, \dots, d\}$,

$$X_i = Y_{i,1} + \dots + Y_{i,n}, \quad (1)$$

where each vector $Y_{i,\cdot}$ has independent marginals, and each vector $Y_{\cdot,j}$ has comonotonous marginals. Since on one hand some $Y_{i,j}$ might be identically zero, as $0 \in \mathcal{G}_{1,1}$, and on the other hand every $Y_{i,j}$ is infinite divisible, by increasing n the model can achieve a wide variety of dependence structures and approach any marginal in \mathcal{G}_1 .

A useful tool for the analysis of these models is the Thorin measure. It can be shown that the cumulant generating function of a random vector in \mathcal{G}_d writes:

$$K(\mathbf{t}) = \ln \mathbb{E} \left(e^{\langle \mathbf{t}, \mathbf{X} \rangle} \right) = - \int_{\mathbb{R}_+^d} \ln(1 - \langle \mathbf{t}, \mathbf{s} \rangle) \nu(\partial \mathbf{s}),$$

where the Thorin measure ν is discrete with n atoms if $\mathbf{X} \in \mathcal{G}_{d,n}$. In this case, atoms and weights of ν are respectively scales and shapes of gamma vectors $Y_{\cdot,j}$. Therefore, the model parameters can be fully summarized by ν , which we would like to estimate.

The deconvolution problem of estimating distributions in $\mathcal{G}_{d,n}$, which is equivalent to the problem of retrieving the measure ν , is an inverse problem known to be numerically challenging, even when

$d = 1, n = 2$. Thus, the current literature contains no estimation procedure for distributions in \mathcal{G}_d or $\mathcal{G}_{d,n}$.

In this work, we investigate the univariate and multivariate estimation of atomic Thorin measures. Through a projection in a Laguerre basis, we provide a deconvolution procedure that produces estimators in $\mathcal{G}_{d,n}$, for any finite d, n , be the distribution given through an exact density or an empirical dataset. To make the estimation viable, we designed fast ad-hoc algorithms to compute Laguerre coefficients of multivariate gamma convolutions, leveraging the clever Faa di Bruno trick from Miatto [5].

Through an analytic combinatorics approach, we construct a regularity condition on the random vector that is necessary and sufficient for the (uniform) exponential decay of the Laguerre coefficients, and derive a consistency result for our estimator. Moreover, under the same regularity condition, the approach provides a series expansion for densities in $\mathcal{G}_{d,n}$, where the current literature provides density expansions only when $d = 1$, and which are known to be unstable, or even dramatically failing for certain parameters ranges, noteworthy those which correspond to projection of log-normal, Pareto or Weibull distributions.

We propose some numerical illustrations of the method, and highlight the flexibility of the obtained model.

Références

- [1] Lennart Bondesson. *Generalized Gamma Convolutions and Related Classes of Distributions and Densities*, volume 76 of *Lecture Notes in Statistics*. Springer New York, New York, NY, 1992.
- [2] Lennart Bondesson. A Class of Probability Distributions that is Closed with Respect to Addition as Well as Multiplication of Independent Random Variables. *Journal of Theoretical Probability*, 28(3):1063–1081, September 2015.
- [3] KC Cherian. A bivariate correlated gamma-type distribution function. *Journal of the Indian Mathematical Society*, 5:133–144, 1941.
- [4] Wissem Jedidi and Thomas Simon. Further examples of GGC and HCM densities. *Bernoulli*, 19(5A):1818–1838, November 2013.
- [5] Filippo M Miatto. Recursive multivariate derivatives of $e(x_1, \dots, x_n)$ of arbitrary order. *CoRR*, 2019.
- [6] Tord Sjödin. On Mixtures of Gamma Distributions, Distributions with Hyperbolically Monotone Densities and Generalized Gamma Convolutions (GGC). *arXiv:1806.03926 [math]*, June 2018.
- [7] Olof Thorin. On the infinite divisibility of the Pareto distribution. *Scandinavian Actuarial Journal*, 1977(1):31–40, January 1977.
- [8] Olof Thorin. On the infinite divisibility of the lognormal distribution. *Scandinavian Actuarial Journal*, 1977(3):121–148, March 1977.