

Propositional Logic Syntactic Sugar

$\varphi \Rightarrow \psi := (\neg \varphi \vee \psi) \wedge (\neg \psi \vee \varphi)$ $\varphi \rightarrow \psi := \neg \varphi \vee \psi$
 $\varphi \oplus \psi := (\varphi \wedge \neg \psi) \vee (\psi \wedge \neg \varphi)$ $\varphi \bar{\wedge} \psi := \neg(\varphi \wedge \psi)$
 $(\alpha \Rightarrow \beta | \gamma) := (\neg \alpha \vee \beta) \wedge (\alpha \vee \gamma)$ $\varphi \bar{\vee} \psi := \neg(\varphi \vee \psi)$
Distributivity: $a \wedge (b \vee c) = (a \wedge b) \vee (a \wedge c)$
 $a \vee (b \wedge c) = (a \vee b) \wedge (a \vee c)$
CNF: from truth table, take minterms that are 0.
Each minterm is built as an OR of the negated variables. E.g., $(0, 0, 1) \rightarrow (x \vee y \vee \neg z)$.

Satisfiability, Validity and Equivalence

$SAT(\varphi) := \neg VALID(\neg \varphi)$ $\varphi \Leftrightarrow \psi := VALID(\varphi \leftrightarrow \psi)$
 $VALID(\varphi) := (\varphi \Rightarrow 1)$ $SAT(\varphi) := \neg(\varphi \Leftrightarrow 0)$.

Sequent Calculus:

— **Validity** of ϕ : start with $\{\} \vdash \phi$; ϕ is valid iff $\Gamma \cap \Delta \neq \{\}$ FOR ALL leaves.
— **Satisfiability** of ϕ : start with $\{\phi\} \vdash \{\}$; ϕ is satisfiable iff $\Gamma \cap \Delta = \{\}$ for AT LEAST ONE leaf.
— var is true, if $x \in \Gamma$; false otherwise; "don't care", if variable doesn't appear.

OPER.	LEFT	RIGHT
NOT	$\frac{\neg \phi, \Gamma \vdash \Delta}{\Gamma \vdash \phi, \Delta}$	$\frac{\Gamma \vdash \neg \phi, \Delta}{\phi, \Gamma \vdash \Delta}$
AND	$\frac{\phi \wedge \psi, \Gamma \vdash \Delta}{\phi, \psi, \Gamma \vdash \Delta}$	$\frac{\Gamma \vdash \phi \wedge \psi, \Delta}{\Gamma \vdash \phi, \Delta \quad \Gamma \vdash \psi, \Delta}$
OR	$\frac{\phi \vee \psi, \Gamma \vdash \Delta}{\phi, \Gamma \vdash \Delta \quad \psi, \Gamma \vdash \Delta}$	$\frac{\Gamma \vdash \phi \vee \psi, \Delta}{\Gamma \vdash \phi, \psi, \Delta}$

Resolution Calculus

$\frac{\neg x \vee y \quad \{x\} \cup C_2}{\{y\} \cup C_2}$
To prove unsatisfiability of given clauses in CNF: If we reach $\{\}$, the formula is unsatisfiable. E.g., $\{\{a\}, \{\neg a, b\}, \{\neg b\}\}$, we get: $\{a\} + \{\neg a, b\} \rightarrow \{b\}$; $\{b\} + \{\neg b\} \rightarrow \{\}$ (unsatisfiable).
To prove validity, prove UNSAT of negated formula.
Davis Putnam Procedure - proves SAT; To prove validity: prove unsatisfiability of negated formula.
(1) Compute Linear Clause Form
(*Don't forget to create the last clause $\{x_n\}$*)


(2) Last variable has to be \perp (true) \rightarrow find implied variables.
(3) For remaining variables: assume values and compute newly implied variables.
(4) If contradiction reached: backtrack.

Linear Clause Forms (Computes CNF) -

Bottom up (inside out) in the syntax tree: convert "operators and variables" into new variable. E.g., $\neg a \vee b$ becomes $x_1 \leftrightarrow \neg a$; $x_2 \leftrightarrow x_1 \vee b$. Use rules below to find CNF. Create last clause $\{X_n\}$

$x \leftrightarrow \neg y \Leftrightarrow (\neg x \vee \neg y) \wedge (x \vee y)$
 $x \leftrightarrow y_1 \wedge y_2 \Leftrightarrow (\neg x \vee y_1) \wedge (\neg x \vee y_2) \wedge (x \vee \neg y_1 \vee \neg y_2)$
 $x \leftrightarrow y_1 \vee y_2 \Leftrightarrow (\neg x \vee y_1 \vee y_2) \wedge (x \vee \neg y_1) \wedge (x \vee \neg y_2)$
 $x \leftrightarrow y_1 \leftrightarrow y_2 \Leftrightarrow (x \vee y_1 \vee y_2) \wedge (x \vee \neg y_1 \vee \neg y_2) \wedge (\neg x \vee y_1 \vee \neg y_2) \wedge (\neg x \vee \neg y_1 \vee y_2)$
 $x \leftrightarrow y_1 \oplus y_2 \Leftrightarrow (x \vee \neg y_1 \vee y_2) \wedge (x \vee y_1 \vee \neg y_2) \wedge (\neg x \vee y_1 \vee y_2) \wedge (\neg x \vee \neg y_1 \vee \neg y_2)$

Quotient: Bisimulation of the kripke with itself
Product - $\mathcal{K}_1 \times \mathcal{K}_2 = (\mathcal{V}_1 \cup \mathcal{V}_2, \varphi_I \wedge \psi_I, \varphi_R \wedge \psi_R)$
Quantif. $\exists x.\varphi := [\varphi]_x^1 \vee [\varphi]_x^0$ $\forall x.\varphi := [\varphi]_x^1 \wedge [\varphi]_x^0$
Tarski-Knaster Theorem: $\mu \nu :=$ starts $\perp \rightarrow$ least fixpoint $\spadesuit \nu :=$ starts $\top \rightarrow$ greatest fixpoint



```
Apply(⊙, BddNode a, b)
int m; BddNode h, l;
if isLeaf(a)&isLeaf(b)
  then
    return Eval(⊙, label(a), label(b));
else
  m=max{label(a), label(b)};
  (a0, a1):=Ops(a, m);
  (b0, b1):=Ops(b, m);
  h:=Apply(⊙, a1, b1);
  l:=Apply(⊙, a0, b0);
  return CreateNode(m, h, l)
end;
end
```

```
ITE(BddNode i, j, k)
int m; BddNode h, l;
if i = 0 then return k
elseif i=1 then
  return j
elseif j=k then
  return k
else
  m = max{label(i), label(j), label(k)};
  (i0, i1):=Ops(i, m);
  (j0, j1):=Ops(j, m);
  (k0, k1):=Ops(k, m);
  l:=ITE(i0, j0, k0);
  h:=ITE(i1, j1, k1);
  return CreateNode(m, h, l)
end;
end
```

```
Restrict(Φ, β)
if β=0
  return 0
elseif
  Φ ∈ {0, 1} ∨ (β = 1)
  then
    return Φ
else
  m=max{label(β), label(Φ)};
  (Φ0, Φ1):=Ops(Φ, m);
  (β0, β1):=Ops(β, m)
  if β0=0
    return Restrict(Φ1, β1)
  elseif β1=0
    return Restrict(Φ0, β0)
  elseif m=label(Φ)
    return CreateNode(m, Restrict(Φ1, β1), Restrict(Φ0, β0))
  else
    return Restrict(Φ, Apply(∨, β0, β1))
endif;
endif;
end
```

Ops(v, m)
x:=label(v);
if m=degree(x)
 return (low(v), high(v))
else return(v, v)
end;
end

```
Exists(BddNode e, ϕ)
if isLeaf(ϕ)VisLeaf(e)
  return ϕ;
elseif label(e)>label(ϕ)
  return Exist(high(e), ϕ)
elseif label(e)=label(ϕ)
  h=Exist(high(e), high(ϕ))
  l=Exist(high(e), low(ϕ))
  return Apply(∨, l, h)
else (label(e)<label(ϕ))
  h:=Exists(e, high(ϕ))
  l:=Exists(e, low(ϕ))
  return CreateNode(label(ϕ), h, l)
endif;
end function.
```

ZDD: If positive cofactor = 0, redirect edge to negative cofactor.
If variable not in the formula, add with both edges pointing to same node.

FDD: Positive Davio Decomposition. (Keep both edges to 1 if happens!)

$\varphi = [\varphi]_x^0 \vee x \wedge [\varphi \partial x]$
 $(\partial \varphi / \partial x) := [\varphi]_x^0 \vee [\varphi]_x^1$

Local Model Checking

$\frac{s \vdash \Phi, \varphi \wedge \psi}{\{s\} \vdash \Phi, \varphi} \wedge$	$\frac{s \vdash \Phi, \varphi \vee \psi}{\{s\} \vdash \Phi, \varphi} \vee$
$\frac{s \vdash \Phi, \varphi}{\{s_1 \vdash \Phi, \varphi\} \dots \{s_n \vdash \Phi, \varphi\}} \wedge$	$\frac{s \vdash \Phi, \varphi}{\{s_1 \vdash \Phi, \varphi\} \dots \{s_n \vdash \Phi, \varphi\}} \vee$
$\frac{s \vdash \Phi, \varphi}{\{s'_1 \vdash \Phi, \varphi\} \dots \{s'_n \vdash \Phi, \varphi\}} \wedge$	$\frac{s \vdash \Phi, \varphi}{\{s'_1 \vdash \Phi, \varphi\} \dots \{s'_n \vdash \Phi, \varphi\}} \vee$
$\frac{s \vdash \mu x. \varphi}{s \vdash \Phi, \varphi}$	$\frac{s \vdash \nu x. \varphi}{s \vdash \Phi, \varphi}$
$\frac{s \vdash \Phi, \varphi}{\{s_1 \dots s_n\} = suc_{\exists}^R(s)}$	$\frac{s \vdash \Phi, \varphi}{\{s_1 \dots s_n\} = pre_{\exists}^R(s)}$

If $(s, \mu x. \varphi)$ repeats $\rightarrow 1$ If $(s, \nu x. \varphi)$ repeats $\rightarrow 0$
Predecessor and Successor
 $\diamond := pre_{\exists}^R(Q) := \exists x_1', \dots, x_n'. \varphi_{\mathcal{R}} \wedge [\varphi Q]_{x_1', \dots, x_n'}^{x_1', \dots, x_n'}$
 $\bar{\diamond} := suc_{\exists}^R(Q) := [\exists x_1, \dots, x_n. \varphi_{\mathcal{R}} \wedge \varphi Q]_{x_1', \dots, x_n'}^{x_1, \dots, x_n}$

$\square = pre_{\forall}^R(Q) := \forall x_1', \dots, x_n'. \varphi_{\mathcal{R}} \rightarrow [\varphi Q]_{x_1', \dots, x_n'}^{x_1', \dots, x_n'}$
 $\bar{\square} := suc_{\forall}^R(Q) := [\forall x_1, \dots, x_n. \varphi_{\mathcal{R}} \rightarrow \varphi Q]_{x_1', \dots, x_n'}^{x_1, \dots, x_n}$
Automata types: G \rightarrow Safety; F \rightarrow Liveness;
FG \rightarrow Persistence/Co-Buchi; GF \rightarrow Fairness/Buchi.

Automaton Determinization

NDet_G \rightarrow Det_G: 1.Remove all states/edges that do not satisfy acceptance condition; 2.Use Subset construction (Rabin-Scott); 3.Acceptance condition will be the states where $\{\}$ is never reached.
{NDet_F (partial) or NDet_{prefix}} \rightarrow Det_{FG}: Breakpoint Construction.
NDet_F (total) \rightarrow Det_F: Subset Construction.
NDet_{FG} \rightarrow Det_{FG}: Breakpoint Construction.
NDet_{GF} \rightarrow {Det_{Rabin} or Det_{streett}}: Safra
*** Rabin-Scott Subset Construction** Acceptance condition:set of states containing acceptance states.
*** Breakpoint Construction** 1. Each state is composed by two components **2.** Initial state first component is a set of all initial states, and second component is the empty set. Ex.: $(\mathcal{I}, \{\})$. **3.** a successor for a state (Q, Q_f) is generated as follows:

$\left\{ \begin{array}{ll} \text{If } Q_f = \{\} & (suc_{\exists}^{R_a}(Q), (suc_{\exists}^{R_a}(Q) \cap F)) \\ \text{Otherwise} & (suc_{\exists}^{R_a}(Q), (suc_{\exists}^{R_a}(Q_f) \cap F)) \end{array} \right.$
4. Acceptance states are states where $Q_f \neq \{\}$.
Boolean Operations on ω -Automata
Complement
 $\neg A_{\forall}(Q, \mathcal{I}, \mathcal{R}, F) = A_{\exists}(Q, \mathcal{I}, \mathcal{R}, \neg F)$
 $\neg A_{\exists}(Q, \mathcal{I}, \mathcal{R}, F) = A_{\forall}(Q, \mathcal{I}, \mathcal{R}, \neg F)$

Conjunction

$(A_{\exists}(Q_1, \mathcal{I}_1, \mathcal{R}_1, F_1) \wedge A_{\exists}(Q_2, \mathcal{I}_2, \mathcal{R}_2, F_2)) = A_{\exists}(Q_1 \cup Q_2, \mathcal{I}_1 \wedge \mathcal{I}_2, \mathcal{R}_1 \wedge \mathcal{R}_2, F_1 \wedge F_2)$

Disjunction

$(A_{\exists}(Q_1, \mathcal{I}_1, \mathcal{R}_1, F_1) \vee A_{\exists}(Q_2, \mathcal{I}_2, \mathcal{R}_2, F_2)) = A_{\exists}\left(\begin{array}{l} Q_1 \cup Q_2 \cup \{q\}, \\ (\neg q \wedge \mathcal{I}_1) \vee (q \wedge \mathcal{I}_2), \\ (\neg q \wedge \mathcal{R}_1 \wedge \neg q') \vee (q \wedge \mathcal{R}_2 \wedge q'), \\ (\neg q \wedge F_1) \vee (q \wedge F_2) \end{array}\right)$

If both automata are totally defined,
 $(A_{\exists}(Q_1, \mathcal{I}_1, \mathcal{R}_1, F_1) \vee A_{\exists}(Q_2, \mathcal{I}_2, \mathcal{R}_2, F_2)) = A_{\exists}(Q_1 \cup Q_2, \mathcal{I}_1 \wedge \mathcal{I}_2, \mathcal{R}_1 \wedge \mathcal{R}_2, F_1 \vee F_2)$
Eliminate Nesting - Acceptance condition **must** be an automata of the same type
 $A_{\exists}(Q^1, \mathcal{I}_1^1, \mathcal{R}_1^1, A_{\exists}(Q^2, \mathcal{I}_1^2, \mathcal{R}_1^2, F_1)) = A_{\exists}(Q^1 \cup Q^2, \mathcal{I}_1^1 \wedge \mathcal{I}_1^2, \mathcal{R}_1^1 \wedge \mathcal{R}_1^2, F_1))$

Boolean Operations of G

(1) $\neg G\varphi = F\neg\varphi$ (2) $G\varphi \wedge G\psi = G[\varphi \wedge \psi]$
(3) $G\varphi \vee G\psi = A_{\exists}(\{p, q\}, p \wedge q, [p' \leftrightarrow p \wedge \varphi] \wedge [q' \leftrightarrow q \wedge \psi], G[p \vee q])$

Boolean Operations of F

(1) $\neg F\varphi = G\neg\varphi$ (2) $F\varphi \vee F\psi = F[\varphi \vee \psi]$
(3) $F\varphi \wedge F\psi = A_{\exists}(\{p, q\}, \neg p \wedge \neg q, [p' \leftrightarrow p \vee \varphi] \wedge [q' \leftrightarrow q \vee \psi], F[p \wedge q])$

Boolean Operations of FG

(1) $\neg FG\varphi = GF\neg\varphi$ (2) $FG\varphi \wedge FG\psi = FG[\varphi \wedge \psi]$
(3) $FG\varphi \vee FG\psi = A_{\exists}(\{q\}, \neg q, q' \leftrightarrow (q \Rightarrow \psi | \neg \varphi), FG[\neg q \vee \psi])$

Boolean Operations of GF

(1) $\neg GF\varphi = FG\neg\varphi$ (2) $GF\varphi \vee GF\psi = GF[\varphi \vee \psi]$
(3) $GF\varphi \wedge GF\psi = A_{\exists}(\{q\}, \neg q, q' \leftrightarrow (q \Rightarrow \neg \psi | \varphi), GF[q \wedge \psi])$

Transformation of Acceptance Conditions
Reduction of G
 $G\varphi = A_{\exists}(\{q\}, q, \varphi \wedge q \wedge q', Fq)$
 $G\varphi = A_{\exists}(\{q\}, q, q' \leftrightarrow q \wedge \varphi, FGq)$
 $G\varphi = A_{\exists}(\{q\}, q, q' \leftrightarrow q \wedge \varphi, GFq)$

Reduction of F

$F\varphi$ can **not** be expressed by $NDet_G$
 $F\varphi = A_{\exists}(\{q\}, \neg q, q' \leftrightarrow q \vee \varphi, FGq)$
 $F\varphi = A_{\exists}(\{q\}, \neg q, q' \leftrightarrow q \vee \varphi, GFq)$
Reduction of FG
 $FG\varphi$ can **not** be expressed by $NDet_G$
 $FG\varphi = A_{\exists}(\{q\}, \neg q, q \rightarrow \varphi \wedge q', Fq)$

$$FG\varphi = A_{\exists} \left(\begin{array}{l} \{p, q\}, \quad \neg p \wedge \neg q, \\ \left[(p \rightarrow p') \wedge (p' \rightarrow p \vee \neg q) \wedge \right. \\ \left. (q' \leftrightarrow (p \wedge \neg q \vee \neg \varphi) \vee (p \wedge q)) \right] \end{array} \right),$$
$$FG\varphi = A_{\exists} \left(\begin{array}{l} \{p, q\}, \quad \neg p \wedge \neg q, \\ \left[(p \rightarrow p') \wedge (p' \rightarrow p \vee \neg q) \wedge \right. \\ \left. (q' \leftrightarrow (p \wedge \neg q \vee \neg \varphi) \vee (p \wedge q)) \right] \end{array} \right),$$

Temporal Logics Beware of Finite Paths
E and A quantify over infinite paths.
 $A\varphi$ holds on every state that has no infinite path;
 $E\varphi$ is false on every state that has no infinite path;
 $A0$ holds on states with only finite paths;
 $E1$ is false on state with only finite paths;
 $\square 0$ holds on states with no successor states;
 $\diamond 1$ holds on states with successor states.
 $F\varphi = \varphi \vee XF\varphi$ $G\varphi = \varphi \wedge XG\varphi$
 $[\varphi U \psi] = \psi \vee (\varphi \wedge X[\varphi U \psi])$
 $[\varphi B \psi] = \neg \psi \wedge (\varphi \vee X[\varphi B \psi])$
 $[\varphi W \psi] = (\psi \wedge \varphi) \vee (\neg \psi \wedge X[\varphi W \psi])$

Negation Normal Form

$\neg(\varphi \wedge \psi) = \neg\varphi \vee \neg\psi$ $\neg(\varphi \vee \psi) = \neg\varphi \wedge \neg\psi$
 $\neg\neg\varphi = \varphi$ $\neg X\varphi = X\neg\varphi$
 $\neg G\varphi = F\neg\varphi$ $\neg F\varphi = G\neg\varphi$
 $\neg[\varphi U \psi] = [(\neg\varphi) \bar{B} \psi]$ $\neg[\varphi \bar{U} \psi] = [(\neg\varphi) B \psi]$
 $\neg[\varphi B \psi] = [(\neg\varphi) \bar{U} \psi]$ $\neg[\varphi \bar{B} \psi] = [(\neg\varphi) U \psi]$
 $\neg A\varphi = E\neg\varphi$ $\neg E\varphi = A\neg\varphi$
 $\neg \bar{X}\varphi = \bar{X}\neg\varphi$ $\neg \bar{X}\varphi = \bar{X}\neg\varphi$
 $\neg \bar{G}\varphi = \bar{F}\neg\varphi$ $\neg \bar{F}\varphi = \bar{G}\neg\varphi$
 $\neg[\varphi \bar{U} \psi] = [(\neg\varphi) \bar{B} \psi]$ $\neg[\varphi \bar{U} \psi] = [(\neg\varphi) \bar{B} \psi]$
 $\neg[\varphi \bar{B} \psi] = [(\neg\varphi) \bar{U} \psi]$ $\neg[\varphi \bar{B} \psi] = [(\neg\varphi) \bar{U} \psi]$

LTL Syntactic Sugar: analog for past operators
 $G\varphi = \neg[1 \bar{U} (\neg\varphi)]$ $F\varphi = [1 \bar{U} \varphi]$
 $[\varphi W \psi] = \neg[(\neg\varphi \vee \neg\psi) \bar{U} (\neg\varphi \wedge \psi)]$ $(\neg\psi \text{ holds until } \varphi \wedge \psi)$
 $[\varphi W \psi] = [(\neg\psi) \bar{U} (\varphi \wedge \psi)]$
 $[\varphi B \psi] = [(\neg\psi) \bar{U} (\varphi \wedge \neg\psi)]$ $(\psi \text{ can't hold when } \varphi \text{ holds})$
 $[\varphi \bar{B} \psi] = [(\neg\psi) \bar{U} (\neg\varphi \wedge \neg\psi)]$
 $[\varphi U \psi] = [(\neg\psi) \bar{U} (\neg\varphi \wedge \neg\psi)]$
 $[\varphi U \psi] = [\varphi \bar{U} \psi] \vee G\varphi$
 $[\varphi \bar{U} \psi] = \neg[(\neg\psi) U (\neg\varphi \wedge \neg\psi)]$
 $[\varphi \bar{U} \psi] = \neg[(\neg\psi) W (\varphi \rightarrow \psi)]$
 $[\varphi \bar{U} \psi] = \neg[(\neg\varphi) B \psi]$ $(\varphi \text{ doesn't matter when } \psi \text{ holds})$
 $[\varphi \bar{U} \psi] = [\psi B (\neg\varphi \wedge \neg\psi)]$

CTL* Modelchecking to LTL model checking
Let's φ_i be a pure path formula (without path quantifiers), Ψ be a propositional formula, abbreviate subformulas $E\varphi$ and $A\psi$ working bottom-up the syntax tree to obtain the following

normal form: $\phi = \text{let } \begin{bmatrix} x_1 = A\varphi_1 \\ \vdots \\ x_n = A\varphi_n \end{bmatrix} \text{ in } \Psi \text{ end}$

Use LTL model checking to compute $Q_i := \llbracket A\varphi_i \rrbracket_{\mathcal{K}_{i-1}}$, where $\mathcal{K}_0 := \mathcal{K}$ and \mathcal{K}_{i+1} is obtained from \mathcal{K}_i by labelling the states Q_i with x_i . Finally compute $\llbracket \Psi \rrbracket_{\mathcal{K}_n}$

CTL Syntactic Sugar: analog for past operators

Existential Operators

$$EF\varphi = E[1 \ \underline{U} \ \varphi]$$

$$EG\varphi = E[\varphi \ \underline{U} \ 0]$$

$$E[\varphi \ U \ \psi] = E[\varphi \ \underline{U} \ \psi] \vee EG\varphi$$

$$E[\varphi \ B \ \psi] = E[(\neg\psi) \ \underline{U} \ (\varphi \wedge \neg\psi)] \vee EG\neg\psi$$

$$E[\varphi \ B \ \psi] = E[(\neg\psi) \ \underline{U} \ (\varphi \wedge \neg\psi)]$$

$$E[\varphi \ \underline{B} \ \psi] = E[(\neg\psi) \ \underline{U} \ (\varphi \wedge \neg\psi)]$$

$$E[\varphi \ \underline{B} \ \psi] = E[(\neg\psi) \ \underline{U} \ (\varphi \wedge \neg\psi)]$$

$$E[\varphi \ \underline{W} \ \psi] = E[(\neg\psi) \ \underline{U} \ (\varphi \wedge \psi)] \vee EG\neg\psi$$

$$E[\varphi \ W \ \psi] = E[(\neg\psi) \ \underline{U} \ (\varphi \wedge \psi)]$$

$$E[\varphi \ \underline{W} \ \psi] = E[(\neg\psi) \ \underline{U} \ (\varphi \wedge \psi)]$$

Universal Operators

$$AX\varphi = \neg EX\neg\varphi$$

$$AG\varphi = \neg E[1 \ \underline{U} \ \neg\varphi]$$

$$AF\varphi = \neg EG\neg\varphi$$

$$AF\varphi = \neg E[(\neg\varphi) \ U \ 0]$$

$$A[\varphi \ U \ \psi] = \neg E[(\neg\psi) \ \underline{U} \ (\neg\varphi \wedge \neg\psi)]$$

$$A[\varphi \ \underline{U} \ \psi] = \neg E[(\neg\psi) \ \underline{U} \ (\neg\varphi \wedge \neg\psi)] \wedge \neg EG\neg\psi$$

$$A[\varphi \ \underline{U} \ \psi] = \neg E[(\neg\psi) \ \underline{U} \ (\neg\varphi \wedge \neg\psi)]$$

$$A[\varphi \ B \ \psi] = \neg E[(\neg\varphi) \ \underline{U} \ \psi]$$

$$A[\varphi \ \underline{B} \ \psi] = \neg E[(\neg\varphi) \ \underline{U} \ \psi]$$

$$A[\varphi \ \underline{B} \ \psi] = \neg E[(\neg\varphi \vee \psi) \ \underline{U} \ \psi] \wedge \neg EG(\neg\varphi \vee \psi)$$

$$A[\varphi \ W \ \psi] = \neg E[(\neg\psi) \ \underline{U} \ (\neg\varphi \wedge \psi)]$$

$$A[\varphi \ W \ \psi] = \neg E[(\neg\psi) \ \underline{U} \ (\neg\varphi \wedge \psi)] \wedge \neg EG\neg\psi$$

$$A[\varphi \ \underline{W} \ \psi] = \neg E[(\neg\psi) \ \underline{U} \ (\neg\varphi \wedge \psi)]$$

CTL to μ -Calculus($\Phi_{inf} = \nu y. \Diamond y$)

STRONG: op= μ / WEAK: op= ν

$$EX\varphi = \Diamond(\Phi_{inf} \wedge \varphi)$$

$$EG\varphi = \nu x. \varphi \wedge \Diamond x$$

$$EF\varphi = \mu x. \Phi_{inf} \wedge \varphi \vee \Diamond x$$

$$E[\varphi U \psi] = \text{op } x. (\Phi_{inf} \wedge \psi) \vee \varphi \wedge \Diamond x$$

$$E[\varphi B \psi] = \text{op } x. \neg\psi \wedge (\Phi_{inf} \wedge \varphi \vee \Diamond x)$$

$$AX\varphi = \Box(\Phi_{inf} \rightarrow \varphi)$$

$$AG\varphi = \nu x. (\Phi_{inf} \rightarrow \varphi) \wedge \Box x$$

$$AF\varphi = \mu x. \varphi \vee \Box x$$

$$A[\varphi U \psi] = \text{op } x. \psi \vee (\Phi_{inf} \rightarrow \varphi) \wedge \Box x$$

$$A[\varphi B \psi] = \text{op } x. (\Phi_{inf} \rightarrow \neg\psi) \wedge (\varphi \vee \Box x)$$

CTL* to CTL - Existential Operators (weak = str)

$$EX\varphi = EXE\varphi$$

$$EF\varphi = EFE\varphi \quad EFG\varphi \equiv EFEG\varphi$$

$$E[\varphi \ W \ \psi] = E[(E\varphi) \ W \ \psi]$$

$$E[\psi \ U \ \varphi] = E[\psi \ U \ E(\varphi)]$$

$$E[\varphi \ B \ \psi] = E[(E\varphi) \ B \ \psi]$$

obs. $EGF\varphi \neq EGEF\varphi \rightarrow$ can't be converted

CTL* to CTL - Universal Operators (weak = str)

$$AX\varphi = AXA\varphi$$

$$AG\varphi = AGA\varphi$$

$$A[\varphi \ W \ \psi] = A[(A\varphi) \ W \ \psi]$$

$$A[\varphi \ U \ \psi] = A[A(\varphi) \ U \ \psi]$$

$$A[\psi \ B \ \varphi] = A[\psi \ B \ (E(\varphi))]$$

Weak Equivalences

$$*[\varphi U \psi] := [\varphi \underline{U} \ \psi] \vee G\varphi \quad *[\varphi B \psi] := [\varphi \underline{B} \ \psi] \vee G\neg\psi$$

$$*\text{same to past version} \quad [\varphi W \psi] := \neg[(\neg\varphi) \underline{W} \ \psi]$$

$$\overleftarrow{X}\varphi := \neg \overleftarrow{X}\neg\varphi \text{ (at } t0 : \text{ weak true, strong false)}$$

Eliminate boolean op. after path quantify

$$[\varphi_1 \ \underline{U} \ \psi_1] \wedge [\varphi_2 \ \underline{U} \ \psi_2] =$$

$$\left[(\varphi_1 \wedge \varphi_2) \ \underline{U} \ \left(\psi_1 \wedge [\varphi_2 \ \underline{U} \ \psi_2] \vee \right) \right]$$

$$[\varphi_1 \ \underline{U} \ \psi_1] \wedge [\varphi_2 \ U \ \psi_2] =$$

$$\left[(\varphi_1 \wedge \varphi_2) \ \underline{U} \ \left(\psi_1 \wedge [\varphi_2 \ U \ \psi_2] \vee \right) \right]$$

$$[\varphi_1 \ U \ \psi_1] \wedge [\varphi_2 \ U \ \psi_2] =$$

$$\left[(\varphi_1 \wedge \varphi_2) \ \underline{U} \ \left(\psi_1 \wedge [\varphi_2 \ U \ \psi_2] \vee \right) \right]$$

TLT to ω -Automata

$$\phi\langle X\varphi \rangle_x \Leftrightarrow \mathcal{A}_\exists(\{q\}, 1, q \leftrightarrow X\varphi, \phi\langle q \rangle_x)$$

$$\phi\langle X\varphi \rangle_x \Leftrightarrow$$

$$\mathcal{A}_\exists(\{q_0, q_1\}, 1, (q_0 \leftrightarrow \varphi) \wedge (q_1 \leftrightarrow Xq_0), \phi\langle q_1 \rangle_x)$$

$$\phi\langle G\varphi \rangle_x \Leftrightarrow$$

$$\mathcal{A}_\exists(\{q\}, 1, q \leftrightarrow \varphi \wedge Xq, \phi\langle q \rangle_x \wedge GF[\varphi \rightarrow q])$$

$$\phi\langle F\varphi \rangle_x \Leftrightarrow$$

$$\mathcal{A}_\exists(\{q\}, 1, q \leftrightarrow \varphi \wedge Xq, \phi\langle q \rangle_x \wedge GF[q \rightarrow \varphi])$$

$$\phi\langle [\varphi \ U \ \psi] \rangle_x \Leftrightarrow$$

$$\mathcal{A}_\exists(\{q\}, 1, q \leftrightarrow \psi \vee \varphi \wedge Xq, \phi\langle q \rangle_x \wedge GF[\varphi \rightarrow q])$$

$$\phi\langle [\varphi \ \underline{U} \ \psi] \rangle_x \Leftrightarrow$$

$$\mathcal{A}_\exists(\{q\}, 1, q \leftrightarrow \psi \vee \varphi \wedge Xq, \phi\langle q \rangle_x \wedge GF[q \rightarrow \psi])$$

$$\phi\langle [\varphi \ \underline{B} \ \psi] \rangle_x \Leftrightarrow$$

$$\mathcal{A}_\exists(\{q\}, 1, q \leftrightarrow \neg\psi \wedge (\varphi \vee Xq), \phi\langle q \rangle_x \wedge GF[q \vee \psi])$$

$$\phi\langle [\varphi \ \underline{B} \ \psi] \rangle_x \Leftrightarrow$$

$$\mathcal{A}_\exists(\{q\}, 1, q \leftrightarrow \neg\psi \wedge (\varphi \vee Xq), \phi\langle q \rangle_x \wedge GF[q \rightarrow \varphi])$$

$$\phi\langle \overleftarrow{X}\varphi \rangle_x \Leftrightarrow \mathcal{A}_\exists(\{q\}, q, Xq \leftrightarrow \varphi, \phi\langle q \rangle_x)$$

$$\phi\langle \overleftarrow{X}\varphi \rangle_x \Leftrightarrow \mathcal{A}_\exists(\{q\}, \neg q, Xq \leftrightarrow \varphi, \phi\langle q \rangle_x)$$

$$\phi\langle \overleftarrow{G}\varphi \rangle_x \Leftrightarrow \mathcal{A}_\exists(\{q\}, q, Xq \leftrightarrow \varphi \wedge q, \phi\langle \varphi \wedge q \rangle_x)$$

$$\phi\langle \overleftarrow{F}\varphi \rangle_x \Leftrightarrow \mathcal{A}_\exists(\{q\}, \neg q, Xq \leftrightarrow \varphi \vee q, \phi\langle \varphi \vee q \rangle_x)$$

$$\phi\langle [\varphi \ \underline{U} \ \psi] \rangle_x \Leftrightarrow$$

$$\mathcal{A}_\exists(\{q\}, q, Xq \leftrightarrow \psi \vee \varphi \wedge q, \phi\langle \psi \vee \varphi \wedge q \rangle_x)$$

$$\phi\langle [\varphi \ \underline{U} \ \psi] \rangle_x \Leftrightarrow$$

$$\mathcal{A}_\exists(\{q\}, \neg q, Xq \leftrightarrow \psi \vee \varphi \wedge q, \phi\langle \psi \vee \varphi \wedge q \rangle_x)$$

$$\phi\langle [\varphi \ \underline{B} \ \psi] \rangle_x \Leftrightarrow$$

$$\mathcal{A}_\exists(\{q\}, q, Xq \leftrightarrow \neg\psi \wedge (\varphi \vee q), \phi\langle \neg\psi \wedge (\varphi \vee q) \rangle_x)$$

$$\phi\langle [\varphi \ \underline{B} \ \psi] \rangle_x \Leftrightarrow$$

$$\mathcal{A}_\exists(\{q\}, \neg q, Xq \leftrightarrow \neg\psi \wedge (\varphi \vee q), \phi\langle \neg\psi \wedge (\varphi \vee q) \rangle_x)$$

Equivalences and Tips

$$[\varphi \underline{U} \ \psi] \equiv \varphi \text{ don't matter when } \psi \text{ hold}$$

$$[\varphi \underline{B} \ \psi] \equiv \psi \text{ can't hold when } \varphi \text{ hold}$$

$$[\varphi W \ \psi] \equiv \neg\psi \text{ hold until } \varphi \wedge \psi$$

$$F[aUb] \equiv Fb \equiv [FaUFb] \equiv [aUFb]$$

$$F[aBb] \equiv F[a \wedge \neg b]$$

$$AEA \equiv A$$

$$GFX \equiv GXF$$

$$FF\varphi \equiv F\varphi$$

$$GG\varphi \equiv G\varphi$$

$$GF\varphi \equiv XGF\varphi \equiv FGF\varphi \equiv GGF\varphi \equiv GFGF\varphi \equiv$$

$$FGGF\varphi$$

$$FG\varphi \equiv XFG\varphi \equiv FFG\varphi \equiv GFG\varphi \equiv GFFG\varphi \equiv$$

$$FGFG\varphi$$

$$E[(aUb) \wedge (cUd)] \equiv$$

$$E[(a \wedge c) \underline{U} (b \wedge E(cUd) \vee d \wedge E(aUb))]$$

\Rightarrow Rules from F apply to E and rules from G to A.

G and μ -calculus (safety property)

$$-[\nu x. \varphi \wedge \Diamond x]_K$$

-Contains states s where an infinite path π starts

with $\forall t. \pi^{(t)} \in [\varphi]_K$

- φ holds always on π

F and μ -calculus (liveness property)

$$-[\mu x. \varphi \vee \Diamond x]_K$$

-Contains states s where a (possibly finite) path π

starts with $\exists t. \pi^{(t)} \in [\varphi]_K$

- φ holds at least once on π

FG and μ -calculus (persistence property)

$$-[\mu y. [\nu x. \varphi \wedge \Diamond x] \vee \Diamond y]_K$$

-Contains states s where an infinite path π starts

with $\exists t1. \forall t2. \pi^{(t1+t2)} \in [\varphi]_K$

- φ holds after some point on π

GF and μ -calculus (fairness property)

$$-[\nu y. [\mu x. (y \wedge \varphi) \vee \Diamond x]]_K$$

-Contains states s where an infinite path π starts

with

$$\forall t1. \exists t2. \pi^{(t1+t2)} \in [\varphi]_K \text{????}t1 + t2 \text{ort} t1 + t0 \text{????}$$

- φ holds infinitely often on π