

Propositional Logic - Syntactic Sugar

$\varphi \Leftrightarrow \psi := (\neg\varphi \vee \psi) \wedge (\neg\psi \vee \varphi)$ $\varphi \rightarrow \psi := \neg\varphi \vee \psi$

$\varphi \oplus \psi := (\varphi \wedge \neg\psi) \vee (\psi \wedge \neg\varphi)$ $\varphi \bar{\wedge} \psi := \neg(\varphi \wedge \psi)$

$(\alpha \Rightarrow \beta | \gamma) := (\neg\alpha \vee \beta) \wedge (\alpha \vee \gamma)$ $\varphi \bar{\vee} \psi := \neg(\varphi \vee \psi)$

Satisfiability, Validity and Equivalence

$\text{SAT}(\varphi) := \neg \text{VALID}(\neg\varphi)$ $\varphi \Leftrightarrow \psi := \text{VALID}(\varphi \leftrightarrow \psi)$

$\text{VALID}(\varphi) := (\varphi \Leftrightarrow 1)$ $\text{SAT}(\varphi) := \neg(\varphi \Leftrightarrow 0)$.

Conjunctive Normal Form: from truth table, take minterms that are 0. Each minterm is built as an OR of the negated variables. E.g.,

$(0, 0, 1) \rightarrow (x \vee y \vee \neg z)$.

Distributivity: $a \wedge (b \vee c) = (a \wedge b) \vee (a \wedge c)$

$a \vee (b \wedge c) = (a \vee b) \wedge (a \vee c)$

Sequent Calculus:

1. Prove validity of ϕ : start with $\{\} \vdash \phi$; ϕ is valid iff $\Gamma \cap \Delta \neq \{\}$ for all leaves; else, counterexample: var is true, if $x \in \Gamma$; false otherwise; "don't care", if variable doesn't appear.
2. Prove satisfiability of ϕ : start with $\{\phi\} \vdash \{\}$; ϕ is satisfiable iff $\Gamma \cap \Delta = \{\}$ for at least one leaf. Satisfying interpretation: same as counterexample.

OPER.	LEFT	RIGHT
NOT	$\neg\phi, \Gamma \vdash \Delta$ $\Gamma \vdash \phi, \Delta$	$\Gamma \vdash \neg\phi, \Delta$ $\neg\phi, \Gamma \vdash \Delta$
AND	$\phi \wedge \psi, \Gamma \vdash \Delta$ $\phi, \psi, \Gamma \vdash \Delta$	$\Gamma \vdash \phi \wedge \psi, \Delta$ $\Gamma \vdash \phi, \Delta$ $\Gamma \vdash \psi, \Delta$
OR	$\phi \vee \psi, \Gamma \vdash \Delta$ $\phi, \Gamma \vdash \Delta$ $\psi, \Gamma \vdash \Delta$	$\Gamma \vdash \phi \vee \psi, \Delta$ $\Gamma \vdash \phi, \psi, \Delta$

Resolution Calculus

$\frac{\{ \neg x \} \cup C_1 \quad \{ x \} \cup C_2}{C_1 \cup C_2}$
To prove unsatisfiability of given clauses in CNF: If we reach $\{\}$, the formula is unsatisfiable. E.g., $\{\{a\}, \{\neg a, b\}, \{\neg b\}\}$, we get:

$\{a\} + \{\neg a, b\} \rightarrow \{b\}$; $\{b\} + \{\neg b\} \rightarrow \{\}$ (unsatisfiable).

To prove validity, prove UNSAT of negated formula.

Linear Clause Forms (Computes CNF)

Bottom up in the syntax tree: convert "operators and variables" into new variable. E.g., $\neg a \vee b$ becomes $x_1 \leftrightarrow \neg a$; $x_2 \leftrightarrow x_1 \vee b$. Use rules below to find CNF.

$$x \leftrightarrow \neg y \Leftrightarrow (\neg x \vee \neg y) \wedge (x \vee y)$$

$$x \leftrightarrow y_1 \wedge y_2 \Leftrightarrow (\neg x \vee y_1) \wedge (\neg x \vee y_2) \wedge (x \vee \neg y_1 \vee \neg y_2)$$

$$(x \vee \neg y_1 \vee \neg y_2)$$

$$x \leftrightarrow y_1 \vee y_2 \Leftrightarrow (\neg x \vee y_1 \vee y_2) \wedge (x \vee \neg y_1) \wedge (x \vee \neg y_2)$$

$$(x \vee \neg y_1) \wedge (x \vee \neg y_2)$$

$$x \leftrightarrow y_1 \rightarrow y_2 \Leftrightarrow (x \vee y_1) \wedge (x \vee \neg y_1 \vee \neg y_2) \wedge (\neg x \vee \neg y_1 \vee y_2)$$

$$(\neg x \vee \neg y_1 \vee y_2)$$

$$x \leftrightarrow (y_1 \leftrightarrow y_2) \Leftrightarrow (x \vee y_1 \vee y_2) \wedge (x \vee \neg y_1 \vee \neg y_2) \wedge (\neg x \vee y_1 \vee \neg y_2) \wedge (\neg x \vee \neg y_1 \vee y_2)$$

$$(\neg x \vee y_1 \vee \neg y_2) \wedge (\neg x \vee \neg y_1 \vee y_2)$$

$$x \leftrightarrow y_1 \oplus y_2 \Leftrightarrow (x \vee \neg y_1 \vee y_2) \wedge (x \vee y_1 \vee \neg y_2) \wedge (\neg x \vee y_1 \vee y_2) \wedge (\neg x \vee \neg y_1 \vee \neg y_2)$$

$$(\neg x \vee y_1 \vee y_2) \wedge (\neg x \vee \neg y_1 \vee \neg y_2)$$

Davis Putnam Procedure - proves SAT; To prove validity: prove unsatisfiability of negated formula. **(1)** Compute Linear Clause Form

(Don't forget to create the last clause $\{x_n\}$) **(2)** Last

variable has to be \perp (true) \rightarrow find implied variables.


(3) For remaining variables: assume values and

compute newly implied variables. **(4)** If

contradiction reached: backtrack.

<pre> Apply(⊙, BddNode a, b) int m; BddNode h, l; if isLeaf(a)&isLeaf(b) then return Eval(⊙, label(a), label(b)); else m=max(label(a),label(b)) (a0,a1):=Ops(a,m); (b0,b1):=Ops(b,m); h:=Apply(⊙,a1,b1); l:=Apply(⊙,a0,b0); return CreateNode(m,h,l) end; </pre>	<pre> Compose(int x, BddNode ψ, α) int m; BddNode h, l; if x>label(ψ) then return ψ; elseif x=label(ψ) then return ITE(α,high(ψ), low(ψ)); else m=max{label(ψ),label(α)}; (α0,α1):=Ops(α, m); (ψ0,ψ1):=Ops(ψ, m); h:=Compose(x,ψ1,α1); l:=Compose(x,ψ0,α0); return CreateNode(m,h,l) endif; end </pre>
<pre> ITE(BddNode i, j, k) int m; BddNode h, l; if i = 0 then return k elseif i=1 then return j elseif j=k then return k else m = max{label(i), label(j),label(k)} (i0,i1):=Ops(i,m); (j0,j1):=Ops(j,m); (k0,k1):=Ops(k,m); l:=ITE(i0,j0,k0); h:=ITE(i1,j1,k1); return CreateNode(m,h,l) end; end </pre>	<pre> Constrain(Φ, β) if β=0 then ret 0 elseif Φ ∈ {0,1} (β = 1) ret Φ else m=max{label(β),label(Φ)} (Φ0,Φ1):=Ops(Φ,m); (β0,β1):=Ops(β,m); if β0=0 ret Constrain(Φ1,β1) elseif β1=0 then ret Constrain(Φ0,β0) else l:=Constrain(Φ0,β0); h:=Constrain(Φ1,β1); ret CreateNode(m,h,l) endif; endif; end </pre>
<pre> Restrict(Φ, β) if β=0 return 0 elseif Φ ∈ {0,1} ∨ (β = 1) return Φ else m=max{label(β),label(Φ)} (Φ0,Φ1):=Ops(Φ,m); (β0,β1):=Ops(β,m) if β0=0 return Restrict(Φ1,β1) elseif β1=0 return Restrict(Φ0,β0) elseif m=label(Φ) return CreateNode(m, Restrict(Φ1,β1), Restrict(Φ0,β0)) else return Restrict(Φ, Apply(v,β0,β1)) endif; endif; end </pre>	<pre> Ops(v,m) x:=label(v); if m=degree(x) return (low(v),high(v)) else return(v, v) end; end </pre> <p>Other Diagrams: TODD ZDD FDD</p> <p>----</p>

Simulation:



given $K_1 = (\mathcal{I}_1, \mathcal{S}_1, \mathcal{R}_1, \mathcal{L}_1)$ and $K_2 = (\mathcal{I}_2, \mathcal{S}_2, \mathcal{R}_2, \mathcal{L}_2)$; $\sigma \subseteq \mathcal{S}_1 \times \mathcal{S}_2$ is a sim. relation between K_1 and K_2 ($K_1 \preceq K_2$) if: **SIM1-** $(s_1, s_2) \in \sigma$ implies $\mathcal{L}_1(s_1) = \mathcal{L}_2(s_2)$; **SIM2-** for $s_1, s'_1 \in \mathcal{S}_1, s_2 \in \mathcal{S}_2$ with $(s_1, s_2) \in \sigma$ and $(s_1, s'_1) \in \mathcal{R}_1$, there must be $s'_2 \in \mathcal{S}_2$ with $(s'_1, s'_2) \in \sigma$ ($s_2, s'_2 \in \mathcal{R}_2$); **SIM3-** for all $s_1 \in \mathcal{I}_1$, there is a $s_2 \in \mathcal{I}_2$ with $(s_1, s_2) \in \sigma$.

Greatest Simulation Relation

$(s_1, s_2) \in \mathcal{H}_0 \Leftrightarrow \mathcal{L}_1(s_1) = \mathcal{L}_2(s_2)$

$(s_1, s_2) \in \mathcal{H}_{i+1} \Leftrightarrow$

$\left(\begin{array}{l} (s_1, s_2) \in \mathcal{H}_i \wedge \\ \forall s'_1 \in \mathcal{S}_1. \exists s'_2 \in \mathcal{S}_2. \\ (s_1, s'_1) \in \mathcal{R}_1 \rightarrow (s_2, s'_2) \in \mathcal{R}_2 \wedge (s'_1, s'_2) \in \mathcal{H}_i \end{array} \right)$

\mathcal{H}_* is the greatest simulation relation if **SIM3:**

$\mathcal{I}_1 \subseteq \{s_1 \in \mathcal{S}_1 | \exists s_2 \in \mathcal{I}_2. (s_1, s_2) \in \mathcal{H}_*\}$

Bisimulation: $\sigma \subseteq \mathcal{S}_1 \times \mathcal{S}_2$ is a bisim. relation

between K_1 and K_2 ($K_1 \approx K_2$) if: **BISIM1-**

$(s_1, s_2) \in \sigma$ implies $\mathcal{L}_1(s_1) = \mathcal{L}_2(s_2)$; **BISIM2a-**

$(s_1, s'_1 \in \mathcal{S}_1, s_2 \in \mathcal{S}_2, (s_1, s_2) \in \sigma, (s_1, s'_1) \in \mathcal{R}_1$, imply that there is $s'_2 \in \mathcal{S}_2$ with $(s'_1, s'_2) \in \sigma$ and $(s_2, s'_2) \in \mathcal{R}_2$; **BISIM2b-** $s_2, s'_2 \in \mathcal{S}_2, s_1 \in \mathcal{S}_1, (s_1, s_2) \in \sigma, (s_2, s'_2) \in \mathcal{R}_2$, imply that there is $s'_1 \in \mathcal{S}_1$ with $(s'_1, s'_2) \in \sigma$ and $(s_1, s'_1) \in \mathcal{R}_1$; **BISIM3a-** for all $s_1 \in \mathcal{I}_1$, there is a $s_2 \in \mathcal{I}_2$ with $(s_1, s_2) \in \sigma$; **BISIM3b-** for all $s_1 \in \mathcal{I}_2$, there is a $s_2 \in \mathcal{I}_2$ with $(s_1, s_2) \in \sigma$.

Greatest Bisimulation Relation (Equivalence)

$(s_1, s_2) \in \mathcal{B}_0 \Leftrightarrow \mathcal{L}_1(s_1) = \mathcal{L}_2(s_2)$

$(s_1, s_2) \in \mathcal{B}_{i+1} \Leftrightarrow$

$\left(\begin{array}{l} (s_1, s_2) \in \mathcal{B}_i \wedge \\ \forall s'_1 \in \mathcal{S}_1. \exists s'_2 \in \mathcal{S}_2. \\ (s_1, s'_1) \in \mathcal{R}_1 \rightarrow (s_2, s'_2) \in \mathcal{R}_2 \wedge (s'_1, s'_2) \in \mathcal{B}_i \\ \forall s'_2 \in \mathcal{S}_2. \exists s'_1 \in \mathcal{S}_1. \\ (s_2, s'_2) \in \mathcal{R}_2 \rightarrow (s_1, s'_1) \in \mathcal{R}_1 \wedge (s'_1, s'_2) \in \mathcal{B}_i \end{array} \right)$

\mathcal{B}_* is the greatest simulation relation if

$\mathcal{I}_1 \subseteq \{s_1 \in \mathcal{S}_1 | \exists s_2 \in \mathcal{I}_2. (s_1, s_2) \in \mathcal{B}_*\}$

$\mathcal{I}_2 \subseteq \{s_2 \in \mathcal{S}_2 | \exists s_1 \in \mathcal{I}_1. (s_1, s_2) \in \mathcal{B}_*\}$

Quotient: given $\mathcal{K} = (\mathcal{I}, \mathcal{S}, \mathcal{R}, \mathcal{L})$ and the


equivalence relation $\sigma \subseteq \mathcal{S} \times \mathcal{S}$; Quotient structure

$\mathcal{K}_{/\sigma} = (\tilde{\mathcal{I}}, \tilde{\mathcal{S}}, \tilde{\mathcal{R}}, \tilde{\mathcal{L}})$: $\tilde{\mathcal{I}} := \{\{s' \in \mathcal{S} | (s, s') \in \sigma\} | s \in \mathcal{I}\}$

$\tilde{\mathcal{S}} := \{\{s' \in \mathcal{S} | (s, s') \in \sigma\} | s \in \mathcal{S}\}$

$(\tilde{s}_1, \tilde{s}_2) \in \tilde{\mathcal{R}}_1 \Leftrightarrow \exists s'_1 \in \tilde{s}_1. \exists s'_2 \in \tilde{s}_2. (s'_1, s'_2) \in \mathcal{R}$

$\tilde{\mathcal{L}}(\tilde{s}) := \mathcal{L}(s)$



Symbolic Product Computation - given

$\mathcal{K}_1 = (\mathcal{V}_1, \varphi_{\mathcal{I}}, \varphi_{\mathcal{R}})$ and $\mathcal{K}_2 = (\mathcal{V}_2, \psi_{\mathcal{I}}, \psi_{\mathcal{R}})$, the

product is: $\mathcal{K}_1 \times \mathcal{K}_2 = (\mathcal{V}_1 \cup \mathcal{V}_2, \varphi_{\mathcal{I}} \wedge \psi_{\mathcal{I}}, \varphi_{\mathcal{R}} \wedge \psi_{\mathcal{R}})$

Quantif. $\exists x. \varphi := [\varphi]_x^1 \vee [\varphi]_x^0 \forall x. \varphi := [\varphi]_x^1 \wedge [\varphi]_x^0$


Predecessor and Successor

$\diamond := pre_{\mathcal{R}}^{\mathcal{R}}(Q) := \exists x'_1, \dots, x'_n. \varphi_{\mathcal{R}} \wedge [\varphi_Q]_{x'_1, \dots, x'_n}^{x_1, \dots, x_n}$

$\diamondsuit := suc_{\mathcal{R}}^{\mathcal{R}}(Q) := \exists x_1, \dots, x_n. \varphi_{\mathcal{R}} \wedge \varphi_Q]_{x'_1, \dots, x'_n}^{x_1, \dots, x_n}$

$\square := pre_{\mathcal{V}}^{\mathcal{R}}(Q) := \forall x'_1, \dots, x'_n. \varphi_{\mathcal{R}} \rightarrow [\varphi_Q]_{x'_1, \dots, x'_n}^{x_1, \dots, x_n}$

$\square \vdash := suc_{\mathcal{V}}^{\mathcal{R}}(Q) := \forall x_1, \dots, x_n. \varphi_{\mathcal{R}} \rightarrow \varphi_Q]_{x'_1, \dots, x'_n}^{x_1, \dots, x_n}$



$pre_{\mathcal{V}}^{\mathcal{R}}(Q = \{S_1, \dots, S_n\})$ for each node n in \mathcal{K} : if n points to a node that is not in Q n $\notin pre_{\mathcal{V}}^{\mathcal{R}}(Q)$ else n $\in pre_{\mathcal{V}}^{\mathcal{R}}(Q)$	$suc_{\mathcal{V}}^{\mathcal{R}}(Q = \{S_1, \dots, S_n\})$ for each node n in \mathcal{K} : if n is pointed by a node that is not in Q n $\notin suc_{\mathcal{V}}^{\mathcal{R}}(Q)$ else n $\in suc_{\mathcal{V}}^{\mathcal{R}}(Q)$
---	---

Tarski-Knaster Theorem: $\mu :=$ starts $\perp \rightarrow$

least fixpoint $\spadesuit \nu :=$ starts $\top \rightarrow$ greatest fixpoint *

Rabin-Scott Subset Construction 1. Initial

state is a set of states containing all the initial

states. **2.** For all transitions of a set of states,

compute the successors and create a set of states

containing all the possible reachable states when

performing that transition. **3.** Acceptance condition

are set of states containing acceptance states.

Local Model Checking

$\frac{s \models \varphi \wedge \psi}{\{s\} \models \varphi \wedge \psi} \wedge$	$\frac{s \models \varphi \vee \psi}{\{s\} \models \varphi \vee \psi} \vee$
$\frac{s \models \varphi \sqsubseteq \psi}{\{s_1 \vdash \varphi\} \dots \{s_n \vdash \varphi\} \wedge} \wedge$	$\frac{s \models \varphi \supset \psi}{\{s_1 \vdash \varphi\} \dots \{s_n \vdash \varphi\} \vee} \vee$
$\frac{s \models \varphi \supset \psi}{\{s'_1 \vdash \varphi\} \dots \{s'_n \vdash \varphi\} \wedge} \wedge$	$\frac{s \models \varphi \supset \psi}{\{s'_1 \vdash \varphi\} \dots \{s'_n \vdash \varphi\} \vee} \vee$
$\frac{s \models \varphi \mu x. \varphi}{s \models \varphi} \mu$	$\frac{s \models \varphi \nu x. \varphi}{s \models \varphi} \nu$
$\frac{s \models \varphi}{s \models \varphi} \mathcal{D}_{\varphi}(\text{replace w. initial form.})$	

$\{s_1 \dots s_n\} = suc_{\mathcal{R}}^{\mathcal{R}}(s)$ and $\{s'_1 \dots s'_n\} = pre_{\mathcal{R}}^{\mathcal{R}}(s)$

Approximations and Ranks

If $(s, \mu x. \varphi)$ repeats \rightarrow return 1	$apx_0(\mu x. \varphi) := 0$
If $(s, \nu x. \varphi)$ repeats \rightarrow return 0	$apx_0(\nu x. \varphi) := 1$
$apx_{n+1}(\mu x. \varphi) := [\varphi]_x^{apx_n(\mu x. \varphi)}$	
$apx_{n+1}(\nu x. \varphi) := [\varphi]_x^{apx_n(\nu x. \varphi)}$	

Automata types: G \rightarrow Safety; F \rightarrow Liveness;

FG \rightarrow Persistence/Co-Buchi; GF \rightarrow Fairness/Buchi.

Automaton Determinization

NDet_G \rightarrow Det_G: 1. Remove all states/edges that do

not satisfy acceptance condition; 2. Use Subset

construction (Rabin-Scott); 3. Acceptance condition

will be the states where $\{\}$ is never reached.

{NDet_F(partial) or NDet_{prefix}} \rightarrow Det_{FG}:

Breakpoint Construction.

NDet_F(total) \rightarrow Det_F: Subset Construction.

NDet_{FG} \rightarrow Det_{FG}: Breakpoint Construction.

NDet_{GF} \rightarrow {Det_{Rabin} or Det_{Streett}}: Safra

Algorithm.

* **Breakpoint Construction 1.** Each state is

composed by two components **2.** Initial state first

component is a set of all initial states, and second

component is the empty set. Ex.: $(\mathcal{I}, \{\})$. **3.** a

successor for a state (Q, Q_f) is generated as follows:

$\left\{ \begin{array}{l} \text{If } Q_f = \{\} \quad (suc_{\mathcal{R}}^{\mathcal{R}a}(Q), (suc_{\mathcal{R}}^{\mathcal{R}a}(Q) \cap \mathcal{F}) \\ \text{Otherwise} \quad (suc_{\mathcal{R}}^{\mathcal{R}a}(Q), (suc_{\mathcal{R}}^{\mathcal{R}a}(Q_f) \cap \mathcal{F}) \end{array} \right.$

4. Acceptance states are states where $Q_f \neq \{\}$.

Boolean Operations on ω -Automata

Complement

$\neg \mathcal{A}_{\forall}(Q, \mathcal{I}, \mathcal{R}, \mathcal{F}) = \mathcal{A}_{\exists}(Q, \mathcal{I}, \mathcal{R}, \neg \mathcal{F})$

$\neg \mathcal{A}_{\exists}(Q, \mathcal{I}, \mathcal{R}, \mathcal{F}) = \mathcal{A}_{\forall}(Q, \mathcal{I}, \mathcal{R}, \neg \mathcal{F})$

Conjunction

$(\mathcal{A}_{\exists}(Q_1, \mathcal{I}_1, \mathcal{R}_1, \mathcal{F}_1) \wedge \mathcal{A}_{\exists}(Q_2, \mathcal{I}_2, \mathcal{R}_2, \mathcal{F}_2)) =$

$\mathcal{A}_{\exists}(Q_1 \cup Q_2, \mathcal{I}_1 \wedge \mathcal{I}_2, \mathcal{R}_1 \wedge \mathcal{R}_2, \mathcal{F}_1 \wedge \mathcal{F}_2)$

Disjunction

$(\mathcal{A}_{\exists}(Q_1, \mathcal{I}_1, \mathcal{R}_1, \mathcal{F}_1) \vee \mathcal{A}_{\exists}(Q_2, \mathcal{I}_2, \mathcal{R}_2, \mathcal{F}_2)) =$

$\mathcal{A}_{\exists}(Q_1 \cup Q_2, \mathcal{I}_1 \wedge \mathcal{I}_2, \mathcal{R}_1 \wedge \mathcal{R}_2, \mathcal{F}_1 \vee \mathcal{F}_2)$

$\mathcal{A}_{\exists} \left(\begin{array}{l} Q_1 \cup Q_2 \cup \{q\}, \\ (\neg q \wedge \mathcal{I}_1) \vee (q \wedge \mathcal{I}_2), \\$

Boolean Operations of F	
(1) $\neg F\varphi = G\neg\varphi$	(2) $F\varphi \vee F\psi = F[\varphi \vee \psi]$
(3) $F\varphi \wedge F\psi = A\exists(\{p, q\}, \neg p \wedge \neg q, [p' \leftrightarrow p \vee \varphi] \wedge [q' \leftrightarrow q \vee \psi], F[p \wedge q])$	
Boolean Operations of FG	
(1) $\neg FG\varphi = GF\neg\varphi$	(2) $FG\varphi \wedge FG\psi = FG[\varphi \wedge \psi]$
(3) $FG\varphi \vee FG\psi = A\exists(\{q\}, \neg q, q' \leftrightarrow (q \Rightarrow \psi \neg\varphi), FG[\neg q \vee \psi])$	

Boolean Operations of GF	
(1) $\neg GF\varphi = FG\neg\varphi$	(2) $GF\varphi \vee GF\psi = GF[\varphi \vee \psi]$
(3) $GF\varphi \wedge GF\psi = A\exists(\{q\}, \neg q, q' \leftrightarrow (q \Rightarrow \neg\psi \varphi), GF[q \wedge \psi])$	

Transformation of Acceptance Conditions

<u>Reduction of G</u>
$G\varphi = A\exists(\{q\}, q, \varphi \wedge q \wedge q', Fq)$
$G\varphi = A\exists(\{q\}, q, q' \leftrightarrow q \wedge \varphi, FGq)$
$G\varphi = A\exists(\{q\}, q, q' \leftrightarrow q \wedge \varphi, GFq)$

<u>Reduction of F</u>
$F\varphi$ can not be expressed by $NDet_G$
$F\varphi = A\exists(\{q\}, \neg q, q' \leftrightarrow q \vee \varphi, FGq)$
$F\varphi = A\exists(\{q\}, \neg q, q' \leftrightarrow q \vee \varphi, GFq)$
<u>Reduction of FG</u>
$FG\varphi$ can not be expressed by $NDet_G$
$FG\varphi = A\exists(\{q\}, \neg q, q \rightarrow \varphi \wedge q', Fq)$

$FG\varphi = A\exists \left(\begin{array}{c} \{p, q\}, \quad \neg p \wedge \neg q, \\ (p \rightarrow p') \wedge (p' \rightarrow p \vee \neg q) \wedge \\ (q' \leftrightarrow (p \wedge \neg q \vee \neg\varphi) \vee (p \wedge q)) \end{array} \right),$	$G\neg q \wedge Fp$
$FG\varphi = A\exists \left(\begin{array}{c} \{p, q\}, \quad \neg p \wedge \neg q, \\ (p \rightarrow p') \wedge (p' \rightarrow p \vee \neg q) \wedge \\ (q' \leftrightarrow (p \wedge \neg q \vee \neg\varphi) \vee (p \wedge q)) \end{array} \right),$	$GF[p \wedge \neg q]$

Temporal Logics *Beware of Finite Paths*
E and A quantify over infinite paths.
A φ holds on every state that has no infinite path;
E φ is false on every state that has no infinite path;
A0 holds on states with only finite paths;
E1 is false on state with only finite paths;
□0 holds on states with no successor states;
◇1 holds on states with successor states.

$F\varphi = \varphi \vee XF\varphi$	$G\varphi = \varphi \wedge XG\varphi$
$[\varphi U \psi] = \psi \vee (\varphi \wedge X[\varphi U \psi])$	
$[\varphi B \psi] = \neg\psi \wedge (\varphi \vee X[\varphi B \psi])$	
$[\varphi W \psi] = (\psi \wedge \varphi) \vee (\neg\psi \wedge X[\varphi W \psi])$	

Negation Normal Form

$\neg(\varphi \wedge \psi) = \neg\varphi \vee \neg\psi$	$\neg(\varphi \vee \psi) = \neg\varphi \wedge \neg\psi$
$\neg\neg\varphi = \varphi$	$\neg X\varphi = X\neg\varphi$
$\neg G\varphi = F\neg\varphi$	$\neg F\varphi = G\neg\varphi$
$\neg[\varphi U \psi] = [(\neg\varphi) \underline{B} \psi]$	$\neg[\varphi \underline{U} \psi] = [(\neg\varphi) B \psi]$
$\neg[\varphi B \psi] = [(\neg\varphi) \underline{U} \psi]$	$\neg[\varphi U \psi] = [(\neg\varphi) \underline{U} \psi]$
$\neg A\varphi = E\neg\varphi$	$\neg E\varphi = A\neg\varphi$
$\neg \overline{X}\varphi = \overline{X}\neg\varphi$	$\neg \overline{X}\varphi = \overline{X}\neg\varphi$
$\neg \overline{G}\varphi = \overline{F}\neg\varphi$	$\neg \overline{F}\varphi = \overline{G}\neg\varphi$
$\neg[\varphi \overline{U} \psi] = [(\neg\varphi) \overline{\underline{B}} \psi]$	$\neg[\varphi \underline{\overline{U}} \psi] = [(\neg\varphi) \overline{\underline{B}} \psi]$
$\neg[\varphi \overline{B} \psi] = [(\neg\varphi) \overline{\underline{U}} \psi]$	$\neg[\varphi \underline{\overline{B}} \psi] = [(\neg\varphi) \overline{\underline{U}} \psi]$

LTL Syntactic Sugar: analog for past operators
 $G\varphi = \neg[1 \underline{U} (\neg\varphi)]$ $F\varphi = [1 \underline{U} \varphi]$

$[\varphi W \psi] = \neg[(\neg\varphi \vee \neg\psi) \underline{U} (\neg\varphi \wedge \psi)]$	
$[\varphi \underline{W} \psi] = [(\neg\psi) \underline{U} (\varphi \wedge \psi)]$ <small>($\neg\psi$ holds until $\varphi \wedge \psi$)</small>	
$[\varphi B \psi] = \neg[(\neg\varphi) \underline{U} \psi]$	
$[\varphi \underline{B} \psi] = [(\neg\psi) \underline{U} (\varphi \wedge \neg\psi)]$ <small>(ψ can't hold when φ holds)</small>	
$[\underline{U} \psi] = \neg[(\neg\psi) \underline{U} (\neg\varphi \wedge \neg\psi)]$	
$[\varphi U \psi] = [\varphi \underline{U} \psi] \vee G\varphi$	

$[\varphi \underline{U} \psi] = \neg[(\neg\psi) U (\neg\varphi \wedge \neg\psi)]$	
$[\varphi \underline{U} \psi] = \neg[(\neg\psi) W (\varphi \rightarrow \psi)]$	
$[\varphi \underline{U} \psi] = [\psi \underline{W} (\varphi \rightarrow \psi)]$	
$[\varphi \underline{U} \psi] = \neg[(\neg\varphi) B \psi]$ <small>(φ doesn't matter when ψ holds)</small>	
$[\varphi \underline{U} \psi] = [\psi \underline{B} (\neg\varphi \wedge \neg\psi)]$	

CTL Syntactic Sugar: analog for past operators

Existential Operators	
$EF\varphi = E[1 \underline{U} \varphi]$	
$EG\varphi = E[\varphi U 0]$	
$E[\varphi U \psi] = E[\varphi \underline{U} \psi] \vee EG\varphi$	
$E[\varphi B \psi] = E[(\neg\psi) \underline{U} (\varphi \wedge \neg\psi)] \vee EG\neg\psi$	
$E[\varphi B \psi] = E[(\neg\psi) U (\varphi \wedge \neg\psi)]$	
$E[\varphi \underline{B} \psi] = E[(\neg\psi) \underline{U} (\varphi \wedge \neg\psi)]$	
$E[\varphi \underline{B} \psi] = E[(\neg\psi) \underline{U} (\varphi \wedge \neg\psi)]$	
$E[\varphi W \psi] = E[(\neg\psi) \underline{U} (\varphi \wedge \psi)] \vee EG\neg\psi$	
$E[\varphi W \psi] = E[(\neg\psi) U (\varphi \wedge \psi)]$	
$E[\varphi \underline{W} \psi] = E[(\neg\psi) \underline{U} (\varphi \wedge \psi)]$	

Universal Operators	
$AX\varphi = \neg EX\neg\varphi$	
$AG\varphi = \neg E[1 \underline{U} \neg\varphi]$	
$AF\varphi = \neg EG\neg\varphi$	
$AF\varphi = \neg E[(\neg\varphi) U 0]$	
$A[\varphi U \psi] = \neg E[(\neg\psi) \underline{U} (\neg\varphi \wedge \neg\psi)]$	
$A[\varphi \underline{U} \psi] = \neg E[(\neg\psi) \underline{U} (\neg\varphi \wedge \neg\psi)] \wedge \neg EG\neg\psi$	
$A[\varphi \underline{U} \psi] = \neg E[(\neg\psi) U (\neg\varphi \wedge \neg\psi)]$	
$A[\varphi B \psi] = \neg E[(\neg\varphi) \underline{U} \psi]$	
$A[\varphi \underline{B} \psi] = \neg E[(\neg\varphi) U \psi]$	
$A[\varphi B \psi] = \neg E[(\neg\varphi \vee \psi) \underline{U} \psi] \wedge \neg EG(\neg\varphi \vee \psi)$	
$A[\varphi W \psi] = \neg E[(\neg\psi) \underline{U} (\neg\varphi \wedge \psi)]$	
$A[\varphi \underline{W} \psi] = \neg E[(\neg\psi) \underline{U} (\neg\varphi \wedge \psi)] \wedge \neg EG\neg\psi$	
$A[\varphi \underline{W} \psi] = \neg E[(\neg\psi) U (\neg\varphi \wedge \psi)]$	

CTL to μ - Calculus ($\Phi_{inf} = \nu y. \Diamond y$)
$EX\varphi = \Diamond(\Phi_{inf} \wedge \varphi)$
$EG\varphi = \nu x. \varphi \wedge \Diamond x$
$EF\varphi = \mu x. \Phi_{inf} \wedge \varphi \vee \Diamond x$
$E[\varphi \underline{U} \psi] = \mu x. (\Phi_{inf} \wedge \psi) \vee \varphi \wedge \Diamond x$
$E[\varphi U \psi] = \nu x. (\Phi_{inf} \wedge \psi) \vee \varphi \wedge \Diamond x$
$E[\varphi \underline{B} \psi] = \mu x. \neg\psi \wedge (\Phi_{inf} \wedge \varphi \vee \Diamond x)$
$E[\varphi B \psi] = \nu x. \neg\psi \wedge (\Phi_{inf} \wedge \varphi \vee \Diamond x)$
$AX\varphi = \Box(\Phi_{inf} \rightarrow \varphi)$
$AG\varphi = \nu x. (\Phi_{inf} \rightarrow \varphi) \wedge \Box x$
$AF\varphi = \mu x. \varphi \vee \Box x$

$A[\varphi U \psi]$	$= \mu x. \psi \vee (\Phi_{inf} \rightarrow \varphi) \wedge \Box x$
$A[\varphi U \psi]$	$= \nu x. \psi \vee (\Phi_{inf} \rightarrow \varphi) \wedge \Box x$
$A[\varphi B \psi]$	$= \mu x. (\Phi_{inf} \rightarrow \neg \psi) \wedge (\varphi \vee \Box x)$
$A[\varphi B \psi]$	$= \nu x. (\Phi_{inf} \rightarrow \neg \psi) \wedge (\varphi \vee \Box x)$

obs. $EGF\varphi \neq EGEF\varphi \rightarrow$ can't be converted

CTL* to CTL	
Universal Operators	
$AX\varphi = AXA\varphi$	
$AG\varphi = AGA\varphi$	
$A[\varphi W \psi] = A[(A\varphi) W \psi]$	
$A[\varphi \underline{W} \psi] = A[(A\varphi) \underline{W} \psi]$	
$A[\varphi U \psi] = A[A(\varphi) U \psi]$	
$A[\varphi \underline{U} \psi] = A[A(\varphi) \underline{U} \psi]$	

$A[\psi B \varphi] = A[\psi B (E(\varphi))]$	
$A[\psi \underline{B} \varphi] = A[\psi \underline{B} (E(\varphi))]$	
Eliminate boolean op. after path quantify	
$[\varphi_1 \underline{U} \psi_1] \wedge [\varphi_2 \underline{U} \psi_2] =$	

$$\left[(\varphi_1 \wedge \varphi_2) \underline{U} \left(\psi_1 \wedge [\varphi_2 \underline{U} \psi_2] \vee \right) \right]$$

$$[\varphi_1 \underline{U} \psi_1] \wedge [\varphi_2 U \psi_2] =$$

$$\left[(\varphi_1 \wedge \varphi_2) \underline{U} \left(\psi_1 \wedge [\varphi_2 U \psi_2] \vee \right) \right]$$

$$[\varphi_1 U \psi_1] \wedge [\varphi_2 U \psi_2] =$$

$$\left[(\varphi_1 \wedge \varphi_2) \underline{U} \left(\psi_1 \wedge [\varphi_2 U \psi_2] \vee \right) \right]$$

CTL* Modelchecking to LTL model checking

Let's φ_i be a pure path formula (without path quantifiers), Ψ be a propositional formula, abbreviate subformulas $E\varphi$ and $A\psi$ working bottom-up the syntax tree to obtain the following

$$\text{normal form: } \Phi = \text{let } \begin{bmatrix} x_1 = A\varphi_1 \\ \vdots \\ x_n = A\varphi_n \end{bmatrix} \text{ in } \Psi \text{ end}$$

Use LTL model checking to compute $Q_i := \llbracket A\varphi_i \rrbracket_{\mathcal{K}_{i-1}}$, where $\mathcal{K}_0 := \mathcal{K}$ and \mathcal{K}_{i+1} is obtained from \mathcal{K}_i by labelling the states Q_i with x_i .

Finally compute $\llbracket \Psi \rrbracket_{\mathcal{K}_n}$

LTL to ω -automata

$$\Phi(X\varphi)_x \Leftrightarrow A\exists(\{q\}, 1, q \leftrightarrow X\varphi, \Phi(q)_x)$$

$$\Phi(X\varphi)_x \Leftrightarrow A\exists(\{q_0, q_1\}, 1, (q_0 \leftrightarrow \varphi) \wedge (q_1 \leftrightarrow Xq_0), \Phi(q_1)_x)$$

$$\Phi(G\varphi)_x \Leftrightarrow A\exists(\{q\}, 1, q \leftrightarrow \varphi \wedge Xq, \Phi(q)_x \wedge GF[\varphi \rightarrow q])$$

$$\Phi(F\varphi)_x \Leftrightarrow A\exists(\{q\}, 1, q \leftrightarrow \varphi \wedge Xq, \Phi(q)_x \wedge GF[q \rightarrow \varphi])$$

$$\Phi([\varphi U \psi])_x \Leftrightarrow A\exists(\{q\}, 1, q \leftrightarrow \psi \vee \varphi \wedge Xq, \Phi(q)_x \wedge GF[\varphi \rightarrow q])$$

$$\Phi([\varphi \underline{U} \psi])_x \Leftrightarrow A\exists(\{q\}, 1, q \leftrightarrow \psi \vee \varphi \wedge Xq, \Phi(q)_x \wedge GF[q \rightarrow \psi])$$

$$\Phi([\varphi B \psi])_x \Leftrightarrow A\exists(\{q\}, 1, q \leftrightarrow \neg\psi \wedge (\varphi \vee Xq), \Phi(q)_x \wedge GF[q \vee \psi])$$

$$\Phi([\varphi \underline{B} \psi])_x \Leftrightarrow A\exists(\{q\}, 1, q \leftrightarrow \neg\psi \wedge (\varphi \vee Xq), \Phi(q)_x \wedge GF[q \rightarrow \varphi])$$

$$\Phi(\overline{X}\varphi)_x \Leftrightarrow A\exists(\{q\}, q, Xq \leftrightarrow \varphi, \Phi(q)_x)$$

$$\Phi(\overline{X}\varphi)_x \Leftrightarrow A\exists(\{q\}, \neg q, Xq \leftrightarrow \varphi, \Phi(q)_x)$$

$$\Phi(\overline{G}\varphi)_x \Leftrightarrow A\exists(\{q\}, q, Xq \leftrightarrow \varphi \wedge q, \Phi(\varphi \wedge q)_x)$$

$$\Phi(\overline{F}\varphi)_x \Leftrightarrow A\exists(\{q\}, \neg q, Xq \leftrightarrow \varphi \vee q, \Phi(\varphi \vee q)_x)$$

$$\Phi([\varphi \overline{U} \psi])_x \Leftrightarrow A\exists(\{q\}, q, Xq \leftrightarrow \psi \vee \varphi \wedge q, \Phi(\psi \vee \varphi \wedge q)_x)$$

$$\Phi([\varphi \underline{\overline{U}} \psi])_x \Leftrightarrow A\exists(\{q\}, \neg q, Xq \leftrightarrow \neg\psi \wedge (\varphi \vee q), \Phi(\neg\psi \wedge (\varphi \vee q))_x)$$

$$\Phi([\varphi \overline{B} \psi])_x \Leftrightarrow A\exists(\{q\}, q, Xq \leftrightarrow \neg\psi \wedge (\varphi \vee q), \Phi(\neg\psi \wedge (\varphi \vee q))_x)$$

$$\Phi([\varphi \underline{\overline{B}} \psi])_x \Leftrightarrow A\exists(\{q\}, \neg q, Xq \leftrightarrow \neg\psi \wedge (\varphi \vee q), \Phi(\neg\psi \wedge (\varphi \vee q))_x)$$

$$\Phi([\varphi \overline{\underline{U}} \psi])_x \Leftrightarrow A\exists(\{q\}, q, Xq \leftrightarrow \psi \vee \varphi \wedge q, \Phi(\psi \vee \varphi \wedge q)_x)$$

$$\Phi([\varphi \underline{\overline{\underline{U}}} \psi])_x \Leftrightarrow A\exists(\{q\}, \neg q, Xq \leftrightarrow \neg\psi \wedge (\varphi \vee q), \Phi(\neg\psi \wedge (\varphi \vee q))_x)$$

$$\Phi([\varphi \overline{\underline{B}} \psi])_x \Leftrightarrow A\exists(\{q\}, q, Xq \leftrightarrow \neg\psi \wedge (\varphi \vee q), \Phi(\neg\psi \wedge (\varphi \vee q))_x)$$

$$\Phi([\varphi \underline{\overline{\underline{B}}} \psi])_x \Leftrightarrow A\exists(\{q\}, \neg q, Xq \leftrightarrow \neg\psi \wedge (\varphi \vee q), \Phi(\neg\psi \wedge (\varphi \vee q))_x)$$

$$\Phi([\varphi \overline{\underline{\underline{U}}} \psi])_x \Leftrightarrow A\exists(\{q\}, q, Xq \leftrightarrow \psi \vee \varphi \wedge q, \Phi(\psi \vee \varphi \wedge q)_x)$$

$$\Phi([\varphi \underline{\overline{\underline{\underline{U}}}} \psi])_x \Leftrightarrow A\exists(\{q\}, \neg q, Xq \leftrightarrow \neg\psi \wedge (\varphi \vee q), \Phi(\neg\psi \wedge (\varphi \vee q))_x)$$

$$\Phi([\varphi \overline{\underline{\underline{\underline{B}}}} \psi])_x \Leftrightarrow A\exists(\{q\}, q, Xq \leftrightarrow \neg\psi \wedge (\varphi \vee q), \Phi(\neg\psi \wedge (\varphi \vee q))_x)$$

$$\Phi([\varphi \underline{\overline{\underline{\underline{\underline{U}}}} \psi])_x \Leftrightarrow A\exists(\{q\}, \neg q, Xq \leftrightarrow \neg\psi \wedge (\varphi \vee q), \Phi(\neg\psi \wedge (\varphi \vee q))_x)$$

$$\Phi([\varphi \overline{\underline{\underline{\underline{\underline{B}}}} \psi])_x \Leftrightarrow A\exists(\{q\}, q, Xq \leftrightarrow \neg\psi \wedge (\varphi \vee q), \Phi(\neg\psi \wedge (\varphi \vee q))_x)$$

$$\Phi([\varphi \underline{\overline{\underline{\underline{\underline{\underline{U}}}} \psi])_x \Leftrightarrow A\exists(\{q\}, \neg q, Xq \leftrightarrow \neg\psi \wedge (\varphi \vee q), \Phi(\neg\psi \wedge (\varphi \vee q))_x)$$

$$\Phi([\varphi \overline{\underline{\underline{\underline{\underline{\underline{B}}}} \psi])_x \Leftrightarrow A\exists(\{q\}, q, Xq \leftrightarrow \neg\psi \wedge (\varphi \vee q), \Phi(\neg\psi \wedge (\varphi \vee q))_x)$$

$$\Phi([\varphi \underline{\overline{\underline{\underline{\underline{\underline{\underline{U}}}} \psi])_x \Leftrightarrow A\exists(\{q\}, \neg q, Xq \leftrightarrow \neg\psi \wedge (\varphi \vee q), \Phi(\neg\psi \wedge (\varphi \vee q))_x)$$

$$\Phi([\varphi \overline{\underline{\underline{\underline{\underline{\underline{\underline{B}}}} \psi])_x \Leftrightarrow A\exists(\{q\}, q, Xq \leftrightarrow \neg\psi \wedge (\varphi \vee q), \Phi(\neg\psi \wedge (\varphi \vee q))_x)$$

$$\Phi([\varphi \underline{\overline{\underline{\underline{\underline{\underline{\underline{\underline{U}}}} \psi])_x \Leftrightarrow A\exists(\{q\}, \neg q, Xq \leftrightarrow \neg\psi \wedge (\varphi \vee q), \Phi(\neg\psi \wedge (\varphi \vee q))_x)$$

$$\Phi([\varphi \overline{\underline{\underline{\underline{\underline{\underline{\underline{\underline{B}}}} \psi])_x \Leftrightarrow A\exists(\{q\}, q, Xq \leftrightarrow \neg\psi \wedge (\varphi \vee q), \Phi(\neg\psi \wedge (\varphi \vee q))_x)$$

$$\Phi([\varphi \underline{\overline{\underline{\underline{\underline{\underline{\underline{\underline{\underline{U}}}} \psi])_x \Leftrightarrow A\exists(\{q\}, \neg q, Xq \leftrightarrow \neg\psi \wedge (\varphi \vee q), \Phi(\neg\psi \wedge (\varphi \vee q))_x)$$

$\neg\exists t. \varphi \in L_{S1S}$	
$\neg\exists p. \varphi \in L_{S1S}$	
where:	
$\neg\tau \in Term_{\Sigma}^{S1S}$	
$\neg\varphi, \psi \in \zeta_{S1S}$	
$\neg t \in V_{\Sigma}$ with $typ_{\Sigma}(t) = \mathbb{N}$	
$\neg p \in V_{\Sigma}$ with $typ_{\Sigma}(p) = \mathbb{N} \rightarrow \mathbb{B}$	
LO2	
first order terms are defined as:	
$\neg t \in V_{\Sigma} typ_{\Sigma}(t) = \mathbb{N} \subseteq Term_{\Sigma}^{LO2}$	
formulas LO2 are defined as:	
$\neg t_1 < t_2 \in L_{LO2}$	
$\neg p^{(t)} \in L_{LO2}$	
$\neg\varphi, \varphi \wedge \psi \in L_{LO2}$	
$\neg\exists t. \varphi \in L_{LO2}$	
$\neg\exists p. \varphi \in L_{LO2}$	
where:	
$\neg t, t_1, t_2 \tau \in V_{\Sigma}$ with $typ_{\Sigma}(t) = typ_{\Sigma}(t_1) = typ_{\Sigma}(t_2) = \mathbb{N}$	
$\neg\varphi, \psi \in \zeta_{LO2}$	
$\neg t \in V_{\Sigma}$ with $typ_{\Sigma}(t) = \mathbb{N}$	
$\neg p \in V_{\Sigma}$ with $typ_{\Sigma}(p) = \mathbb{N} \rightarrow \mathbb{B}$	
function LO2_S1S(Φ)	
case Φ of	
$t_1 < t_2 : \text{return } \exists p. [\forall t. p^{(t)} \rightarrow p^{(SUC(t))}] \wedge \neg p^{(t_1)} \wedge p^{(t_2)} :$	
$p^{(t)} : \text{return } p^{(t)} ;$	
$\neg\varphi : \text{return } \neg LO2_S1S(\varphi) ;$	
$\varphi \wedge \psi : \text{return } LO2_S1S(\varphi) \wedge LO2_S1S(\psi) ;$	
$\exists t. \varphi : \text{return } \exists t. LO2_S1S(\varphi) ;$	
$\exists p. \varphi : \text{return } \exists p. LO2_S1S(\varphi) ;$	
end	
end	
function S1S_LO2(Φ)	
case Φ of	
$p^{(n)} :$	
return $\exists t_0 \dots t_n. p^{(t_n)} \wedge zero(t_0) \wedge \bigwedge_{i=0}^{n-1} succ(ti, ti+1) ;$	
$p^{(t_0+n)} :$	
return $\exists t_1 \dots t_n. p^{(t_n)} \wedge \bigwedge_{i=0}^{n-1} succ(ti, ti+1) ;$	
$\neg\varphi : \text{return } \neg S1S_LO2(\varphi) ;$	</

$\varphi \vee \psi : \mathbf{return} \text{ } ElimFO(\varphi) \vee ElimFO(\psi);$ $\exists t.\varphi : \mathbf{return} \text{ } \exists qt.Sing(qt) \wedge ElimFO(\varphi);$ $\exists p.\varphi : \mathbf{return} \text{ } \exists p.ElimFO(\varphi);$ end end function Tp2Od(t_0, Φ) <i>temporal to LO1</i> case Φ of $is_var(\Phi) : \Psi^{(t_0)};$ $\neg\varphi : \mathbf{return} \text{ } \neg Tp2Od(\varphi);$ $\varphi \wedge \psi : \mathbf{return} \text{ } Tp2Od(\varphi) \wedge Tp2Od(\psi);$ $\varphi \vee \psi : \mathbf{return} \text{ } Tp2Od(\varphi) \vee Tp2Od(\psi);$ $X\varphi : \Psi := \exists t_1.(t_0 < t_1) \wedge (\forall t_2.t_0 < t_2 \rightarrow t_1 \leq t_2) \wedge Tp2Od(t_1, \varphi);$ $[\varphi \underline{U} \psi] : \Psi := \exists t_1.t_0 \leq t_1 \wedge Tp2Od(t_1, \psi) \wedge interval((t_0, 1, t_1, 0), \varphi);$ $[\varphi B \psi] : \Psi := \forall t_1.t_0 \leq t_1 \wedge interval((t_0, 1, t_1, 0), \neg\varphi) \rightarrow Tp2Od(t_1, \neg\psi);$ $\overline{X}\varphi : \Psi := \forall t_1.(t_1 < t_0) \wedge (\forall t_2.t_2 < t_0 \rightarrow t_2 \leq t_1) \rightarrow Tp2Od(t_1, \varphi);$ $\overline{X}\varphi : \Psi := \exists t_1.(t_1 < t_0) \wedge (\forall t_2.t_2 < t_0 \rightarrow t_2 \leq t_1) \wedge Tp2Od(t_1, \varphi);$ $[\varphi \underline{U} \psi] : \Psi := \exists t_1.t_1 \leq t_0 \wedge Tp2Od(t_1, \psi) \wedge interval((t_1, 0, t_0, 1), \varphi);$	$[\varphi \overleftarrow{B} \psi] : \Psi := \forall t_1.t_1 \leq t_0 \wedge interval((t_1, 0, t_0, 1), \neg\varphi) \rightarrow Tp2Od(t_1, \neg\psi);$ end return Ψ end function interval(l, φ) case Φ of $(t_0, 0, t_1, 0) :$ return $\forall t_2.t_0 < t_2 \wedge t_2 < t_1 \rightarrow Tp2Od(t_2, \varphi);$ $(t_0, 0, t_1, 1) :$ return $\forall t_2.t_0 < t_2 \wedge t_2 \leq t_1 \rightarrow Tp2Od(t_2, \varphi);$ $(t_0, 1, t_1, 0) :$ return $\forall t_2.t_0 \leq t_2 \wedge t_2 < t_1 \rightarrow Tp2Od(t_2, \varphi);$ $(t_0, 1, t_1, 1) :$ return $\forall t_2.t_0 \leq t_2 \wedge t_2 \leq 3t_1 \rightarrow Tp2Od(t_2, \varphi);$ end end Temporal Logic Equivalences and Tips $[\varphi \underline{U} \psi] \equiv \varphi \text{ don't matter when } \psi \text{ hold}$ $[\varphi \underline{B} \psi] \equiv \psi \text{ can't hold when } \varphi \text{ hold}$ $[\varphi \underline{W} \psi] \equiv \neg\psi \text{ hold until } \varphi \wedge \psi$ $[\varphi \underline{U} \psi] \equiv [\varphi \underline{U} \psi] \vee G\varphi$ $[a \underline{U} Fb] \equiv Fb$ $F[a \underline{U} b] \equiv Fb \equiv [Fa \underline{U} Fb]$	$[\varphi B \psi] \equiv [\varphi \underline{B} \psi] \vee G\neg\psi$ $F[a \underline{B} b] \equiv F[a \wedge \neg b]$ $[\varphi \underline{W} \psi] \equiv \neg[\neg\varphi \underline{W} \psi]$ $E(\varphi \wedge \psi) \equiv E\varphi \wedge E\psi (in\ general)$ $AEA \equiv A$ $GF(x \vee y) \equiv GFx \vee GFy$ $FF\varphi \equiv F\varphi$ $GG\varphi \equiv G\varphi$ $GF\varphi \equiv XGF\varphi \equiv FGF\varphi \equiv GGF\varphi \equiv GFGF\varphi \equiv FGGF\varphi$ $FG\varphi \equiv XFG\varphi \equiv FFG\varphi \equiv GFG\varphi \equiv GFFG\varphi \equiv FGGFG\varphi$ G and μ-calculus (safety property) $-\nu x.\varphi \wedge \Diamond x]_K$ -Contains states s where an infinite path π starts with $\forall t.\pi^{(t)} \in [\varphi]_K$ $-\varphi$ holds always on π F and μ-calculus (liveness property) $-\mu x.\varphi \vee \Diamond x]_K$ -Contains states s where a (possibly finite) path π starts with $\exists t.\pi^{(t)} \in [\varphi]_K$ $-\varphi$ holds at least once on π FG and μ-calculus (persistence property) $-\mu y.[\nu x.\varphi \wedge \Diamond x] \vee \Diamond y]_K$	-Contains states s where an infinite path π starts with $\exists t_1.\forall t_2.\pi^{(t_1+t_2)} \in [\varphi]_K$ $-\varphi$ holds after some point on π GF and μ-calculus (fairness property) $-\nu y.[\mu x.(y \wedge \varphi) \vee \Diamond x]]_K$ -Contains states s where an infinite path π starts with $\forall t_1.\exists t_2.\pi^{(t_1+t_2)} \in [\varphi]_K$ $-\varphi$ holds infinitely often on π ω - Automaton to LO2 $A_{\exists}(q_1, ..., qn, \psi I, \psi R, \psi F) (input\ automaton)$ $\exists q_1..qn, \Theta LO2(0, \psi I) \wedge (\forall t.\Theta LO2(t, \psi R)) \wedge (\forall t_1 \exists t_2.t_1 < t_2 \wedge \Theta LO2(t_2, \psi F))$ Where $\Theta LO2(t, \Phi)$ is: $-\Theta LO2(t, p) := p(t) \text{ for variable } p$ $-\Theta LO2(t, X\psi) := \Theta LO2(t+1, \psi)$ $-\Theta LO2(t, \neg\psi) := \neg\Theta LO2(t, \psi)$ $-\Theta LO2(t, \varphi \wedge \psi) := \Theta LO2(t, \varphi) \wedge \Theta LO2(t, \psi)$ $-\Theta LO2(t, \varphi \vee \psi) := \Theta LO2(t, \varphi) \vee \Theta LO2(t, \psi)$ Temporal logic set examples -Pure LTL: AFGa -Pure CTL: AGEFa -LTL + CTL: AFa -CTL*: AFGa \vee AGEFa
--	---	---	---