Christian DiPietrantonio
ME 5130: Advanced Heat and Mass Transfer
Homework 02: Mass Diffusion
October 15, 2026

18B.19, Problem Statement: *Oxygen uptake by a bacterial aggregate.* Under suitable circumstances the rate of oxygen metabolism by bacterial cells is very nearly zero order with respect to oxygen concentration. We examine such a case here and focus our attention on a spherical aggregate of cells, which has a radius R. We wish to determine the total rate of oxygen uptake by the aggregate as a function of aggregate size, oxygen mass concentration ρ_0 at the aggregate surface, the metabolic activity of the cells, and the diffusional behavior of the oxygen. For simplicity we consider the aggregate to be homogeneous. We then approximate the metabolic rate by an effective volumetric reaction rate $r_{O_2} = -k_0'''$ and the diffusional behavior by Fick's law, with an effective pseudobinary diffusivity $\mathcal{D}_{O_2 m}$. Because the solubility of oxygen is very low in this system, both convective oxygen transport and transient effects may be neglected.

(a) Show by means of a shell mass balance that the pseudo-steady-state oxygen concentration profile is described by the differential equation

$$\frac{1}{\xi^2}\frac{d}{d\xi}\left(\xi^2\frac{d\chi}{d\xi}\right) = N \tag{18B.19–1}$$

where $\chi = \rho_{O_2}/\rho_0$, $\xi = r/R$, and $N = k_0''' R^2/\rho_0 \mathcal{D}_{O_2 m}$.

(b) There may be an oxygen-free core in the aggregate, if N is sufficiently large, such that $\chi = 0$ for $\xi < \xi_0$. Write sufficient boundary conditions to integrate Eq. 18B.19–1 for this situation. To do this, it must be recognized that both χ and $d\chi/d\xi$ are zero at $\xi = \xi_0$. What is the physical significance of this last statement?

(c) Perform the integration of Eq. 18B.19–1 and show how ξ_0 may be determined.

(d) Sketch the total oxygen uptake rate and ξ_0 as functions of N, and discuss the possibility that no oxygen-free core exists.

Answer: (c)

$$\chi = 1 - \frac{N}{6}\left(1 - \xi^2\right) + \frac{N}{3}\xi_0^3\left(\frac{1}{\xi} - 1\right) \tag{1}$$

for $\xi \geq \xi_0 \geq 0$, where ξ_0 is determined as a function of N from

$$\xi_0^3 - \frac{3}{2}\xi_0^2 + \left(\frac{1}{2} - \frac{3}{N}\right) = 0 \tag{2}$$

18B.19, Solution:
To solve this problem, we can consider the oxygen going into and out of a thin spherical shell between radius r and $r + \Delta r$. The rate of oxygen going in through the outer surface is:

$$-n_{O_{2,r}}\Big|_{r+\Delta r} \cdot 4\pi(r+\Delta r)^2 \tag{3}$$

The rate of oxygen going out of the inner surface is:

$$-n_{O_{2,r}}\Big|_{r} \cdot 4\pi r^2 \tag{4}$$

and the rate of oxygen being produced within the shell is:

$$-k_0''' \cdot 4\pi r^2 \Delta r \tag{5}$$

Therefore, the shell balance can be constructed as:

$$n_{O_{2,r}}\Big|_{r} \cdot 4\pi r^2 - n_{O_{2,r}}\Big|_{r+\Delta r} \cdot 4\pi(r+\Delta r)^2 - k_0''' \cdot 4\pi r^2 \Delta r = 0 \tag{6}$$

Dividing by $4\pi\Delta r$:

$$\begin{aligned} \frac{\left(r^2 n_{O_{2,r}}\right)\Big|_{r} - \left(r^2 n_{O_{2,r}}\right)\Big|_{r+\Delta r}}{\Delta r} - k_0''' r^2 = 0 \\ \frac{\left(r^2 n_{O_{2,r}}\right)\Big|_{r+\Delta r} - \left(r^2 n_{O_{2,r}}\right)\Big|_{r}}{\Delta r} = -k_0''' r^2 \end{aligned} \tag{7}$$

We can now consider the definition of a derivative:

$$\frac{df}{dr} = \lim_{\Delta r \to 0} \frac{f(r+\Delta r) - f(r)}{\Delta r} \tag{8}$$

Therefore, if we take $\lim_{\Delta r \to 0}$, we have:

$$\frac{d}{dr}\left(r^2 n_{O_{2,r}}\right) = -k_0''' r^2 \tag{9}$$

From Fick's law we have:

$$n_{O_{2,r}} = -\mathcal{D}\frac{d\rho_{O_2}}{dr} \tag{10}$$

and can write:

$$\begin{aligned} \frac{d}{dr}\left(r^2\left(-\mathcal{D}\frac{d\rho_{O_2}}{dr}\right)\right) = -k_0''' r^2 \\ \mathcal{D}\frac{d}{dr}\left(r^2\frac{d\rho_{O_2}}{dr}\right) = k_0''' r^2 \end{aligned} \tag{11}$$

Since $\xi = r/R$:

$$r = R\xi \tag{12}$$

$$dr = Rd\xi \tag{13}$$

Therefore:

$$\frac{d}{dr} = \frac{1}{R}\frac{d}{d\xi} \tag{14}$$

and we can write:

$$\begin{aligned} \mathcal{D}\frac{1}{R}\frac{d}{d\xi}\left(R^2\xi^2\frac{1}{R}\frac{d\rho_{O_2}}{d\xi}\right) &= k_0'''R^2\xi^2 \\ \mathcal{D}\frac{d}{d\xi}\left(\xi^2\frac{d\rho_{O_2}}{d\xi}\right) &= k_0'''R^2\xi^2 \end{aligned} \tag{15}$$

and since $\chi = \rho_{O_2}/\rho_0$:

$$\rho_{O_2} = \chi\rho_0 \tag{16}$$

$$\frac{d\rho_{O_2}}{d\xi} = \rho_0\frac{d\chi}{d\xi} \tag{17}$$

therefore:

$$\begin{aligned} \mathcal{D}\frac{d}{d\xi}\left(\xi^2\rho_0\frac{d\chi}{d\xi}\right) &= k_0'''R^2\xi^2 \\ \mathcal{D}\rho_0\frac{d}{d\xi}\left(\xi^2\frac{d\chi}{d\xi}\right) &= k_0'''R^2\xi^2 \\ \frac{1}{\xi^2}\frac{d}{d\xi}\left(\xi^2\frac{d\chi}{d\xi}\right) &= \frac{k_0'''R^2}{\mathcal{D}\rho_0} \end{aligned} \tag{18}$$

and since:

$$N = \frac{k_0'''R^2}{\rho_0\mathcal{D}_{O_2m}} \tag{19}$$

We have:

$$\boxed{\frac{1}{\xi^2}\frac{d}{d\xi}\left(\xi^2\frac{d\chi}{d\xi}\right) = N} \tag{20}$$

If there is an oxygen-free core in the aggregate, the cells use oxygen faster than diffusion can bring it in, so the oxygen must run out before reaching the center such that $\chi = 0$ for $\xi < \xi_0$. Therefore the sphere has two regions, the outer shell where oxygen is present:

$$\xi_0 < \xi \leq 1 \tag{21}$$

And the core, where no oxygen is present:

$$0 \leq \xi \leq \xi_0 \tag{22}$$

At the outer surface, $\rho_{O2} = \rho_0$, therefore:

$$\boxed{\chi = 1 \quad \text{at} \quad \xi = 1} \tag{23}$$

At the edge of the core:

$$\boxed{\chi = 0 \quad \text{at} \quad \xi = \xi_0} \tag{24}$$

$$\boxed{\frac{d\chi}{d\xi} = 0 \quad \text{at} \quad \xi = \xi_0} \tag{25}$$

The physical significance of this is that when $\chi = 0$ at ξ_0, the oxygen runs out exactly at the edge of the core. Furthermore, when $\frac{d\chi}{d\xi} = 0$ at ξ_0, no oxygen is flowing into the core. We know this because from part a we have:

$$n_{O_{2,r}} = -\mathcal{D}\frac{\rho_0}{R}\frac{d\chi}{d\xi} \tag{26}$$

From this we can see that the mass flux is proportional to the slope of the normalized mass concentration with respect to the normalized radius. Therefore, if this slope is zero at $\xi = \xi_0$, then so is the mass flux. We can now integrate Eq. 18B.19–1:

$$\begin{aligned}
\frac{1}{\xi^2}\frac{d}{d\xi}\left(\xi^2\frac{d\chi}{d\xi}\right) &= N \\
\frac{d}{d\xi}\left(\xi^2\frac{d\chi}{d\xi}\right) &= N\xi^2 \\
\int \frac{d}{d\xi}\left(\xi^2\frac{d\chi}{d\xi}\right)d\xi &= \int N\xi^2 d\xi \\
\xi^2\frac{d\chi}{d\xi} &= \frac{N\xi^3}{3} + C_1 \\
\frac{d\chi}{d\xi} &= \frac{N\xi}{3} + \frac{C_1}{\xi^2}
\end{aligned} \tag{27}$$

Now, we know what at $\xi = \xi 0$, $\frac{d\chi}{d\xi} = 0$:

$$\begin{aligned}
0 &= \frac{N}{3}\xi_0 + \frac{C_1}{\xi_0^2} \\
0 &= \frac{N}{3}\xi_0^3 + C_1 \\
C_1 &= -\frac{N}{3}\xi_0^3
\end{aligned} \tag{28}$$

Therefore:

$$\begin{aligned}
\frac{d\chi}{d\xi} &= \frac{N\xi}{3} + \frac{N}{3}\frac{\xi_0^3}{\xi^2} \\
\frac{d\chi}{d\xi} &= \frac{N}{3}\left(\xi - \frac{\xi_0^3}{\xi^2}\right)
\end{aligned} \tag{29}$$

We can now evaluate:

$$\begin{aligned}\int \frac{d\chi}{d\xi} d\xi &= \frac{N}{3} \int \left(\xi - \frac{\xi_0^3}{\xi^2}\right) d\xi \\ \chi &= \frac{N}{3}\left(frac\xi^2 2 + \frac{\xi_0^3}{\xi}\right) + C_2\end{aligned} \tag{30}$$

We also know that at $\xi = \xi_0$, $\chi = 0$:

$$\begin{aligned}0 &= \frac{N}{3}\left(frac\xi_0^2 2 + \frac{\xi_0^3}{\xi_0}\right) + C_2 \\ 0 &= \frac{N}{3}\left(frac\xi_0^2 2 + \xi_0^2\right) + C_2 \\ 0 &= \frac{N}{2}\xi_0^2 + C_2 \\ C_2 &= -\frac{N}{2}\xi_0^2\end{aligned} \tag{31}$$

Thus:

$$\begin{aligned}\chi &= \frac{N}{3}\left(\frac{\xi^2}{2} + \frac{\xi_0^3}{\xi}\right) - \frac{N}{2}\xi_0^2 \\ &= \frac{N}{3}\left(\frac{\xi^2}{2} + \frac{\xi_0^3}{\xi} - \frac{3}{2}\xi_0^2\right)\end{aligned} \tag{32}$$

and Finally, we can write:

$$\boxed{\chi = \frac{N}{3}\left(\frac{\xi^2}{2} + \frac{\xi_0^3}{\xi} - \frac{3}{2}\xi_0^2\right), \qquad \xi_0 < \xi \leq 1} \tag{33}$$

and:

$$\boxed{\chi = 0, \qquad 0 \leq \xi \leq xi_0} \tag{34}$$

To show how ξ_0 may be determined, we can recall that at $\xi = 1$, $\chi = 1$. Therefore:

$$\begin{aligned}1 &= \frac{N}{3}\left(\frac{1}{2} + \xi_0^3 - \frac{3}{2}\xi_0^2\right) \\ \frac{3}{N} &= \frac{1}{2} + \xi_0^3 - \frac{3}{2}\xi_0^2\end{aligned} \tag{35}$$

and thus:

$$\boxed{\xi_0^3 - \frac{3}{2}\xi_0^2 + \left(\frac{1}{2} - \frac{3}{N}\right) = 0} \tag{36}$$

Now, we know that the total oxygen uptake rate can be expressed as:

$$W_{O_2} = -4\pi R^2 n_{O_2,r}\big|_{r=R} \tag{37}$$

Which can be re-written using Fick's law as:

$$W_{O_2} = 4\pi R^2, \mathcal{D} \frac{d\rho_{O_2}}{dr}\Big|_{r=R} \tag{38}$$

Now, we have already shown that:

$$\frac{d\rho_{O_2}}{dr} = \frac{\rho_0}{R}\frac{d\chi}{d\xi} \tag{39}$$

So we have:

$$W_{O_2} = 4\pi R \mathcal{D} \rho_0 \frac{d\chi}{d\xi}\Big|_{\xi=1} \tag{40}$$

18C.2, Problem Statement: *Diffusion and reaction in a partially impregnated catalyst.* Consider a catalytic sphere like that in §18.7, except that the active ingredient of the catalyst is present only in the annular region between $r = \kappa R$ and $r = R$:

In region I $(0 < r < \kappa R)$,	$k_1'' a = 0$	(18C.2–1)
In region II $(\kappa R < r < R)$,	$k_1'' a = \text{constant} > 0$	(18C.2–2)

Such a situation may arise when the active ingredient is put on the particles after pelleting, as is done for many commercial catalysts.

(a) Integrate Eq. 18.7–6 separately for the active and inactive regions. Then apply the appropriate boundary conditions to evaluate the integration constants, and solve for the concentration profile in each region. Give qualitative sketches to illustrate the forms of the profiles.

(b) Evaluate W_{AR}, the total molar rate of conversion of A in a single particle.

18C.2, Solution: