
Disease-causing Mutations Have a Stronger Predictive Effect Than ApoE-e4 on The Development of Alzheimer's Disease

Cartooned by Hong CJ

Lancet Neurol 2018; 17: 434–44



Early-onset Type

Late-onset Type

	Age of onset	Probability of having a genetic mutation	
Affected family members in three generations	<60 years	86%	
Two or more affected first-degree relatives	<61 years	68%	
Two or more affected first-degree relatives	<65 years	15%	(31 fam)
Two or more affected first-degree relatives	>65 years	<1%	(39 fam)


Table 1: **Probability of finding a pathogenic mutation in one of the recognised Alzheimer's disease genes among multiplex families**

https://www.ncbi.nlm.nih.gov/pubmed/23927914

Loy et al. Lancet 2014


Janssen, et al. Neurology 2003
Zekanowski, et al. Exp Neurol 2003

Exceptionally low likelihood of Alzheimer's dementia in APOE2 homozygotes from a 5,000-person neuropathological study



Nat Commun 2020;11(1):667

https://www.nature.com/articles/s41467-019-14279-8#MOESM1



dementia can be caused by multiple neurodegenerative or cerebrovascular pathologies, particularly in older patients. DLB, dementia with Lewy bodies; FTLD, frontotemporal lobar degeneration.

Nature Reviews Disease Primers volume 7, Article number: 33 (2021)