

Section 2 AI Solutions

Connor Leggett

July 2020

1

- **Question:** Three people walk into an elevator. Each of these people chooses a floor randomly and independently between 2 and 10 (inclusive) and presses the button for that floor. What is the probability that the buttons for 3 consecutive floors are pressed?
- **Answer:**
 - There are seven different combinations of consecutive floors: (2, 3, 4), (3, 4, 5), (4, 5, 6), (5, 6, 7), (6, 7, 8), (7, 8, 9), (8, 9, 10).
 - We don't care about the order in which they were pressed though, (3, 4, 5) is equivalent to (4, 5, 3) for example, and there are 6 different ways each group of three can be re-arranged, so there are actually $7 * 6 = 42$ successful worlds.
 - There are 9 possible numbers that each of the three people could choose, meaning there are $9^3 = 729$ possible worlds.
 - We find the probability by dividing successful worlds by total worlds:

$$p = \frac{\text{success}}{\text{total}} = \frac{42}{729} = \boxed{\frac{14}{243} \approx .0576}$$

2

- **Question:** Men who smoke are 23 times more likely to develop lung cancer than men who don't smoke. The probability of a man smoking is .216. What is the probability that a man smokes given, that he has lung cancer?
- **Answer:**
 - Let S be the event of a man smoking
 - Let L be the event of a man having lung cancer.
 - We are told that:

$$* P(L|S) = 23P(L|S^C)$$

$$* P(S) = .216$$

– By Bayes' Rule, we know that:

$$P(S|L) = \frac{P(L|S)P(S)}{P(L)}$$

Substituting our values:

$$P(S|L) = \frac{P(L|S) \cdot .216}{P(L)}$$

Using the law of total probability:

$$P(S|L) = \frac{P(L|S) \cdot .216}{P(L|S)P(S) + P(L|S^C)P(S^C)}$$

Using the fact that we know $P(L|S) = 23P(L|S^C)$:

$$P(S|L) = \frac{23P(L|S^C) \cdot .216}{23P(L|S^C)P(S) + P(L|S^C)P(S^C)}$$

Removing the common term of $P(L|S^C)$:

$$P(S|L) = \frac{23 \cdot .216}{23 \cdot P(S) + P(S^C)}$$

All probabilities must sum to 1:

$$P(S|L) = \frac{23 \cdot .216}{23 \cdot P(S) + (1 - P(S))}$$

Substituting the probability of smoking again:

$$P(S|L) = \frac{23 \cdot .216}{23 \cdot .216 + (1 - .216)}$$

Calculating:

$$\boxed{P(S|L) \approx .864}$$

3

- **Question:** You are on a game show with three doors. There are goats behind two of the doors, and a car behind one of the doors, and your goal is to pick the car. You initially choose a door, and then the host (who knows which doors have what behind them) opens a different door to reveal a goat, and asks if you would like to switch doors. What is the probability of winning when switching?

• **Answer:**

- Let's decide ahead of time that we'll use the switching strategy.
- Let W be the event that you win the game by switching doors.
- Let C be the event that your initial guess was a car. For readability, we'll also let G be the event that your initial guess was a goat.
- By the law of total probability, we can split this into cases:

$$P(W) = P(W|C)P(C) + P(W|G)P(G)$$

- Now, let's look at some of these individual values:
 - * $P(W|C)$: The probability of winning when your initial guess was a car is 0 because we've decided we're switching from our original guess.
 - * $P(C)$: There are two goats and one car, so the probability that your initial guess was a car is $\frac{1}{3}$.
 - * $P(W|G)$: If your original guess was a goat and the host reveals another goat, then when you switch you must be switching to the car. Therefore, this probability is 1.
 - * $P(G)$: There are two goats and one car, so the probability that your initial guess was a goat is $\frac{2}{3}$.
- Now, putting these values together:

$$P(W) = 0 \cdot \frac{1}{3} + 1 \cdot \frac{2}{3} = \boxed{\frac{2}{3}}$$