

A test of desert shrub facilitation via radiotelemetric monitoring of a diurnal lizard

Journal:	<i>Journal of Applied Ecology</i>
Manuscript ID	Draft
Manuscript Type:	Research Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Westphal, Michael; Bureau of Land Management, Central Coast Field Office Noble, Taylor; York University Butterfield, Scott; The Nature Conservancy Iortie, Christopher; York University,
Key-words:	Gambelia sila, Ephedra californica, San Joaquin Desert, plant-animal interactions, endangered species, thermoregulatory behavior, ectotherm

SCHOLARONE™
Manuscripts

1

2

3

4

5

6

7

8

9

10

11

A test of desert shrub facilitation via radiotelemetric monitoring of a diurnal lizard

Michael F. Westphal, US Bureau of Land Management, Central Coast Field Office, 940 2nd Avenue, Marina California 93933 mwestpha@blm.gov (corresponding author)

Taylor Noble, Department of Biology, York University. 4700 Keele St. Toronto, Ontario, Canada. M3J 1P3

H. Scott Butterfield, The Nature Conservancy, 201 Mission St, San Francisco, CA, 94105, USA

Christopher J. Lortie Department of Biology, York University. 4700 Keele St. Toronto, Ontario, Canada. M3J 1P3

Summary

1. Preservation of desert ecosystems is a worldwide conservation priority. Shrubs can play a key role in the structure of desert communities, and can function as foundation species. Understanding desert shrub ecology is therefore an important task in desert conservation. A useful model for the function of shrubs in deserts is ecological facilitation, which explores benefits that shrubs confer on their community. Facilitation has been well developed in the context of shrub-plant interactions but less well studied for plant-animal interactions.
2. We used radiotelemetry to test the hypothesis that a dominant desert shrub facilitates one species of diurnal lizard. We hypothesized that the blunt-nosed leopard lizard *Gambelia sila* would spend some part of its daily activity cycle associated with California jointfir *Ephedra californica*, and that lizard association with shrubs would increase during the afternoon peak temperature period. We relocated lizards three times daily for 24 days and scored whether lizards were within 0.5 meters of a shrub, which we used as an indicator of shrub association. For each relocation we also scored lizard association with a set of predefined microhabitat features. We also scored lizard behavior according to a set of predefined behavioral traits. We constructed home ranges following the minimum convex polygon method and generated estimates of shrub density and relative shrub area within each home range polygon.
3. We obtained 1190 datapoints from a sample of 27 lizards. We found that lizards were associated with open sites significantly more often than with shrubs but were associated with shrubs more than predicted by percent shrub area within their home ranges. Lizards were associated significantly more often under shrubs during the afternoon peak temperature period, and lizards were observed cooling under shrubs significantly more often. The frequency of association of individual lizards with shrubs was not correlated with the density of shrubs within their home range.

4. *Synthesis and Applications.* Shrubs can be considered as a component of high-quality habitat for ectothermic desert vertebrates for the purposes of restoration and management. Furthermore, radiotelemetry provides a novel methodological approach for assessing shrub-animal facilitative interactions within desert communities.

Keywords: *Gambelia sila*, *Ephedra californica*, San Joaquin Desert, plant-animal interactions, endangered species, thermoregulatory behavior, ectotherm

Introduction

Deserts are highly distinct ecosystems that contribute significantly to global biodiversity and global ecosystem function. The conversion and loss of desert habitat is therefore a global biodiversity crisis requiring immediate intervention, including conservation of remaining undisturbed habitat and restoration of degraded desert (Hannah et al. 1995, Cook et al. 2004, Hoekstra et al. 2005, Kéfi et al. 2007, Mouat et al. 2008, Bachelet et al. 2016, Westphal et al. 2016). Identifying the drivers of ecological health in desert communities will be a crucial component of such interventions. Shrubs can maintain the diversity of desert plant communities (Flores & Jurado 2003) and are predicted to play significant roles in the thermal ecology of desert ectotherms (Sears et al 2016, Basson et al. 2017). Shrubs can also facilitate ectotherm populations in the face of climate change (Adolph 1990, Kearney, Shine and Porter 2009, Sinervo et al 2010, Sears and Angilletta 2015, Sears et al 2016).

Ecological facilitation theory provides a roadmap for describing and predicting the beneficial interactions of shrubs with other organisms within their communities (Bruno et al. 2003, McIntire & Fajardo 2014, Filazzola and Lortie 2014, Bulleri et al 2016, Filazzola et al. 2017). Using facilitation theory, Filazzola et al (2017) extended the exploration of the beneficial interactions between desert shrubs

and vertebrates, and found that one species of shrub provided facilitative benefits to a target species of lizard. We sought to confirm and add depth to their findings using radiotelemetry tracking of the same target species. . Radio telemetry is a well-tested and powerful tool that allows the longitudinal tracking of individual animals throughout their daily behavioral cycles (McGowan et al 2017) and enables the direct observation of habitat interactions and behaviors. We used radio telemetry study to test and refine our understanding of the beneficial interaction of shrubs with lizards. To our knowledge, incorporating radiotelemetry into a facilitation study is a novel use of the method.

We sought to test the hypothesis that shrubs facilitated lizards by providing thermoregulatory opportunity. We predicted that lizards would associate with shrubs for a meaningful proportion of their daily activity cycle; that shrub association would increase in the afternoon when daytime temperatures peak (Filazzola, Sotomayor, and Lortie, 2017); and that lizard association with shrubs would be correlated with thermoregulatory behaviors. The results of our study confirm the application of radiotelemetry to ecological facilitation studies and the application of such studies to the description of beneficial interactions between shrubs and vertebrate ectotherms.

Materials and Methods

Study site.-- The study was conducted on the Elkhorn Plain within Carrizo Plain National Monument (San Luis Obispo County, California, USA, 35.1914° N, 119.7929° W) (Fig. 1) within the San Joaquin Desert ecosystem (Germano et al 2011). Average annual precipitation within the Monument ranges from 15 cm in the southeast to 25 cm in the northwest (Hijmans et al. 2005). The Elkhorn Plain is located within the Monument on an elevated plain separated from the main valley floor of the Carrizo Plain by the San Andreas Fault (Germano et al. 1994). The area has been heavily invaded by non-native annual grasses including *Bromus madritensis*, *Erodium cicutarium*, and *Hordeum murinum* (Schiffman 1994, Stout et al. 2014, Gurney et al. 2015) but still provides habitat for endemic keystone species such as the giant kangaroo rat *Dipodomys ingens* (Bean et al. 2014). California jointfir, *Ephedra californica* was

the dominant shrub at our study site. The blunt-nosed leopard lizard, *Gambelia sila*, was well documented on the study (Germano, Smith and Tabor, 2007).

Study species .— *E. californica*, a basal gymnosperm in the Gnetophyta division, is a large, slow-growing woody shrub restricted to arid environments in western North America (Sawyer, Keeler-Wolf & Evens 2009). Although the genus has a worldwide distribution and is represented by over a dozen species in the desert southwest of North America, *E. californica* is the only species that occurs in the San Joaquin Valley, where it is locally considered rare and sensitive (Sawyer, Keeler-Wolf & Evens 2009) and has been documented to be a foundation species in the San Joaquin desert community (Hawbecker 1951, Lortie et al. 2017, Lortie, Filazzola & Westphal 2017). *Gambelia sila* is a state and federally listed endangered species endemic to the San Joaquin Valley and restricted to San Joaquin Desert habitat (Germano & Williams 1992, U.S. Fish and Wildlife Service 1998, Warrick, Kato, & Rose 1998, Germano et al. 2011, Germano & Rathbun 2016). *Gambelia sila* are diurnal and mainly insectivorous though they may eat smaller lizard species on occasion (Warrick et al. 1998, Germano, Smith & Tabor, 2007). Though *G. sila* can bury themselves and will occasionally dig primitive burrows, they mostly utilize abandoned burrows of other animals such as *D. ingens* (Fields, Coffin & Gosz 1999, Prugh et al. 2012). Adult *G. sila* are inactive in burrows for much of the year, emerging only from late March or April through July (U.S. Fish and Wildlife Service 1998, Warrick, Kato, & Rose 1998, Germano et al. 2016). During the active season, *G. sila* will also spend the night underground in burrows and may return to a burrow during the day if the temperature becomes too hot or cold (Warrick, Kato, and Rose 1998, Germano and Rathbun 2016).

Experimental design.-- *Gambelia sila* individuals were located during foot and vehicle surveys and captured using a pole and noose. Individuals were collared following the method of Germano & Rathbun (2016). VHF radio transmitters (Holohil model BD-2, frequency 151-152 MHz, battery life 8-16 weeks, Holohil Systems Ltd., Carp, ON, Canada) were attached to a small beaded chain collar using jewelry wire and epoxy, and the collars were then fastened around the lizard's neck. *Gambelia sila* were

kept overnight to ensure the collar was fitted correctly and did not irritate or harm the animal, and then were then released at their capture site. Collars weighed 1.6-2.2 grams (depending on the size of chain needed for the lizard's neck), and we ensured that the weight of the collar did not exceed between 5% and 10% of the body mass of the individual.

In the first two days following release, all captured *G. sila* individuals were relocated (i.e. repeatedly sighted using radio telemetry) several times between to ensure that the lizards were successfully adjusting to the collars and that impacts to their behavior and survival were minimal. We looked for any negative effects the collar had on the lizards, such as impacts on movement or any other deviation from normal lizard behaviors. *Gambelia sila* were then formally surveyed for 24 consecutive days. Surveys were conducted on each lizard 3 times a day. Two of these daily surveys were conducted during daylight hours, when lizards were typically active above ground. One survey was conducted before noon and one was conducted after noon. The third survey was conducted during the night when lizards are inactive below ground. The 'night survey' was conducted before 7:30 AM or after 7:30 PM on each day.

Gambelia sila were relocated using a 3-element Yagi antenna and Model R-100 telemetry receiver (Communications Specialists, Inc., Orange, CA, USA). Once found, a location was taken for each lizard using a Garmin 64st GPS unit (Garmin Ltd., Olathe, KS, USA) and a laser range-finder (Bushnell Outdoor Products, Overland Park, KS, USA). Habitat was categorized as whether a lizard was within 0.5 meters of a shrub (shrub) or not (open) (henceforth, the "shrub association zone"), and behavior was scored from a suite of predetermined behavioral syndromes (Table S1). Disturbance from the observer to the lizard was kept to a minimum for each observation to avoid influencing behavior and habitat selection. At the completion of the study all collars were removed from the lizards.

Analyses.-- Analyses were conducted in R (version 3.3.2). Habitat association was analyzed using a generalized linear model (Bolker et al. 2009) with the multcomp package (Hothorn, Bretz & Westfall

2008). Behavioral data were analyzed with a multinomial logistic regression using the nnet package that accounts for the multiple levels of nominal outcomes of the observations (Venables and Ripley 2002). Home range size was calculated using a 95% Minimum Convex Polygon (MCP) estimation (Mohr 1947) using the adehabitatHR package (Calenge 2006). MCPs were visualized in two dimensions in R.

Shrub density was calculated by visually counting individual shrubs within each lizard's MCP using aerial photographs (Google Earth, image taken December 20, 2016, accessed November 2017) and dividing that number by the area in square meters of the MCP. We calculated a standardized measure of shrub association zone area using on-the-ground measurements of a large number of randomly chosen shrubs in the study area, from which we calculated an average radius for each shrub following the method of Filazzola et al. (2017) and to which we added the 0.5m association criterion described above. We calculated the area of each shrub association zone using the formula πr^2 and then took the average across the sample. We multiplied this standardized shrub association zone area by the number of shrubs counted in each MCP to obtain an estimate of the percent area of an MCP subsumed by shrub association zones.

R code used for this project can be found at <https://cjlortie.github.io/Carrizo.telemetry>.

Results

A total of 27 lizards were relocated more than 5 instances cumulatively either in the AM or the PM across the sampling period. On a given day, the median total number of relocations was 22 with a maximum of 27 and a minimum of 1 relocation. There were a total of 1190 relocations and MCPs generally did not overlap (Fig 1). Mean female MCP area was 1.87 ha +/- 0.53 se. Mean male MCP area was 5.14 ha +/- 2.15 se. The difference in MCP area between males and females was not significant ($P < \text{Chi } 0.095920$). Gender was initially included as a factor in all other analyses but no relevant effects were significant (not reported), therefore gender was subsequently removed from the remaining analyses.

Habitat:-- The frequency of lizard observation differed significantly between habitat types (Fig. 2, Table 1, $p < 0.01$). Lizards were observed in the open on an average of 18.8 days and in shrubs an average of 10.5 days. Shrub association frequencies of individual lizards ranged from 0 to 0.63 with a mean of 0.23 ± 0.035 se (Table S2). Observations of lizards within open habitat did not differ between different times of day, but observations of lizards associating with shrubs differed significantly between morning and afternoon with lizards being found more frequently at shrubs in the afternoon (Table 1, $p = 0.0252$).

Behavior:—Behavior differed significantly between habitat types (Fig. 2, Table 2, $p < 0.0001$). Lizards were observed cooling under shrubs significantly more than other habitat types (Fig. 2, Table 2, $p < 0.0001$). Lizards were also observed avoiding predators under shrubs more frequently than at other microhabitat types (Table 3, $p < 0.0001$). The predators that lizards were observed avoiding in this study were all aerial predators (either ravens or raptor species). Burrowing and interacting occurred significantly less often under shrubs ($p < 0.0001$). Other types of behavior such as sunning, hunting, or active observation did not differ significantly between habitat types. Observed behavior also differed significantly between different times of day e.g. lizards were more frequently observed sunning in the morning in both habitat types compared to the afternoon and more often burrowing and avoiding predators in the afternoon (Fig. 2, Table 2, $p < 0.001$).

Shrub use as a function of shrub density and area.-- Shrub use by individual lizards did not vary significantly as a function of shrub density within that lizard's home range (Fig 3.) Percent of MCP areas subsumed by shrub association zones ranged from 1% to 15% with an average of 5% of total surface area, and frequency of shrub use by lizards was significantly higher than predicted by the percent of MCP area subsumed by shrubs ($Z = -4.714$ from a Wilcoxon Signed ranks test, $p < 0.001$).

Discussion

Shrubs are foundation species in many ecosystems due to the facilitative benefits that they provide to both plant and animal species (Filazzola & Lortie 2014, Lortie, Filazzola & Sotomayor 2015). We hypothesized that *E. californica* facilitates *G. sila* by providing thermoregulatory benefits. Our finding that *G. sila* was associated with *E. californica* an average of 10 out of 24 days, and that individual lizards were associated with shrubs more than predicted by shrub area within their home ranges, support our hypothesis. Our hypothesis is also supported by the observed significant shift towards shrub association in the afternoon during peak daytime temperatures. Our observation that shrub use was not correlated with shrub density suggests that lizards are actively choosing shrubs over open habitats rather than as a consequence of shrubs being more densely distributed in their home ranges. The observed association of *G. sila* with shrubs is consistent with results of studies of thermoregulatory behavior of lizards (Sears et al. 2016, Vickers et al 2016, Basson et al. 2017) and suggest that shrubs facilitate *G. sila* by providing shade. Shrubs can buffer the extremes of multiple environmental conditions such as temperature, wind, and solar radiation, creating a moderate microclimate under their canopy (Kerr et al. 2004, Pugnaire 2010). At the landscape scale, the presence of shrubs and their pattern of distribution (i.e. clumped vs. dispersed) will affect lizard thermoregulatory behavior and can be crucial to an ectotherm's thermoregulatory efficiency (Sears et al 2016, Basson et al 2017). Sources of shade are particularly important for ectotherms, which must maintain body temperature through behavior (Huey 1974, Huey and Slatkin 1976, Díaz and Cabezas-Díaz 2004, Kerr et al. 2004). Visual concealment from predators and physical protection is also important (Fields et al. 1999, Anderson et al. 2010, Filazzola et al. 2017). Shrubs may therefore provide important mechanisms of facilitation for *G. sila*. Our results suggest an important mechanism (shrub restoration) for the management of desert ectotherms such as *G. sila*, and provide support for radiotelemetry as a viable method for studying ecological facilitation.

Shrub use by *G. sila* was addressed in one previous paper that also used radiotelemetry. Germano & Rathbun (2016) employed *post hoc* tests to answer the question of whether shrubs are important components of *G. sila* habitat. One test depended on an assumption based on Schoepf et al 2015 that

home ranges were resource-based (ie shrub-limited) and would thus be smaller in the presence of high quality habitat (= shrubs), while another test sought to bound the amount of shrub habitat present in lizard home ranges away from a null expectation. The authors found no effect of shrubs on home range size but did find more shrubs present within lizard home ranges than predicted. Our *a priori* approach (ie taking direct observations of association with shrubs) provided evidence that lizards actively seek out shrubs rather than randomly encountering them during their daily activity. The lack of a correlation between individual shrub use and shrub density suggests that a threshold presence of shrubs may be sufficient to provide thermoregulatory opportunity, therefore a strong correlation between absolute number of shrubs within a home range and home range size would not be predicted. This conclusion is further supported by the Germano and Rathbun's (2016) result that home ranges tended to include more shrub habitat than predicted by the study-site wide prediction, i.e. it is likely beneficial that SOME shrubs be available within the daily activity theater of individual lizards. Our results therefore confirm and are consistent with the results from Germano & Rathbun (2016). Germano and Rathbun (2016) also provide a caveat against overestimating the importance of shrubs to *G. sila* by noting that *G. sila* occurs in places that lack shrubs. Given the variation that we observed in lizard shrub association within one population, it is not surprising that entire populations can persist in relatively shrubless areas. Although heritability of thermoregulatory response in lizards is still undescribed, heritable variation in propensity to use shrubs would predispose a population to adapt either to the presence of shrubs or the absence of shrubs (Logan, Cox & Calsbeek 2014). More to the point, where population-scale variation exists in the predisposition to use shrubs, such as we found in this study, it would be reasonable to propose that shrubs be made available to those lizards that are predisposed to associated with shrubs. The net effect would be to optimize the habitat available for that population. Such optimization may be crucial to impart population resilience to climate change (Sinervo et al 2010, Sears et al 2016). Additionally, structured and/or heterogeneous habitats are becoming increasingly recognized as important to achieve individual-scale thermoregulatory optimization for lizards (Clusella-Trullas & Chown 2014, Basson et al 2016, Sears et al 2016).

Conclusions

Our results document the benefits of shrubs to vertebrate ectotherms in desert communities, including endangered species such as *G. sila*, thus providing guidance for land managers evaluating habitat preservation and restoration designs. We also advance methodology by demonstrating the utility of combining ecological facilitation theory with radiotelemetry. It should be noted that our study was not intended to test the hypothesis that *G. sila* require shrubs per se. Rather, we designed our study to ask whether shrubs provide benefits to *G. sila*, and found evidence to support our hypothesis. In our view this subtle divergence in focus and outcome demonstrates the power of taking an ecological facilitation approach to community interactions.

Author’s Contributions

MFW, CL and CB acquired funding for the project; MW, CJL, SB and TN conceived the study; TN and MW collected the data; CL, TN and MW analysed the data; MW and TN led the writing of the manuscript. Noble participated in design, fieldwork and writing. All authors contributed critically to the drafts and gave final approval for publication.

Data Accessibility

Data are available at <https://cjlortie.github.io/Carrizo.telemetry>

Acknowledgements

R. Seymour, J. Hurl, L. Saslaw, R. Cooper, K. Sharum, E. Gruber all provided assistance. D. Germano allowed us to collar lizards under his permit, for which we thank him.

References

252

253 Adolph, S. C. (1990) 'Influence of behavioral thermoregulation on microhabitat use by two
254 *Sceloporus* lizards', *Ecology*, pp. 315–327. doi: 10.2307/1940271.

255 Bachelet, D. et al. (2016) 'Climate change effects on southern California deserts', *Journal of*
256 *Arid Environments*, 127, pp. 17–29. doi: 10.1016/j.jaridenv.2015.10.003.

257 Basson, C. H. et al. (2017) 'Lizards paid a greater opportunity cost to thermoregulate in a less
258 heterogeneous environment', *Functional Ecology*, 31(4), pp. 856–865. doi:
259 10.1111/1365-2435.12795.

260 Bean, W. T. et al. (2014) 'Species distribution models of an endangered rodent offer conflicting
261 measures of habitat quality at multiple scales', *Journal of Applied Ecology*, 51(4), pp.
262 1116–1125. doi: 10.1111/1365-2664.12281.

263 Bolker, B. M. et al. (2009) 'Generalized linear mixed models: a practical guide for ecology and
264 evolution', *Trends in ecology & evolution*, 24(3), pp. 127–35. doi:
265 10.1016/j.tree.2008.10.008.

266 Bruno, J. F., Stachowicz, J. J. and Bertness, M. D. (2003) 'Inclusion of facilitation into
267 ecological theory', *Trends in Ecology and Evolution*, pp. 119–125. doi: 10.1016/S0169-
268 5347(02)00045-9.

269 Bulleri, F. et al. (2016) 'Facilitation and the niche: Implications for coexistence, range shifts and
270 ecosystem functioning', *Functional Ecology*, 30(1), pp. 70–78. doi: 10.1111/1365-
271 2435.12528.

- 272 Calenge, C. (2006) 'The package adehabitat for the R software: a tool for the analysis of space
273 and habitat use by animals' *Ecological modelling*, 197, 516–519
- 274 Clusella-Trullas, S. and Chown, S. L. (2014) 'Lizard thermal trait variation at multiple scales: A
275 review', *Journal of Comparative Physiology B: Biochemical, Systemic, and*
276 *Environmental Physiology*, pp. 5–21. doi: 10.1007/s00360-013-0776-x.
- 277 Cook, E. R. et al. (2004) 'Long-term aridity changes in the western United States.', *Science*
278 (New York, N.Y.), 306(5698), pp. 1015–8. doi: 10.1126/science.1102586.
- 279 Díaz, J. A. and Cabezas-Díaz, S. (2004) 'Seasonal variation in the contribution of different
280 behavioural mechanisms to lizard thermoregulation', *Functional Ecology*, 18(6), pp. 867–
281 875. doi: 10.1111/j.0269-8463.2004.00916.x.
- 282 Fields, M., Coffin, D. and Gosz, J. (1999) 'Burrowing activities of kangaroo rats and patterns in
283 plant species dominance at a shortgrass steppe-desert grassland ecotone', *Journal of*
284 *Vegetation Science*, 10(1), pp. 123–130. doi: 10.2307/3237167.
- 285 Filazzola, A. and Lortie, C. J. (2014) 'A systematic review and conceptual framework for the
286 mechanistic pathways of nurse plants', *Global Ecology and Biogeography*, pp. 1335–
287 1345. doi: 10.1111/geb.12202.
- 288 Filazzola, A. et al. (2017) 'Non-trophic interactions in deserts: Facilitation, interference, and an
289 endangered lizard species', *Basic and Applied Ecology*, 20, pp. 51–61. doi:
290 10.1016/j.baae.2017.01.002.
- 291 Filazzola, A., Sotomayor, D. A. and Lortie, C. J. (2017) 'Modelling the niche space of desert
292 annuals needs to include positive interactions', *Oikos*. doi: 10.1111/oik.04688.

- 293 Flores, J., & Jurado, E. (2003). Are nurse-protégé interactions more common among plants from
294 arid environments? *Journal of Vegetation Science*, 14, 911–916.
- 295 Germano, D. J. and Williams, D. F. (1992) ‘Recovery of the blunt-nosed leopard lizard: past
296 efforts, present knowledge, and future opportunities’, *Transactions of the Western*
297 *Section of the Wildlife Society*, 28, pp. 38–47.
- 298 Germano D.J., Smith P.T., Tabor S.P. 2007. Food habits of the blunt-nosed leopard lizard
299 (*Gambelia sila*). *The Southwestern Naturalist*. 52(2):318-323.
- 300 Germano, D. J. and Williams, D. F. (2005) ‘Population ecology of blunt-nosed leopard lizards in
301 high elevation foothill habitat’, *Journal of Herpetology*, 39(1), pp. 1–18. doi:
302 10.1670/0022-1511(2005)039[0001:PEOBLL]2.0.CO;2.
- 303 Germano, D. J. et al. (2011) ‘The San Joaquin Desert of California: ecologically misunderstood
304 and overlooked’, *Natural Areas Journal*, 31(2), pp. 138–147. doi: 10.3375/043.031.0206.
- 305 Germano D.J., Rathbun G.B. (2016). ‘Home range and habitat use by blunt-nosed leopard lizards
306 in the southern San Joaquin Desert of California.’, *Journal of Herpetology*. 50(3):429-
307 434.
- 308 Gurney, C. M., Prugh, L. R. and Brashares, J. S. (2015) ‘Restoration of native plants is reduced
309 by rodent-caused soil disturbance and seed removal’, *Rangeland Ecology and*
310 *Management*, 68(4), pp. 359–366. doi: 10.1016/j.rama.2015.05.001.
- 311 Hannah, L., Carr, J. L. and Lankerani, A. (1995) ‘Human disturbance and natural habitat: a
312 biome level analysis of a global data set’, *Biodiversity and Conservation*, 4, pp. 128–155.
313 doi: 10.1007/BF00137781.

- 314 Hawbecker A.C. (1951). 'Small mammal relationships in an *Ephedra* community', Journal of
315 Mammalogy. 32(1), pp. 50-60.
- 316 Hijmans, R. J. et al. (2005) 'Very high resolution interpolated climate surfaces for global land
317 areas', International Journal of Climatology, 25(15), pp. 1965–1978. doi:
318 10.1002/joc.1276.
- 319 Hoekstra, J. M. et al. (2005) 'Confronting a biome crisis: Global disparities of habitat loss and
320 protection', Ecology Letters, 8(1), pp. 23–29. doi: 10.1111/j.1461-0248.2004.00686.x.
- 321 Hothorn, T., Bretz, F. and Westfall, P. (2008) 'Simultaneous inference in general parametric
322 models', Biometrical Journal, pp. 346–363. doi: 10.1002/bimj.200810425.
- 323 Huey, R. B. (1974) 'Behavioral thermoregulation in lizards: importance of associated costs',
324 Science, 184(140), pp. 1001–1003. doi: 10.1126/science.184.4140.1001.
- 325 Huey, R. B. and Slatkin, M. (1976) 'Cost and benefits of lizard thermoregulation', The Quarterly
326 Review of Biology, 51(3), pp. 363–384. doi: 10.1086/409470.
- 327 Kearney, M., Shine, R. and Porter, W. P. (2009) 'The potential for behavioral thermoregulation
328 to buffer "cold-blooded" animals against climate warming', Proceedings of the National
329 Academy of Sciences, 106(10), pp. 3835–3840. doi: 10.1073/pnas.0808913106.
- 330 Kéfi, S. et al. (2007) 'Spatial vegetation patterns and imminent desertification in Mediterranean
331 arid ecosystems', Nature, 449(7159), pp. 213–217. doi: 10.1038/nature06111.
- 332 Kerr, G. D. and Bull, C. M. (2004) 'Microhabitat use by the scincid lizard *Tiliqua rugosa*:
333 Exploiting natural temperature gradients beneath plant canopies', Journal of Herpetology,
334 38(4), pp. 536–545. doi: 10.1670/82-04A.

- 335 Logan, M. L., Cox, R. M. and Calsbeek, R. (2014) 'Natural selection on thermal performance in
336 a novel thermal environment', *Proceedings of the National Academy of Sciences of the*
337 *United States of America*, (19), pp. 1–5. doi: 10.1073/pnas.1404885111.
- 338 Lortie, C. J., Filazzola, A. and Sotomayor, D. A. (2016) 'Functional assessment of animal
339 interactions with shrub-facilitation complexes: A formal synthesis and conceptual
340 framework', *Functional Ecology*, 30(1), pp. 41–51. doi: 10.1111/1365-2435.12530.
- 341 Lortie C.J., Liczner A., Filazzola A., Noble T., Gruber E., Westphal M.F. 2017. 'The Groot
342 Effect: plant facilitation and desert shrub regrowth following extensive damage' *Ecology*
343 *and Evolution* 2017:1–10 doi: 10.1002/ece3.3671
- 344 Lortie C, Filazzola A, Westphal M. (2017): 'The foundation species effect of *Ephedra*
345 *californica*.' Knowledge Network for Biocomplexity. doi:10.5063/F1VM49D1.
- 346 McGowan, J. et al. (2017) 'Integrating research using animal-borne telemetry with the needs of
347 conservation management', *Journal of Applied Ecology*. doi: 10.1111/1365-2664.12755.
- 348 McIntire, E. J. B. and Fajardo, A. (2014) 'Facilitation as a ubiquitous driver of biodiversity',
349 *New Phytologist*, 201(2), pp. 403–416. doi: 10.1111/nph.12478.
- 350 Mouat D.A., Lancaster J.M. 2008. Drylands in Crisis. *Environmental Change and Human*
351 *Security*. 67-80.
- 352 Mohr, C. O. (1947) 'Table of Equivalent Populations of North American Small Mammals',
353 *American Midland Naturalist*, 37(1), p. 223. doi: 10.2307/2421652.

- 354 Prugh, L.R. and Brashares, J. S. (2012) 'Partitioning the effects of an ecosystem engineer:
355 Kangaroo rats control community structure via multiple pathways', *Journal of Animal*
356 *Ecology*, 81(3), pp. 667–678. doi: 10.1111/j.1365-2656.2011.01930.x.
- 357 Pugnaire F.I. (ed.) (2010). 'Positive Plant Interactions and Community Dynamics.' CRC Press,
358 Boca Raton, FL, USA.
- 359 Sawyer J., Keeler-Wolf T and Evens J. 2009. A manual of California vegetation, 2nd ed.
360 California Native Plant Society. Sacramento. 1300 p.
- 361 Schiffman, P.M. (1994) 'Promotion of exotic weed establishment by endangered giant kangaroo
362 rats (*Dipodomys ingens*) in a California grassland', *Biodiversity and Conservation*, 3(6),
363 pp. 524–537. doi: 10.1007/BF00115158.
- 364 Schoepf, I., Schmol, G., Keonig, B., Pillay, N., Schradin, C. (2015). 'Manipulation of
365 population density and food availability affects home range sizes of African Striped
366 Mouse females', *Animal Behaviour* 99:53–60.
- 367 Sears, M.W. and Angilletta, M. J. (2015) 'Costs and benefits of thermoregulation revisited: Both
368 the heterogeneity and spatial Structure of temperature drive energetic costs', *The*
369 *American Naturalist*, 185(4), pp. E94–E102. doi: 10.1086/680008.
- 370 Sears, M.W. et al. (2016) 'Configuration of the thermal landscape determines thermoregulatory
371 performance of ectotherms', *Proceedings of the National Academy of Sciences*, 113(38),
372 pp. 10595–10600. doi: 10.1073/pnas.1604824113.
- 373 Sinervo, B. et al. (2010) 'Erosion of lizard diversity by climate change and altered thermal
374 niches', *Science*, 328(5980), pp. 894–899. doi: 10.1126/science.1184695.

- 375 Stout D., Buck-Diaz J., Taylor S., Evens J. (2014). Vegetation mapping and accuracy assessment
376 report for Carrizo Plain National Monument. California Native Plants Society. Available:
377 https://www.cnps.org/cnps/vegetation/pdf/carrizo-mapping_rpt2013.pdf. Accessed Jan.
378 21th, 2017.
- 379 U.S. Fish and Wildlife Service. 1998. Recovery plan for upland species of the San Joaquin
380 Valley, California. Portland, OR. 1998; 1-319.
- 381 Venables W. N., Ripley B. D. 2002. Modern Applied Statistics with S. Fourth Edition. Springer,
382 New York. ISBN 0-387-95457-0
- 383 Vickers, M., Manicom, C. and Schwarzkopf, L. (2011) 'Extending the cost-benefit model of
384 thermoregulation: high-yemperature environments', *The American Naturalist*, 177(4), pp.
385 452–461. doi: 10.1086/658150.
- 386 Warrick, G. D., Kato, T. T. and Rose, B. R. (1998) 'Microhabitat Use and Home Range
387 Characteristics of Blunt-Nosed Leopard Lizards', *Journal of Herpetology*, 32(2), pp. 183–
388 191. doi: 10.2307/1565295.
- 389 Westphal, M. F. et al. (2016) 'Contemporary drought and future effects of climate change on the
390 endangered blunt-nosed leopard lizard, *Gambelia sila*', *PLoS ONE*, 11(5). doi:
391 10.1371/journal.pone.0154838.

392

393

394

395

396

Tables

Table 1: Generalized linear model for habitat associated with relocated *G. sila*, with degrees of freedom, deviance, and p-values.

Generalized linear model					
Factor	Df	Deviance	P-value		
habitat	1	88.33	< 0.0001		
Time class	1	2.901	0.1		
habitat:time.class	1	5.281	0.01		
Post Hoc, least squared means					
contrast	estimate	SE	df	z.ratio	p.value
open,AM-shrub,AM	0.769229	0.102934	NA	7.473	<.0001
open,AM-open,PM	-0.01848	0.067966	NA	-0.272	0.993
open,AM-shrub,PM	0.44597	0.085189	NA	5.235	<.0001
shrub,AM-open,PM	-0.78771	0.102727	NA	-7.668	<.0001
shrub,AM-shrub,PM	-0.32326	0.11485	NA	-2.815	0.0252
open,PM-shrub,PM	0.464446	0.084938	NA	5.468	<.0001

Table 2: Multinomial logistic regression for observations of *G. sila* behaviors associated with shrubs.

	shrub		Time.class	
Factor	z	P-value	z	P-value
avoiding.predators	6.61E+01	<0.0001	4.60E+07	<0.0001
burrowing	-1.88E+07	<0.0001	2.71E+01	<0.0001
cooling	8.80E+00	<0.0001	1.65E+00	9.91E-02
hunting	8.27E-01	0.4084232	-1.94E+00	5.23E-02
interacting	-1.74E+01	<0.0001	-8.19E-01	4.13E-01
observing	1.14E+00	0.2534383	-8.04E-01	4.21E-01
sunning	6.02E-01	0.5468632	-6.51E+00	7.67E-11

Figures

Figure 1: Left: location of *G. sila* radiotelemetry study. Top left: Location of study area within California. Top right: aerial photograph of study site overlain with sample home ranges calculated using a 95% minimum convex polygon (MCP) estimate, for each individual. Bottom: Aerial image depicting all home ranges of lizards in the study. Different individuals are indicated by different colors.




Figure 2: Plot of *G. sila* behaviors with respect to habitat and time. Lizards engaged significantly more often in cooling behaviors when under shrubs during afternoon temperature peak.





Figure 3: Plots of shrub density on the weighted *G. sila* associations with shrubs.



Supporting Information

Table S1: Behavior classification table for lizard observations.

Classification	Observed behavior
avoiding predators	Moving (most often running) away from predators
burrowing	Actively digging a burrow, or burying itself. This classification was only used if the lizards was actively creating its own burrow, it was not used if a pre-existing burrow was utilized.
cooling	Lizard moving into, or remaining still in shade. Lizard would typically sit upright in shade with front legs extended and rear toes pointed up and off the ground. Occasionally the tail would be lifted off the ground as well.
hunting	Actively stalking or attempting to catch prey. Usually comprised of a slow stalking of an insect and then a sudden burst of speed.
interacting	Interacting with another lizard including either genders of the same species as well as members of other lizard species such as whiptail lizards (<i>Aspidocelis tigrinum</i>).
observing	Actively observing environment (eg moving head or body to track motion).
underground	Lizard underground, behavior could not be otherwise be determined.
sunning	Lizard in sun, not moving. Most often either low to ground, with lower body touching ground or sitting upright with head and shoulders up and rear toes pointed out. Eyes often closed or squinted.

485 **Table S2.** Frequencies of shrub association of individual lizard, density of shrubs within individual home ranges, and percent of area of individual
486 Minimum Convex Polygons (MCP) subsumed by shrub association zones for radiotracked *Gambelia sila* on Elkhorn Plain in 2016.

487

Lizard ID	Total obs.	Shrub obs.	Shrub association frequency	MCP area m ²	Number of shrubs	Shrub density	Shrub assoc area m ²	% Shrub assoc area within MCP
180	59	0	0.000	7509	8	0.001	73.491	0.010
320	5	0	0.000	2	0	0.000	0.000	0.000
360	5	0	0.000	8	0	0.000	0.000	0.000
500	50	0	0.000	803	0	0.000	0.000	0.000
900	9	0	0.000	354	1	0.003	9.186	0.026
939	12	0	0.000	9	0	0.000	0.000	0.000
780	55	5	0.091	8190	50	0.006	459.318	0.056
220	56	7	0.125	57400	104	0.002	955.381	0.017
740	31	4	0.129	5636	7	0.001	64.304	0.011
919	47	8	0.170	40998	124	0.003	1139.108	0.028
381	17	3	0.176	5533	92	0.017	845.144	0.153
439	55	10	0.182	3597	44	0.012	404.200	0.112
760	52	10	0.192	4356	10	0.002	91.864	0.021
660	50	11	0.220	5500	22	0.004	202.100	0.037
960	54	12	0.222	18920	48	0.003	440.945	0.023
520	53	12	0.226	4120	5	0.001	45.932	0.011
820	55	14	0.255	3802	59	0.016	541.995	0.143
840	55	14	0.255	10693	50	0.005	459.318	0.043
240	57	15	0.263	21786	75	0.003	688.976	0.032
979	50	16	0.320	13723	91	0.007	835.958	0.061
0	46	15	0.326	20331	88	0.004	808.399	0.040
800	28	11	0.393	7136	25	0.004	229.659	0.032
540	50	20	0.400	15908	81	0.005	744.095	0.047
420	50	21	0.420	4488	10	0.002	91.864	0.020
680	56	24	0.429	51618	138	0.003	1267.717	0.025
860	51	25	0.490	12759	81	0.006	744.095	0.058
460	50	26	0.520	3838	52	0.014	477.690	0.124
717	56	36	0.643	39292	226	0.006	2076.116	0.053