

# How to critically read ecological meta-analyses

Christopher J. Lortie,<sup>a\*</sup> Gavin Stewart,<sup>b</sup> Hannah Rothstein<sup>c</sup>  
and Joseph Lau<sup>d</sup>

**Meta-analysis offers ecologists a powerful tool for knowledge synthesis. Albeit a form of review, it also shares many similarities with primary empirical research. Consequently, critical reading of meta-analyses incorporates criteria from both sets of approaches particularly because ecology is a discipline that embraces heterogeneity and broad methodologies. The most important issues in critically assessing a meta-analysis initially include transparency, replicability, and clear statement of purpose by the authors. Specific to ecology, more so than other disciplines, tests of the same hypothesis are generally conducted at different study sites, have variable ecological contexts (i.e., seasonality), and use very different methods. Clear reporting and careful examination of heterogeneity in ecological meta-analyses is thus crucial. Ecologists often also test similar hypotheses with different species, and in these meta-analyses, the reader should expect exploration of phylogenetic dependencies. Finally, observational studies not only provide the substrate for potential current manipulative experiments in this discipline but also form an important body of literature historically for synthesis. Sensitivity analyses of observational versus manipulative experiments when aggregated in the same ecological meta-analysis are also frequent and appropriate. This brief conceptual review is not intended as an instrument to rate meta-analyses for ecologists but does provide the appropriate framing for those purposes and directs the reader to ongoing developments in this direction in other disciplines. Copyright © 2013 John Wiley & Sons, Ltd.**

**Keywords:** ecology; criteria; guidelines; interpretation; meta-analysis; reading; and synthesis

## 1. Introduction

We propose that progress in ecology is best promoted by using a range of nonexclusive methods including data synthesis based on formal meta-analyses, systematic reviews, conceptual and theoretical papers, and combinations of large and small experimental studies. 'Conceptual evolution' can occur through detailed experimental tests that refine and hone hypotheses thereby increasing the depth of our understanding of a natural system or set of ideas describing patterns or predicting processes (Paine, 2002). Synthesis in ecology can however also be achieved via the integration of different concepts (Ford, 2000, Ford and Ishii, 2001) and quantitative syntheses of studies testing related ideas or hypotheses (Arnquist and Wooster, 1995). It may seem obvious that science can progress effectively through both avenues, but the majority of ecological research in recent decades has focused on experimental tests of hypotheses. Similar to other mature/maturing disciplines, ecology is now in a position to capitalize on these intensive efforts, because the literature has accrued adequate capacity in breadth and depth on many dominant ideas (Cadotte *et al.*, 2012, Castellanos and Verdu, 2012, Hampton *et al.*, 2013, Jennions *et al.*, 2013a, 2013b). Ecology is thus ripe for a profound shift in focus to use reviews as a means to develop ideas, explore conceptual evolution, seek general conclusions, and assess validity. This paradigm shift is not a new idea in evidence-based medicine (Higgins and Green, 2006) nor in ecology (Arnquist and Wooster, 1995, Carpenter *et al.*, 2009, Gurevitch and Hedges, 1993, Gurevitch *et al.*, 1992, Hampton *et al.*, 2013, Sidlauskas *et al.*, 2009), but the reviews in ecology are shifting from narrative descriptions to systematic reviews and meta-analyses (Cadotte *et al.*, 2012, Chaudhary *et al.*, 2010, Pullin and Stewart, 2006, Vetter *et al.*,

<sup>a</sup>Department of Biology, York University, 4700 Keele St., Toronto, ON M3J1P3, Canada

<sup>b</sup>Centre for Reviews and Dissemination, University of York, York, UK

<sup>c</sup>Department of Management, Baruch College—CUNY, 1 Bernard Baruch Way, New York, NY 10010, USA

<sup>d</sup>Center for Clinical Evidence Synthesis, Tufts Medical Center, 800 Washington Street, Box 63, Boston, MA 02111, USA

\*Correspondence to: Christopher J. Lortie, Department of Biology, York University, 4700 Keele St., Toronto, ON M3J1P3, Canadas.

E-mail: lortie@yorku.ca

A brief note on reading ecological meta-analyses.

2013). These forms of synthesis provide quantitative, replicable insights that can accelerate progress within ecology by providing the reader with a general framework of research completed to date and ideally insights into future directions.

Meta-analysis is an effective tool for synthesizing independent research efforts, comparing the relative success of treatments associated with groups of studies, testing whether mean treatment effects are significantly different than zero, and testing whether the effects are homogeneous or heterogeneous among and within groups or categories of studies. Unfortunately in ecology, the semantics of synthesis, particularly the use of the term meta-analysis, varies significantly (Vetter *et al.*, 2013). Herein, we define an ecological meta-analysis as a review that includes statistical analysis of strength of evidence within and between the studies summarized (Koricheva and Gurevitch, 2013, Vetter *et al.*, 2013). In other disciplines, systematic reviews frequently include not only a summary of the population of studies included in the review and analyses of the literature but statistical analyses of the strength of evidence. The emerging trend in ecology is to decouple the use of the term systematic review from meta-analysis and reserve the latter for reviews that include effect sizes and statistical analysis of within-study evidence. This is an appropriate and convenient distinction for ecology that we would like to further propagate for readers. Meta-analyses thus offer readers in ecology additional capacity to assess attributes of a hypothesis such as testability, generality, consistency, accuracy, and bias in the studies published on a topic at the time of analysis. In contrast to experimental hypothesis testing, the only attribute of a hypothesis perhaps not best tested via meta-analysis is direct falsification. Meta-analyses can be used to summarize evidence for or against an ecological hypothesis. However, when the evidence in a meta-analysis fails to support a hypothesis, this does not necessarily constitute rejection of the hypothesis as useful or predictive in ecology *per se* given that there is a wide range of hypotheses in this discipline ranging from purely descriptive to more formally falsifiable when associated with sets of manipulative experiments (i.e., trials are uncommon in ecology). Failure of a meta-analysis to support a hypothesis may also be a product of the particular set of studies used in the synthesis (Lortie and Callaway, 2006). Nonetheless, this synthesis tool is an important shift toward 'effective thinking' in weighing the strength of ecological ideas, that is, considering effect sizes (Jennions *et al.*, 2013a, 2013b, Nakagawa and Cuthill, 2007) and this ongoing shift away from *p*-values is an important disciplinary transition (Sterne and Smith, 2001). In summary, a meta-analysis is not an experiment and does not test hypotheses but is a means to explore the strength of evidence associated with hypotheses. This is a critical clarification for ecology because there are often large collections of studies documenting only pattern, randomized controlled trials are not used, and even the exact replication of experiments or general protocols is unfortunately relatively infrequent (Kelly, 2006). This trajectory is not unlike progress made in other disciplines such as evidence-based medicine, but ecologists may face unique or at least more frequent subsets of challenges specific to the study of diverse natural systems.

The effective execution of meta-analyses is comprehensively described in a recent handbook for ecologists (Koricheva *et al.*, 2013) and in a succinct methods paper (Harrison, 2011). A recent special issue in the journal 'Evolutionary Ecology' (Nakagawa and Poulin, 2012) also details trends in publications including a meta-analysis of meta-analyses (Castellanos and Verdu, 2012) and the demonstration that meta-analyses are increasing in both frequency and complexity in ecology (Cadotte *et al.*, 2012). However, a simple guide or outline for readers is lacking for ecologists. Only a handful of summaries are available along these lines but for clinicians (Leucht *et al.*, 2009, Ried, 2006) or medical practitioners in general (Russo, 2007), and these deal primarily with synthesis of randomized controlled trials. Epistemologically and empirically, ecology is applying this approach to very different forms of experiments and datasets. Herein, we provide a broad-stroke summary of the general principles/philosophy for the reader to aid in assessing a meta-analysis and summarize a few of the emerging issues specific to ecology. The goal is to provide a general heuristic for the ecological practitioner that has not yet done a meta-analysis.

## 2. Reader guidelines

### 2.1. General principles

**2.1.1. Literature and scope.** In reading ecological meta-analyses, the form of evidence used in the synthesis should be assessed to ensure that the review possesses the capacity to describe the ecological process or hypothesis of interest. Surrogate measures of fitness, stress, population regulation, performance, and sometimes even diversity are common in primary-research ecological studies. This does not weaken the discipline in any way. It is simply a product of the wide-ranging patterns and processes associated with the study of complex and diverse study systems that include different species and categories of drivers. Ecological synthesis efforts include similar decisions. A clear statement of purpose by the authors is necessary and defines the scope of the review and the datasets aggregated. A meta-analysis is still a review and must tell a story (Humphrey, 2011). In spite of its statistics, it is nonetheless a simplification of primary research, and in a discipline that rarely replicates experiments and infrequently uses exact methodologies (Vetter *et al.*, 2013), ecological review stories have considerable potential as an explanatory tool but can also be very easy to misinterpret when the natural history of a system is neglected or as the degree of abstraction is overextended (Lortie and Callaway, 2006). The ecological reader should thus initially confirm that the meta-analysis includes the salient elements needed for any good study such as transparency, replicability, and a clear statement of purpose by the authors (Text Box 1). There are numerous very well articulated current ecological meta-analyses to this effect, including a meta-analysis on climate and litter quality effects on decomposition rates across biomes that provided a map of all

studies used in analyses, a table of study selection criteria, and a clear statement of handling nonindependence issues when a study included more than one location (Garcia-Closas *et al.*, 2013). The savvy reader should thus look for scope of the search, choice of relevant studies, and reporting of the methods used to combine studies (Text Box 1a). There are formal evaluation tools available for syntheses of clinical research such as MOOSE (Stroup *et al.*, 2000) and a quality standard checklist for ecological meta-analysts as well (Rothstein *et al.*, 2013). These instruments need not be used by the reader in any/every instance, but PRISMA (Moher *et al.*, 2009) style tables or visualizations showing how the final population of studies included in the meta-analysis was decided should be provided to the reader (sample flowchart provided in PRISMA statement here: <http://www.prisma-statement.org/2.1.4%20-%20PRISMA%20Flow%202009%20Diagram.pdf>). Inclusion of this aspect of the synthesis workflow provides a degree of transparency (studies were not selected at random or preferentially), the opportunity for replication (the reader could repeat the search on Web of Knowledge or Scopus to confirm that a similar set of studies is generated), and evidence for the purpose of the meta-analysis (excluded studies were outside the statement of scope provided by the authors). The extent to which a meta-analysis can generate robust results thus depends not only on the quality of the synthesis but also on the scope and quality of the included studies. Publication bias is discussed at great length in the technical literature associated with meta-analysis (Koricheva, 2003, Leimu and Koricheva, 2004, Moller and Jennions, 2001, Peters *et al.*, 2007, Tomkins and Kotiaho, 2004), and the reader should expect at least some examination to ensure that the studies included were not unduly skewed to only those that reported positive findings (Jennions *et al.*, 2013a, 2013b). A very thorough exemplar of effective handling of bias is described appropriately in an ecological meta-analysis testing for shifts in plant interactions with increasing environmental stress (He *et al.*, 2013). The authors clearly describe all sets of tests applied to explore bias and inform the reader of decisions associated with exclusions when data were removed. The capacity for a reader to assess the reported statistical evidence will be facilitated by consideration of these three general criteria (transparency, replicability, and purpose) and will reduce the likelihood that the story becomes an oversimplification of the underlying ecology associated with the synthesis effort.

Text Box 1. General guidelines for ecological meta-analysis readers.

## Overarching Principles

Transparency  
Replicability  
Statement of purpose

## A. Literature & Scope

### General Heuristic

Scope of search  
Choice of relevant studies  
Representativeness

### Specifics

Defined inclusion/exclusion criteria for identification of relevant (evidence) studies  
Reasons for inclusion/exclusion documented for each study  
Inclusion/exclusion controlled & listed excluded studies in appendix  
Assessment of study quality/validity: design, context, scale, and taxa  
Data extraction methodology documented and repeatable  
Reporting of aggregation methods across studies  
Estimation of publication bias

## B. Results & Interpretation

### Heuristic

Larger context of evidence framed & interpreted  
Variation effectively explored  
Ecology of system included, i.e. generalizable results

### Specifics

Reported number of studies (N) relative to number of effect size estimates (n)  
Investigation of sources of variation including heterogeneity  
Conducted sub-group analyses or meta-regression  
Partial reporting of covariates in studies listed  
Alternative response variables explored  
Identification evidence gaps & proposed future designs and/or sample sizes  
Common ecological drivers tested (latitude, climate, etc.)  
Appropriate effect sizes calculated & statistical methods applied

Sources: Pullin, A. S., and G. B. Stewart. 2006. *Guidelines for Systematic Review in Conservation and Environmental Management*. *Conservation biology* 10.1111/j.1523-1739.2006.00485.x.  
Russo, M. W. 2007. *How to review a meta-analysis*. *Gastroenterology & Hepatology* 3: 637-642.

It is useful to consider two elements of study validity when interpreting the purpose proposed for ecological meta-analyses. The first element, internal validity, relates to whether a study answers its research question using methods that are free from bias (Gates, 2002, Juni *et al.*, 1999, Treadwell *et al.*, 2007). The reader could consider the experimental design or sampling accuracy of the studies included in the meta-analysis (Text box 1a). Tight experimental control tends to result in studies with high internal validity. The second element, external validity, relates to the generalizability of the research question. Studies with high external validity (i.e., have a broad scope in spatial scale or are not very taxon specific) have high generalizability; thus, the use of appropriate spatial and temporal scales may be important elements of external validity (Text Box 1a). This issue is handled particularly well in an environmental meta-analysis that tested multiple hypotheses for sex differences in mercury contamination in birds (Robinson *et al.*, 2012). The scope of inference or interpretations was tightly controlled, and primary versus ancillary hypotheses were interpreted independently. Ecology often involves tradeoffs between internal and external validity with reductionist approaches having high internal validity and holistic approaches increasing external validity but decreasing internal validity. These tradeoffs are not specific to synthesis.

Formal study inclusion criteria have been proposed and discussed extensively for meta-analysis because of its quantitative nature relative to narrative reviews (Moher *et al.*, 1995, Moher *et al.*, 1996) and for ecology (Cote *et al.*, 2013, Gates, 2002). The primary purpose of inclusion criteria is to ensure that the patterns detected within a meta-analysis are representative of the ecological processes. Effective interpretation of a meta-analysis should at least include a brief inspection of the studies listed. Hence, a useful meta-analysis should provide a list of studies for the reader, preferably in the main body of the paper and not in the appendix. This reporting approach also has the additional benefit of formally crediting the studies used in main citation list of the meta-analysis. A perfect ecological example to this end examining whether similarity between invasive and native species explained resistance clearly listed all studies and how each study tested the hypotheses in question directly in the paper prior to the presentation of the main synthetic findings (Price and Partel, 2013). Similar to primary research studies, most readers should also acquaint herself/himself with the study species or geography of the study to place the findings reported into the appropriate context. Because studies are the substrate of meta-analyses, the same principle applies; the reader should ensure that the synthetic ecosystem is appropriate. This does not imply that authors should not seek to aggregate studies from different ecosystems or use studies with relatively small sample sizes, but that several general principles are relevant for ecologists. The sole criteria for the interpretation of a good ecological study should not be its *p*-value but rather an assessment of whether a given study has the capacity to detect the treatment effect or ecological process of interest. In meta-analyses, the capacity of the primary study to test for an effect is called the prestudy odds that a true effect can be detected (Wacholder *et al.*, 2004), and this is a judgment made by the author. If possible, the reader should consider whether this decision corresponds with their judgment. Importantly, it has been shown that for ecology, there is a strong tendency to cite primary research papers with larger effect sizes regardless of data quality (Barto and Rillig, 2012). Cumulatively adding single studies does little to increase the capacity of a meta-analysis to detect treatment effects (Ioannidis *et al.*, 1998, Ioannidis and Lau, 2001), but a major strength of meta-analysis is nonetheless that many small or even low-quality studies can be combined to provide a comprehensive overview of a hypothesis when each independent experimental test may be equivocal. The reader should be prepared for such surprises but also ensure that the studies used match the scope and scale of the ecological process.

**2.1.2. Results.** Interpreting the magnitude of effect sizes differs from the interpretation of significance levels of statistical tests within a given study. The purpose of any meta-analysis is to compare standardized data across studies, but the interpretation of relevance of the meta-analysis takes place in a much larger context (Walker *et al.*, 2008). The reader needs a frame of reference to assess the outcome of a meta-analysis, and several options are available to generate context. These include comparison with other meta-analyses, translation of effect sizes to other metrics or direct contrasts to contexts or specific groups that readers can comprehend readily (Lipsey and Wilson, 2001). For example, a very general delineation was made from 300 meta-analyses of small versus large effect-sizes in the social sciences (in terms of standardized mean difference), with 'small' being less than 0.2 (i.e., the means of the experimental and control groups differ by 0.2 standard deviations) and 'large' being 0.8 or greater (Lipsey and Wilson, 2001) similar to the original levels proposed for individual studies (Cohen, 1992). A similar exercise was conducted for ecology and evolutionary biology, and it generated 44 published examples (that met particular criteria including reporting effect sizes, etc.), with a mean number of data sets of  $5.3 \pm 1.0$  (Jennions and Moller, 2002) and mean effect sizes ranging from Pearson  $r = 0.180$  to  $0.193$  and Hedges'  $d = 0.631$  to  $0.721$  (Moller and Jennions, 2002). As a starting point, this is an excellent example of an opportunity to both calibrate and estimate the efficacy of sets of ecological approaches. Other researchers reported in a summary note that most effect sizes in ecology and evolution are less than  $d = 0.3$  (Kotiaho and Tomkins, 2002). In general, however, interpretation of the magnitude of mean effect sizes in ecology, at least at this point in time, conforms to the coarse benchmarks for other disciplines with  $d = 0.2$  as small,  $0.5$  as moderate, and  $0.8$  as large, and as such, we propose that readers adopt a similar perspective on assigning relevance. Nonetheless, meta-analytical statistical relevance does not necessarily map directly onto biological significance, and smaller mean effect sizes may be highly relevant and ecologically important in complex, diffuse natural systems (Text Box 1b). Given the increasing frequency of meta-analyses (Cadotte *et al.*, 2012, Chaudhary *et al.*, 2010) and reporting of effect sizes


in primary studies in addition to significance tests, readers can nonetheless now begin to calibrate strength of evidence frameworks for ecology

Finally, the reader should also consider the breadth of confidence intervals when considering the magnitude of mean effect size because variation is such an important aspect of ecology in general. Forest plots are the most common and basic element of most, if not all, meta-analyses in ecology (Lortie *et al.*, 2013). These plots depict both mean effect size estimates and the variances thereby providing a rapid assessment tool for the reader of patterns in relative variation. For instance,  $d=0.6$  is a large effect but if its 95% confidence interval varies extensively, one or two extra studies can change its statistical significance from 'different from 0' to 'not significantly different' (Figure 1, group b). A very clear treatment of variance in forest plots was provided in a meta-analysis testing how insects sense olfactory patches by providing summary effect size estimates for insects from different foraging categories with confidence intervals for each group but also by providing confidence intervals associated with the slope of fit across groups (Andersson *et al.*, 2013). The reader may wish to consider the following general questions: is the number of studies very low (also often provided in good forest plots), is there an important source of variation which has been ignored in the analysis, is there partial reporting of important covariates in some studies and not others which could be added to the meta-analysis, and is there another response variable available to assess whether the patterns of variability associated with the confidence interval is a property of this particular biological system or the response variable selected (Text Box 1b). The reader should also assess whether the authors explored potential dependencies between moderators, provided more than one is available, that is, the importance of latitude in plant community studies or plant traits in interaction studies (Michalet *et al.*, 2013). In many respects, ecology is about variation. Hence, confidence intervals and moderators provide the reader an opportunity to not only more soundly infer the state of research but to assess whether additional studies are needed and if the primary research studies are best categorized into the groupings commonly adopted.

## 2.2. Particularly relevant issues for ecology

**2.2.1. Heterogeneity.** The more diverse the designs used in a meta-analysis, the more likely the synthetic conclusions are to be false (Ioannidis, 2005). Ecologists frequently use very different methods to test hypotheses because we measure populations, communities, ecosystem properties, and organisms. There is also a strong bias against publishing replicated experiments (Vetter *et al.*, 2013). Arguably, this is strength because we can attack hypotheses from various angles, but the reader must cautiously interpret the conclusions of meta-analyses, taking into consideration the diversity of the study set included. In meta-analyses, this is termed heterogeneity (Higgins and Thompson, 2002). There is both methodological heterogeneity (i.e., qualitative) and statistical heterogeneity (Higgins and Green, 2011, Higgins *et al.*, 2003). The reader should expect some treatment and discussion of patterns in both forms, and a common statistical metric,  $I^2$ , which ranges from 0% to 100% has accepted benchmarks of 15% as low, 50% as moderate, and 75% as high (Higgins and Green, 2011). The combination of very different groups, species, processes, and places is integral to predictive and applied ecology (Stewart, 2010). There are three sets of issues the reader should consider with respect to heterogeneity: model, low-quality studies, and sensitivity.

The choice of statistical model (fixed or random) should match the *a priori* purpose of the meta-analysis, that is, is this treatment effective, are these groups different, has this hypothesis been successfully tested, and so on. The



**Figure 1.** A simple forest plot common in ecological meta-analyses. The axes can be flipped, and in most cases, one axis provides the scale of the effect size estimate calculated, and the other axis shows the different groups contrasted. The overall grand mean for the statistical model can also be shown (not provided here). Confidence intervals calculated in various forms, often bootstrapped 95% estimates, are provided. This facilitates an easy visual summary of the differences between and within the groups examined. The sample size is also commonly provided either beside each point estimate as shown here or on the axis in parentheses.



reader should inspect degree of fit, amount of heterogeneity explained, and appropriateness of the statistic generated in satisfying the purpose of the meta-analysis (Text Box 2a). Furthermore, some ecologists have emphasized that different models and metrics may be appropriate depending on the type of variation associated with the set of studies (Osenberg *et al.*, 1999), although some aspects of this recommendation are controversial. As developed in the previous text, definition of the scope of review and evaluation of the respective evidence provides a robust means to assess whether the interpretations proposed by the authors are balanced and reasonable. In some instances, single summary estimates or models may not be adequate to encompass the range of natural variability associated with an ecological hypothesis and may warrant presentation of multiple estimates and models to satisfactorily explain the context dependence (Text Box 2a). A 'great' meta-analysis may focus more extensively on exploring the variation between studies and associated implications (Humphrey, 2011) and less on rejection of a method or hypothesis.

Text Box 2. Frequent considerations in ecological meta-analysis

#### **A. Heterogeneity**

##### Heuristic

Statistical model  
Low-quality studies  
Sensitivity

##### Specifics

Degree of fit of statistical model  
Heterogeneity reported & statistically tested  
Heterogeneity within & between groups interpreted & explanations proposed  
Alternative models explored  
Sign consistency & changes addressed  
Observational versus mensurative methods contrasted  
Studies coded whether directly tested question or reported associated data

#### **B. Phylogenetics**

##### Heuristic

Inclusion of many different species  
Phylogenetic signal & size treated as factors  
Size of dataset relates to nonindependence

##### Specifics

Fixed versus random effects models tested or justified  
Number of species included in the meta-analyses provided  
Tree balance, distribution of nodes, & reporting of phylogenetic correlations  
Phylogenetic signal examined as a form of nonindependence bias  
Alternative statistical approaches explored  
Functional classifications considered

*Sources: (1) Stewart, G. 2010. Meta-analysis in applied ecology. Biology Letters 6: 78-81. (2) Lajeunesse, M. et al. 2013. Phylogenetic nonindependence and meta-analysis. Handbook of meta-analysis in ecology and evolution: 284-299. (3) Chamberlain, S.A. et al. 2012. Does phylogeny matter? Assessing the impact of phylogenetic information in ecological meta-analysis. Ecology Letters 15: 627-636.*

In addition to 'true' heterogeneity among studies or groups of studies, in some instances, the studies may be of 'low quality' for various reasons, may be diverse in methodology, and vary in standards. Nevertheless, it may still be valuable to do a systematic review or even preliminary meta-analysis with the intention of qualitatively describing the body of studies (Treadwell *et al.*, 2007). In this situation, the primary purpose could be to determine net sign, sign differences between groups, or compare levels of variation (Text Box 2a). In ecology, many sets of studies may fall into this qualitative category, and the purpose can be to assess sign, bias, consistency, and generality of outcomes. The ecological reader should expect these synthetic efforts to clearly differentiate between 'no significant mean effect' and 'no differences', depending on the purpose of the meta-analysis (Higgins and Green, 2011). Interpretations rejecting support for an ecological process or hypothesis may not be correct if the evidence loaded into the meta-analysis has low capacity to test the ideas. As such, readers can also use magnitude, sign, sign changes, and comparison between groups as well as formal analysis of heterogeneity to assess meta-analyses.

Different categories of studies are also very common in ecology. In some instances, this can introduce heterogeneity, and the reader should be alert to effective exploration such as sensitivity analyses (i.e., repeated

runs of meta-analysis with and without sets of studies) and contrasts of important categories (Text Box 2a). There is a strong legacy of natural history and pattern analyses in ecology. Consequently, the reader of ecological meta-analyses should also expect different categories of studies provided by authors. Most ecological meta-analyses in the last 5 years provide this information. This not only provides the reader with the capacity to infer research gaps but also provides the substrate for the sensitivity analyses. One of the most common groupings in ecological meta-analyses is observational versus mensurative experiments (Cote and Jennions, 2013, Gates, 2002). Differences between these coarse groups provide the reader with direct insights into how an ecological topic is explored to date and opportunities for novel research. Importantly, this same principle can be applied to contrast sets of methodologies, measures, or group of hypotheses depending on the extent the topic has been explored (Lamarque *et al.*, 2011, Robinson *et al.*, 2012). Heterogeneity is thus an instructive opportunity to explore scaling-up in ecology (Stewart, 2010) and consistency of various methodologies applied.

**2.2.2. Phylogenetics.** Similar to heterogeneity, groups of species can introduce both significant patterns of variation and nonindependence in meta-analyses. This topic is described succinctly in the meta-analysis handbook (Lajeunesse *et al.*, 2013). Application of models to include phylogenetic dependence have also been developed in general (Hadfield and Nakagawa, 2009, Nakagawa and Santos, 2012), and a specific tool for phylogenetic meta-analyses is also available (Lajeunesse, 2011). Conducting a meta-analysis and extracting the data needed to calculate effect sizes are not trivial. Adding evolutionary relationships of the taxa included is also a challenge (Lajeunesse *et al.*, 2013). Nonetheless, this is an important issue. A reanalysis of 30 published meta-analyses to include phylogenetic dependencies dramatically altered the overall pooled effect size of the syntheses in nearly 50% of the fixed-effect analyses and shifted effects from not significantly different from 0 to significance in up to 40% of the datasets (Chamberlain *et al.*, 2012). If possible, inclusion of even coarse but accurate trees in meta-analyses can improve synthesis (Lajeunesse *et al.*, 2013). General principles for the reader are to include as many species as possible in meta-analysis, to consider phylogenetic signal and size as potentially important factors, and to report size of dataset in general because this can generate non-independence issues (Text Box 2b). Important specific applications for the reader to be aware of also include fixed versus random effects models (Borenstein *et al.*, 2010), number of species included in the meta-analyses (Root *et al.*, 2003), tree balance (Blum *et al.*, 2006) or phylodynamics (Poon *et al.*, 2013) statistically contrasting shape, distribution of nodes, reporting of phylogenetic correlations, phylogenetic signal as a form of bias, and alternative statistical approaches (Text Box 2b). The reader should also examine classification of species into functional groups to ensure that these data aggregations are appropriate and meaningful (Maestre and Cortina, 2004). A recent excellent ecological example sorted species into both life history groups and exotic versus native species (He *et al.*, 2013). Phylogenetics added to all forms ecological syntheses will however inject an important and often necessary dimension of evolution to the discipline.

### 3. Summary

To summarize, we propose that there are both several general and specific heuristics that ecological readers should consider in using meta-analyses to discover broad patterns and reach general conclusions about a topic. These include general guidelines associated with meta-analyses such as literature and scope (transparency, replicability, and purpose) and critical appraisal of the results (focus on strength of evidence and not statistical significance). In framing the ecological relevance of a synthesis endeavor, readers should assess heterogeneity, sensitivity, and phylogenetics. Readers should also look for clear delineation of how the strength of evidence was used, reporting and exploration of variability in evidence and sign, and careful use of groups to determine the importance of moderators, covariates, and subgroups. Ecology often takes place in less controlled environments (relative to some other scientific fields) and incorporates many aspects of natural variation in the field, sometimes using very diverse methods to test a hypothesis. Ecology is now in a position to practically apply many forms of synthesis to examine important issues. Interpretations derived from meta-analyses within this domain have the opportunity to not only enhance predictive ecology but also facilitate exploration at larger and novel scales.

### Acknowledgements

This review was funded by a National Center for Ecological Analysis and Synthesis working group grant.

### References

- Andersson P, Lofstedt C, Hamback PA. 2013. How insects sense olfactory patches—the spatial scaling of olfactory information. *Oikos*.
- Arnquist G, Wooster D. 1995. Meta-analysis: synthesizing research findings in ecology and evolution. *Trends in Ecology and Evolution* **10**: 236–240.

- Barto EK, Rillig M. 2012. Dissemination biases in ecology: effect sizes matter more than quality. *Oikos* **121**: 228–235.
- Blum MGB, Francois O, Janson S. 2006. The mean, variance and limiting distribution of two statistics sensitive to phylogenetic tree balance. *The Annals of Applied Probability* **16**: 2195–2214.
- Borenstein M, Hedges LV, Higgins JPT, Rothstein HR. 2010. A basic introduction to fixed-effect and random-effects models for meta-analysis. *Research Synthesis Methods* **1**: 97–111.
- Cadotte MW, Mehrkens LR, Menge DNL. 2012. Gauging the impact of meta-analysis on ecology. *Evolutionary Ecology* **26**: 1153–67.
- Carpenter SR, Armbrust VE, Arzberger PW, Chapin FS, Elser JJ, Hackett EJ, Ives AR, Kareiva PM, Leibold MA, Lundberg P, Mangel M, Merchant N, Murdoch WM, Palmer MA, Peters D, Pickett S, Smith KK, Wall DH, Zimmerman AS. 2009. Accelerate synthesis in ecology and environmental sciences. *Bioscience* **59**: 699–701.
- Castellanos MC, Verdu M. 2012. Meta-analysis of meta-analyses in plant evolutionary ecology. *Evolutionary Ecology* **26**: 1187–1196.
- Chamberlain SA, Hovick SM, Dibble CJ, Rasmussen NL, Van Allen BG, Maitner BS, Ahern JR, Bell-Dereske LP, Roy CL, Meza-Lopez M, Carrillo J, Siemann E, Lajeunesse MJ, Whitney KD 2012. Does phylogeny matter? Assessing the impact of phylogenetic information in ecological meta-analysis. *Ecology Letters* **15**: 627–636.
- Chaudhary VB, Walters LL, Bever JD 2010. Advancing synthetic ecology: a database system to facilitate complex ecological meta-analyses. *Bulletin of the Ecological Society of America* **91**: 235–43.
- Cohen J. 1992. A power primer. *Psychological Bulletin* **112**: 155–159.
- Cote IM, Jennions MD. 2013. The procedure of meta-analysis in a nutshell. In *Handbook of Meta-analysis in Ecology and Evolution*, Koricheva J, Gurevitch J, Mengersen K (eds.). Princeton University Press; 14–26.
- Cote IM, Curtis PS, Rothstein HR, Stewart G 2013. Gathering data: searching literature and criteria selection. In *Handbook of Meta-analysis in Ecology and Evolution*, Koricheva J, Gurevitch J, Mengersen K (eds). Princeton University Press: Oxford; 37–51.
- Ford ED. 2000. *Scientific Method for Ecological Research*. Cambridge University Press: Cambridge.
- Ford ED, Ishii H. 2001. The method of synthesis in ecology. *Oikos* **93**: 153–160.
- García-Closas M, Maestre F, Kattge J, Wall DH 2013. Climate and litter quality differently modulate the effects of soil fauna on litter decomposition across biomes. *Ecology Letters* **16**: 1045–1053.
- Gates S. 2002. Review of methodology of quantitative reviews using meta-analysis in ecology. *Journal of Animal Ecology* **71**: 547–557.
- Gurevitch J, Hedges LV 1993. Meta-analysis: combining the results of independent experiments. In *Design and analysis of ecological experiments*, Gurevitch J (ed.). Chapman and Hall: New York.
- Gurevitch J, Morrow LL, Wallace A, Walsh, JS. 1992. A meta-analysis of competition in field experiments. *American Naturalist* **140**: 539–572.
- Hadfield JD, Nakagawa S. 2009. General quantitative genetic methods for comparative biology: phylogenies, taxonomies and multi-trait models for continuous and categorical characters. *Journal of Evolutionary Biology* **23**: 494–508.
- Hampton SE, Strasser CA, Tewksbury JJ, Gram WK, Budden AE, Batcheller AL, Duke CS, Porter JH. 2013. Big data and the future of ecology. *Frontiers in Ecology & the Environment* **11**: 156–162.
- Harrison F. 2011. Getting started with meta-analysis. *Methods in Ecology and Evolution* **2**: 1–10.
- He Q, Bertness MD, Altieri AH. 2013. Global shifts towards positive species interactions with increasing environmental stress. *Ecology Letters* **16**: 695–706.
- Higgins JPT, Green S. 2006. *Cochrane Handbook for Systematic Reviews of Interventions* 4.2.6. The Cochrane Library 4.
- Higgins JPT, Green S. 2011. *Cochrane handbook for systematic reviews of interventions* 5.1.0. The Cochrane Library 5.1.
- Higgins JPT, Thompson SG. 2002. Quantifying heterogeneity in a meta-analysis. *Statistics in Medicine* **21**: 1539–1558.
- Higgins JPT, Thompson SG, Deeks JJ, Altman DG. 2003. Measuring inconsistency in meta-analyses. *British Medical Journal* **327**: 557–560.
- Humphrey SE. 2011. What does a great meta-analysis look like? *Organizational Psychology Review* **1**: 99–103.
- Ioannidis PA. 2005. Why most published findings are false. *Plos Medicine* **2**: e124.
- Ioannidis PA, Lau J. 2001. Evolution of treatment effects over time: empirical insight from recursive metaanalyses. *Proceedings of the National Academy of Science* **98**: 831–836.
- Ioannidis PA, Cappelleri JC, Lau J. 1998. Issues in comparisons between meta-analyses and large trials. *Journal of American Medical Association* **279**: 1089–1093.
- Jennions MD, Moller AP. 2002. Relationships fade with time: a meta-analysis of temporal trends in publication in ecology and evolution. *Proceedings of the Royal Society of London B* **269**: 43–48.
- Jennions MD, Lortie CJ, Koricheva J. 2013a. Role of meta-analyses in interpreting the scientific literature. In *Handbook of Meta-analysis in Ecology and Evolution*, Koricheva J, Gurevitch J, Mengersen K (eds). Princeton University Press; 364–380.
- Jennions MD, Lortie CJ, Rosenberg MS, Rothstein HR. 2013b. Publication and related biases. In *Handbook of Meta-analysis in Ecology and Evolution*, Koricheva J, Gurevitch J, Mengersen K (eds). Princeton University Press; 207–236.



- Juni P, Witschi A, Bloch R, Egger M. 1999. The hazards of scoring the quality of clinical trials for meta-analysis. *Journal of the American Medical Association* **282**: 1054–1060.
- Kelly CD. 2006. Replicating empirical research in behavioral ecology: how and why it should be done but rarely ever is. *The Quarterly Review of Biology* **81**: 221–236.
- Koricheva J 2003. Non-significant results in ecology: a burden or a blessing in disguise? *Oikos* **102**: 397–401.
- Koricheva J, Gurevitch J. 2013. Place of meta-analysis among other methods of research synthesis. In *Handbook of Meta-analysis in Ecology and Evolution*, Koricheva J, Gurevitch J, Mengersen K. (eds). Princeton University Press; 3–13.
- Koricheva J, Gurevitch J, Mengersen K. 2013. *Handbook of Meta-analysis in Ecology and Evolution*. Princeton University Press.
- Kotiaho JS, Tomkins JL. 2002. Meta-analysis, can it ever fail? *Oikos* **96**: 551–553.
- Lajeunesse M. 2011. phyloMeta: a program for phylogenetic comparative analyses with meta-analysis. *Bioinformatics* **27**: 2603–2604.
- Lajeunesse M, Rosenberg MS, Jennions MD. 2013. Phylogenetic nonindependence and meta-analysis. In *Handbook of Meta-analysis in Ecology and Evolution*, Koricheva J, Gurevitch J, Mengersen K (eds). Princeton University Press; 284–299.
- Lamarque LJ, Delzon S, Lortie CJ. 2011. Tree invasions: a comparative test of the dominant hypotheses and predictor traits. *Biological Invasions* **13**: 1969–1989.
- Leimu R, Koricheva J. 2004. Cumulative meta-analysis: a new tool for detection of temporal trends and publication bias in ecology. *Proceedings of the Royal Society of London B* **271**: 1961–1966.
- Leucht S, Kissling W, Davis JM. 2009. How to read and understand and use systematic reviews and meta-analyses. *Acta Psychiatrica Scandinavica* **119**: 443–450.
- Lipsey MW, Wilson DB. 2001. *Practical Meta-analysis*. Sage Publications Inc: Thousand Oaks, California.
- Lortie CJ, Callaway RM. 2006. Re-analysis of meta-analysis: a test of the stress-gradient hypothesis. *Journal of Ecology* **94**: 7–16.
- Lortie CJ, Lau J, Lajeunesse M. 2013. Graphical presentation of results. In *Handbook of meta-analysis in ecology and evolution*, Koricheva J, Gurevitch J, Mengersen K (eds). Princeton University Press; 339–347.
- Maestre FT, Cortina J. 2004. Do positive interactions increase with abiotic stress? A test from a semi-arid steppe. *Proc. R. Soc. Lond. B* **271**: s331–s333.
- Michalet R, Schob C, Lortie CJ, Brooker RW, Callaway RM. 2013. Partitioning net interactions among plants along altitudinal gradients to study community responses to climate change. *Functional Ecology*. doi: 10.1111/1365-2435.12136.
- Moher D, Jadad AR, Nichol G, Penman M, Tugwell P, Walsh S. 1995. Assessing the quality of randomized controlled trials: an annotated bibliography of scales and checklists. *Controlled Clinical Trials* **16**: 62–73.
- Moher D, Jadad AR, Tugwell P. 1996. Assessing the quality of randomized controlled trials: current issues and future directions. *International Journal of Technology Assessment in Health Care* **12**: 195–208.
- Moher D, Liberati A, Tetzlaff J, Altman DG, Group TP. 2009. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *BMJ* **339**: b2535.
- Moller AP, Jennions MD. 2001. Testing and adjusting for publication bias. *Trends in Ecology and Evolution* **16**: 580–586.
- Moller AP, Jennions MD. 2002. How much variance can be explained by ecologists and evolutionary biologists? *Oecologia* **132**: 492–500.
- Nakagawa S, Cuthill IC. 2007. Effect size, confidence interval and statistical significance: a practical guide for biologists. *Biological Reviews* **82**: 591–605.
- Nakagawa S, Poulin R. 2012. meta-analytic insights into evolutionary ecology: an introduction and synthesis. *Evolutionary Ecology* **26**: 1085–1099.
- Nakagawa S, Santos ES. 2012. Methodological issues and advances in biological meta-analysis. *Evolutionary Ecology* **26**: 1253–1274.
- Osenberg CW, Sarnelle O, Cooper SD, Holt RD. 1999. Resolving ecological questions through meta-analysis: goals, metrics, and models. *Ecology* **80**: 1105–1117.
- Paine RT. 2002. Advances in ecological understanding: by Kuhnian revolution or conceptual evolution. *Ecology* **83**: 1553–1559.
- Peters JL, Sutton AJ, Jones DR, Abrams KR, Rushton L. 2007. Performance of the trim and fill method in the presence of publication bias and between-study heterogeneity. *Statistics in Medicine* **26**: 4544–4562.
- Poon AFY, Walker LW, Murray H, McCloskey RM, Harrigan PR, Liang RH. 2013. Mapping the shapes of phylogenetic trees from human and zoonotic RNA viruses. *PLoS ONE* **8**: e78122.
- Price J, Partel M. 2013. Can limiting similarity increase invasion resistance? A meta-analysis of experimental studies. *Oikos* **122**: 649–656.
- Pullin AS, Stewart GB 2006. Guidelines for Systematic Review in Conservation and Environmental Management. *Conservation Biology* **20**: 1647–1656.
- Ried K 2006. Interpreting and understanding meta-analysis graphs. *Australian Family Physician* **35**: 635–638.
- Robinson SA, Lajeunesse M, Forbes MR. 2012. Sex differences in mercury contamination of birds: testing multiple hypotheses with meta-analysis. *Environmental Science & Technology* **46**: 7094–7101.

- Root TL, Price JT, Hall KR, Schneider SH, Rosenzweig C, Pounds JA. 2003. Fingerprints of global warming on wild animals and plants. *Nature* **421**: 57–60.
- Rothstein HR, Lortie CJ, Stewart G. 2013. Quality standards for research syntheses. In *Handbook of Meta-analysis in Ecology and Evolution*, Koricheva J, Gurevitch J, Mengersen, K (eds). Princeton University Press; 323–228.
- Russo MW. 2007. How to review a meta-analysis. *Gastroenterology & Hepatology* **3**: 637–642.
- Sidlauskas B, Ganapathy G, Jenkins KP, Lapp H, McCall LW, Price S, Scherle R, Spaeth PA, Kidd DM. 2009. Linking big: the continuing promise of evolutionary synthesis. *Evolution* **64**: 871–880.
- Sterne JAC, Smith GD. 2001. Sifting the evidence—what’s wrong with significance tests? *British Medical Journal* **322**: 226–231.
- Stewart G. 2010. Meta-analysis in applied ecology. *Biology Letters* **6**: 78–81.
- Stroup DF, Berlin JA, Morton SC, *et al.*. 2000. Meta-analysis of observational studies in epidemiology: a proposal for reporting. *JAMA* **283**: 2008–2012.
- Tomkins JL, Kotiaho JS. 2004. Publication bias in meta-analysis: seeing the wood for the trees. *Oikos* **104**: 194–196.
- Treadwell JR, Tregar SJ, Reston JT, Turkelson CM. 2007. A system for rating the stability and strength of medical evidence. *BMC Medical Research Methodology* **6**: 52.
- Vetter D, Rucker G, Storch I. 2013. Meta-analysis: A need for well-defined usage in ecology and conservation biology. *Ecosphere* **4**: 74.
- Wacholder S, Chancok S, Garcia-Closas M, El ghormli L, Rothman N. 2004. Assessing the probability that a positive report is false: an approach for molecular epidemiology studies. *National Cancer Institute* **96**: 434–442.
- Walker E, Hernandez AV, Kattan MW. 2008. Meta-analysis: its strengths and limitations. *Cleveland Clinic Journal of Medicine* **75**: 431–439.