
Supplementary information

**A SARS-CoV-2 protein interaction map
reveals targets for drug repurposing**

In the format provided by the
authors and unedited

A list of authors and affiliations appears at the end of the paper

A SARS-CoV-2 Protein Interaction Map Reveals Targets for Drug-Repurposing

Supplementary Information

Supplementary Discussion

In this study we capture and identify 332 high confidence host-pathogen protein-protein interactions (PPIs) between SARS-CoV-2 and human proteins. Below in the Supplemental Discussion we provide what we hope is a thorough, well-sourced, review of the principal interactions for each bait. We have focused on prey proteins involved in comorbidities (e.g. cancer; lung and heart function and disease) and those that are connected to other viral infections or viral protein interactions. When applicable, we have indicated those prey proteins that exhibit host and restriction factor activities. Along these lines, we have characterized evolutionary properties of all prey proteins and identified 40 that evolved under recurrent positive selection in simian primates, thus highlighting a subset of prey proteins that are potentially hijacked or targeted by different pathogens (**Supplementary Table 3**).

All SARS-CoV-2 protein and gene functions described in the subnetwork appendices provided here, including the text below and the text found in the individual bait subnetwork figures, are based on the functions of homologous genes from other coronavirus species. These are mainly from SARS-CoV and MERS-CoV, but when available and relevant other related viruses were used to provide insight into function. The SARS-CoV-2 proteins in this study were designed and researched based on the gene alignments provided by Chan et. al. 2020¹ and Wu et. al. 2020². Though we are reasonably sure the genes here are well annotated, we want to note that not every protein has been verified to be expressed or functional during SARS-CoV-2 infections, either *in vitro* or *in vivo*. In an effort to be as comprehensive and transparent as possible, we are reporting the subnetworks of these functionally unverified proteins along with the other SARS-CoV-2 proteins. In such cases, we have made notes within the text and on the corresponding subnetwork figures, and would advise that more caution be taken when examining these proteins and their molecular interactions. Due to practical limits in our sample preparation and data collection process, we were unable to generate data for proteins corresponding to Nsp3, Orf7b, and Nsp16. Therefore these three genes have been left out of the following literature review of the SARS-CoV-2 proteins and PPIs identified in this study.

To facilitate visualization and understanding we have provided zoomed views of individual bait subnetworks. An interactive version of these networks, including relevant drug and functional information, can be found at the following website: <http://kroganlab.ucsf.edu/network-maps>. Along with our partners at Zoic Labs, we are providing this fully interactive network and the corresponding data tables as a free service to the public. Given the urgent and unprecedented nature of the current crisis and global pandemic, we hope that this will be of significant use to both the scientific and global communities.

STRUCTURAL PROTEINS (S, E, M, N)

S


Function*: Spike (S) is a surface glycoprotein that is responsible for binding and fusion with the host membrane.

- Classified as a class I fusion protein.

- Has 2 subunits that need to be processed by cellular protease TMPRSS2. S1 mediates receptor (ACE2) binding whereas S2 mediates fusion.

Similarity to SARS-CoV

Identity: 76.3% Similarity: 87.0%



Protein

Palmitoylation:

GOLGA7-ZDHHC5 is a protein acyl-transferase (PAT) complex that may play a role in Spike palmitoylation.

E


Function*: Envelope (E) protein plays a central role in virus morphogenesis and assembly.

- Acts as a viroporin, assembling in host membranes and forming pentameric protein-lipid pores that allow ion transport.

- Binds to protein M. Co-expression of M and E is sufficient for VLP formation and release. Lack of E reduces viral titers about 20-fold.

Similarity to SARS-CoV

Identity: 94.7% Similarity: 96.1%



BRD2/BRD4:

Bromodomain extra terminal (BET) proteins are implicated as epigenetic factors that regulate genes crucial for cell cycle progression, inflammation and immune response.

M

Function*: Membrane (M) protein is the major driver for virus assembly and budding.

- Exists as a dimer in two major conformations, long and compact, which together determine the membrane curvature and spike density.

- M-M, M-S and M-N protein interactions contribute to virus assembly.

Similarity to SARS-CoV

Identity: 90.5% Similarity: 96.4%


ATP6V1A: Affects Dengue, West Nile and Influenza A H1N1 virus infections.

Mitochondrial Metabolism

Solute Transport

REEP6: Involved in IL-8 secretion, which plays a role in recruitment of neutrophils in lungs of patients with sepsis-induced ARDS.

ER Morphology





PITRM1: Differentially expressed in RSV-infected human small airway epithelial cells.

BZW2: A known host restriction factor for Dengue virus infection.

ANO6: Deficiency leads to severe T cell exhaustion and the inability of the host to control viral burden

TUBGCP2: Induced by EBV infection and is a known host dependency factor for Influenza A virus.



STRUCTURAL PROTEINS (S, E, M, N)

N

Function*: Nucleocapsid (N) protein binds to the RNA genome.

G3BP1 and G3BP2: G3BP1 and G3BP2 are core structural components of stress granules (SGs) which are broadly refractory to replication of viruses. Viruses have evolved diverse mechanisms, such as direct cleavage of G3BPs (poliovirus, FMDV) or sequestration away from other granule components (SmFV, VV, TMCV, SAFV-2, Mengovirus, DENV, JEV, Ebola virus) to prevent granule formation. Coronaviridae like MERS and IBV also possess specific mechanisms to abrogate SG assembly. Certain viruses utilize G3BPs to promote their replication cycle, a function that is almost exclusively extra-granular.


CK2 (CSNK2A2 and CSNK2B): CSNK2A2 and CSNK2B are subunits of the tetrameric Casein Kinase 2. CK2 phosphorylates G3BPs and disassembles and/or inhibits the formation of stress granules. The activity of CK2 is thus presumptively proviral. CK2 is inhibited at sub-nanomolar concentrations by an orally bioavailable molecule, Silmitasertib.

LARP1: LARP1 is a major effector of the mTOR pathway, suppressing translation of terminal oligopyrimidine mRNAs. LARP1 binds the N protein in a variety of viruses (e.g. IBV, IAV). LARP1 knockdown decreases DENV viral titers, while inhibition of mTOR (e.g. with rapamycin) impairs MERS-CoV replication and exerts immunosuppressant functions.

MOV10: MOV10 is a 5' to 3' RNA helicase that interacts with UPF1 and binds 3'UTRs. Its antiviral functions are independent of the helicase activity, and often through IFN stimulation. MOV10 exhibits P-body dependent antiviral activity by binding the N protein and preventing its nuclear localization.


PABPC1/4: Poly-A binding proteins are involved in many steps of mRNA processing. Viral factors cause PABPC1/4 to shuttle into the nucleus, causing mRNA hyperadenylation and nuclear retention. In a complex with LARP1 and RyDEN, PABPC1/4 can promote virulence.

UPF1: UPF1 is a RNA helicase that functions in the nonsense-mediated mRNA decay (NMD) pathway. Murine Hepatitis Virus (MHV) mRNAs are subjected to NMD, while the MHV N protein has been shown to have NMD inhibitory functions.



Similarity to SARS-CoV

Identity: 90.5%	Similarity: 94.3%
------------------------	--------------------------



NON-STRUCTURAL PROTEINS

Nsp1


Function*: Nsp1 antagonizes interferon induction to suppress host antiviral response.

- Overexpression of Nsp1 in A549 cells increases production of pro-inflammatory chemokines CCL5, CXCL10, and CCL3.

- Can also inhibit host gene expression by binding to ribosomes and modifying host mRNAs.

Similarity to SARS-CoV

Identity: 84.4% **Similarity:** 91.1%



DNA Polymerase α Complex: Regulates the activation of type I interferons through cytosolic RNA-DNA synthesis and primes DNA replication in the nucleus.

PKP2 (Plakophilin): Binds cadherins and intermediate fibers, crucial for desmosome formation.

COLGALT1: Required for galactosylation of collagen IV and VI to form the collagen triple helix.


Nsp2

Function*: Nsp2 is translated as part of a single protein along with Nsp3 and may serve as an adaptor for Nsp3.

-While not essential for viral replication, deletion of Nsp2 diminishes viral growth and RNA synthesis.

Similarity to SARS-CoV

Identity: 68.3% **Similarity:** 82.9%



Translation Repression

SLC27A2: Previously described as a Dengue virus restriction factor.

Endosomal Transport

Nsp4

Function*: Nsp4 forms a complex with Nsp3 and Nsp6.

- Together, these proteins are predicted to nucleate and anchor viral replication complexes on double-membrane vesicles in the cytoplasm.


Similarity to SARS-CoV

Identity: 80.0% **Similarity:** 90.8%

TIM Complex: Involved in the import and insertion of hydrophobic membrane proteins into the mitochondrial inner membrane. Regulates import of transmembrane proteins into the inner mitochondrial membrane.

ALG11: Mannosyltransferase involved in the synthesis of core oligosaccharide.

IDE: Involved in antigen presentation via MHC class I.



SARS-CoV-2 Viral Protein (26)
 Human Protein
 Drug Target
 Human-Human Protein-Protein Interaction

MIST Score
 Spectral Count


Protein Complex
 Biological Process

NON-STRUCTURAL PROTEINS

Nsp5

Function*: Nsp5 is the 3C-like protease.

-Cleaves the viral polyprotein.



HDAC2: Deacetylates lysines at the N-terminus of histones. Reduced levels of HDAC2 leads to increased transcription of inflammatory genes. Low HDAC2 expression contributes to disease severity in COPD patients.

Similarity to SARS-CoV

Identity: 96.1% Similarity: 98.7%

Nsp5_C145A

Function*: Nsp5_C145A is a catalytically dead mutant of the Nsp5 3C-like protease.

-The catalytic residues of SARS-CoV Nsp5 align to H41 and C145 of SARS-CoV2 Nsp5.



GPX1: Glutathione peroxidase 1 is an antioxidant enzyme helps regulate the redox state of the cell, mitochondrial function, and apoptosis. It has been linked to a number of diseases that increase the risk of severe SARS-CoV-2 comorbidity including cancer, diabetes, and cardiovascular disease.

Similarity to SARS-CoV

Identity: 96.1% Similarity: 98.7%


Nsp6

Function*: Nsp6 limits autophagosome expansion.

- Nsp6 may favor SARS-CoV infection by compromising the ability of autophagosomes to deliver lysosomes for degradation

-Complexes with Nsp3 and Nsp4 to form double-membrane vesicles that anchor viral replication complexes.

Ion Transport: Components of the Mitochondrial Complex V co-purify with Nsp6. This complex regenerates ATP from ADP.




ATP6AP1: Subunit of the vacuolar ATP synthase protein pump. Dysregulation results in impaired vesicle acidification and intracytoplasmic granules, resulting in a range of pathologies including an immunodeficiency syndrome and granular cell tumors. Identified as a potential host factor for IAV, WNV, and DENV.

SIGMAR1: Sigma Receptor 1 is an ER chaperone protein that modulates calcium signaling through its interaction with the IP3 receptor. It is targeted by a number of existing drugs including Chloroquine and haloperidol.

Similarity to SARS-CoV

Identity: 87.2% Similarity: 94.8%



REPLICATION COMPLEX (Nsp7, Nsp8, Nsp12)

Nsp7

Function*: The Nsp7-Nsp8 complex is part of a unique multimeric RNA-dependent RNA replicase capable of both de novo initiation and primer extension.


- Forms the primase in complex with Nsp8.

Similarity to SARS-CoV

Identity: 98.8% **Similarity:** 100.0%

ACSL3: Induction results in increased acyl-CoA synthesis that is essential for providing prostaglandin.

MOGS: Cleaves the distal alpha 1,2-linked glucose residue from the N-linked oligosaccharide precursor in a highly specific manner.



LMAN2: Intracellular lectin involved in the early secretory pathway. May interact with O-linked glycans and N-acetyl-D-galactosamine.

Nsp8

Function*: Nsp8 forms a primase in complex with Nsp7.


-Eight Nsp8 proteins complex with eight Nsp7 proteins to form a a hexadecameric structure surrounding dsRNA.

Similarity to SARS-CoV

Identity: 97.5% **Similarity:** 99.0%

Exosome: Degrades ssRNA in a 3' to 5' direction and is involved in homeostatic degradation of host RNA as well as antiviral immunity.

Signal Recognition Particle: Binds to signal recognition peptides and mediates insertion of proteins into the endoplasmic reticulum.



Nsp12

Function*: Nsp12 is the RNA-dependent RNA polymerase (RdRp).

- Nsp12 contains a large two-domain N-terminus of little known function and a canonical RdRp domain in the C-terminus.


- Nsp7-Nsp8 heterodimer binds the RdRp domain of Nsp12.

Similarity to SARS-CoV

Identity: 96.4% **Similarity:** 98.3%

RIPK1: Triggers cell death by apoptosis or necrosis as an active regulatory kinase. Regulates inflammatory signaling and inhibits cell death as a scaffold protein.

Spliceosome: Removes introns from pre-mRNA. SLU7 is essential for the second catalytic step of pre-mRNA splicing.



◆ SARS-CoV-2 Viral Protein (26)
● Human Protein
● Drug Target
— Human-Human Protein-Protein Interaction

1.0
0.7
181
2
MIST Score
Spectral Count

● Protein Complex
● Biological Process

NON-STRUCTURAL PROTEINS

Nsp9

Function*: Nsp9 is an essential single-stranded RNA binding protein.


- Shown to interact with the replication complex (Nsp7, Nsp8, and Nsp12).
- Binds to both DNA and RNA but preferentially binds single-stranded RNA.

DCAF7: Potential substrate adaptor for CUL4 E3 Ligase; interacts with adenovirus E1A innate immune response and depresses IFN stimulated genes.

NEK9: Serine/threonine kinase shown as an adenovirus dependency factor.

Similarity to SARS-CoV

Identity: 97.3% Similarity: 98.2%



Nuclear Pore Complex: Several components of the NPC are targeted by viruses in an effort to block or promote nuclear transport beneficial to virus replication.

Fibrillin: Fibrillin-1 and -2 (FBN1/2) are extracellular matrix glycoproteins involved in elastin fiber and respiratory organ development.

Nsp10

Function*: Nsp10 is a zinc-finger protein essential for replication.

- Has been implicated in negative-strand RNA synthesis.
- Acts as a stimulatory factor for Nsp16 to execute its methyltransferase activity.


GFER: FAD-dependent sulfhydryl oxidase that regenerates the redox-active disulfide bonds in CHCHD4/MIA40.

ERGIC1: Plays a possible role in transport between the endoplasmic reticulum and Golgi.

GRPEL1: Essential component of the PAM complex, required for the translocation of transit peptide-containing proteins from the inner membrane into the mitochondrial matrix in an ATP-dependent manner.

Similarity to SARS-CoV

Identity: 97.1% Similarity: 99.3%



AP2 Clathrin: Nsp10 may hijack the clathrin machinery and endocytosis host proteins.

Nsp11

Function*: It is unclear if Nsp11 encodes a functional viral protein.


- Short peptide (13 amino acids) at the end of Orf1a.

Similarity to SARS-CoV

Identity: 84.6% Similarity: 92.3%



TBCA: Involved in the early steps of the tubulin folding pathway




CAPPING ENZYMES (Nsp13, Nsp14)

Nsp13

Function*: Nsp13 is a helicase and triphosphatase that initiates the first step in viral mRNA capping.

- Nsp13, along with Nsp14 and Nsp16, installs the cap structure onto viral mRNA in the cytoplasm instead of the nucleus where host mRNA is capped.



Protein Kinase A Signaling: PKA signaling coordinates multiple steps of membrane transport and may be hijacked for viral benefit. AKAP9 is a large scaffolding protein that localizes to the Golgi and centrosomes.


Similarity to SARS-CoV

Identity: 99.8% Similarity: 100.0%

Nsp14

Function*: Nsp14 is a bifunctional enzyme encoding both an exonuclease and a SAM dependent methyltransferase domain.

- The exonuclease domain corrects mutations that arise during genome replication.
- The SAM dependent methyltransferase domain facilitates capping of viral mRNA.



IPDH2: Catalyzes the conversion of isosine 5' phosphate (IMP) ultimately to guanine 5' monophosphate for de novo synthesis of guanine nucleotides. Nsp14's interaction with IMPDH2 may reflect an interplay with purine nucleotide metabolism.

Similarity to SARS-CoV


Identity: 95.1% Similarity: 98.7%

Nsp15

Function*: Nsp15 has uridine-specific endoribonuclease (endoU) activity and is essential for viral RNA synthesis.

- endoUs shown to: (i) have endonucleolytic activity; (ii) cleave 3' of pyrimidines, preferring uridine > cytidine; and (iii) release reaction products with 2'-3'-cyclic phosphate and 5'-OH ends.
- Shown to form homohexamers composed of a dimer of trimers.

NUTF2: Mediates the import of GDP-bound RAN from the cytoplasm into the nucleus, thus indirectly plays a more general role in cargo receptor-mediated nucleocytoplasmic transport.




RNF41: E3 ubiquitin-protein ligase that promotes TRIF-dependent production of type I interferon and inhibits infection with vesicular stomatitis virus.

ARF6: GTP-binding protein involved in protein trafficking. Regulates endocytic recycling and cytoskeleton remodeling.

Similarity to SARS-CoV

Identity: 88.7% Similarity: 95.7%



OPEN READING FRAMES

Orf3a


Function*: Orf3a is not essential for replication but contributes to pathogenesis. It is packaged into virions.

- Mediates trafficking of Spike (S protein) by providing an ER/Golgi retention signal.
- Induces elevation of IL-1 β secretion and activates NF- κ B and the NLRP3 inflammasome by promoting TRAF3-dependent ubiquitination of p105 and ASC.
- Expression of Orf3a induces apoptosis in viral infection and cell line models.

Similarity to SARS-CoV

Identity: 72.4% **Similarity:** 85.1%

ALG5: Involved in N-linked protein glycosylation. Identified as a host factor for IAV replication.



CLCC1: An intracellular chloride channel implicated in ER stress. Interacts physically with RSV and druggable protein SLC5A13.

HMOX1: Key enzyme in heme catabolism. Has shown a cytoprotective and anti-inflammatory effect both in pulmonary pathologies and viral pathogenesis.

HOPS Complex: VPS11 and VPS39 are members of the HOPS and CORVET complexes, respectively, which coordinate fusion of the lysosome with the endosome and autophagosome.

Orf3b

Function*: Orf3b is shown to be an interferon antagonist and is involved in pathogenesis.

Similarity to SARS-CoV

Identity: 7.1% **Similarity:** 9.5%



STOML2: Stimulates cardiolipin biosynthesis and binds and stabilizes prohibitin. Both STOML2 and prohibitin have been shown to be host dependency factors for Enterovirus (EV)71.


Orf6

Function*: Orf6 is a type 1 interferon antagonist. Expression of Orf6 suppresses the induction of interferon and interferon signaling pathways.

- C-terminal region of SARS-CoV Orf6 interacts with the nuclear import protein, karyopherin alpha-2, sequestering it in the cytoplasm. This prevents import of STAT1, activator of interferon response genes, into the nucleus.

Similarity to SARS-CoV

Identity: 66.7% **Similarity:** 85.7%



SARS-CoV-2 Viral Protein (26)
 Human Protein
 Drug Target
 Human-Human Protein-Protein Interaction

MIST Score
 Spectral Count

Protein Complex
 Biological Process

OPEN READING FRAMES

Orf7a


Function*: Orf7a may play a role in pathogenesis via its role in virus-induced apoptosis.

- ΔOrf7a SARS-CoV is still able to release virions at similar levels as wild-type virus.

Similarity to SARS-CoV

Identity: 85.2% Similarity: 90.2%

HEATR3: Involved in ribosomal protein transport. Implicated in mediating NF-κB signaling.



MDN1: Nuclear chaperone that is critical for the export of pre-60s ribosome subunits.

Orf8


Function*: Orf8 is an accessory protein not essential for virus replication in vitro and in vivo.

- Previously shown to be a recombination hotspot, one of the most rapidly evolving regions among SARS-CoV genomes.

- SARS-CoV Orf8b was shown to induce ER stress and activate NLRP3 inflammasomes.

Similarity to SARS-CoV

Identity: 28.5% Similarity: 45.3%




Orf9b

Function*: Orf9b is an accessory protein synthesized from an alternative complete reading frame within the viral N gene.


- Targets the mitochondrial-associated adaptor molecule MAVS signalosome by utilizing PCBP2 and E3 ligase AIP4, resulting in the degradation of MAVS and therefore limiting host cell interferon responses.

Similarity to SARS-CoV

Identity: 72.4% Similarity: 84.7%



TOMM70: Receptor that accelerates the import of all mitochondrial precursor proteins. TOMM70 interacts with MAVS protein upon virus infection.



OPEN READING FRAMES

Orf9c

Function*: Orf9c is a short polypeptide (70 amino acids) dispensable for viral replication. There is no data yet providing evidence that the protein is expressed during SARS-CoV-2 infection.

ACC1: ABCC1 (also MRP1) is a multifunctional ATP-binding cassette protein involved in controlling the efflux of drugs in cells. MRP1 is a well-known viral host factor and physical interactor of both IAV and WNV proteins. Further, it has been implicated in disease progression of pneumonia and COPD, as well as drug resistance in the lung.


F2LR1: The protein product of F2LR1, PAR2, is a protease-activated receptor that has a cytoprotective and inflammatory role in the pathogenesis of viral infection and progression of pulmonary disease.

BCS1L: BCS1L is a mitochondrial chaperone located in the inner mitochondrial membrane. Loss of BCS1L is associated with severe clinical disorders such as GRACILE and Bjornstad Syndrome.

Similarity to SARS-CoV

Identity: 74.0% Similarity: 78.1%

Electron Transport: Orf9c interacts with four proteins in the mitochondrial respiratory electron transport chain complex I, which has demonstrated roles in TLR/IL-1 signaling and mediating inflammation.




GPI-Anchor Biosynthesis: GPAA1 is essential for GPI-anchoring of precursor proteins, while PIGS and PIGO are involved in GPI-synthesis.

Orf10

Function*: Orf10 codes for a peptide only 38 amino acids long. There is no data yet providing evidence that the protein is expressed during SARS-CoV-2 infection.


- Does not have a homolog in SARS-CoV.

Cul2 Complex: Commonly hijacked by viral proteins to ubiquitinate and degrade viral restriction factors. The CUL2^{ZYG11B} E3 ligase targets substrates with exposed N-terminal glycines for degradation



Similarity to SARS-CoV

Identity: NA Similarity: NA



SARS-CoV Structural Proteins

SARS-CoV-2 Spike/S Protein

Spike (S protein) is the viral surface protein that mediates viral entry through its interaction with cellular host factor angiotensin-converting enzyme 2 (ACE2)³. The full-length S protein is 141 kDa (1273 amino acids (aa)) and encodes a N-terminal signal peptide, a transmembrane region (aa 1214-1237), and a patch of cysteine residues (aa 1235-1254) which are predicted to be palmitoylated (protein S-acylation of cysteine residues)^{4,5}. In previous studies, mutation of these cytosolic cysteine residues inhibited fusogenicity of S, and suggest that targeting the s-acylation modifications via acyl-transferase inhibitors could potentially be a therapeutic strategy for inhibiting coronavirus infection⁶.

In our interactome, we demonstrate high-confidence binding of SARS-CoV-2 Spike to the GOLGA7-ZDHHC5 complex, a protein acyl-transferase (PAT) complex. GOLGA7 characteristically localizes to the Golgi apparatus⁷ but the GOLGA7-ZDHHC5 complex has been found at the plasma membrane⁸. While this interaction is exciting and could potentially point to a targetable enzymatic activity responsible for Spike palmitoylation, further experiments need to be done to fully clarify the functional role of this complex and the consequence of its inhibition. ZDHHC5 is highly expressed in neuronal tissue and is reported to have a role in hippocampal function⁹. In addition, ZDHHC5 is shown to palmitoylate G protein-coupled receptors¹⁰ and activate nucleotide oligomerization domain (NOD)-like receptors 1 and 2 (NOD1/2)¹¹ which recognize pathogenic peptidoglycans and activate immune signaling¹¹. GOLGA7 is also a general adaptor for ZDHHC proteins and, in complex with these acyl-transferase proteins, can regulate additional processes including the palmitoylation of Ras¹². It should also be noted that the ZDHHC5-GOLGA7 complex is required for the non-apoptotic cell death phenotype attributed to the synthetic oxime CIL56, indicating a role for this complex in the cell death pathway⁸. Thus, we would caution against long-term inhibition of this enzymatic complex as a general treatment strategy.

Though identified just below our cutoffs, one of the lower confidence hits for Spike was ATP1A1, a protein at the plasma membrane that promotes viral entry and replication of related coronaviruses (M-CoV, FCoV, MERS-CoV), and a number of other viruses including Ebola, Lassa, live-attenuated Junín virus, and respiratory syncytial virus (RSV)^{13–16}. Interestingly, this protein is from the same ATPase family as ATP1B1, a high-confidence interactor of M protein.

SARS-CoV-2 E Protein

The SARS-CoV Envelope (E) protein, a small integral membrane protein, is important for virus production, assembly, intracellular trafficking, morphogenesis and virulence^{17,18}. The transmembrane domain of E has amphipathic properties and can oligomerize to form cation-permeant ion channels¹⁹. The C-terminal cytoplasmic domain contains a targeting signal for protein localization to the Golgi complex^{20,21} and a PDZ binding motif important for virulence²². Interestingly, E protein is expressed at very high levels inside the infected cells, but only a small portion is incorporated in virions and the rest is largely distributed in intracellular membranes between ER and Golgi compartments indicating an important role of E protein in manipulating the cellular environment²³. Increased cellular stress, unfolded protein response, apoptosis, and heightened host immune response were all observed in infected cells lacking E protein¹⁸. In addition, E has also been implicated in inflammasome activation and consequential inflammation in the lung parenchyma¹⁸.

In our study, we identify six high-confidence host protein interactions with E protein: AP31B, BRD2, BRD4, CWC27, SLC44A2, and ZC3H18. BRD2 and BRD4 are bromodomain extra terminal (BET) family

proteins that bind acetylated chromatin and activate transcription²⁴. BRD4 is a co-activator of NF- κ B and facilitates the transcription of NF- κ B-dependent inflammatory response²⁵. Sequestration of BRD4 by E protein may represent a means for SARS-CoV-2 to protect against the host immune response. Previously it's been shown that inhibition of BRD4 activity in primary lung epithelial cells results in diminished innate immune response after challenge with poly(I:C), a viral pattern that simulates acute RNA virus infections²⁶. In addition, the interaction of BRD4 with Bovine Papillomavirus E2 protein tethers viral DNA to host mitotic chromosomes ensuring viral persistence in infected cells²⁷. BRD2 is a transcriptional regulator that binds hyperacetylated histones and is implicated in a variety of cellular processes^{28,29}. Though functionally similar, BRD2 and BRD4 are shown to regulate different transcriptional programs³⁰. Given that bromodomain proteins and their ability to regulate transcription are implicated in the life cycle of several viruses^{27,28,31,32}, the observed interactions of E protein with BRD2 and BRD4 present exciting avenues to pursue. Future research is still needed to fully establish the structural basis of these interactions and the resulting functional implication for SARS-CoV-2 infection.

E protein also interacts with host factors involved in protein and mRNA trafficking, important for potentially carrying out many of E protein's functions. AP3B1 encodes the beta-1 subunit of adaptor protein complex 3 (AP-3), that is involved in signal-mediated protein sorting from Golgi membranes to endosomal-lysosomal organelles^{33,34}. Mutations in AP3B1 affect protease activity in endosomes and produce pathology-associated defects with aberrant transmembrane lysosomal protein trafficking³⁵. AP-3 depletion resulted in decreased localization of the human immunodeficiency virus (HIV)-1 major structural protein Gag from plasma membrane and late endosome, suggesting that intact AP-3 is required for HIV-1 particle production and release³⁴. Also, we have previously shown that AP-3 is targeted by the NS5 protein of West Nile virus³⁶. Thus it is possible that SARS-CoV-2 may also utilize AP-3 for particle assembly, transport, and release.

Other notable high-confidence host protein interactions with E protein include CWC27 and SLC44A2. Both SLC44A2 and CWC27 are proteins implicated in other diseases, but have yet to be characterized in relation to viral infections.

SARS-CoV-2 M Protein

Membrane (M) protein, the most abundant protein in coronavirus particles, is a type III transmembrane glycoprotein located in the virus envelope. M protein manipulates cellular membranes to bring viral and host factors together and therefore is the main driver for virus assembly and budding processes. M-M, M-S and M-N protein interactions all contribute to this assembly process³⁷. In the virion, M exists as a dimer in two major conformations (long and compact), which together determine the membrane curvature and spike density^{38,39}. SARS-CoV M protein is known to affect various host processes in a manner beneficial to viral replication and infectivity⁴⁰. M suppresses key inflammatory molecules (e.g. NF- κ B and Cox-2)⁴¹ and counteracts host viral defenses by inhibiting type I interferon production⁴². SARS-CoV M protein modulates apoptosis by interfering with PDK1–PKB/Akt signaling. This interaction between M protein and PDK1 could be targeted as a plausible therapeutic approach to modulate the pro-apoptotic properties attributed to coronavirus infection^{43–45}.

In this study, we identified 30 high-confidence physical interactors of SARS-CoV-2 M protein. Transmembrane domain-mediated oligomerization of M protein at ER-Golgi intermediate compartment (ERGIC) membranes drives the assembly of coronavirus which buds into the lumen of the ERGIC⁴⁶. Therefore, it was not surprising to find many SARS-CoV-2 M protein interactors were transmembrane proteins including mitochondrial proteases (PITRM1, PMPCA, PMPCB), ATPases (ATP6V1A, ATP1B1) and kinases (AKAP8L, COQ8B, FASTKD5). In addition, we identify known host dependency (ATP6V1A, ATP1B1, TUBGCP2, RTN4) and restriction (BZW2) factors of other viruses. ATP6V1A is a dependency factor for Dengue virus⁴⁷, West Nile

virus⁴⁷ and Influenza A H1N1 virus⁴⁸, ATP1B1 is involved in viral autophagy⁴⁹ and has a role in Sindbis virus (SINV) infection⁵⁰, and TUBGCP2 is a host dependency factor for Influenza A virus^{51,52}. RTN4 is a reticulon protein involved in the formation and stabilization of endoplasmic reticulum (ER) tubules^{53,54}. Reticulon proteins play crucial roles in viral RNA replication, compartment formation, and function⁵⁵ and are known to protect ER membrane integrity during polyomavirus SV40 infection⁵⁶. RTN4 is also involved in the formation of the *Legionella pneumophila* containing vacuoles⁵⁷. In a functional RNAi screen, BWZ2 was identified as a host restriction factor for dengue virus (DENV) infection⁵⁸. BZW2 is a translation repressor known to regulate the PI3K/AKT/mTOR signaling pathway⁵⁹⁻⁶¹. Recently, BZW2 was identified as an oncogene implicated in a number of cancers including lung adenocarcinoma, where overexpression was negatively correlated with overall disease free survival⁶⁰⁻⁶⁴. Studies suggest that BZW2 overexpression could also contribute to drug resistance in cancer cells, including resistance to rapamycin⁶¹.

The mechanism behind acute respiratory distress in severe SARS-CoV-2 infections is not understood but the host immune response is likely to determine the pathogenesis to some extent. We found that two related host factors REEP5 and REEP6 copurify with SARS-CoV-2 M protein. REEP5 and REEP6 can refine CXCR1-mediated cellular responses and lung cancer progression⁶⁵. REEP6 is also involved in IL-8 secretion⁶⁶. IL-8 is the major chemoattractant for neutrophils and is implicated to have a major role in the recruitment of neutrophils to the lungs in patients with sepsis-induced acute respiratory distress syndrome⁶⁷. Another high-confidence M protein interactor ANO6 is also very exciting from the perspective of viral pathogenesis. ANO6 is required to curb excessive T cell responses in chronic viral infections. ANO6-deficient T cells are hyperactivated during the early phase of infection, exhibiting increased proliferation and cytokine production. This overactivation ultimately leads to severe T cell exhaustion and the inability of the host to control viral burden⁶⁸. Interestingly, ANO6 is localized in late endosomes and SARS-CoV is also targeted to endosomes⁶⁹. Mice deficient for a related protein ANO1 suffer from tracheomalacia and die shortly after birth because of respiratory failure; ANO6 is highly expressed in the respiratory system and its involvement in respiratory functions is likely⁷⁰ making it an interesting candidate for future mechanistic studies in the context of SARS-CoV-2 infection.

SARS-CoV-2 N Protein

SARS-CoV nucleocapsid (N) protein is an essential structural protein that binds to viral genomic RNA (gRNA) in virions. N protein dimerization and its association with viral gRNA are crucial for viral assembly^{71,72}. It has been shown that the N protein binds to both intracellular gRNA and subgenomic RNA (sgRNA), suggesting functions in viral transcription and translation⁷³. N protein also has implications in a variety of cellular responses, including stress granule formation and host translation shutoff⁷⁴, inhibition of nonsense mediated mRNA decay (NMD)⁷⁵ and cell cycle regulation⁷⁶.

The SARS-CoV-2 N interactome includes 15 high-confidence host protein interactions, many of which are host mRNA binding proteins, including stress granule (SG)-related factors (G3BP1/2, MOV10, CK2 subunits, and PABP proteins), mRNA decay factors (MOV10 and UPF1), mTOR translational repressors (LARP1), and protein kinases (CK2 subunits). Several N protein interactors including G3BP1, G3BP2, LARP1, MOV10, PABPC1 and PABPC4 have also been detected in a mouse hepatitis virus interactome⁷⁷ and in two previous nucleocapsid protein interactomes from infectious bronchitis virus⁷⁸ and Influenza A virus⁷⁹. Numerous viruses have evolved diverse mechanisms to abrogate SG assembly. Poliovirus specifically cleaves G3BP to facilitate translation of viral mRNAs⁸⁰. Foot and Mouth Disease virus cleaves G3BP and inhibits SG formation⁸¹. Several other viruses including Semliki Forest virus, vaccinia virus, cardioviruses, dengue virus, Japanese encephalitis virus, and Ebola virus sequester G3BP to inhibit SG formation and to promote virus replication^{82,83,84,85,86,87,88}. Amongst *Coronaviridae*, MERS virus employs its dsRNA-binding 4a protein to inhibit

SG formation and promote viral replication⁸⁹ whereas avian infectious bronchitis virus (IBV) infection prevents the formation of SGs via yet-unknown mechanisms⁹⁰.

N protein also interacts with subunits of the broad-spectrum Casein Kinase 2 (CK2) complex (CSNK2A2 and CSNK2B). The N protein of other *Coronaviridae* has been shown to be phosphorylated at sites predicted to be substrates for CK2⁹¹, and there is a NetPhos predicted CK2 phosphorylation site in N. CK2 has previously been shown to inhibit granule formation or promote granule disassembly in a G3BP phosphorylation-dependent manner⁹². CK2 inhibition sequesters SARS-CoV N protein in the nucleus, away from the G3BP subunits of SGs⁷⁶. It is possible that the SARS-CoV-2 N protein inhibits the formation of cellular SGs, potentially by mediating the phosphorylation of G3BP by CK2. siRNA-mediated knockdown of either CSNK2A2 or CSNK2B reduced viral replication of SARS-CoV, indicating that CK2 may have a proviral function⁹³. These findings suggest that the CK2 inhibitor Silmitasertib (also known as CX-4945) could be effective in slowing SARS-CoV-2 replication. Silmitasertib is a sub-nanomolar inhibitor of CK2 currently in phase II trials for indications such as multiple myeloma or metastatic basal cell carcinoma⁹⁴. Potential antiviral effects of Silmitasertib via inhibition of CK2 and subsequent increase in the formation of antiviral SGs merits further investigation.

UPF1 is an RNA helicase functioning in the nonsense-mediated mRNA decay (NMD) pathway. In common with other virus species in the order *Nidovirales*, SARS-CoV-2 produces a nested set of sgRNAs sharing the same 3'UTR but are different in the number of ORFs contained⁹⁵. In principle, only the first ORF in the 5' end is translated. Thus, many of the subgenomic mRNAs have long 3' UTRs and are potentially targeted for NMD. A previous study showed that Murine Hepatitis Virus (MHV) mRNAs were subjected to NMD⁷⁵. Transfection of plasmids containing MHV N protein had an NMD inhibitory function and prevented MHV mRNA from rapid decay. Based on the observation that SARS-CoV-2 shares a similar feature of nested sgRNAs, we suspect it is also subjected to NMD and that its N protein may have similar NMD inhibitory function. It is also possible that UPF1 is involved in the programmed ribosomal frameshifting during SARS-CoV-2 translation given its discovered function in suppressing nonsense mutations⁹⁶.

LARP1 is an RNA binding protein, which is known to regulate protein synthesis as well as modulate mTOR pathway⁹⁷. LARP interacts with actively translating ribosomes via another N protein interactor, PABPC1, also a regulator of mTOR pathway⁹⁸. Activation of the mTOR pathway is advantageous for a broad spectrum of virus species, as it counteracts the host cell response by inhibiting autophagy and apoptosis⁹⁹. Therefore inhibition of mTOR activity has been proven useful to counteract viral infection and replication. For instance, Sirolimus (rapamycin) has been used to reduce MERS-CoV infection^{100,101} and alleviate H1N1 pneumonia and acute respiratory failure¹⁰². Therefore, mTOR pathway represents a potential therapeutic target for SARS-CoV-2 as well.

PABPC1 and PABPC4 are poly-A binding proteins, involved in both nuclear and cytoplasmic mRNA processing¹⁰³. They are known to shuttle to the nucleus via interaction with Nsp1 proteins from different viruses (including SARS-CoV), down-regulating gene expression via hyperadenylation and nuclear retention of mRNAs¹⁰⁴. In a complex with LARP1 and RyDEN, PABPC1/4 have been shown to promote DENV replication¹⁰⁵. PABPC4 has been described as a potential biomarker for primary lung adenocarcinoma¹⁰⁶.

Moloney Leukemia Virus 10 Protein (MOV10) is a host cytoplasmic 5' to 3' RNA helicase that interacts with UPF1 and binds 3'UTRs¹⁰⁷. It exhibits antiviral functions independent of its helicase activity towards PRRSV, DENV and Influenza A viruses, through IFN stimulation^{108,109}. Additionally, MOV10 exhibits P-body dependent antiviral activity by binding to nucleocapsid proteins of other viruses and preventing their nuclear localization^{110,111}.

SARS-CoV Non-Structural Proteins

SARS-CoV-2 Nsp1

In SARS-CoV, Nsp1 is likely dispensable for CoV RNA synthesis¹¹², but may play specific roles in the interaction of the virus with the innate immune response via directly antagonizing IFN induction¹¹³. In addition, overexpression of Nsp1 in lung epithelial cells (A549) increases the production of the chemokines CCL5, CXCL10 and CCL3 30-200-fold compared with mock-transfected cells or cells expressing Nsp5, suggesting that Nsp1 may contribute to the inflammatory phenotype of SARS-CoV and SARS-CoV-2 pathology¹¹⁴.

In our interactome, SARS-CoV-2 Nsp1 interacts with six host proteins. Four of these host proteins form the DNA polymerase alpha complex (POLA1, POLA2, PRIM1, and PRIM2). The DNA polymerase alpha complex was recently shown to modulate the type I interferons through cytosolic RNA:DNA synthesis¹¹⁵, raising a possibility that SARS-CoV-2 Nsp1 may bind to the DNA polymerase alpha complex in the cytosol and modulate its activity to antagonize the innate immune response. Alternatively but not exclusively, Nsp1 may also interfere with the canonical DNA replication function of the complex, causing DNA replication stress and ATR activation¹¹⁶. Along this line, treatment with ATR inhibitors significantly reduced viral RNA replication for avian infectious bronchitis virus (IBV) coronavirus, but it remains elusive how ATR activation promotes viral replication¹¹⁶. One other protein interacting with Nsp1 is PKP2, an Influenza A virus PB1 protein interactor that restricts Influenza A virus replication¹¹⁷ and therefore may also act as a restriction factor for SARS-CoV-2.

SARS-CoV-2 Nsp2

Nsp2 is highly variable among *Coronaviridae*, and while not essential for viral replication, deletion of Nsp2 in SARS-CoV diminishes viral growth and RNA synthesis^{118,119}. Nsp2 is translated as part of a single protein along with Nsp3 and may serve as an adaptor for Nsp3¹¹⁹. The SARS-CoV-2 Nsp2 protein has a stabilizing mutation in the endosome-associated-protein-like domain, and is thought to have implication in virus pathogenesis¹²⁰.

In this study, we identified seven high-confidence host protein interactions of Nsp2. Among these are two endosomal proteins, FKBP15 and WASHC4, that are known to regulate endosome transport¹²¹⁻¹²³. It is known that SARS-CoV is translocated to endosomes after cell entry³ and interaction with endosomal transport proteins may reflect a mechanism by which SARS-CoV-2 regulates this process. Nsp2 also interacts with translational repressors EIF4E2 and GIGYF2, and disruption of the EIF4E2-GIGYF2 complex leads to increased translation¹²⁴. Nsp2 may bind to EIF4E2 and GIGYF2 to modulate translation of host and viral mRNAs. Nsp2 was also shown to interact with the acyl-CoA synthetase SLC27A2, a previously described Dengue virus restriction factor¹²⁵. Nsp2 interacts with chaperone protein and guanine nucleotide exchange factor RAP1GDS1/SmgGDS, which is known to promote the malignant phenotype in non-small cell lung carcinoma (NSCLC) by regulating cell proliferation, migration, and NF-κB transcriptional activity¹²⁶. Activation of NF-κB in pulmonary epithelial cells induces transcription of a variety of proteins that protect the cells from the inflammatory process, suggesting that RAP1GDS1/SmgGDS could be implicated in lung inflammation during SARS-CoV-2 infection¹²⁷.

SARS-CoV-2 Nsp4

Nsp4 is a non-structural, transmembrane protein that forms a complex with Nsp3 and Nsp6, and is involved in double membrane vesicle (DMV) formation¹²⁸. Together, Nsp3, Nsp4, and Nsp6 are predicted to nucleate and anchor viral replication complexes on DMVs in the cytoplasm¹²⁹. Nsp4 is likely essential, as loss of the Nsp3-Nsp4 interaction eliminated viral replication using an infectious cDNA clone and replicon system of SARS-CoV¹³⁰.

We identified eight high-confidence host protein interactions with Nsp4. One such interactor NUP210, is a nuclear pore membrane glycoprotein involved in nuclear pore assembly, fusion, spacing and structural integrity^{131,132}. Although RNA viruses replicate in the cytoplasm, they still target the nuclear pore complex (NPC) to improve viral replication and transmission. Certain RNA viruses such as rhinoviruses and polioviruses inhibit active nuclear import by the proteolytic degradation of NUPs, such as NUP62, NUP153, and NUP98, and thereby weaken the host cell's immune response against the virus¹³³. SARS-CoV-1 protein nsp1 is known to disrupt the localization of NUP93 from the NPC deregulating host cell functions. SARS-CoV-2 Nsp4 interactor Nup210 identified in this study forms the ring structure of the NPC along with NUP93¹³⁴, and it is therefore likely that Nsp4-NUP210 interaction also affects SARS-CoV-2 pathology by NPC deregulation.

Nsp4 also interacts with Insulin degrading enzyme (IDE), which is involved in intercellular signaling through the cellular breakdown of diverse signaling peptides^{135,136}. IDE is also involved in antigen processing through the production of an antigenic peptide that is presented to cytotoxic T lymphocytes by MHC class I¹³⁷. Interestingly, IDE acts as an entry receptor for varicella-zoster virus (VZV), where VZV glycoprotein E interacts with IDE through its extracellular domain^{138,139}.

Viruses are known to target components of mitochondrial proteins to enhance their own replication¹⁴⁰. Nsp4 interacts with various mitochondrial proteins linked to transport, including TIMM10, TIMM10B, TIMM9 and TIMM29, members of the TIMM22 complex. TIMM22 facilitates the import and insertion of multi-pass transmembrane proteins into the mitochondrial inner membrane, with TIMM9 additionally involved in protein homodimerization and chaperone binding¹⁴¹. TIMM10 and TIMM29 are known host dependency factors in Zika virus infection^{142,143}. Another Nsp4 interactor DNAJC11 is also involved in mitochondrial function. DNAJC11 is required for mitochondrial inner membrane assembly and functions through involvement with the MICOS complex and the MOM sorting assembly machinery complex¹⁴⁴. Other Nsp4 interactor ALG11, a mannosyltransferase catalyzing oligosaccharide linkage^{145,146} has no known links to viral infections.

SARS-CoV-2 Nsp5/Nsp5_C145

Nsp5 encodes the coronavirus main protease (M^{pro}) responsible for cleaving itself and the other subunits from the polyproteins Orf1a and Orf1ab¹⁴⁷. As these proteins include the replicase machinery, Nsp5 M^{pro} is essential for all coronaviruses, and indeed is functionally and structurally conserved throughout the order *Nidovirales*¹⁴⁸. Many protease PPIs are transient as substrates are further cleaved and processed, making it harder to capture and identify interacting proteins by native affinity purification mass spectrometry (AP-MS) experiments. To address this issue, we carry out pull-down experiments on both wild-type and catalytically inactive protease proteins. Here, we have used both the wild-type and C145A mutant Nsp5 proteins as baits. We have called C145A catalytically inactive based on sequence alignments, as the catalytic residues of SARS-CoV align to H41 and C145 in our SARS-CoV-2 construct. However, we have not yet tested its activity *in vitro* or *in vivo*, and suggest researchers keep this in mind when examining the functional significance of high confidence host interactors.

Catalytically active Nsp5, but not the C145A mutant, interacts with Histone Deacetylase 2 (HDAC2). HDACs deacetylate lysines on the N-termini of histones creating epigenetic markers for transcriptionally silent genes (HDACs involved in disease reviewed in ¹⁴⁹). HDAC2 is a class I deacetylase that regulates gene expression important for several processes including hormone signaling, cytokine signaling, embryonic development, as well as heart, muscle, and neuronal function¹⁴⁹. Specifically, HDAC2 is known to regulate genes involved in inflammatory response, especially in the pulmonary context where low HDAC2 expression contributes to increased disease severity and glucocorticoid resistance in patients with asthma¹⁵⁰ and chronic obstructive pulmonary disease¹⁵¹. Cigarette smoke is also shown to down-regulate HDAC2 expression levels through ubiquitylation and proteasomal degradation, with overexpression of the deubiquitinase USP17 restoring HDAC2 levels¹⁵². In addition to its connection to lung diseases, HDAC2 can act as either a restriction or dependency factor for various viruses. For example, during infection with Influenza A virus, HDAC2 is targeted for proteasomal degradation, and it was shown that siRNA knockdown of HDAC2 increases IAV replication and growth¹⁵³. In contrast, during human respiratory syncytial virus (RSV) infection HDAC2 expression is increased and treatment with HDAC inhibitors limited RSV replication and alleviated airway inflammation in mouse models of the disease, indicating HDAC2 as a dependency factor for RSV¹⁵⁴. General HDAC inhibitors have also shown antiviral activity against human adenovirus where treatment repressed viral gene expression and viral replication¹⁵⁵. While HDAC inhibitors represent a potential strategy for treatment of viral infection, more research is required to determine HDAC2 role in SARS-CoV-2 infection. It is clear that HDAC2 has different roles during infection with different pathogens, and its role in inflammation response can be modulated on an individual patient level depending on their history and genetic background.

Catalytically inactive Nsp5_C145A interacts with two host proteins: TRMT1 and GPX1. TRMT1 is a tRNA methylase that has recently been linked to intellectual disorders¹⁵⁶ but has no other known functions or links to viral infections. The strongest interactor of the C145A Nsp5 mutant is GPX1, a glutathione peroxidase antioxidant enzyme that catalyzes the reduction of peroxides like H_2O_2 to less harmful substances like water¹⁵⁷. GPX1 is a complex regulator that not only helps control the redox state of the cell, but also mediates a number of cellular processes including mitochondrial function and apoptosis¹⁵⁷. The enzyme is transcriptionally regulated by a variety of factors including NFkB and has been linked to inflammation^{158,157}. In addition, GPX1 has been linked to a number of human diseases that increase the risk of comorbidity with SARS-CoV-2 infection including cancer, diabetes, and cardiovascular disease (reviewed in detail in ¹⁵⁷). Interestingly, there is evidence suggesting GPX1 is regulated by oestrogens contributing to a higher expression of GPX1 in females over males¹⁵⁹. Taken together, it's possible that GPX1 could play a role in the severity of disease in different individuals, though more research is needed to clarify its interaction with Nsp5 as a substrate or interactor, and determine its role during infection. As we learn more about COVID-19 and individual risk factors, we can more accurately assess the clinical relevance of specific interactors.

Given we typically utilize catalytically dead mutants to stabilize protease interactions with substrates, it is not surprising that we identify more interactors for the C145A Nsp5 mutant. The fact that HDAC2 specifically interacts with wild-type Nsp5 may suggest that this interaction is unusually stable, and perhaps that the C145A mutation may interfere with this interaction.

SARS-CoV-2 Nsp6

Nsp6 complexes with two other transmembrane proteins, Nsp3 and Nsp4, to form double membrane vesicles (DMVs), anchoring viral replication complexes inside^{160–162}. Chemical inhibition and targeted mutagenesis studies have shown that these DMVs are crucial to viral replication and are formed early after viral entry into the cell^{163–166}. The most well-characterized function of Nsp6 is limiting autophagosome expansion, which likely benefits the virus by preventing its components from being sent to the lysosome for

degradation¹⁶⁷. In the context of IBV it was shown that the Nsp6-mediated generation of autophagosomes can be induced by chemical inhibition of the mTOR pathway¹⁶⁷. While this behavior has been shown for SARS-CoV Nsp6 and related viruses¹⁶⁷, its exact mechanism and whether SARS-CoV-2 Nsp6 will function similarly remains uncertain.

In this study, we identified four high confidence interactors of SARS-CoV-2 Nsp6 (ATP5MG, ATP6AP1, ATP13A3, and SIGMAR1). Notably, three out of four of these prey proteins are subunits of different ATP synthases, suggesting that Nsp6 may be manipulating the metabolic program of the cell through ATP synthases of different organelles. As viral infection requires a new surplus of energy, it is unsurprising that viruses hijack host metabolic machinery to carry out a wide variety of cellular functions^{168–171}. Specifically, ATPases are crucial for budding by multiple viruses, including HIV-1, Influenza A virus (IAV), and Ebola^{172–174}. One such target is vacuolar ATPases (vATPases), which are responsible for modulating endo-lysosomal acidification and thus a number of processes like membrane trafficking and protein degradation¹⁷⁵. In the context of IAV infection, vATPase ATP6V1A was identified as a key host dependency factor for viral replication across multiple RNAi knockdown screens^{48,176}. Influenza A virus is believed to leverage this interaction to lower the pH of the lysosomal environment to accelerate the process of viral uncoating in the early stages of viral infection^{177,178}. ATP6AP1, a subunit of the vATPase protein pump and high-confidence physical interactor of Nsp6 in this study, has also been described as a host factor for a number of viruses, including IAV, West Nile virus (WNV), and DENV^{47,48,176,179,180}. Further, dysregulation of ATP6AP1 leads to impaired vesicle acidification and intracytoplasmic granules, resulting in multiple clinical pathologies including an immunodeficiency syndrome and granular cell tumors^{181,182}. Given the known role of ATP6V1A as an IAV host factor, the mechanism of SARS-CoV-2 Nsp6 could leverage its interaction with ATP6AP1 similarly to speed replication in DMV complexes.

ATP13A3 is a poorly understood cation-transporting P-type subunit of the mitochondrial ATPase that has recently been implicated in both host innate immunity to pathogens and pulmonary pathologies. During Herpes simplex virus (HSV)-1 infection, ATP13A3 was shown to be down regulated at the plasma membrane of infected cells, but its expression could be rescued by the deletion of a specific HSV-1 protein pUL56, suggesting that it may be specifically targeted as a host factor in the context of some viral infections¹⁸³. Furthermore, ATP13A3 was identified in the plasma membrane of primary CD4+ T cells infected with Vpu-deficient and Nef-deficient HIV-1, and was differentially expressed in CD8+ T cells during IAV infection¹⁸⁴. More broadly, ATP13A3 expression was found to be a significant variable in determining interindividual response to lipopolysaccharide sensing in human dendritic cells¹⁸⁵. It was recently reported that loss of function of *ATP13A3* results in disruption of polyamine homeostasis in pulmonary arterial endothelial cells, leading to endothelial dysfunction and ultimately pulmonary arterial hypertension^{186,187}. Notably, ATP13A3 is also amenable to direct drug targeting.

Sigma Receptor 1 (SIGMAR1) is an ER chaperone protein that modulates calcium signaling through its interaction with the IP3 receptor (IP3R-3) and adaptor protein ankyrin B (ANK2)¹⁸⁸. While SIGMAR1 has no known role in viral pathogenesis or pulmonary pathologies, it is targeted by a number of potential SARS-CoV-2 drug candidates highlighted in this study, including approved drugs chloroquine and haloperidol, as well as other preclinical drugs such as RS-PPCC and PB28.

SARS-CoV RNA polymerase: Nsp7, Nsp8 and Nsp12

SARS-CoV Nsp7, Nsp8 and Nsp12 proteins form the RNA-dependent RNA polymerase (RdRp)^{189,190}. Eight Nsp7 proteins bind stoichiometrically to eight Nsp8 proteins to form a hexadecameric barrel-shaped structure thought to surround double-stranded RNA¹⁹¹. The catalytic activity of the Nsp7-Nsp8 complex is much

weaker than that of Nsp12, though this macromolecular complex is primer-independent and RNA products of Nsp7-Nsp8 complex are typically ≤ 6 bases. Thus it is believed that the Nsp7-Nsp8 complex functions as a primase, generating short RNA primers for Nsp12's more processive, primer-dependent RNA polymerase activity^{192,193}.

Nsp12 encodes a canonical, primer-dependent RdRp domain in its C-terminus^{192,194}, and is essential for viral RNA synthesis. A Nsp7-Nsp8 heterodimer binds to and stabilizes loops in the polymerase domain of Nsp12, and is thought to facilitate interactions between Nsp12 and RNA synthesis/processing machinery¹⁹⁰. Nsp12 contains an unusually large N-terminal extension with a conserved, essential nucleotidyltransferase domain¹⁹⁵ and protein interface domain that binds a second Nsp8¹⁹⁰. The full functions of these domains in virus replication or fitness and host biology are not known. Nsp12 may have additional roles outside of the RdRp complex mediated through interactions with the uncharacterized N-terminus^{190,192,195} and potentially other virus proteins including Nsp5, Nsp8 and Nsp9¹⁹⁶.

The structure of SARS-CoV-2 Nsp7/Nsp8/Nsp12 complex closely mimics that of SARS-CoV¹⁹⁷, and the complex members bear 98.8%, 97.5%, and 96.4% genetic similarity at the codon level respectively, suggesting that function will be conserved.

SARS-CoV-2 Nsp7

We identified 32 high-confidence host protein interactions with Nsp7. These interactors include multiple Rab GTPases with various roles in exocytic and endocytic membrane trafficking, mitochondrial proteins such as cytochromes, and several other factors that were previously identified as host interactors of other viruses.

Rab GTPases act as molecular switches that regulate dynamic networks and complexes involved in vesicle trafficking¹⁹⁸. Among the Rab GTPases identified as Nsp7 interactors are Rab14, Rab1a, Rab7a, Rab5c and Rab8a. Rab14 regulates the membrane recycling pathway from endosomes to the plasma membrane which is required for ADAM protease trafficking and regulation of cell-cell junctions¹⁹⁹. Rab14 was previously characterized to be required for HIV-1 envelope glycoprotein particle incorporation²⁰⁰. Rab1a regulates vesicular protein transport from the ER to the Golgi and was identified to be required for production of extracellular enveloped virions of vaccinia virus (VACV)²⁰¹, as well as for particle assembly of classical swine fever virus (CSFV)²⁰². Rab7a and Rab5c are essential for Hepatitis B virus (HBV) infection²⁰³ with Rab7a implicated in promoting virus entry in early stages²⁰³ and restricting exocytic virion release in late stages²⁰⁴. Rab8a was shown to be an important regulator of HIV-1 trafficking to the virological synapses²⁰⁵ which enables delivery of the virus to CD4+ T-cells.

The ACSL3-LPIAT1 fusion protein is a known cancer gene and ACSL3 is highly expressed in human lung cancer²⁰⁶. ACSL3 induction results in prostaglandin production by increasing acyl-CoA synthesis, which is required for the maintenance of non-small cell lung cancer²⁰⁷ and mutant KRAS lung cancer tumorigenesis *in vivo*²⁰⁶. Besides the function of ACSL3 in lung cancer, it is known to interact with several viral proteins and is required for poliovirus replication^{208–210}.

MOGS is a Nsp7 interactor that has previously been shown to interact with various pathogen proteins, including HIV Vpr and gp120, and *Mycobacterium tuberculosis* (Mtb) Ppe11^{209,211,212}. This gene is well-characterized in the context of a congenital disorder of glycosylation. Interestingly, patients with genetic defects in MOGS manifest decreased susceptibility to viral infections²¹³, providing a potential target for broad spectrum therapy of viral infections.

SARS-CoV-2 Nsp8

We find that SARS-CoV-2 Nsp8 interacts with 23 host proteins that are enriched for a number of functions and complexes, including the exosome complex, ribosome biogenesis, and regulation of protein translation. In our dataset, we identify four (EXOSC2, EXOSC3, EXOSC5, and EXOSC8) members of the exosome complex as high confidence Nsp8 interactors, with an additional six subunits just narrowly missing our stringent threshold (EXOSC1, EXOSC4, EXOSC6, EXOSC7, EXOSC9, and EXOSC10). The eukaryotic RNA exosome complex plays a critical role in RNA processing, degradation, and quality control. It consists of 10-11 subunits and functions to remove unstable transcripts including cryptic unstable transcripts, antisense RNA, and unstable bidirectional transcripts/enhancer RNAs^{215,216,217}. Its barrel-shaped catalytically inactive core is made up of nine subunits that form the 'cap' and 'ring' subcomplexes, all of which are identified in our Nsp8 pull-down to varying degrees of confidence. In addition, we identify (below our threshold) one of the putative catalytic subunits, EXOSC10^{215,217}. In drosophila and human cells the RNA exosome complex broadly restricts a number of viral infections including vesicular stomatitis virus (VSV), Sindbis virus (SINV), and Rift Valley fever virus (RVFV)²¹⁴. However, other studies have shown that the RNA exosome can act as a dependency factor. For instance, H1N1 influenza A virus (IAV) polymerase PA subunit binds to subunits of the exosome, which were then shown to promote viral ribogenesis and IAV growth²¹⁵. Given that IAV PA and SARS-CoV-2 Nsp8 are part of viral polymerase complexes, it is tempting to speculate host cell RNA exosome complexes function as dependency factors in a similar manner during both IAV and SARS-CoV-2 infection.

LARP7, a member of the 7SK snRNP RNA complex, interacts with WNV, ZIKV capsid, and HIV-1 Tat in such a way as to compete with viral proteins for the transcriptional elongator pTEFb. The interaction of LARP7 and coronavirus primase points to a mechanism by which the virus promotes elongation of its vRNA.

Interestingly, we find that Nsp8 interacts with proteins just under our scoring threshold found to be differentially regulated or expressed during acute lung disease secondary to pathogen infection, including CEBPZ, DAP3, NKRF and ZN512. NKRF for example is upregulated in the circulating monocytes and alveolar macrophages of patients with active pulmonary TB, and NKRF serves as an endogenous repressor for IP-10 and IL-8 synthesis to hinder host from robust response to Mtb infection^{216,217}.

SARS-CoV-2 Nsp12

In our study, SARS-CoV-2 Nsp12 interacts with 20 high confidence host proteins. Consistent with Nsp12 RdRp activity, eight host protein interactors are RNA binding factors involved at multiple steps of RNA processing and regulation. These host proteins could facilitate long-range RNA interactions that occur during genome replication and discontinuous transcription²¹⁸, or mediate viral protein translation. Of note, Nsp12 interacts with three proteins involved in pre-mRNA splicing: A-kinase anchor protein 8 (AKAP8)²¹⁹; and spliceosome components pre-mRNA-splicing factor SLU7 (SLU7)²²⁰ and peptidyl-prolyl cis-trans isomerase-like 3 (PPIL3). Notably, siRNA knockdown of SLU7 inhibits early stages of HIV-1 replication²²¹. Nsp12 also interacts with La-related protein 4B (LARP4B), a cytoplasmic RNA binding protein that promotes mRNA translation and interacts with PABPC1 and RACK1 kinase, potentially connecting 3' mRNA factors with translation machinery²²². PABP, a 3' poly(A) tail binding protein, is a cis acting element on coronavirus RNA that is essential for bovine coronavirus replication^{223,224}. Nsp12 could be targeting LARP4B as a bridge to recruit members of the PABP family for efficient RNA replication and translation. Nsp12 interacts with two RNA binding proteins that localize to and nucleate formation of stress granules: ubiquitin-associated protein 2 (UBAP2) and ubiquitin-associated protein 2-like (UBAP2L)²²⁵. Interestingly, UBAP2L interaction is also identified in SARS-CoV-2 Nsp9 pull-downs, although it falls below our scoring threshold. Coronavirus species

mouse hepatitis virus (MHV) Nsp9 has been shown to interact with Nsp12¹⁹⁶. This may suggest a potential role for SARS-CoV-2 Nsp12 and Nsp9 in coordinating host and/or viral RNA regulation (see also Nsp9), as stress granules are sites of RNA storage and intermediate stages between translation and mRNA decay²²⁶. Additionally, since MERS-CoV has been shown to inhibit stress granule formation to promote viral replication⁸⁹, SARS-CoV-2 Nsp12 may interact with and sequester UBAP2L as a potential mechanism to inhibit stress granule nucleation and promote virus replication (see also N protein for role of stress granules).

Although coronavirus genome replication and transcription occurs in the cytoplasm, SARS-CoV-2 Nsp12 interacts with five nuclear DNA-related factors: transcription factors CREB-regulated transcription coactivator 3 (CRTC3), transcription factor 12 (TCF12) and zinc finger protein 318 (ZNF318); chromatin factor SBNO1; and AKAP8 which regulates histone methylation and gene expression²²⁷. Nsp12 shows DNA-dependent activity and can synthesize nucleotides from a DNA template *in vitro*^{192,193}. Nsp12 may have novel roles in chromatin and transcription regulation, although the advantage to the virus is unclear.

Nsp12 also interacts with host proteins that regulate inflammatory signaling and apoptotic pathways. One of the top Nsp12 interactors is receptor-interacting serine/threonine-protein kinase 1 (RIPK1). As an active regulatory kinase, RIPK1 triggers cell death by apoptosis or necroptosis; as a scaffold independent of its kinase activity, RIPK1 regulates inflammatory signaling and inhibits cell death^{228–230}. Many diverse viral and bacterial proteins interact with and modify RIPK1 to modulate host defense pathways, cytokine signaling and host cell death^{231–233}. For example, HIV-1 protease cleaves and inactivates RIPK1, which then impairs host defense pathways²³³. Importantly, RIPK1 is a druggable target, and many inhibitors are being tested as anti-inflammatory treatments for neurodegenerative diseases and cancer²³⁴. However, given the two faces of RIPK1 regulating apoptosis for pathogen clearance²³² or stimulating cytokine production as part of the inflammatory response²³⁰, the usefulness of drugs targeting RIPK1 as a treatment for SARS-CoV-2 will depend on future research identifying which pathway is engaged by Nsp12. In addition, Nsp12 interactor AKAP8 binds and shuttles caspase 3 to the nucleus as part of caspase-mediated proteolysis and apoptosis²³⁵. These interactions suggest additional novel roles for Nsp12 outside of canonical RdRp activity in regulating host inflammation and cell death, perhaps mediated through the large uncharacterized N-terminal domain of Nsp12.

SARS-CoV-2 Nsp9

In SARS-CoV and related coronaviruses, Nsp9 is an essential non-structural protein that binds to RNA and DNA, with a preference for single-stranded RNA^{236–239}. The function of Nsp9 is not well annotated, though in SARS-CoV it is believed to interact with Nsp8 and the viral replication complex (Nsp7, Nsp8, and Nsp12)^{239,240}. In our hands, SARS-CoV-2 Nsp9 interacts with 16 high confidence host proteins, with many having been shown in previous studies to regulate nuclear transport, transcription, and mRNA degradation in response to various viral infections. Unexpectedly, Nsp9 also demonstrates some strong interactions with extracellular matrix proteins involved in elastin formation, lung development, and lung injury and repair.

Nsp9 interacts with Nup62, Nup58, and Nup54, the three components of the Nup62 subcomplex which forms the channel of the nuclear pore complex (NPC)^{241–245}. In addition, Nsp9 interacts with nucleoporins on the cytoplasmic side of the nuclear pore complex (i.e. NUP214 and NUP88) but none from the nucleoplasmic side²⁴⁶ indicating a cytoplasmic role of Nsp9 at the NPC²⁴⁶. Many viruses have been shown to exploit cellular nuclear transport machinery to block host-related transport or promote viral transport in a manner beneficial to the virus^{247–260}. Nup62 in particular interacts with a number of viral proteins including Epstein Barr Virus (EBV) BGLF4 protein kinase²⁶¹, human papillomavirus (HPV)16 and HPV8 E7 proteins^{262,263}, and HIV-1 IN²⁶⁴. Several positive-sense single strand RNA viruses (e.g. poliovirus, enterovirus (EV)71, rhinovirus, and cardioviruses) target Nup62 either through viral protease-directed cleavage^{250–256} or induced hyperphosphorylation^{257–259} in order to limit nuclear transport. Vaccinia virus, a DNA poxvirus, replicates in

cytoplasmic virus factories that recruit G3BP1 and Nup62²⁶⁰. Nup54 interacts with Influenza A virus polymerase and was shown to be important for virus replication and transcriptional activity²⁶⁵. In addition to the nucleoporins, Nsp9 interacts with MIB1, a RING-type E3 ubiquitin ligase shown to act as a dependency factor during adenovirus infection²⁶⁶. Adenoviruses are non-enveloped DNA viruses that utilize the NPC to dock and deliver genomic cargo into the nucleus, and it has been shown that MIB1 mediates the delivery of viral DNA through the NPC²⁶⁶.

In addition to nuclear transport machinery, Nsp9 interacts with three transcription regulators, Nek9, DCAF7 and eIF4H. Nek9 is a serine/threonine kinase that regulates mitotic progression²⁶⁷. It has been shown as an adenovirus dependency factor promoting viral growth, interacting with E1A protein to silence the expression of certain host genes, and was demonstrated to colocalize at adenovirus replication centers²⁶⁸. DCAF7 is a potential substrate adaptor for CUL4-DDB1 E3 ligase²⁶⁹ that has also been shown to interact with adenovirus E1A protein to suppress innate immune response and depresses IFN stimulated genes (ISGs)²⁷⁰. In combination with eIF4A, eIF4H was shown to interact with HSV virus host shut off proteins, with eIF4A shown to help degrade mRNA and switch from host to viral gene expression^{271–274}. Taken together, these interactors suggest a potential function of Nsp9 in the inhibition of host mRNA expression potentially to limit the expression of ISGs.

Somewhat unexpectedly, Nsp9 interacts with several proteins involved in lung development, lung injury and repair, and lung cancers. The most abundant interactor in the Nsp9 interactome is Fibrillin-2 (FBN2), an extracellular matrix glycoprotein involved in elastin fiber and respiratory organ development²⁷⁵. Additional related interactors were identified, albeit with lower abundance. Fibrillin-1 (FBN1) and Fibulin-5 (FBLN5) are both extracellular matrix proteins, also involved in elastin fibre development. Mutations in FBN1 cause Marfan syndrome (MFS), a connective tissue disease that can result in early morbidity and mortality mainly caused by aortic aneurysm and rupture, though additional clinical manifestations include lung complications²⁷⁶. FBLN5 is indicated to serve a role during lung injury and repair, is frequently silenced in lung cancer, and has been shown to suppress cancer cell invasion^{277–282}.

The Nsp9 interactome reveals a potential function for the protein not only in viral replication, but potentially in regulating nuclear transport, though it is unclear if this regulation would block host cell transport or promote viral transport. In addition, the interaction with transcription regulators indicates a potential role in inhibition of host cell transcription and potentially host shut off. And finally, the unexpected interaction with fibrillins-1 and -2, and fibulin-5 could implicate Nsp9 as a potential complicating factor during disease pathogenesis, and may point to additional molecular reasons underlying SARS-CoV-2 complications in the lungs.

SARS-CoV-2 Nsp10

Nsp10 contains two Zn-finger motifs, binds nucleic acids non-specifically, and has been implicated in minus-strand RNA synthesis, thus performing an essential role in viral replication^{283,284}. A unique feature for SARS-CoV is that Nsp16 requires Nsp10 as a stimulatory factor to execute its methyltransferase activity²⁸⁵. Nsp10 on its own forms a dodecameric homomeric complex²⁸⁶ which excludes its interaction interfaces with Nsp14 and Nsp16, meaning that Nsp10 could have at least two different functional quaternary structures.

In our map, we identify five high-confidence host protein interactions with Nsp10, including two subunits of the clathrin adaptor protein complex 2 (AP-2), AP2A2 and AP2M1. This interaction is reminiscent of Nef proteins from Human and Simian Immunodeficiency Viruses (HIV and SIV) which are shown to bind the AP-2 complex through a canonical AP-2 recognition acidic dileucine motif ([RQED]XXXL[LIV])²⁸⁷. HIV and SIV are thought to use the interaction between Nef and AP-2 to hijack the clathrin machinery and endocytose host proteins such as CD4 and MHC-I²⁸⁸. Interestingly, coronavirus Nsp10 proteins do not appear to contain the

canonical AP-2 binding motifs (either the acidic dileucine motif [RQED]XXXL[LIV] for binding AP2A2 nor YXXΦ for binding AP2M1).

In addition to the AP-2 complex, Nsp10 interacts with GFER, ERGIC1, and GRPEL1. GFER was identified as a host dependency factor for West Nile and Dengue virus infections⁴⁷. ERGIC1 and GRPEL do not have known roles in viral pathogenesis. ERGIC1 plays a possible role in transport between endoplasmic reticulum and Golgi²⁸⁹; and GRPEL1 participates in the translocation of transit peptide-containing proteins from the inner membrane into the mitochondrial matrix²⁹⁰.

SARS-CoV-2 Nsp11

SARS-CoV Nsp11 encodes a short peptide that is 13 aa in length. It is not clear whether Nsp11 encodes a functional protein. In our map, we identify only one high-confidence host protein as an interactor of Nsp11. Tubulin-specific chaperone A (TBCA) is a tubulin-folding protein and is known to be involved in the early step of the tubulin folding pathway²⁹¹.

SARS-CoV Capping Enzymes: Nsp13, Nsp14 and Nsp16

The m7GpppN (N = any nucleotide) cap of mRNA promotes translation. SARS-CoV Nsp13, Nsp14, and Nsp16 encode enzymes that install cap structure onto mRNA^{292,293}. The pathway of cap synthesis on nascent viral mRNA is thought to be similar to capping of cellular mRNA, but occurs in the cytoplasm instead of the nucleus^{292,294,295}.

SARS-CoV-2 Nsp13

Nsp13 is essential for SARS-CoV viral RNA synthesis²⁹⁶, and SARS-CoV-2 Nsp13 shares 100% amino acid similarity with SARS-CoV. Nsp13 is a helicase/triphosphatase, and triphosphate cleavage initiates the first step in mRNA capping²⁹⁷. Nsp13 hydrolyzes the gamma phosphate of nascent mRNA and the resulting diphosphate is then converted to a GpppN RNA by a yet-to-be identified guanylyl transferase.

SARS-CoV-2 Nsp13 was found to interact with 40 host proteins with a broad range of cellular processes and complexes from Protein Kinase A (PKA) signalling, to the Golgi apparatus, to multiple members of microtubules and centrosomes. SARS-CoV-2 Nsp13 showed a strong interaction with Giantin (GOLGB1), which has previously been shown to interact with HSV-1 UL37²⁹⁸ and the Tick-borne encephalitis virus (TBEV) replicon²⁹⁹. Interestingly, in macaque experiments, expression of the HIV protein Nef resulted in increased GOLGB1 levels and led to Golgi disruption and specific pulmonary vasculopathies³⁰⁰.

Unexpectedly, we found that SARS-CoV-2 Nsp13 pulled down both the regulatory (PRKAR2A and PRKAR2B) and the catalytic (PRKACA) subunits of PKA as well as the A-kinase anchoring protein AKAP9 and the phosphodiesterase interacting protein PDE4DIP. AKAP9 (also called AKAP450) is a large scaffolding protein that localizes to the Golgi apparatus and centrosomes^{301,302} where it assembles multiple signaling proteins (e.g., PKA and PDE4D) that control microtubule organization³⁰³, polarized secretion³⁰⁴, Golgi morphology^{304,305}, ciliogenesis³⁰⁶, directional cell migration³⁰⁷ and cell cycle progression³⁰². Importantly, the activities of PKA signaling complexes have been implicated in multiple membrane transport steps^{308–311}, suggestive of a role for SARS-CoV-2 Nsp13 in hijacking the host secretory pathway for viral benefit. Also notably, a pool of AKAP9 relocates to RNA stress granules upon treatment with arsenite where it forms a complex with G3BP and CCAR1 and regulates stress granule size and composition³¹². PKA, being a kinase, is also targetable by small molecule inhibitors or peptides that target the AKAP-PKA binding interface.

An additional SARS-CoV-2 Nsp13 interaction partner was the endosomal transport protein ERC1. Knockout of ERC1 causes a significant decrease in dengue virus replication, and interestingly a similar phenotype was observed with additional SARS-CoV-2 Nsp13 pulldown target GOLGA2³¹³. The NS3 protein of hepatitis C virus binds to ERC1 and may mediate the pathogenesis of HCV³¹⁴, and knockdown of ERC1 significantly decreases human cytomegalovirus (HCMV) viral production³¹⁵. Of particular interest, ERC1 has previously been identified as a potential drug target in dengue virus infection³¹⁶.

Lastly, Nsp13 interacts with transducin-like enhancer of split (TLE) family members TLE1, 3 and 5.

SARS-CoV-2 Nsp14

Nsp14 is a bifunctional enzyme. It encodes an exonuclease (exo) domain that corrects mutations that arise during genome replication³¹⁷. In addition, a separate domain of Nsp14 functions as a SAM dependent methyltransferase (MTase) that generates the N-7 Guanosine of the m7GpppN cap on viral mRNA. High-resolution crystal structures of SARS-CoV Nsp10/Nsp14 suggest exo and MTase activity is stimulated by Nsp10 through an allosteric mechanism³¹⁸.

There are three host factors that copurify with Nsp14 with all three being druggable targets. GLA is an alpha galactosidase implicated in Fabry Disease³¹⁹. Migalastat, a pharmacological chaperone, targets GLA through the inhibition of alpha-glucosidase and glycosylation, increasing its lysosomal activity³²⁰. SIRT5 is a mitochondrial protein linked to metabolism and aging that removes malonyl, succinyl, acetyl, and glutaryl groups on lysines of target proteins^{321–324}. Numerous compounds target SIRT5, including HDAC inhibitors³²⁵. IMPDH2 catalyzes the conversion of isosine 5' phosphate (IMP) to xanthine 5'-phosphate (XMP) which is then converted into guanine 5' monophosphate for *de novo* synthesis of guanine nucleotides³²⁶. It is tempting to speculate that the copurification of Nsp14 and IMPDH2 reflects an interplay between Nsp14 activities and purine nucleotide metabolism, though more experiments need to be done to verify this. Merimepodib, a nucleoside analog and broad spectrum antiviral, is among the compounds that target IMPDH2³²⁷.

SARS-CoV-2 Nsp15

In SARS-CoV, Nsp15 has uridine-specific endoribonuclease (endoU) activity, and is essential for viral RNA synthesis³²⁸. The Nsp15-associated endoU domain is one of the most conserved proteins among CoVs and related viruses, suggesting important functions in the viral replicative cycle. The endoUs were shown (i) to have endonucleolytic activity, (ii) to cleave 3' of pyrimidines, preferring uridine over cytidine, and (iii) to release reaction products with 2'-3'-cyclic phosphate and 5'-OH ends. Deletion of a conserved domain for this enzymatic activity led to loss of viral RNA generation measured by RT-PCR³²⁸. Nsp15 is also known to be critical for evasion of host dsRNA sensors in macrophages³²⁹. Nsp15 proteins were shown to form homohexamers composed of a dimer of trimers³³⁰.

In our map, Nsp15 interacts with three host proteins. NUFT2 mediates the import of GDP-bound RAN from the cytoplasm to the nucleus and indirectly plays a general role in cargo receptor-mediated nucleocytoplasmic transport^{331,332}. ARF6 is a GTP-binding protein involved in protein trafficking that regulates endocytic recycling and cytoskeleton remodeling^{333–337}. ARF6 has been shown to play a role in the activation of cholera toxin³³⁸. RNF41 acts as E3 ubiquitin-protein ligase and promotes TRIF-dependent production of type I interferon; it has been shown to inhibit vesicular stomatitis virus infection³³⁹.

SARS-CoV Open Reading Frames

SARS-CoV-2 Orf3a

Orf3a is the largest (274 aa) group-specific Orf in the SARS-CoV-2 genome and encodes a type IIIa integral membrane protein. It is thought to be non-essential but has multiple key functions in viral pathogenesis from mediating the trafficking of SARS-CoV Spike (S protein), to inducing apoptosis and inflammation during infection^{340–345}. Specifically, SARS-CoV Orf3a induces pro-IL-1 β expression and protein maturation through the TRAF3-dependent ubiquitination of ASC and p105, ultimately activating NF- κ B and the NLRP3 inflammasome³⁴¹. In addition, Orf3a has been shown to upregulate the secretion of fibrinogen in lung epithelial cells, a process responsible for the induction of cytokine storm, particularly in the respiratory tract³⁴⁶. Within the cell, Orf3a localizes to the perinuclear region and plasma membrane, forming punctae throughout the cytoplasm as it complexes with cellular factors^{340,347,348}. In the rER and Golgi, Orf3a has been shown to cause substantial ER stress during infection³⁴⁹. A Yxx Φ motif present in the C-terminal cytoplasmic domain of Orf3a is required for delivery to the plasma membrane, from which Orf3a is internalized and traffics through the endocytic pathway to lysosomes³⁵⁰. Orf3a can form tetramers that are proposed to act as cation-permeant ion channels³⁵¹. In this study, Orf3a pulled down eight high-confidence host protein interactors (ALG5, ARL6IP6, CLCC1, HMOX1, SUN2, TRIM59, VPS11, VPS39) with functional enrichments in autophagy (HMOX1, VPS11, VPS39) and organelle localization (HMOX1, VPS11, SUN2). VPS11 and VPS39 serve as members of the HOPS and CORVET complexes, respectively, which coordinate fusion of the lysosome with the endosome and autophagosome^{352,353}.

HMOX1 is a key enzyme in heme catabolism. This oxygenase cleaves free heme to produce iron, biliverdin and carbon monoxide, providing a cytoprotective effect to cells as excess free heme is shown to induce apoptosis³⁵⁴. In turn, heme cleavage elicits a cascade of physiological events, most notably the induction of anti-inflammatory cytokines IL-10 and IL-1RA^{355,356}. Given these features of HMOX1 and its high expression in the lungs, it has been implicated in a broad range of disease states, including diabetes, heart failure, lung carcinoma and chronic obstructive pulmonary disease (COPD)^{357,358}. In the context of infection, upregulation of HMOX1 has been shown to have a protective effect against the oxidative stress produced during various pathogen infections, including viruses like HIV, DENV, HCV, and IAV, as well as parasites and *Mycobacterium* species^{359,360}. As severe inflammatory response in the respiratory tract is a key clinical feature of SARS-CoV-2 infection, further exploration of HMOX1 could elucidate mechanisms of SARS-CoV-2 pathogenesis and a path forward for treatments.

Orf3a also strongly interacts with, CLCC1, an intracellular chloride channel that is localized to the ER and Golgi apparatus and is ubiquitously expressed in numerous tissues including the lung³⁶¹. Loss of *CLCC1* results in ER stress and disruption of the protein folding capacity of the ER, leading to misfolded protein accumulation and well-characterized retinal cell dysfunction in the clinic^{362,363}. Notably, both CLCC1 and SARS-CoV Orf3a are localized to the ER membrane^{340,361}. In addition, expression of *CLCC1* has been correlated with volume of adipose tissue in HIV-infected men³⁶⁴. RSV NS1 protein also interacts with CLCC1, though the function of the interaction is unknown³⁶⁵. Interestingly, CLCC1 also interacts with solute carrier superfamily member SLC15A3, a druggable peptide/histidine transporter and known interferon-stimulated gene that is regulated by TLR-activation and contributes to TLR4-mediated inflammation in macrophages and epithelial cells^{366,367}. While SLC15A3 was not identified as an interactor of Orf3a, we identified interactions between SARS-CoV-2 proteins and other members of the solute carrier superfamily including SLC25A17 (SARS-CoV-2 Orf9b and Orf9c) and SLC6A15 (SARS-CoV-2 Orf9c, M and Nsp6), although these interactions

fall below our scoring thresholds. This may suggest that Orf3a may play a role in modulating host innate immunity via an indirect interaction with CLCC1 as part of a larger protein complex or pathway.

In addition to HMOX1 and CLCC1, Orf3a also interacts with known viral host factors and interactors. ARL6IP6, a transmembrane protein responsible for ADP ribosylation, is known to physically interact with both HPV-31 E5 and WNV NS4b, and was differentially expressed in CD8+ T cells from HIV+ progressors on HAART^{36,368,369}. ALG5, an ER-localized glycosyltransferase involved in N-glycan biosynthesis, has been characterized as a cellular host factor for IAV replication in multiple genome-wide CRISPR-Cas9 knockout screening efforts, along with other ALG family proteins^{51,52,36,368,369}.

SARS-CoV-2 Orf3b

SARS-CoV-2 Orf3b encodes a 168 aa protein that is not well conserved (9.5% codon similarity as compared to SARS-CoV Orf3b). While Orf3b is thought to be non-essential^{1,342}, it has been shown to be an IFN antagonist in both SARS-CoV and SL-CoV (bat)^{370,371} and to subsequently be involved in pathogenesis¹. The only confident interaction partner detected for Orf3b was the mitochondrial protein STML2/STOML2. STOML2 stimulates cardiolipin biosynthesis and recruits and stabilizes prohibitin. Both STOML2 and its interactor prohibitin have been shown to be host dependency factors for EV71 neuropathogenesis³⁷². STOML2 forms large complexes with the i-AAA protease YME1L and the rhomboid protease PARL at the inner mitochondrial membrane, which regulate key proteins of the mitochondrial stress response such as PGAM5 and PINK1³⁷³. STOML2 may directly be involved in regulating T-cell-mediated immune responses by modulating T-cell receptor activation³⁷⁴.

SARS-CoV-2 Orf6

SARS-CoV-2 Orf6 encodes a 7.3kDa protein that is dispensable for virus replication, but affects viral production³⁷⁵. Orf6 functions as a type I IFN antagonist and suppresses IFN induction and IFN signalling pathways. It is localized to the ER/Golgi membrane in infected cells, where it binds to and disrupts nuclear import complex formation by tethering karyopherin alpha 2 and karyopherin beta 1 to the membrane. This results in the loss of nuclear import of the interferon signalling responsive transcription factor STAT1^{371,376} and therefore blocks the expression of STAT1-activated genes establishing an antiviral state. In SARS-CoV infected cells, Orf6 complexes with Orf9b³⁷⁷. SARS-CoV-2 Orf6 was found to interact with three host proteins: NUP98, RAE1, and MTCH1. As described in the main text, NUP98-RAE1 is an interferon-inducible mRNA nuclear export complex that is targeted by multiple viruses including VSV, IAV, KSHV, and Polio^{378,379}. Orf6 also interacts with mitochondrial carrier homologs 1 (MTCH), which is known to regulate apoptosis by modulating activity of the mitochondrial permeability transition pore³⁸⁰ and therefore could be involved in virus-induced apoptosis.

SARS-CoV-2 Orf7a

SARS-CoV Orf7 is divided into two open reading frames, designated Orf7a and Orf7b. Orf7a (also known as U122) encodes a 122-amino-acid protein and contains a compact seven-stranded β -stack similar in structure to members of the immunoglobulin superfamily³⁸¹. The role of Orf7a in apoptosis was highlighted by an increase in caspase-3 protease activity that resulted in a significant induction of apoptosis³⁸². Orf7a expression has also been shown to downregulate cyclin D3, resulting in the accumulation of retinoblastoma protein (Rb) and ultimately cell cycle arrest in G0/G1 phase³⁸³. Orf7a traffics to different locations throughout the cell, including the perinuclear region, the cytoplasm, and the plasma membrane through its type-1

transmembrane domain^{382–384}. Further, Orf7a has been shown to colocalize with ER and ER-Golgi intermediate compartment (ERGIC) markers during infection³⁸⁴. Interestingly, this cellular localization coincides with that of high-confidence protein interactors MDN1 and HEATR3, as well as a factor that was just below our MIST threshold, TNPO1. While it did not satisfy our stringent scoring criteria, TNPO1, or transportin-1, is worth mentioning due to its demonstrated role in other viral infections such as Influenza A virus, HIV-1 and Hepatitis C virus, where it plays a crucial role in mediating nuclear transport of viral proteins and protein complexes^{385–388}. HEATR3 has also been shown to activate the NF- κ B pathway via NOD-2, which has been implicated in a pro-inflammatory response during Crohn's disease^{389,390} and therefore Orf7a may target HEATR3 to modulate inflammatory response upon SARS-CoV-2 infection.

SARS-CoV-2 Orf8

Orf8 is an accessory protein and is not essential for virus replication *in vitro* and *in vivo*^{342,391}. It is one of the most rapidly evolving regions among SARS-CoV genomes and was previously shown to be a recombination hotspot^{392–394}. Pairwise comparison of amino acid sequences showed that SARS-CoV-2 Orf8 exhibited 45.3% sequence similarity with SARS-CoV. Orf8 in human isolates from the 2003 epidemic contained a signature 29-nucleotide deletion compared to all civet and bat SARS-related CoVs, which causes the split of full-length Orf8 into two small proteins: 8a and 8b³⁹⁵. Orf8 from SARS-CoV-2 encodes a single polypeptide and lacks the aggregation motif VLVVL present in SARS-CoV Orf8b, which was shown to induce ER stress and activate NLRP3 inflammasomes³⁹⁶. Further, Orf8b protein has been shown to be modified by N-linked glycosylation on N81 residue, which protects Orf8ab protein from proteasomal degradation³⁹⁷. This novel Orf8 likely encodes a secreted protein and has an N-glycosylation site at N78, within the consensus sequence NYT.

We identified 47 high-confidence host protein interactions with Orf8. Several Orf8 interactors are involved in ER stress and ER-associated degradation (ERAD) pathway, including UDP-glucose/glycoprotein glucosyltransferase 2 (UGGT2), ER degradation enhancing alpha-mannosidase like protein 3 (EDEM3), OS9³⁹⁸, N-glycanase 1 (NGLY1), and FAD-dependent oxidoreductase domain-containing protein 2 (FOXRED2)^{399–402}. The ERAD pathway targets unassembled glycoproteins for ubiquitylation and proteasomal degradation. EDEM3 has been shown to increase ubiquitylation of HCV envelope proteins via direct physical interaction and consequently reduce viral production⁴⁰³. OS9 and ERLEC1 proteins are also known to be targeted by other virus-encoding proteins from Dengue virus²¹⁰, HIV²⁰⁹, WNV³⁶, HPV and KSHV⁴⁰⁴, suggesting common molecular mechanisms of infection and proliferation used by these different pathogens.

Infection with the SARS virus results in severe inflammation in the lungs, which can lead to respiratory distress and fibrosis during the late stages of infection⁴⁰⁵. Fibroblast activation and overexpression of collagen are two important aspects of the pathogenesis of lung fibrosis. Interestingly, we identified a number of Orf8 interactors implicated in pulmonary fibrogenesis including FKBP10⁴⁰⁶, GDF15⁴⁰⁷, NEU1⁴⁰⁸ and IL17RA⁴⁰⁹. In addition, the expression levels of ADAMTS1⁴¹⁰ and HS6ST2⁴¹¹ are modulated during lung inflammation and fibrosis and are identified as Orf9b interactors. Growth differentiation factor 15 (GDF15) is a fibroblast-inhibiting cytokine that inhibits the growth and activation of lung fibroblasts by inactivating the TGF- β /Smad pathway, suggesting this cytokine could be a potential therapeutic for ameliorating interstitial lung fibrosis during severe SARS infection. FK506-binding protein 10 (FKBP10) is a collagen chaperone and inhibition attenuates expression of profibrotic mediators and effectors⁴⁰⁶, suggesting that this protein can be targeted to reduce virus-induced lung fibrosis. Activation of the pro-inflammatory cytokine receptor IL17RA in lung tissues is an important host defense mechanism upon fungal, bacterial and viral infections, but its overactivation increases collagen secretion and exacerbates pulmonary fibrosis⁴⁰⁹. Inhibition of IL17RA signaling promotes resolution of

pulmonary inflammation and fibrosis in *in vivo* models⁴¹² and may therefore serve as a therapeutic strategy to reduce lung fibrosis during SARS infection.

Though below our stringent scoring threshold, one additional interesting protein identified in Orf8 pull-downs was the cellular guanyl transferase RNGTT/Mce1. This interaction was significant (>0.05 BFDR) but just below our MIST threshold (MIST score = 0.649). Previous studies suggest RNGTT can exist in the cytoplasm^{413,414}, it is possible that Orf8 recruits RNGTT to viral mRNA to add G to nascent mRNAs after they are acted on by Nsp13 to make GpppN mRNA that is subsequently acted on by Nsp14 and Nsp16 (see section on SARS-CoV Capping Enzymes).

SARS-CoV-2 Orf9b

Orf9b is an accessory protein synthesized from an alternative complete reading frame within the viral N gene, which encodes for a 98-aa long protein. Orf9b has been shown to be expressed in SARS-CoV-infected cells and antibodies against Orf9b were detected in the sera from convalescent-phase SARS patients^{415,416}, however the function of Orf9b is largely unknown. It is known that Orf9b can passively diffuse into the nucleus and is actively exported via Crm1-mediated nucleocytoplasmic export⁴¹⁷. In addition, Orf9b localizes to mitochondria and causes mitochondrial elongation by inducing ubiquitination-mediated proteasomal degradation of the main pro-fission factor dynamin-like protein 1 (DRP1)⁴¹⁸. Orf9b targets the mitochondrial-associated adaptor molecule MAVS signalosome by utilizing PCBP2 and the HECT domain-containing E3 ligase AIP4, resulting in the degradation of MAVS and therefore limiting host cell interferon responses⁴¹⁸.

We found that SARS-CoV-2 Orf9b interacts with 11 human proteins, including with a mitochondrial import receptor, translocase of outer membrane 70 (TOM70). TOM70 is known to interact with MAVS protein upon RNA virus infection and it acts as a critical adaptor linking MAVS to TBK1/IRF3, resulting in the activation of IRF-3⁴¹⁹. TOM70 also makes a dynamic protein complex with HSP90/IRF3/BAX and mediates virus-induced apoptosis⁴²⁰. Though more studies need to be done to fully flesh out this interaction, it is possible SARS-CoV-2 Orf9b may target TOM70 to modulate IRF3-mediated gene expression or apoptosis upon virus infection. Another mitochondrial protein identified as interacting with Orf9b is BCL2-associated athanogene 5 (BAG5). BAG5 inhibits mitophagy of damaged mitochondria by suppressing recruitment of Parkin to the sites of damage⁴²¹. Several viruses trigger Parkin-dependent mitophagy to promote persistent infection and impair the innate immune response⁴²². SARS-CoV-2 Orf9b might act similarly by antagonizing the function of BAG5.

In addition to mitochondrial proteins, SARS-CoV-2 Orf9b was also found to interact with CHMP2A, a member of the endosomal sorting complex required for transport (ESCRT)-III machinery⁴²³. CHMP2A was shown to contribute to the budding of a variety of viruses, including HIV⁴²⁴, equine infectious anemia virus (EIAV)⁴²⁵, and murine leukemia virus⁴²⁵, suggesting a critical role for virus release. Other Orf9b interactors of interest include microtubule affinity-regulating kinases MARK1, MARK2, and MARK3. These proteins are involved in regulating microtubule dynamics and phosphorylation of tau⁴²⁶, and MARK2 was also shown to regulate HIV trafficking through phosphorylation of FEZ1⁴²⁷.

SARS-CoV-2 Orf9c

SARS-CoV-2 Orf9c (referred to as Orf9B in Wu et. al.) encodes a short polypeptide that is 70 aa in length². There is some debate over whether Orf9c encodes a functional protein, or what its function would be, as Orf9c is thought to be dispensable for virus replication⁴²⁸⁻⁴³⁰. Therefore it is unclear how clinically relevant molecular interactions for this bait would be in the context of coronavirus infection. Keeping that in mind, we are able to express and purify Orf9c in HEK293T/17 cells, identifying 26 high confidence human protein

interactors of diverse functional enrichments. These functions include mitochondrial respiratory chain complex assembly (NDUFB9, NDUFAF1, ACAD9, ECSIT, BCS1L)^{431–433}, GPI-anchor biosynthesis (PIGO, PIGS, GPAA1)⁴³⁴, and regulation of I-kappa β kinase and NF-kappa β signaling (NLRX1, F2RL1, NDFIP2).

F2RL1 is implicated in a variety of cellular processes related to the pathogenesis of respiratory viruses and pulmonary disease, including NF κ B activation, cooperativity with Toll-like receptors, innate immune recruitment and activation, and acute lung inflammation^{435–439}. In the context of Influenza A virus (IAV) pathogenesis of monocytes and macrophages, F2RL1 activation protects against viral infection through an IFN-gamma-mediated mechanism^{440,441}. Importantly, F2RL1 is the target of four known pharmacologic agents AC-55541, AZ8838, GB110 and Z3451. Another interactor linked to pulmonary function is NLRX1, an attenuator of IAV-induced inflammation⁴⁴². During IAV infection, NLRX1 promotes type I IFN signaling and macrophage survival⁴⁴². It is an essential moderator of macrophage immunity, as it senses the extent of viral replication and maintains a protective balance between antiviral immunity and excessive inflammation within the lungs⁴⁴².

In our study, Orf9c is also shown to interact with MRP1 (encoded by *ABCC1*), a multifunctional ATP-binding cassette protein that, among other diverse functions, controls the ATP-dependent efflux of drugs from the cell. It has been implicated in multidrug resistance, viral pathogenesis, and pulmonary disease, and is directly targetable by FDA-approved pharmacologic agents daunorubicin and mitoxantrone⁴⁴³. MRP1 has a demonstrated role in both HIV and CMV biology, is found to be differentially expressed in a polarized subset of macrophages during HIV-1 infection, and is associated with CMV latency^{444–446}. Interestingly, HIV-1 protease inhibitors saquinavir, zidovudine, and zalcitabine are substrates of MRP1, though it was found that MRP1 did not affect the antiviral activity of these drugs in cell lines⁴⁴⁷. The role of MRP1 in determining the severity of diseases (e.g., COPD, pneumonia, and lung carcinoma) and multidrug resistance in the lung has been well-characterized, which could be particularly relevant given the clinical manifestation and treatment of ARDS and pneumonia during SARS-CoV-2 infection^{448–451}.

SARS-CoV-2 Orf10

SARS-CoV-2 Orf10 codes for a peptide only 38 aa long and does not have a homolog in SARS-CoV. There is no data yet providing evidence that the protein is expressed during SARS-CoV-2 infection, however we found that upon expression in HEK293T/17 cells Orf10 interacts with nine host proteins. Among these are multiple members of the Cullin RING E3 ligase 2 (CRL2) complex, including CUL2, ELOB, ELOC, RBX1 and ZYG11B. Cullin RING E3 ligases play a central role in viral infections, since they are commonly hijacked by viral proteins to ubiquitinate and degrade viral restriction factors. CRL2 has been previously found to be targeted by poxviral ANK/BC via a C-terminal BC box domain resulting in potent suppression of inflammatory cytokines production, including interferon⁴⁵². Similarly, HPV16 E7 protein binds to an active CRL2 complex, and this association correlates with the ability of HPV16 E7 to transform cells⁴⁵³. ZYG11B, a substrate adapter of CUL2, is the highest scoring hit in the Orf10 interactome indicating that Orf10 might bind to the assembled CUL2^{ZYG11B} complex. Interestingly, ZYG11B targets substrates with exposed N-terminal glycines for degradation⁴⁵⁴. Orf10 contains an N-terminal glycine but does not have lysine residues, suggesting a few possible models: (1) Orf10 hijacks the CUL2^{ZYG11B} complex for ubiquitination and degradation of restriction factors, or (2) Orf10 blocks CUL2^{ZYG11B} and prevents the ubiquitination of its targets, or (3) Orf10 is targeted by CUL2^{ZYG11B} for degradation through N-terminal ubiquitination (see main text **Figure 4**).

FUTURE DIRECTIONS AND AVAILABLE RESOURCES

As we've described above, a number of SARS-CoV-2 proteins are known to assemble and function in multi-viral subunit complexes. It is likely that co-expression of specific proteins forming these complexes could identify and implicate additional clinically relevant interactions within the host. At the time of this publication, we are in the process of generating a library of untagged transient expression constructs suitable for co-transfection with the tagged library generated and used in this study. In parallel we are also integrating endogenous affinity tags into SARS-CoV-2 replicons and full-length recombinant SARS-CoV-2 genomes to study interactions in the context of infection. Performing affinity purifications in the context of infection will also address concerns regarding protein overexpression, which often allows the identification of weaker or lower abundance interactions, but can lead to false positive or less clinically relevant identifications. Replication of our HEK293T overexpression AP-MS experiments is currently ongoing in other cell types, including cultured and primary lung epithelial cells. Comparison and integration of AP-MS overexpression data from multiple cell lines under different cellular contexts (e.g. co-expression with SARS-CoV-2 proteins or replicons) will help clarify and prioritize high-confidence host interactions for further follow-up studies. Given the urgent and acute global need for SARS-CoV-2 research, we will make these resources available to the scientific community upon completion, as we have done in this study.

Supplementary Methods

Sequence Analysis of Overlapping Coding Regions (Extended Data Figure 1a). The alignment of 2,784 SARS-CoV-2 sequences (obtained from GISAID on April 4, 2020) was performed with MAFFT v7.450 with default settings using Geneious Prime® 2019.1.1 software.

Western Blotting (Extended Data Figure 1b). Transfected cell lysate (IP input samples) was mixed with SDS sample buffer and beta-mercaptoethanol, heated to 95°C for 5 minutes, then run on a 26-well Criterion TGX gel (BioRad) until the dye front was 1.5 cm from the bottom of the gel. Protein was transferred to PVDF membrane, and all western blotting steps were performed in PBS containing 5% milk and 1% Tween-20. Anti-strep antibody (Qiagen 34850) was used at a concentration of 1:2,500, and goat anti-mouse HRP conjugate (BioRad) was used at a concentration of 1:20,000. Membranes were washed with PBS prior to addition of HRP substrate and signal detection on film.

Pearson correlation and clustering analysis of SARS-CoV-2 affinity purification-mass spectrometry (AP-MS) Data (Extended Data Figure 2). All MS runs were compared and clustered using standard artMS (<https://github.com/biodavidjm/artMS>) procedures on observed feature intensities computed by MaxQuant. Pearson's pairwise correlations between MS runs are clustered according to similar correlation patterns.

PFAM domain enrichment analysis (Extended Data Figure 4). The enrichment of individual PFAM domains (or PFAM clans)⁴⁵⁵ was calculated with a hypergeometric test where success is defined as number of domains, and the number of trials is the number of individual preys pulled-down with each viral bait. The population values were the numbers of individual PFAM domains and clans in the human proteome. To make sure that the p-values that signify enrichment were meaningful, we only considered PFAM domains that have been pulled-down at least three times with any SARS-CoV-2 protein, and which occur in the human proteome at least five times.

Expression analysis of interacting genes (Extended Data Figure 5). We used GTEx (version 8, median gene-level transcripts per million (TPM) by tissue), which consisted of 17382 samples (578 lung samples)⁴⁵⁶ to examine the mRNA expression of all interacting proteins (n=323). The comparison gene group was all RefSeq genes (n=24,491). The lung expression values represent the median expression of each gene across the GTEx lung samples. The lung enrichment values are calculated by dividing the median expression of each gene in lung tissue by the median expression of each gene across all tissues (including lung). A value of greater than one indicates that the gene expression is enriched in lung tissue. Values were plotted on a log10 scale. All figures and statistics were produced in Python3 and code and reference tables can be found at: (https://github.com/stephaniewanko/Fraser_Lab/tree/master/QCRG_COVID19).

Nsp5 main protease (3CLpro) cleavage prediction (Extended Data Figure 6). We used sequence specificity data for SARS Nsp5⁴⁵⁷ (98.7% identical to SARS-CoV-2 nsp5) and NetCorona⁴⁵⁸ to predict cleavage sites within interacting factors. PDB ID: 1UJ1 served as template for peptide docking which was performed using the predicted P4-P1 residues (BioLuminate, Schrödinger, LLC). Illustration of the docked model was generated in PyMol (Schrödinger, LLC).

Orf6 consensus sequence analysis (Extended Data Figure 7). Orf6 sequence homologs were identified using the BLAST tool⁴⁵⁹ (accession number YP_009724394.1), run with the default settings: gap opening and extension costs of 11 and 1, respectively, BLOSUM62 as the scoring matrix, and an e-value threshold of 10. The search yielded 34 homologous sequences. The multiple sequence alignment was visualized using the

MView web server: <https://www.ebi.ac.uk/Tools/msa/mview/>⁴⁶⁰ and the WebLogo server (<https://weblogo.berkeley.edu/logo.cgi>)⁴⁶¹.

Analysis of off-target activities for characteristic Sigma receptor ligands (Extended Data Figure 11). Radioligand binding assays were performed as previously described⁴⁶². Detailed experimental protocols are available on the NIMH PDSP website (<https://pdspdb.unc.edu/html/tutorials/UNC-CH%20Protocol%20Book.pdf>).

Evolutionary analysis (Supplementary Table 3). For each gene, we obtained a human ORF sequence, choosing the splice isoform with the longest ORF. We used this ORF as query in a blastn search⁴⁶³ of NCBI's NR database and for each non-human primate species we collected the blast hit with the highest bit score, filtering out matches of <60% identity or <100bp alignment length, and ignoring database sequences that are >20kb long or have no annotated ORF. We also blasted each primate hit to a collection of all human genes, to ensure all sequences are reciprocal best hits (a proxy for true orthology, albeit imperfect). We extracted ORFs from each primate match, and aligned orthologous sequences using MACSE⁴⁶⁴, treating the human sequence as 'reliable' and the other primate sequences as 'less reliable' (parameters: -fs_l 10 and -stop_l 10). We then manually inspected and, if necessary, edited all alignments to remove unreliable sequence segments, because gene predictions found in NR sometimes contain erroneous exons. In a few cases, manual inspection led us to select a different splice isoform, because the longest ORF occasionally included one very poorly-alignable exon. We used phym⁴⁶⁵ to estimate a phylogeny for each alignment (parameters: -m GTR --pinv e --alpha e -f e). The alignment and phylogeny were then used as input for PAML's codeml algorithm⁴⁶⁶ comparing the neutral/purifying model 8a (where dN/dS for codons follows a beta distribution with values between 0 and 1, with an extra class of sites with dN/dS fixed at 1) with model 8 that allows a subset of codons to have dN/dS > 1 (parameters: codon frequency F3x4, estimate kappa, initial kappa 2, initial omega 0.4, ncatG 10, cleandata 0). We performed a likelihood ratio test⁴⁶⁶ to obtain a p-value, by comparing twice the difference in log-likelihoods with the chi-squared distribution with 1 degree of freedom. After running all 332 analyses, we used the Benjamini-Hochberg procedure⁴⁶⁷ to control the false-discovery rate.

Supplementary References

1. Chan, J. F.-W. *et al.* Genomic characterization of the 2019 novel human-pathogenic coronavirus isolated from a patient with atypical pneumonia after visiting Wuhan. *Emerg. Microbes Infect.* **9**, 221–236 (2020).
2. Wu, F. *et al.* A new coronavirus associated with human respiratory disease in China. *Nature* (2020) doi:10.1038/s41586-020-2008-3.
3. Du, L. *et al.* The spike protein of SARS-CoV--a target for vaccine and therapeutic development. *Nat. Rev. Microbiol.* **7**, 226–236 (2009).
4. Xie, Y. *et al.* GPS-Lipid: a robust tool for the prediction of multiple lipid modification sites. *Sci. Rep.* **6**, 28249 (2016).
5. Ren, J. *et al.* CSS-Palm 2.0: an updated software for palmitoylation sites prediction. *Protein Eng. Des. Sel.* **21**, 639–644 (2008).
6. Petit, C. M. *et al.* Palmitoylation of the cysteine-rich endodomain of the SARS-coronavirus spike glycoprotein is important for spike-mediated cell fusion. *Virology* **360**, 264–274 (2007).
7. Ohta, E. *et al.* Identification and characterization of GCP16, a novel acylated Golgi protein that interacts with GCP170. *J. Biol. Chem.* **278**, 51957–51967 (2003).
8. Ko, P.-J. *et al.* A ZDHHC5-GOLGA7 Protein Acyltransferase Complex Promotes Nonapoptotic Cell Death. *Cell Chem Biol* **26**, 1716–1724.e9 (2019).
9. Li, Y. *et al.* DHHC5 interacts with PDZ domain 3 of post-synaptic density-95 (PSD-95) protein and plays a role in learning and memory. *J. Biol. Chem.* **285**, 13022–13031 (2010).
10. Kokkola, T. *et al.* Somatostatin receptor 5 is palmitoylated by the interacting ZDHHC5 palmitoyltransferase. *FEBS Lett.* **585**, 2665–2670 (2011).
11. Lu, Y. *et al.* Palmitoylation of NOD1 and NOD2 is required for bacterial sensing. *Science* **366**, 460–467 (2019).
12. Swarthout, J. T. *et al.* DHHC9 and GCP16 constitute a human protein fatty acyltransferase with specificity for H- and N-Ras. *J. Biol. Chem.* **280**, 31141–31148 (2005).
13. Burkard, C. *et al.* ATP1A1-mediated Src signaling inhibits coronavirus entry into host cells. *J. Virol.* **89**,

4434–4448 (2015).

14. Lingemann, M. *et al.* The alpha-1 subunit of the Na⁺,K⁺-ATPase (ATP1A1) is required for macropinocytic entry of respiratory syncytial virus (RSV) in human respiratory epithelial cells. *PLoS Pathog.* **15**, e1007963 (2019).
15. García-Dorival, I. *et al.* Elucidation of the Ebola virus VP24 cellular interactome and disruption of virus biology through targeted inhibition of host-cell protein function. *J. Proteome Res.* **13**, 5120–5135 (2014).
16. Iwasaki, M. *et al.* Interactome analysis of the lymphocytic choriomeningitis virus nucleoprotein in infected cells reveals ATPase Na⁺/K⁺ transporting subunit Alpha 1 and prohibitin as host-cell factors involved in the life cycle of mammarenaviruses. *PLoS Pathog.* **14**, e1006892 (2018).
17. Tan, Y.-J., Lim, S. G. & Hong, W. Characterization of viral proteins encoded by the SARS-coronavirus genome. *Antiviral Res.* **65**, 69–78 (2005).
18. DeDiego, M. L. *et al.* Coronavirus virulence genes with main focus on SARS-CoV envelope gene. *Virus Res.* **194**, 124–137 (2014).
19. Schoeman, D. & Fielding, B. C. Coronavirus envelope protein: current knowledge. *Viol. J.* **16**, 69 (2019).
20. Du, Y., Zuckermann, F. A. & Yoo, D. Myristoylation of the small envelope protein of porcine reproductive and respiratory syndrome virus is non-essential for virus infectivity but promotes its growth. *Virus Res.* **147**, 294–299 (2010).
21. Li, Y., Surya, W., Claudine, S. & Torres, J. Structure of a conserved Golgi complex-targeting signal in coronavirus envelope proteins. *J. Biol. Chem.* **289**, 12535–12549 (2014).
22. Jimenez-Guardeño, J. M. *et al.* Identification of the Mechanisms Causing Reversion to Virulence in an Attenuated SARS-CoV for the Design of a Genetically Stable Vaccine. *PLoS Pathog.* **11**, e1005215 (2015).
23. Venkatagopalan, P., Daskalova, S. M., Lopez, L. A., Dolezal, K. A. & Hogue, B. G. Coronavirus envelope (E) protein remains at the site of assembly. *Virology* **478**, 75–85 (2015).
24. Tang, X. *et al.* BET bromodomain proteins mediate downstream signaling events following growth factor stimulation in human lung fibroblasts and are involved in bleomycin-induced pulmonary fibrosis. *Mol.*

Pharmacol. **83**, 283–293 (2013).

25. Huang, B., Yang, X.-D., Zhou, M.-M., Ozato, K. & Chen, L.-F. Brd4 coactivates transcriptional activation of NF-kappaB via specific binding to acetylated RelA. *Mol. Cell. Biol.* **29**, 1375–1387 (2009).
26. Tian, B. *et al.* Selective Antagonists of the Bronchiolar Epithelial NF-κB-Bromodomain-Containing Protein 4 Pathway in Viral-Induced Airway Inflammation. *Cell Rep.* **23**, 1138–1151 (2018).
27. You, J., Croyle, J. L., Nishimura, A., Ozato, K. & Howley, P. M. Interaction of the bovine papillomavirus E2 protein with Brd4 tethers the viral DNA to host mitotic chromosomes. *Cell* **117**, 349–360 (2004).
28. Taniguchi, Y. The Bromodomain and Extra-Terminal Domain (BET) Family: Functional Anatomy of BET Paralogous Proteins. *Int. J. Mol. Sci.* **17**, (2016).
29. LeRoy, G., Rickards, B. & Flint, S. J. The double bromodomain proteins Brd2 and Brd3 couple histone acetylation to transcription. *Mol. Cell* **30**, 51–60 (2008).
30. Hsu, S. C. *et al.* The BET Protein BRD2 Cooperates with CTCF to Enforce Transcriptional and Architectural Boundaries. *Mol. Cell* **66**, 102–116.e7 (2017).
31. Boehm, D., Conrad, R. J. & Ott, M. Bromodomain proteins in HIV infection. *Viruses* **5**, 1571–1586 (2013).
32. McBride, A. A. & Jang, M. K. Current understanding of the role of the Brd4 protein in the papillomavirus lifecycle. *Viruses* **5**, 1374–1394 (2013).
33. Robinson, M. S. & Bonifacino, J. S. Adaptor-related proteins. *Curr. Opin. Cell Biol.* **13**, 444–453 (2001).
34. Liu, L. *et al.* Defective HIV-1 particle assembly in AP-3-deficient cells derived from patients with Hermansky-Pudlak syndrome type 2. *J. Virol.* **86**, 11242–11253 (2012).
35. Jung, J. *et al.* Identification of a homozygous deletion in the AP3B1 gene causing Hermansky-Pudlak syndrome, type 2. *Blood* **108**, 362–369 (2006).
36. Li, M. *et al.* Identification of antiviral roles for the exon-junction complex and nonsense-mediated decay in flaviviral infection. *Nat Microbiol* **4**, 985–995 (2019).
37. Arndt, A. L., Larson, B. J. & Hogue, B. G. A conserved domain in the coronavirus membrane protein tail is important for virus assembly. *J. Virol.* **84**, 11418–11428 (2010).
38. Neuman, B. W. *et al.* A structural analysis of M protein in coronavirus assembly and morphology. *J. Struct.*

- Biol.* **174**, 11–22 (2011).
39. Perrier, A. *et al.* The C-terminal domain of the MERS coronavirus M protein contains a trans-Golgi network localization signal. *J. Biol. Chem.* **294**, 14406–14421 (2019).
 40. Pacciarini, F. *et al.* Persistent replication of severe acute respiratory syndrome coronavirus in human tubular kidney cells selects for adaptive mutations in the membrane protein. *J. Virol.* **82**, 5137–5144 (2008).
 41. Fang, X. *et al.* The membrane protein of SARS-CoV suppresses NF-kappaB activation. *J. Med. Virol.* **79**, 1431–1439 (2007).
 42. Siu, K.-L. *et al.* Severe acute respiratory syndrome coronavirus M protein inhibits type I interferon production by impeding the formation of TRAF3.TANK.TBK1/IKKepsilon complex. *J. Biol. Chem.* **284**, 16202–16209 (2009).
 43. Chan, C.-M., Ma, C.-W., Chan, W.-Y. & Chan, H. Y. E. The SARS-Coronavirus Membrane protein induces apoptosis through modulating the Akt survival pathway. *Arch. Biochem. Biophys.* **459**, 197–207 (2007).
 44. Tsoi, H. *et al.* The SARS-coronavirus membrane protein induces apoptosis via interfering with PDK1-PKB/Akt signalling. *Biochem. J* **464**, 439–447 (2014).
 45. Tan, Y.-J., Lim, S. G. & Hong, W. Regulation of cell death during infection by the severe acute respiratory syndrome coronavirus and other coronaviruses. *Cell. Microbiol.* **9**, 2552–2561 (2007).
 46. Appenzeller-Herzog, C. & Hauri, H.-P. The ER-Golgi intermediate compartment (ERGIC): in search of its identity and function. *J. Cell Sci.* **119**, 2173–2183 (2006).
 47. Krishnan, M. N. *et al.* RNA interference screen for human genes associated with West Nile virus infection. *Nature* **455**, 242–245 (2008).
 48. Karlas, A. *et al.* Genome-wide RNAi screen identifies human host factors crucial for influenza virus replication. *Nature* **463**, 818–822 (2010).
 49. Orvedahl, A. *et al.* Image-based genome-wide siRNA screen identifies selective autophagy factors. *Nature* **480**, 113–117 (2011).
 50. Ooi, Y. S., Stiles, K. M., Liu, C. Y., Taylor, G. M. & Kielian, M. Genome-wide RNAi screen identifies novel

- host proteins required for alphavirus entry. *PLoS Pathog.* **9**, e1003835 (2013).
51. Han, J. *et al.* Genome-wide CRISPR/Cas9 Screen Identifies Host Factors Essential for Influenza Virus Replication. *Cell Rep.* **23**, 596–607 (2018).
 52. Li, B. *et al.* Genome-wide CRISPR screen identifies host dependency factors for influenza A virus infection. *Nat. Commun.* **11**, 164 (2020).
 53. Wang, S., Tukachinsky, H., Romano, F. B. & Rapoport, T. A. Cooperation of the ER-shaping proteins atlastin, lunapark, and reticulons to generate a tubular membrane network. *Elife* **5**, (2016).
 54. Yamamoto, Y., Yoshida, A., Miyazaki, N., Iwasaki, K. & Sakisaka, T. Arl6IP1 has the ability to shape the mammalian ER membrane in a reticulon-like fashion. *Biochem. J* **458**, 69–79 (2014).
 55. Diaz, A., Wang, X. & Ahlquist, P. Membrane-shaping host reticulon proteins play crucial roles in viral RNA replication compartment formation and function. *Proc. Natl. Acad. Sci. U. S. A.* **107**, 16291–16296 (2010).
 56. Chen, Y.-J., Williams, J. M., Arvan, P. & Tsai, B. Reticulon protects the integrity of the ER membrane during ER escape of large macromolecular protein complexes. *J. Cell Biol.* **219**, (2020).
 57. Haenssler, E., Ramabhadran, V., Murphy, C. S., Heidtman, M. I. & Isberg, R. R. Endoplasmic Reticulum Tubule Protein Reticulon 4 Associates with the Legionella pneumophila Vacuole and with Translocated Substrate Ceg9. *Infect. Immun.* **83**, 3479–3489 (2015).
 58. Hafirassou, M. L. *et al.* A Global Interactome Map of the Dengue Virus NS1 Identifies Virus Restriction and Dependency Host Factors. *Cell Rep.* **21**, 3900–3913 (2017).
 59. Singh, C. R. *et al.* Mechanisms of translational regulation by a human eIF5-mimic protein. *Nucleic Acids Res.* **39**, 8314–8328 (2011).
 60. Cheng, D.-D. *et al.* Downregulation of BZW2 inhibits osteosarcoma cell growth by inactivating the Akt/mTOR signaling pathway. *Oncol. Rep.* **38**, 2116–2122 (2017).
 61. Jin, X., Liao, M., Zhang, L., Yang, M. & Zhao, J. Role of the novel gene BZW2 in the development of hepatocellular carcinoma. *J. Cell. Physiol.* (2019) doi:10.1002/jcp.28331.
 62. Sato, K. *et al.* Novel oncogene 5MP1 reprograms c-Myc translation initiation to drive malignant phenotypes in colorectal cancer. *EBioMedicine* **44**, 387–402 (2019).

63. Wang, S., Bai, W., Huang, J., Lv, F. & Bai, H. Prognostic significance of BZW2 expression in lung adenocarcinoma patients. *Int. J. Clin. Exp. Pathol.* **12**, 4289–4296 (2019).
64. Gao, H. *et al.* BZW2 gene knockdown induces cell growth inhibition, G1 arrest and apoptosis in muscle-invasive bladder cancers: A microarray pathway analysis. *J. Cell. Mol. Med.* **23**, 3905–3915 (2019).
65. Park, C. R. *et al.* The accessory proteins REEP5 and REEP6 refine CXCR1-mediated cellular responses and lung cancer progression. *Sci. Rep.* **6**, 39041 (2016).
66. Warner, N., Burberry, A., Pliakas, M., McDonald, C. & Núñez, G. A genome-wide small interfering RNA (siRNA) screen reveals nuclear factor- κ B (NF- κ B)-independent regulators of NOD2-induced interleukin-8 (IL-8) secretion. *J. Biol. Chem.* **289**, 28213–28224 (2014).
67. Strieter, R. M., Kunkel, S. L., Keane, M. P. & Standiford, T. J. Chemokines in lung injury: Thomas A. Neff Lecture. *Chest* **116**, 103S–110S (1999).
68. Hu, Y. *et al.* Scramblase TMEM16F terminates T cell receptor signaling to restrict T cell exhaustion. *J. Exp. Med.* **213**, 2759–2772 (2016).
69. Wang, H. *et al.* SARS coronavirus entry into host cells through a novel clathrin- and caveolae-independent endocytic pathway. *Cell Res.* **18**, 290–301 (2008).
70. Ehlen, H. W. A. *et al.* Inactivation of anoctamin-6/Tmem16f, a regulator of phosphatidylserine scrambling in osteoblasts, leads to decreased mineral deposition in skeletal tissues. *J. Bone Miner. Res.* **28**, 246–259 (2013).
71. He, R. *et al.* Analysis of multimerization of the SARS coronavirus nucleocapsid protein. *Biochem. Biophys. Res. Commun.* **316**, 476–483 (2004).
72. Nelson, G. W., Stohlman, S. A. & Tahara, S. M. High affinity interaction between nucleocapsid protein and leader/intergenic sequence of mouse hepatitis virus RNA. *J. Gen. Virol.* **81**, 181–188 (2000).
73. Baric, R. S. *et al.* Interactions between coronavirus nucleocapsid protein and viral RNAs: implications for viral transcription. *J. Virol.* **62**, 4280–4287 (1988).
74. Raaben, M., Groot Koerkamp, M. J. A., Rottier, P. J. M. & de Haan, C. A. M. Mouse hepatitis coronavirus

- replication induces host translational shutoff and mRNA decay, with concomitant formation of stress granules and processing bodies. *Cell. Microbiol.* **9**, 2218–2229 (2007).
75. Wada, M., Lokugamage, K. G., Nakagawa, K., Narayanan, K. & Makino, S. Interplay between coronavirus, a cytoplasmic RNA virus, and nonsense-mediated mRNA decay pathway. *Proc. Natl. Acad. Sci. U. S. A.* **115**, E10157–E10166 (2018).
76. Surjit, M. *et al.* The severe acute respiratory syndrome coronavirus nucleocapsid protein is phosphorylated and localizes in the cytoplasm by 14-3-3-mediated translocation. *J. Virol.* **79**, 11476–11486 (2005).
77. V'kovski, P. *et al.* Determination of host proteins composing the microenvironment of coronavirus replicase complexes by proximity-labeling. *Elife* **8**, (2019).
78. Emmott, E. *et al.* The cellular interactome of the coronavirus infectious bronchitis virus nucleocapsid protein and functional implications for virus biology. *J. Virol.* **87**, 9486–9500 (2013).
79. Sun, N. *et al.* Proteomics Analysis of Cellular Proteins Co-Immunoprecipitated with Nucleoprotein of Influenza A Virus (H7N9). *Int. J. Mol. Sci.* **16**, 25982–25998 (2015).
80. White, J. P. & Lloyd, R. E. Poliovirus unlinks TIA1 aggregation and mRNA stress granule formation. *J. Virol.* **85**, 12442–12454 (2011).
81. Visser, L. J. *et al.* Foot-and-Mouth Disease Virus Leader Protease Cleaves G3BP1 and G3BP2 and Inhibits Stress Granule Formation. *J. Virol.* **93**, (2019).
82. Panas, M. D. *et al.* Sequestration of G3BP coupled with efficient translation inhibits stress granules in Semliki Forest virus infection. *Mol. Biol. Cell* **23**, 4701–4712 (2012).
83. Katsafanas, G. C. & Moss, B. Colocalization of transcription and translation within cytoplasmic poxvirus factories coordinates viral expression and subjugates host functions. *Cell Host Microbe* **2**, 221–228 (2007).
84. Borghese, F. & Michiels, T. The leader protein of cardioviruses inhibits stress granule assembly. *J. Virol.* **85**, 9614–9622 (2011).
85. Ward, A. M. *et al.* Quantitative mass spectrometry of DENV-2 RNA-interacting proteins reveals that the DEAD-box RNA helicase DDX6 binds the DB1 and DB2 3' UTR structures. *RNA Biol.* **8**, 1173–1186 (2011).

86. Bidet, K., Dadlani, D. & Garcia-Blanco, M. A. G3BP1, G3BP2 and CAPRIN1 are required for translation of interferon stimulated mRNAs and are targeted by a dengue virus non-coding RNA. *PLoS Pathog.* **10**, e1004242 (2014).
87. Katoh, H. *et al.* Japanese encephalitis virus core protein inhibits stress granule formation through an interaction with Caprin-1 and facilitates viral propagation. *J. Virol.* **87**, 489–502 (2013).
88. Nelson, E. V. *et al.* Ebola Virus Does Not Induce Stress Granule Formation during Infection and Sequesters Stress Granule Proteins within Viral Inclusions. *J. Virol.* **90**, 7268–7284 (2016).
89. Nakagawa, K., Narayanan, K., Wada, M. & Makino, S. Inhibition of Stress Granule Formation by Middle East Respiratory Syndrome Coronavirus 4a Accessory Protein Facilitates Viral Translation, Leading to Efficient Virus Replication. *J. Virol.* **92**, (2018).
90. Brownsword, M. J., Doyle, N., Brocard, M., Locker, N. & Maier, H. J. Infectious bronchitis virus regulates cellular stress granule signaling. *bioRxiv* 819482 (2019) doi:10.1101/819482.
91. Chen, H. *et al.* Mass spectroscopic characterization of the coronavirus infectious bronchitis virus nucleoprotein and elucidation of the role of phosphorylation in RNA binding by using surface plasmon resonance. *J. Virol.* **79**, 1164–1179 (2005).
92. Reineke, L. C. *et al.* Casein Kinase 2 Is Linked to Stress Granule Dynamics through Phosphorylation of the Stress Granule Nucleating Protein G3BP1. *Mol. Cell. Biol.* **37**, (2017).
93. de Wilde, A. H. *et al.* A Kinome-Wide Small Interfering RNA Screen Identifies Proviral and Antiviral Host Factors in Severe Acute Respiratory Syndrome Coronavirus Replication, Including Double-Stranded RNA-Activated Protein Kinase and Early Secretory Pathway Proteins. *J. Virol.* **89**, 8318–8333 (2015).
94. Siddiqui-Jain, A. *et al.* CX-4945, an orally bioavailable selective inhibitor of protein kinase CK2, inhibits prosurvival and angiogenic signaling and exhibits antitumor efficacy. *Cancer Res.* **70**, 10288–10298 (2010).
95. Taiaroa, G. *et al.* Direct RNA sequencing and early evolution of SARS-CoV-2. *bioRxiv* 2020.03.05.976167 (2020) doi:10.1101/2020.03.05.976167.
96. Culbertson, M. R., Underbrink, K. M. & Fink, G. R. Frameshift suppression *Saccharomyces cerevisiae*. II.

- Genetic properties of group II suppressors. *Genetics* **95**, 833–853 (1980).
97. Philippe, L., van den Elzen, A. M. G., Watson, M. J. & Thoreen, C. C. Global analysis of LARP1 translation targets reveals tunable and dynamic features of 5' TOP motifs. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 5319–5328 (2020).
98. McKinney, C., Yu, D. & Mohr, I. A new role for the cellular PABP repressor Paip2 as an innate restriction factor capable of limiting productive cytomegalovirus replication. *Genes Dev.* **27**, 1809–1820 (2013).
99. Le Sage, V., Cinti, A., Amorim, R. & Mouland, A. J. Adapting the Stress Response: Viral Subversion of the mTOR Signaling Pathway. *Viruses* **8**, (2016).
100. Dylla, J. *et al.* Middle East Respiratory Syndrome and Severe Acute Respiratory Syndrome: Current Therapeutic Options and Potential Targets for Novel Therapies. *Drugs* **77**, 1935–1966 (2017).
101. Kindrachuk, J. *et al.* Antiviral potential of ERK/MAPK and PI3K/AKT/mTOR signaling modulation for Middle East respiratory syndrome coronavirus infection as identified by temporal kinome analysis. *Antimicrob. Agents Chemother.* **59**, 1088–1099 (2015).
102. Wang, C.-H. *et al.* Adjuvant treatment with a mammalian target of rapamycin inhibitor, sirolimus, and steroids improves outcomes in patients with severe H1N1 pneumonia and acute respiratory failure. *Crit. Care Med.* **42**, 313–321 (2014).
103. Kühn, U. & Wahle, E. Structure and function of poly(A) binding proteins. *Biochim. Biophys. Acta* **1678**, 67–84 (2004).
104. Kumar, G. R. & Glaunsinger, B. A. Nuclear import of cytoplasmic poly(A) binding protein restricts gene expression via hyperadenylation and nuclear retention of mRNA. *Mol. Cell. Biol.* **30**, 4996–5008 (2010).
105. Suzuki, Y. *et al.* Characterization of RyDEN (C19orf66) as an Interferon-Stimulated Cellular Inhibitor against Dengue Virus Replication. *PLoS Pathog.* **12**, e1005357 (2016).
106. Hsu, C.-H. *et al.* Identification and Characterization of Potential Biomarkers by Quantitative Tissue Proteomics of Primary Lung Adenocarcinoma. *Mol. Cell. Proteomics* **15**, 2396–2410 (2016).
107. Gregersen, L. H. *et al.* MOV10 Is a 5' to 3' RNA helicase contributing to UPF1 mRNA target degradation by translocation along 3' UTRs. *Mol. Cell* **54**, 573–585 (2014).

108. Cuevas, R. A. *et al.* MOV10 Provides Antiviral Activity against RNA Viruses by Enhancing RIG-I-MAVS-Independent IFN Induction. *J. Immunol.* **196**, 3877–3886 (2016).
109. Balinsky, C. A. *et al.* IRAV (FLJ11286), an Interferon-Stimulated Gene with Antiviral Activity against Dengue Virus, Interacts with MOV10. *J. Virol.* **91**, (2017).
110. Zhao, K. *et al.* MOV10 inhibits replication of porcine reproductive and respiratory syndrome virus by retaining viral nucleocapsid protein in the cytoplasm of Marc-145 cells. *Biochem. Biophys. Res. Commun.* **504**, 157–163 (2018).
111. Li, J. *et al.* MOV10 sequesters the RNP of influenza A virus in the cytoplasm and is antagonized by viral NS1 protein. *Biochem. J* **476**, 467–481 (2019).
112. Ziebuhr, J. The coronavirus replicase. *Curr. Top. Microbiol. Immunol.* **287**, 57–94 (2005).
113. Wathelet, M. G., Orr, M., Frieman, M. B. & Baric, R. S. Severe acute respiratory syndrome coronavirus evades antiviral signaling: role of nsp1 and rational design of an attenuated strain. *J. Virol.* **81**, 11620–11633 (2007).
114. Law, A. H. Y., Lee, D. C. W., Cheung, B. K. W., Yim, H. C. H. & Lau, A. S. Y. Role for nonstructural protein 1 of severe acute respiratory syndrome coronavirus in chemokine dysregulation. *J. Virol.* **81**, 416–422 (2007).
115. Starokadomskyy, P. *et al.* DNA polymerase- α regulates the activation of type I interferons through cytosolic RNA:DNA synthesis. *Nat. Immunol.* **17**, 495–504 (2016).
116. Xu, L. H., Huang, M., Fang, S. G. & Liu, D. X. Coronavirus infection induces DNA replication stress partly through interaction of its nonstructural protein 13 with the p125 subunit of DNA polymerase δ . *J. Biol. Chem.* **286**, 39546–39559 (2011).
117. Wang, L. *et al.* Comparative influenza protein interactomes identify the role of plakophilin 2 in virus restriction. *Nat. Commun.* **8**, 13876 (2017).
118. Graham, R. L., Sims, A. C., Brockway, S. M., Baric, R. S. & Denison, M. R. The nsp2 replicase proteins of murine hepatitis virus and severe acute respiratory syndrome coronavirus are dispensable for viral replication. *J. Virol.* **79**, 13399–13411 (2005).

119. Graham, R. L., Sims, A. C., Baric, R. S. & Denison, M. R. The nsp2 proteins of mouse hepatitis virus and SARS coronavirus are dispensable for viral replication. *Adv. Exp. Med. Biol.* **581**, 67–72 (2006).
120. Angeletti, S. *et al.* COVID-2019: The role of the nsp2 and nsp3 in its pathogenesis. *J. Med. Virol.* (2020) doi:10.1002/jmv.25719.
121. Viklund, I.-M. *et al.* WAFL, a new protein involved in regulation of early endocytic transport at the intersection of actin and microtubule dynamics. *Exp. Cell Res.* **315**, 1040–1052 (2009).
122. Derivery, E. *et al.* The Arp2/3 activator WASH controls the fission of endosomes through a large multiprotein complex. *Dev. Cell* **17**, 712–723 (2009).
123. Jia, D. *et al.* WASH and WAVE actin regulators of the Wiskott-Aldrich syndrome protein (WASP) family are controlled by analogous structurally related complexes. *Proc. Natl. Acad. Sci. U. S. A.* **107**, 10442–10447 (2010).
124. Morita, M. *et al.* A novel 4EHP-GIGYF2 translational repressor complex is essential for mammalian development. *Mol. Cell. Biol.* **32**, 3585–3593 (2012).
125. Fusco, D. N. *et al.* HELZ2 Is an IFN Effector Mediating Suppression of Dengue Virus. *Front. Microbiol.* **8**, 240 (2017).
126. Tew, G. W. *et al.* SmgGDS regulates cell proliferation, migration, and NF-kappaB transcriptional activity in non-small cell lung carcinoma. *J. Biol. Chem.* **283**, 963–976 (2008).
127. Lawrence, T. The nuclear factor NF-kappaB pathway in inflammation. *Cold Spring Harb. Perspect. Biol.* **1**, a001651 (2009).
128. Oudshoorn, D. *et al.* Expression and Cleavage of Middle East Respiratory Syndrome Coronavirus nsp3-4 Polypeptide Induce the Formation of Double-Membrane Vesicles That Mimic Those Associated with Coronaviral RNA Replication. *MBio* **8**, (2017).
129. Beachboard, D. C., Anderson-Daniels, J. M. & Denison, M. R. Mutations across murine hepatitis virus nsp4 alter virus fitness and membrane modifications. *J. Virol.* **89**, 2080–2089 (2015).
130. Sakai, Y. *et al.* Two-amino acids change in the nsp4 of SARS coronavirus abolishes viral replication. *Virology* **510**, 165–174 (2017).

131. Cohen, M., Feinstein, N., Wilson, K. L. & Gruenbaum, Y. Nuclear pore protein gp210 is essential for viability in HeLa cells and *Caenorhabditis elegans*. *Mol. Biol. Cell* **14**, 4230–4237 (2003).
132. Borlido, J. *et al.* Nuclear pore complex-mediated modulation of TCR signaling is required for naïve CD4⁺ T cell homeostasis. *Nat. Immunol.* **19**, 594–605 (2018).
133. Jamali, T., Jamali, Y., Mehrbod, M. & Mofrad, M. R. K. Nuclear pore complex: biochemistry and biophysics of nucleocytoplasmic transport in health and disease. *Int. Rev. Cell Mol. Biol.* **287**, 233–286 (2011).
134. Gomez, G. N., Abrar, F., Dodhia, M. P., Gonzalez, F. G. & Nag, A. SARS coronavirus protein nsp1 disrupts localization of Nup93 from the nuclear pore complex. *Biochem. Cell Biol.* **97**, 758–766 (2019).
135. Affholter, J. A., Hsieh, C. L., Francke, U. & Roth, R. A. Insulin-degrading enzyme: stable expression of the human complementary DNA, characterization of its protein product, and chromosomal mapping of the human and mouse genes. *Mol. Endocrinol.* **4**, 1125–1135 (1990).
136. Vekrellis, K. *et al.* Neurons regulate extracellular levels of amyloid beta-protein via proteolysis by insulin-degrading enzyme. *J. Neurosci.* **20**, 1657–1665 (2000).
137. Parmentier, N. *et al.* Production of an antigenic peptide by insulin-degrading enzyme. *Nat. Immunol.* **11**, 449–454 (2010).
138. Li, Q., Ali, M. A. & Cohen, J. I. Insulin degrading enzyme is a cellular receptor mediating varicella-zoster virus infection and cell-to-cell spread. *Cell* **127**, 305–316 (2006).
139. Li, Q., Krogmann, T., Ali, M. A., Tang, W.-J. & Cohen, J. I. The amino terminus of varicella-zoster virus (VZV) glycoprotein E is required for binding to insulin-degrading enzyme, a VZV receptor. *J. Virol.* **81**, 8525–8532 (2007).
140. Williamson, C. D., DeBiasi, R. L. & Colberg-Poley, A. M. Viral product trafficking to mitochondria, mechanisms and roles in pathogenesis. *Infect. Disord. Drug Targets* **12**, 18–37 (2012).
141. Mühlenbein, N., Hofmann, S., Rothbauer, U. & Bauer, M. F. Organization and function of the small Tim complexes acting along the import pathway of metabolite carriers into mammalian mitochondria. *J. Biol. Chem.* **279**, 13540–13546 (2004).
142. Sun, X. *et al.* Transcriptional Changes during Naturally Acquired Zika Virus Infection Render Dendritic

- Cells Highly Conducive to Viral Replication. *Cell Rep.* **21**, 3471–3482 (2017).
143. Wells, M. F. *et al.* Genome-wide screens in accelerated human stem cell-derived neural progenitor cells identify Zika virus host factors and drivers of proliferation. *bioRxiv* 476440 (2018) doi:10.1101/476440.
144. Ioakeimidis, F. *et al.* A splicing mutation in the novel mitochondrial protein DNAJC11 causes motor neuron pathology associated with cristae disorganization, and lymphoid abnormalities in mice. *PLoS One* **9**, e104237 (2014).
145. Rind, N. *et al.* A severe human metabolic disease caused by deficiency of the endoplasmatic mannosyltransferase hALG11 leads to congenital disorder of glycosylation-Ip. *Hum. Mol. Genet.* **19**, 1413–1424 (2010).
146. Thiel, C. *et al.* Improved diagnostics lead to identification of three new patients with congenital disorder of glycosylation-Ip. *Hum. Mutat.* **33**, 485–487 (2012).
147. Neuman, B. W., Chamberlain, P., Bowden, F. & Joseph, J. Atlas of coronavirus replicase structure. *Virus Res.* **194**, 49–66 (2014).
148. Ziebuhr, J., Snijder, E. J. & Gorbalenya, A. E. Virus-encoded proteinases and proteolytic processing in the Nidovirales. *J. Gen. Virol.* **81**, 853–879 (2000).
149. Krämer, O. H. HDAC2: a critical factor in health and disease. *Trends Pharmacol. Sci.* **30**, 647–655 (2009).
150. Kim, R. Y. *et al.* MicroRNA-21 drives severe, steroid-insensitive experimental asthma by amplifying phosphoinositide 3-kinase-mediated suppression of histone deacetylase 2. *J. Allergy Clin. Immunol.* **139**, 519–532 (2017).
151. Barnes, P. J. Role of HDAC2 in the pathophysiology of COPD. *Annu. Rev. Physiol.* **71**, 451–464 (2009).
152. Song, H. *et al.* USP17-mediated deubiquitination and stabilization of HDAC2 in cigarette smoke extract-induced inflammation. *Int. J. Clin. Exp. Pathol.* **8**, 10707–10715 (2015).
153. Nagesh, P. T., Hussain, M., Galvin, H. D. & Husain, M. Histone Deacetylase 2 Is a Component of Influenza A Virus-Induced Host Antiviral Response. *Front. Microbiol.* **8**, 1315 (2017).
154. Feng, Q. *et al.* Histone deacetylase inhibitors suppress RSV infection and alleviate virus-induced airway inflammation. *Int. J. Mol. Med.* **38**, 812–822 (2016).

- 155.Saha, B. & Parks, R. J. Histone Deacetylase Inhibitor Suberoylanilide Hydroxamic Acid Suppresses Human Adenovirus Gene Expression and Replication. *J. Virol.* **93**, (2019).
- 156.Dewe, J. M., Fuller, B. L., Lentini, J. M., Kellner, S. M. & Fu, D. TRMT1-Catalyzed tRNA Modifications Are Required for Redox Homeostasis To Ensure Proper Cellular Proliferation and Oxidative Stress Survival. *Mol. Cell. Biol.* **37**, (2017).
- 157.Lubos, E., Loscalzo, J. & Handy, D. E. Glutathione peroxidase-1 in health and disease: from molecular mechanisms to therapeutic opportunities. *Antioxid. Redox Signal.* **15**, 1957–1997 (2011).
- 158.Zhou, L. Z., Johnson, A. P. & Rando, T. A. NF kappa B and AP-1 mediate transcriptional responses to oxidative stress in skeletal muscle cells. *Free Radic. Biol. Med.* **31**, 1405–1416 (2001).
- 159.Borrás, C. *et al.* 17beta-oestradiol up-regulates longevity-related, antioxidant enzyme expression via the ERK1 and ERK2[MAPK]/NFkappaB cascade. *Aging Cell* **4**, 113–118 (2005).
- 160.Angelini, M. M., Akhlaghpour, M., Neuman, B. W. & Buchmeier, M. J. Severe acute respiratory syndrome coronavirus nonstructural proteins 3, 4, and 6 induce double-membrane vesicles. *MBio* **4**, (2013).
- 161.Oostra, M. *et al.* Topology and membrane anchoring of the coronavirus replication complex: not all hydrophobic domains of nsp3 and nsp6 are membrane spanning. *J. Virol.* **82**, 12392–12405 (2008).
- 162.Lundin, A. *et al.* Targeting membrane-bound viral RNA synthesis reveals potent inhibition of diverse coronaviruses including the middle East respiratory syndrome virus. *PLoS Pathog.* **10**, e1004166 (2014).
- 163.Perlman, S. & Netland, J. Coronaviruses post-SARS: update on replication and pathogenesis. *Nat. Rev. Microbiol.* **7**, 439–450 (2009).
- 164.Hagemeijer, M. C., Rottier, P. J. M. & de Haan, C. A. M. Biogenesis and dynamics of the coronavirus replicative structures. *Viruses* **4**, 3245–3269 (2012).
- 165.Neuman, B. W. How the double spherules of infectious bronchitis virus impact our understanding of RNA virus replicative organelles. *mBio* vol. 4 e00987–13 (2013).
- 166.Knoops, K. *et al.* SARS-coronavirus replication is supported by a reticulovesicular network of modified endoplasmic reticulum. *PLoS Biol.* **6**, e226 (2008).
- 167.Cottam, E. M., Whelband, M. C. & Wileman, T. Coronavirus NSP6 restricts autophagosome expansion.

- Autophagy* **10**, 1426–1441 (2014).
168. Munger, J., Bajad, S. U., Collier, H. A., Shenk, T. & Rabinowitz, J. D. Dynamics of the cellular metabolome during human cytomegalovirus infection. *PLoS Pathog.* **2**, e132 (2006).
169. Anand, S. K. & Tikoo, S. K. Viruses as modulators of mitochondrial functions. *Adv. Virol.* **2013**, 738794 (2013).
170. Baratta, M. G. Virus-mediated hijack of one-carbon metabolism. *Nature reviews. Cancer* vol. 19 486 (2019).
171. Fontaine, K. A., Sanchez, E. L., Camarda, R. & Lagunoff, M. Dengue virus induces and requires glycolysis for optimal replication. *J. Virol.* **89**, 2358–2366 (2015).
172. Garrus, J. E. *et al.* Tsg101 and the vacuolar protein sorting pathway are essential for HIV-1 budding. *Cell* **107**, 55–65 (2001).
173. Licata, J. M. *et al.* Overlapping motifs (PTAP and PPEY) within the Ebola virus VP40 protein function independently as late budding domains: involvement of host proteins TSG101 and VPS-4. *J. Virol.* **77**, 1812–1819 (2003).
174. Gorai, T. *et al.* F1Fo-ATPase, F-type proton-translocating ATPase, at the plasma membrane is critical for efficient influenza virus budding. *Proc. Natl. Acad. Sci. U. S. A.* **109**, 4615–4620 (2012).
175. Forgac, M. Vacuolar ATPases: rotary proton pumps in physiology and pathophysiology. *Nat. Rev. Mol. Cell Biol.* **8**, 917–929 (2007).
176. König, R. *et al.* Human host factors required for influenza virus replication. *Nature* **463**, 813–817 (2010).
177. Banerjee, I., Yamauchi, Y., Helenius, A. & Horvath, P. High-content analysis of sequential events during the early phase of influenza A virus infection. *PLoS One* **8**, e68450 (2013).
178. Ochiai, H., Sakai, S., Hirabayashi, T., Shimizu, Y. & Terasawa, K. Inhibitory effect of bafilomycin A1, a specific inhibitor of vacuolar-type proton pump, on the growth of influenza A and B viruses in MDCK cells. *Antiviral Res.* **27**, 425–430 (1995).
179. Brass, A. L. *et al.* The IFITM proteins mediate cellular resistance to influenza A H1N1 virus, West Nile virus, and dengue virus. *Cell* **139**, 1243–1254 (2009).

180. Sessions, O. M. *et al.* Discovery of insect and human dengue virus host factors. *Nature* **458**, 1047–1050 (2009).
181. Jansen, E. J. R. *et al.* ATP6AP1 deficiency causes an immunodeficiency with hepatopathy, cognitive impairment and abnormal protein glycosylation. *Nat. Commun.* **7**, 11600 (2016).
182. Pareja, F. *et al.* Loss-of-function mutations in ATP6AP1 and ATP6AP2 in granular cell tumors. *Nat. Commun.* **9**, 3533 (2018).
183. Soh, T. K. *et al.* Herpes simplex virus-1 pUL56 degrades GOPC to alter the plasma membrane proteome. *bioRxiv* 729343 (2019) doi:10.1101/729343.
184. Matheson, N. J. *et al.* Cell Surface Proteomic Map of HIV Infection Reveals Antagonism of Amino Acid Metabolism by Vpu and Nef. *Cell Host Microbe* **18**, 409–423 (2015).
185. Lee, M. N. *et al.* Common genetic variants modulate pathogen-sensing responses in human dendritic cells. *Science* **343**, 1246980 (2014).
186. Gräf, S. *et al.* Identification of rare sequence variation underlying heritable pulmonary arterial hypertension. *Nat. Commun.* **9**, 1416 (2018).
187. Liu Bin *et al.* Abstract 14378: ATP13A3 Loss of Function Disrupts Polyamine Homeostasis in Pulmonary Arterial Endothelial Cells. *Circulation* **140**, A14378–A14378 (2019).
188. Hayashi, T. & Su, T. P. Regulating ankyrin dynamics: Roles of sigma-1 receptors. *Proc. Natl. Acad. Sci. U. S. A.* **98**, 491–496 (2001).
189. Subissi, L. *et al.* One severe acute respiratory syndrome coronavirus protein complex integrates processive RNA polymerase and exonuclease activities. *Proc. Natl. Acad. Sci. U. S. A.* **111**, E3900–9 (2014).
190. Kirchdoerfer, R. N. & Ward, A. B. Structure of the SARS-CoV nsp12 polymerase bound to nsp7 and nsp8 co-factors. *Nat. Commun.* **10**, 2342 (2019).
191. Zhai, Y. *et al.* Insights into SARS-CoV transcription and replication from the structure of the nsp7-nsp8 hexadecamer. *Nat. Struct. Mol. Biol.* **12**, 980–986 (2005).
192. te Velthuis, A. J. W., Arnold, J. J., Cameron, C. E., van den Worm, S. H. E. & Snijder, E. J. The RNA

- polymerase activity of SARS-coronavirus nsp12 is primer dependent. *Nucleic Acids Res.* **38**, 203–214 (2010).
193. te Velthuis, A. J. W., van den Worm, S. H. E. & Snijder, E. J. The SARS-coronavirus nsp7+nsp8 complex is a unique multimeric RNA polymerase capable of both de novo initiation and primer extension. *Nucleic Acids Res.* **40**, 1737–1747 (2012).
194. Gorbalenya, A. E. *et al.* The palm subdomain-based active site is internally permuted in viral RNA-dependent RNA polymerases of an ancient lineage. *J. Mol. Biol.* **324**, 47–62 (2002).
195. Lehmann, K. C. *et al.* Discovery of an essential nucleotidylating activity associated with a newly delineated conserved domain in the RNA polymerase-containing protein of all nidoviruses. *Nucleic Acids Res.* **43**, 8416–8434 (2015).
196. Brockway, S. M., Clay, C. T., Lu, X. T. & Denison, M. R. Characterization of the expression, intracellular localization, and replication complex association of the putative mouse hepatitis virus RNA-dependent RNA polymerase. *J. Virol.* **77**, 10515–10527 (2003).
197. Gao, Y. *et al.* Structure of RNA-dependent RNA polymerase from 2019-nCoV, a major antiviral drug target. *bioRxiv* 2020.03.16.993386 (2020) doi:10.1101/2020.03.16.993386.
198. Gurkan, C. *et al.* Large-scale profiling of Rab GTPase trafficking networks: the membrome. *Mol. Biol. Cell* **16**, 3847–3864 (2005).
199. Linford, A. *et al.* Rab14 and its exchange factor FAM116 link endocytic recycling and adherens junction stability in migrating cells. *Dev. Cell* **22**, 952–966 (2012).
200. Qi, M. *et al.* Rab11-FIP1C and Rab14 direct plasma membrane sorting and particle incorporation of the HIV-1 envelope glycoprotein complex. *PLoS Pathog.* **9**, e1003278 (2013).
201. Pechenick Jowers, T. *et al.* RAB1A promotes Vaccinia virus replication by facilitating the production of intracellular enveloped virions. *Virology* **475**, 66–73 (2015).
202. Lin, J. *et al.* Rab1A is required for assembly of classical swine fever virus particle. *Virology* **514**, 18–29 (2018).
203. Macovei, A., Petrareanu, C., Lazar, C., Florian, P. & Branza-Nichita, N. Regulation of hepatitis B virus

- infection by Rab5, Rab7, and the endolysosomal compartment. *J. Virol.* **87**, 6415–6427 (2013).
204. Inoue, J. *et al.* HBV secretion is regulated through the activation of endocytic and autophagic compartments mediated by Rab7 stimulation. *J. Cell Sci.* **128**, 1696–1706 (2015).
205. Bayliss, R., Wheeldon, J., Caucheteux, S. M., Niessen, C. M. & Piguet, V. Identification of host trafficking genes required for HIV-1 virological synapse formation in dendritic cells. *J. Virol.* (2020) doi:10.1128/JVI.01597-19.
206. Padanad, M. S. *et al.* Fatty Acid Oxidation Mediated by Acyl-CoA Synthetase Long Chain 3 Is Required for Mutant KRAS Lung Tumorigenesis. *Cell Rep.* **16**, 1614–1628 (2016).
207. Saliakoura, M. *et al.* The ACSL3-LPIAT1 signaling drives prostaglandin synthesis in non-small cell lung cancer. *Oncogene* (2020) doi:10.1038/s41388-020-1196-5.
208. Nchoutmboube, J. A. *et al.* Increased long chain acyl-CoA synthetase activity and fatty acid import is linked to membrane synthesis for development of picornavirus replication organelles. *PLoS Pathog.* **9**, e1003401 (2013).
209. Jäger, S. *et al.* Global landscape of HIV-human protein complexes. *Nature* **481**, 365–370 (2011).
210. Shah, P. S. *et al.* Comparative Flavivirus-Host Protein Interaction Mapping Reveals Mechanisms of Dengue and Zika Virus Pathogenesis. *Cell* **175**, 1931–1945.e18 (2018).
211. Ramage, H. R. *et al.* A combined proteomics/genomics approach links hepatitis C virus infection with nonsense-mediated mRNA decay. *Mol. Cell* **57**, 329–340 (2015).
212. Penn, B. H. *et al.* An Mtb-Human Protein-Protein Interaction Map Identifies a Switch between Host Antiviral and Antibacterial Responses. *Mol. Cell* **71**, 637–648.e5 (2018).
213. Sadat, M. A. *et al.* Glycosylation, hypogammaglobulinemia, and resistance to viral infections. *N. Engl. J. Med.* **370**, 1615–1625 (2014).
214. Molleston, J. M. *et al.* A conserved virus-induced cytoplasmic TRAMP-like complex recruits the exosome to target viral RNA for degradation. *Genes Dev.* **30**, 1658–1670 (2016).
215. Rialdi, A. *et al.* The RNA Exosome Syncs IAV-RNAPII Transcription to Promote Viral Ribogenesis and Infectivity. *Cell* **169**, 679–692.e14 (2017).

216. Huang, K.-H., Wang, C.-H., Lin, C.-H. & Kuo, H.-P. NF- κ B repressing factor downregulates basal expression and mycobacterium tuberculosis induced IP-10 and IL-8 synthesis via interference with NF- κ B in monocytes. *J. Biomed. Sci.* **21**, 71 (2014).
217. Huang, K.-H. *et al.* NF- κ B repressing factor inhibits chemokine synthesis by peripheral blood mononuclear cells and alveolar macrophages in active pulmonary tuberculosis. *PLoS One* **8**, e77789 (2013).
218. Sawicki, S. G. & Sawicki, D. L. Coronaviruses use discontinuous extension for synthesis of subgenome-length negative strands. *Adv. Exp. Med. Biol.* **380**, 499–506 (1995).
219. Hu, J. *et al.* AKAP95 regulates splicing through scaffolding RNAs and RNA processing factors. *Nat. Commun.* **7**, 13347 (2016).
220. Chua, K. & Reed, R. Human step II splicing factor hSlu7 functions in restructuring the spliceosome between the catalytic steps of splicing. *Genes Dev.* **13**, 841–850 (1999).
221. Konig, R. *et al.* Global analysis of host-pathogen interactions that regulate early-stage HIV-1 replication. *Cell* **135**, 49–60 (2008).
222. Schäffler, K. *et al.* A stimulatory role for the La-related protein 4B in translation. *RNA* **16**, 1488–1499 (2010).
223. Lin, Y. J., Liao, C. L. & Lai, M. M. Identification of the cis-acting signal for minus-strand RNA synthesis of a murine coronavirus: implications for the role of minus-strand RNA in RNA replication and transcription. *J. Virol.* **68**, 8131–8140 (1994).
224. Spagnolo, J. F. & Hogue, B. G. Host protein interactions with the 3' end of bovine coronavirus RNA and the requirement of the poly(A) tail for coronavirus defective genome replication. *J. Virol.* **74**, 5053–5065 (2000).
225. Cirillo, L. *et al.* UBAP2L Forms Distinct Cores that Act in Nucleating Stress Granules Upstream of G3BP1. *Curr. Biol.* **30**, 698–707.e6 (2020).
226. Kedersha, N. *et al.* Stress granules and processing bodies are dynamically linked sites of mRNP remodeling. *J. Cell Biol.* **169**, 871–884 (2005).
227. Jiang, H. *et al.* Regulation of transcription by the MLL2 complex and MLL complex-associated AKAP95.

- Nat. Struct. Mol. Biol.* **20**, 1156–1163 (2013).
- 228.Cho, Y. S. *et al.* Phosphorylation-driven assembly of the RIP1-RIP3 complex regulates programmed necrosis and virus-induced inflammation. *Cell* **137**, 1112–1123 (2009).
- 229.Holler, N. *et al.* Fas triggers an alternative, caspase-8-independent cell death pathway using the kinase RIP as effector molecule. *Nat. Immunol.* **1**, 489–495 (2000).
- 230.Weinlich, R. & Green, D. R. The two faces of receptor interacting protein kinase-1. *Mol. Cell* **56**, 469–480 (2014).
- 231.Chaudhary, P. M., Jasmin, A., Eby, M. T. & Hood, L. Modulation of the NF-kappa B pathway by virally encoded death effector domains-containing proteins. *Oncogene* **18**, 5738–5746 (1999).
- 232.Dondelinger, Y. *et al.* Serine 25 phosphorylation inhibits RIPK1 kinase-dependent cell death in models of infection and inflammation. *Nat. Commun.* **10**, 1729 (2019).
- 233.Wagner, R. N., Reed, J. C. & Chanda, S. K. HIV-1 protease cleaves the serine-threonine kinases RIPK1 and RIPK2. *Retrovirology* **12**, 74 (2015).
- 234.Sheridan, C. Death by inflammation: drug makers chase the master controller. *Nat. Biotechnol.* **37**, 111–113 (2019).
- 235.Kamada, S., Kikkawa, U., Tsujimoto, Y. & Hunter, T. A-kinase-anchoring protein 95 functions as a potential carrier for the nuclear translocation of active caspase 3 through an enzyme-substrate-like association. *Mol. Cell. Biol.* **25**, 9469–9477 (2005).
- 236.Miknis, Z. J. *et al.* Severe acute respiratory syndrome coronavirus nsp9 dimerization is essential for efficient viral growth. *J. Virol.* **83**, 3007–3018 (2009).
- 237.Egloff, M.-P. *et al.* The severe acute respiratory syndrome-coronavirus replicative protein nsp9 is a single-stranded RNA-binding subunit unique in the RNA virus world. *Proc. Natl. Acad. Sci. U. S. A.* **101**, 3792–3796 (2004).
- 238.Ponnusamy, R., Moll, R., Weimar, T., Mesters, J. R. & Hilgenfeld, R. Variable oligomerization modes in coronavirus non-structural protein 9. *J. Mol. Biol.* **383**, 1081–1096 (2008).
- 239.Sutton, G. *et al.* The nsp9 replicase protein of SARS-coronavirus, structure and functional insights.

- Structure* **12**, 341–353 (2004).
- 240.von Brunn, A. *et al.* Analysis of intraviral protein-protein interactions of the SARS coronavirus ORFeome. *PLoS One* **2**, e459 (2007).
- 241.Sharma, A., Solmaz, S. R., Blobel, G. & Melčák, I. Ordered Regions of Channel Nucleoporins Nup62, Nup54, and Nup58 Form Dynamic Complexes in Solution. *J. Biol. Chem.* **290**, 18370–18378 (2015).
- 242.Dewangan, P. S., Sonawane, P. J., Chouksey, A. R. & Chauhan, R. The Nup62 Coiled-Coil Motif Provides Plasticity for Triple-Helix Bundle Formation. *Biochemistry* **56**, 2803–2811 (2017).
- 243.Ulrich, A., Partridge, J. R. & Schwartz, T. U. The stoichiometry of the nucleoporin 62 subcomplex of the nuclear pore in solution. *Mol. Biol. Cell* **25**, 1484–1492 (2014).
- 244.Solmaz, S. R., Blobel, G. & Melčák, I. Ring cycle for dilating and constricting the nuclear pore. *Proc. Natl. Acad. Sci. U. S. A.* **110**, 5858–5863 (2013).
- 245.Chug, H., Trakhanov, S., Hülsmann, B. B., Pleiner, T. & Görlich, D. Crystal structure of the metazoan Nup62•Nup58•Nup54 nucleoporin complex. *Science* **350**, 106–110 (2015).
- 246.Beck, M. & Hurt, E. The nuclear pore complex: understanding its function through structural insight. *Nat. Rev. Mol. Cell Biol.* **18**, 73–89 (2017).
- 247.Wang, B., Zhang, X. & Zhao, Z. Validation-based insertional mutagenesis for identification of Nup214 as a host factor for EV71 replication in RD cells. *Biochem. Biophys. Res. Commun.* **437**, 452–456 (2013).
- 248.Di Nunzio, F. *et al.* Human nucleoporins promote HIV-1 docking at the nuclear pore, nuclear import and integration. *PLoS One* **7**, e46037 (2012).
- 249.Malik, P. *et al.* Herpes simplex virus ICP27 protein directly interacts with the nuclear pore complex through Nup62, inhibiting host nucleocytoplasmic transport pathways. *J. Biol. Chem.* **287**, 12277–12292 (2012).
- 250.Park, N., Skern, T. & Gustin, K. E. Specific cleavage of the nuclear pore complex protein Nup62 by a viral protease. *J. Biol. Chem.* **285**, 28796–28805 (2010).
- 251.Watters, K. *et al.* Differential Disruption of Nucleocytoplasmic Trafficking Pathways by Rhinovirus 2A Proteases. *J. Virol.* **91**, (2017).
- 252.Zhang Y.-Z. *et al.* [The 2A protease of enterovirus 71 cleaves nup62 to inhibit nuclear transport]. *Bing Du*

Xue Bao **29**, 421–425 (2013).

253. Watters, K. & Palmenberg, A. C. Differential processing of nuclear pore complex proteins by rhinovirus 2A proteases from different species and serotypes. *J. Virol.* **85**, 10874–10883 (2011).
254. Walker, E. J. *et al.* Rhinovirus 3C protease facilitates specific nucleoporin cleavage and mislocalisation of nuclear proteins in infected host cells. *PLoS One* **8**, e71316 (2013).
255. Castelló, A., Izquierdo, J. M., Welnowska, E. & Carrasco, L. RNA nuclear export is blocked by poliovirus 2A protease and is concomitant with nucleoporin cleavage. *J. Cell Sci.* **122**, 3799–3809 (2009).
256. Park, N., Katikaneni, P., Skern, T. & Gustin, K. E. Differential targeting of nuclear pore complex proteins in poliovirus-infected cells. *J. Virol.* **82**, 1647–1655 (2008).
257. Porter, F. W. & Palmenberg, A. C. Leader-induced phosphorylation of nucleoporins correlates with nuclear trafficking inhibition by cardioviruses. *J. Virol.* **83**, 1941–1951 (2009).
258. Bardina, M. V. *et al.* Mengovirus-induced rearrangement of the nuclear pore complex: hijacking cellular phosphorylation machinery. *J. Virol.* **83**, 3150–3161 (2009).
259. Porter, F. W., Brown, B. & Palmenberg, A. C. Nucleoporin phosphorylation triggered by the encephalomyocarditis virus leader protein is mediated by mitogen-activated protein kinases. *J. Virol.* **84**, 12538–12548 (2010).
260. Khuperkar, D. *et al.* Selective recruitment of nucleoporins on vaccinia virus factories and the role of Nup358 in viral infection. *Virology* **512**, 151–160 (2017).
261. Chang, C.-W. *et al.* Epstein-Barr virus protein kinase BGLF4 targets the nucleus through interaction with nucleoporins. *J. Virol.* **86**, 8072–8085 (2012).
262. Eberhard, J., Onder, Z. & Moroianu, J. Nuclear import of high risk HPV16 E7 oncoprotein is mediated by its zinc-binding domain via hydrophobic interactions with Nup62. *Virology* **446**, 334–345 (2013).
263. Onder, Z. & Moroianu, J. Nuclear import of cutaneous beta genus HPV8 E7 oncoprotein is mediated by hydrophobic interactions between its zinc-binding domain and FG nucleoporins. *Virology* **449**, 150–162 (2014).
264. Ao, Z. *et al.* Contribution of host nucleoporin 62 in HIV-1 integrase chromatin association and viral DNA

- integration. *J. Biol. Chem.* **287**, 10544–10555 (2012).
265. Tafforeau, L. *et al.* Generation and comprehensive analysis of an influenza virus polymerase cellular interaction network. *J. Virol.* **85**, 13010–13018 (2011).
266. Bauer, M. *et al.* The E3 Ubiquitin Ligase Mind Bomb 1 Controls Adenovirus Genome Release at the Nuclear Pore Complex. *Cell Rep.* **29**, 3785–3795.e8 (2019).
267. Fry, A. M., O'Regan, L., Sabir, S. R. & Bayliss, R. Cell cycle regulation by the NEK family of protein kinases. *J. Cell Sci.* **125**, 4423–4433 (2012).
268. Jung, R., Radko, S. & Pelka, P. The Dual Nature of Nek9 in Adenovirus Replication. *J. Virol.* **90**, 1931–1943 (2016).
269. Jin, J., Arias, E. E., Chen, J., Harper, J. W. & Walter, J. C. A family of diverse Cul4-Ddb1-interacting proteins includes Cdt2, which is required for S phase destruction of the replication factor Cdt1. *Mol. Cell* **23**, 709–721 (2006).
270. Zemke, N. R. & Berk, A. J. The Adenovirus E1A C Terminus Suppresses a Delayed Antiviral Response and Modulates RAS Signaling. *Cell Host Microbe* **22**, 789–800.e5 (2017).
271. Feng, P., Everly, D. N., Jr & Read, G. S. mRNA decay during herpes simplex virus (HSV) infections: protein-protein interactions involving the HSV virion host shutoff protein and translation factors eIF4H and eIF4A. *J. Virol.* **79**, 9651–9664 (2005).
272. Doecker, R. C., Hsu, W.-L., Saffran, H. A. & Smiley, J. R. Herpes simplex virus virion host shutoff protein is stimulated by translation initiation factors eIF4B and eIF4H. *J. Virol.* **78**, 4684–4699 (2004).
273. Sarma, N., Agarwal, D., Shiflett, L. A. & Read, G. S. Small interfering RNAs that deplete the cellular translation factor eIF4H impede mRNA degradation by the virion host shutoff protein of herpes simplex virus. *J. Virol.* **82**, 6600–6609 (2008).
274. Teo, C. S. H. & O'Hare, P. A bimodal switch in global protein translation coupled to eIF4H relocalisation during advancing cell-cell transmission of herpes simplex virus. *PLoS Pathog.* **14**, e1007196 (2018).
275. Yin, W. *et al.* Fibrillin-2 is a key mediator of smooth muscle extracellular matrix homeostasis during mouse tracheal tubulogenesis. *Eur. Respir. J.* **53**, (2019).

- 276.Uriarte, J. J. *et al.* Early Impairment of Lung Mechanics in a Murine Model of Marfan Syndrome. *PLoS One* **11**, e0152124 (2016).
- 277.Kuang, P.-P. *et al.* Coordinate expression of fibulin-5/DANCE and elastin during lung injury repair. *Am. J. Physiol. Lung Cell. Mol. Physiol.* **285**, L1147–52 (2003).
- 278.Jean, J.-C., Eruchalu, I., Cao, Y. X. & Joyce-Brady, M. DANCE in developing and injured lung. *Am. J. Physiol. Lung Cell. Mol. Physiol.* **282**, L75–82 (2002).
- 279.Dabovic, B. *et al.* Function of latent TGF β binding protein 4 and fibulin 5 in elastogenesis and lung development. *J. Cell. Physiol.* **230**, 226–236 (2015).
- 280.Brandsma, C.-A. *et al.* A large lung gene expression study identifying fibulin-5 as a novel player in tissue repair in COPD. *Thorax* **70**, 21–32 (2015).
- 281.Albig, A. R. & Schiemann, W. P. Fibulin-5 antagonizes vascular endothelial growth factor (VEGF) signaling and angiogenic sprouting by endothelial cells. *DNA Cell Biol.* **23**, 367–379 (2004).
- 282.Chen, X. *et al.* Fibulin-5 inhibits Wnt/ β -catenin signaling in lung cancer. *Oncotarget* **6**, 15022–15034 (2015).
- 283.Joseph, J. S. *et al.* Crystal structure of nonstructural protein 10 from the severe acute respiratory syndrome coronavirus reveals a novel fold with two zinc-binding motifs. *J. Virol.* **80**, 7894–7901 (2006).
- 284.Siddell, S., Sawicki, D., Meyer, Y., Thiel, V. & Sawicki, S. Identification of the mutations responsible for the phenotype of three MHV RNA-negative ts mutants. *Adv. Exp. Med. Biol.* **494**, 453–458 (2001).
- 285.Chen, Y. *et al.* Biochemical and structural insights into the mechanisms of SARS coronavirus RNA ribose 2'-O-methylation by nsp16/nsp10 protein complex. *PLoS Pathog.* **7**, e1002294 (2011).
- 286.Su, D. *et al.* Dodecamer structure of severe acute respiratory syndrome coronavirus nonstructural protein nsp10. *J. Virol.* **80**, 7902–7908 (2006).
- 287.Buffalo, C. Z. *et al.* Structural Basis for Tetherin Antagonism as a Barrier to Zoonotic Lentiviral Transmission. *Cell Host Microbe* **26**, 359–368.e8 (2019).
- 288.Buffalo, C. Z., Iwamoto, Y., Hurley, J. H. & Ren, X. How HIV Nef Proteins Hijack Membrane Traffic To Promote Infection. *J. Virol.* **93**, (2019).

289. Breuza, L. *et al.* Proteomics of endoplasmic reticulum-Golgi intermediate compartment (ERGIC) membranes from brefeldin A-treated HepG2 cells identifies ERGIC-32, a new cycling protein that interacts with human Erv46. *J. Biol. Chem.* **279**, 47242–47253 (2004).
290. Choglay, A. A., Chapple, J. P., Blatch, G. L. & Cheetham, M. E. Identification and characterization of a human mitochondrial homologue of the bacterial co-chaperone GrpE. *Gene* **267**, 125–134 (2001).
291. Tian, G. & Cowan, N. J. Tubulin-specific chaperones: components of a molecular machine that assembles the α/β heterodimer. *Methods Cell Biol.* **115**, 155–171 (2013).
292. Sevajol, M., Subissi, L., Decroly, E., Canard, B. & Imbert, I. Insights into RNA synthesis, capping, and proofreading mechanisms of SARS-coronavirus. *Virus Res.* **194**, 90–99 (2014).
293. Chen, Y. & Guo, D. Molecular mechanisms of coronavirus RNA capping and methylation. *Viol. Sin.* **31**, 3–11 (2016).
294. Gu, M. & Lima, C. D. Processing the message: structural insights into capping and decapping mRNA. *Curr. Opin. Struct. Biol.* **15**, 99–106 (2005).
295. Ramanathan, A., Robb, G. B. & Chan, S.-H. mRNA capping: biological functions and applications. *Nucleic Acids Res.* **44**, 7511–7526 (2016).
296. Enjuanes, L., Almazán, F., Sola, I. & Zuñiga, S. Biochemical aspects of coronavirus replication and virus-host interaction. *Annu. Rev. Microbiol.* **60**, 211–230 (2006).
297. Menachery, V. D., Debbink, K. & Baric, R. S. Coronavirus non-structural protein 16: evasion, attenuation, and possible treatments. *Virus Res.* **194**, 191–199 (2014).
298. Desai, P., Sexton, G. L., Huang, E. & Person, S. Localization of herpes simplex virus type 1 UL37 in the Golgi complex requires UL36 but not capsid structures. *J. Virol.* **82**, 11354–11361 (2008).
299. Miorin, L., Maiuri, P., Hoenninger, V. M., Mandl, C. W. & Marcello, A. Spatial and temporal organization of tick-borne encephalitis flavivirus replicated RNA in living cells. *Virology* **379**, 64–77 (2008).
300. Sehgal, P. B. *et al.* Golgi dysfunction is a common feature in idiopathic human pulmonary hypertension and vascular lesions in SHIV-nef-infected macaques. *Am. J. Physiol. Lung Cell. Mol. Physiol.* **297**, L729–37 (2009).

301. Rivero, S., Cardenas, J., Bornens, M. & Rios, R. M. Microtubule nucleation at the cis-side of the Golgi apparatus requires AKAP450 and GM130. *EMBO J.* **28**, 1016–1028 (2009).
302. Terrin, A. *et al.* PKA and PDE4D3 anchoring to AKAP9 provides distinct regulation of cAMP signals at the centrosome. *J. Cell Biol.* **198**, 607–621 (2012).
303. Wu, J. *et al.* Molecular Pathway of Microtubule Organization at the Golgi Apparatus. *Dev. Cell* **39**, 44–60 (2016).
304. Wang, Z., Zhang, C. & Qi, R. Z. A newly identified myomegalin isoform functions in Golgi microtubule organization and ER-Golgi transport. *J. Cell Sci.* **127**, 4904–4917 (2014).
305. Witczak, O. *et al.* Cloning and characterization of a cDNA encoding an A-kinase anchoring protein located in the centrosome, AKAP450. *EMBO J.* **18**, 1858–1868 (1999).
306. Greer, Y. E. *et al.* Casein kinase 1δ functions at the centrosome and Golgi to promote ciliogenesis. *Mol. Biol. Cell* **25**, 1629–1640 (2014).
307. Hurtado, L. *et al.* Disconnecting the Golgi ribbon from the centrosome prevents directional cell migration and ciliogenesis. *J. Cell Biol.* **193**, 917–933 (2011).
308. Subramanian, A. *et al.* Auto-regulation of Secretory Flux by Sensing and Responding to the Folded Cargo Protein Load in the Endoplasmic Reticulum. *Cell* **176**, 1461–1476.e23 (2019).
309. Tenorio, M. J., Luchsinger, C. & Mardones, G. A. Protein kinase A activity is necessary for fission and fusion of Golgi to endoplasmic reticulum retrograde tubules. *PLoS One* **10**, e0135260 (2015).
310. Huang, Z.-X., Wang, H., Wang, Y.-M. & Wang, Y. Novel mechanism coupling cyclic AMP-protein kinase A signaling and golgi trafficking via Gyp1 phosphorylation in polarized growth. *Eukaryot. Cell* **13**, 1548–1556 (2014).
311. Cancino, J. *et al.* Control systems of membrane transport at the interface between the endoplasmic reticulum and the Golgi. *Dev. Cell* **30**, 280–294 (2014).
312. Kolobova, E. *et al.* Microtubule-dependent association of AKAP350A and CCAR1 with RNA stress granules. *Exp. Cell Res.* **315**, 542–555 (2009).
313. Khadka, S. *et al.* A physical interaction network of dengue virus and human proteins. *Mol. Cell. Proteomics*

- 10**, M111.012187 (2011).
- 314.Hidajat, R. *et al.* Hepatitis C virus NS3 protein interacts with ELKS- $\{\delta\}$ and ELKS- $\{\alpha\}$, members of a novel protein family involved in intracellular transport and secretory pathways. *J. Gen. Virol.* **86**, 2197–2208 (2005).
- 315.McCormick, D., Lin, Y.-T. & Grey, F. Identification of Host Factors Involved in Human Cytomegalovirus Replication, Assembly, and Egress Using a Two-Step Small Interfering RNA Screen. *MBio* **9**, (2018).
- 316.Amemiya, T., Gromiha, M. M., Horimoto, K. & Fukui, K. Drug repositioning for dengue haemorrhagic fever by integrating multiple omics analyses. *Sci. Rep.* **9**, 523 (2019).
- 317.Eckerle, L. D. *et al.* Infidelity of SARS-CoV Nsp14-exonuclease mutant virus replication is revealed by complete genome sequencing. *PLoS Pathog.* **6**, e1000896 (2010).
- 318.Ma, Y. *et al.* Structural basis and functional analysis of the SARS coronavirus nsp14-nsp10 complex. *Proc. Natl. Acad. Sci. U. S. A.* **112**, 9436–9441 (2015).
- 319.Lukas, J. *et al.* Functional and Clinical Consequences of Novel α -Galactosidase A Mutations in Fabry Disease. *Hum. Mutat.* **37**, 43–51 (2016).
- 320.McCafferty, E. H. & Scott, L. J. Migalastat: A Review in Fabry Disease. *Drugs* **79**, 543–554 (2019).
- 321.Peng, C. *et al.* The first identification of lysine malonylation substrates and its regulatory enzyme. *Mol. Cell. Proteomics* **10**, M111.012658 (2011).
- 322.Du, J. *et al.* Sirt5 is a NAD-dependent protein lysine demalonylase and desuccinylase. *Science* **334**, 806–809 (2011).
- 323.Tan, M. *et al.* Lysine glutarylation is a protein posttranslational modification regulated by SIRT5. *Cell Metab.* **19**, 605–617 (2014).
- 324.Yang, X. *et al.* SHMT2 Desuccinylation by SIRT5 Drives Cancer Cell Proliferation. *Cancer Res.* **78**, 372–386 (2018).
- 325.Fischer, F. *et al.* Sirt5 deacylation activities show differential sensitivities to nicotinamide inhibition. *PLoS One* **7**, e45098 (2012).
- 326.Carr, S. F., Papp, E., Wu, J. C. & Natsumeda, Y. Characterization of human type I and type II IMP

- dehydrogenases. *J. Biol. Chem.* **268**, 27286–27290 (1993).
- 327.Hedstrom, L., Liechti, G., Goldberg, J. B. & Gollapalli, D. R. The antibiotic potential of prokaryotic IMP dehydrogenase inhibitors. *Curr. Med. Chem.* **18**, 1909–1918 (2011).
- 328.Almazán, F., DeDiego, M. L., Galán, C., Alvarez, E. & Enjuanes, L. Identification of essential genes as a strategy to select a SARS candidate vaccine using a SARS-CoV infectious cDNA. *Adv. Exp. Med. Biol.* **581**, 579–583 (2006).
- 329.Deng, X. *et al.* Coronavirus nonstructural protein 15 mediates evasion of dsRNA sensors and limits apoptosis in macrophages. *Proc. Natl. Acad. Sci. U. S. A.* **114**, E4251–E4260 (2017).
- 330.Xu, X. *et al.* New antiviral target revealed by the hexameric structure of mouse hepatitis virus nonstructural protein nsp15. *J. Virol.* **80**, 7909–7917 (2006).
- 331.Paschal, B. M. & Gerace, L. Identification of NTF2, a cytosolic factor for nuclear import that interacts with nuclear pore complex protein p62. *J. Cell Biol.* **129**, 925–937 (1995).
- 332.Steggerda, S. M., Black, B. E. & Paschal, B. M. Monoclonal antibodies to NTF2 inhibit nuclear protein import by preventing nuclear translocation of the GTPase Ran. *Mol. Biol. Cell* **11**, 703–719 (2000).
- 333.Pasqualato, S., Ménétrey, J., Franco, M. & Cherfils, J. The structural GDP/GTP cycle of human Arf6. *EMBO Rep.* **2**, 234–238 (2001).
- 334.Selyunin, A. S. *et al.* The assembly of a GTPase-kinase signalling complex by a bacterial catalytic scaffold. *Nature* **469**, 107–111 (2011).
- 335.Ha, V. L. *et al.* ASAP3 is a focal adhesion-associated Arf GAP that functions in cell migration and invasion. *J. Biol. Chem.* **283**, 14915–14926 (2008).
- 336.D'Souza-Schorey, C. & Stahl, P. D. Myristoylation is required for the intracellular localization and endocytic function of ARF6. *Exp. Cell Res.* **221**, 153–159 (1995).
- 337.Fang, Z. *et al.* Proteomic identification and functional characterization of a novel ARF6 GTPase-activating protein, ACAP4. *Mol. Cell. Proteomics* **5**, 1437–1449 (2006).
- 338.O'Neal, C. J., Jobling, M. G., Holmes, R. K. & Hol, W. G. J. Structural basis for the activation of cholera toxin by human ARF6-GTP. *Science* **309**, 1093–1096 (2005).

- 339.Wang, C. *et al.* The E3 ubiquitin ligase Nrdp1 'preferentially' promotes TLR-mediated production of type I interferon. *Nat. Immunol.* **10**, 744–752 (2009).
- 340.Tan, Y.-J. *et al.* A novel severe acute respiratory syndrome coronavirus protein, U274, is transported to the cell surface and undergoes endocytosis. *J. Virol.* **78**, 6723–6734 (2004).
- 341.Siu, K.-L. *et al.* Severe acute respiratory syndrome coronavirus ORF3a protein activates the NLRP3 inflammasome by promoting TRAF3-dependent ubiquitination of ASC. *FASEB J.* **33**, 8865–8877 (2019).
- 342.Yount, B. *et al.* Severe acute respiratory syndrome coronavirus group-specific open reading frames encode nonessential functions for replication in cell cultures and mice. *J. Virol.* **79**, 14909–14922 (2005).
- 343.McBride, R. & Fielding, B. C. The role of severe acute respiratory syndrome (SARS)-coronavirus accessory proteins in virus pathogenesis. *Viruses* **4**, 2902–2923 (2012).
- 344.Padhan, K., Minakshi, R., Towheed, M. A. B. & Jameel, S. Severe acute respiratory syndrome coronavirus 3a protein activates the mitochondrial death pathway through p38 MAP kinase activation. *J. Gen. Virol.* **89**, 1960–1969 (2008).
- 345.Yue, Y. *et al.* SARS-Coronavirus Open Reading Frame-3a drives multimodal necrotic cell death. *Cell Death Dis.* **9**, 904 (2018).
- 346.Tan, Y.-J. *et al.* The severe acute respiratory syndrome coronavirus 3a protein up-regulates expression of fibrinogen in lung epithelial cells. *J. Virol.* **79**, 10083–10087 (2005).
- 347.Narayanan, K., Huang, C. & Makino, S. SARS coronavirus accessory proteins. *Virus Res.* **133**, 113–121 (2008).
- 348.Ito, N. *et al.* Severe acute respiratory syndrome coronavirus 3a protein is a viral structural protein. *J. Virol.* **79**, 3182–3186 (2005).
- 349.Minakshi, R. *et al.* The SARS Coronavirus 3a protein causes endoplasmic reticulum stress and induces ligand-independent downregulation of the type 1 interferon receptor. *PLoS One* **4**, e8342 (2009).
- 350.Minakshi, R. & Padhan, K. The YXXΦ motif within the severe acute respiratory syndrome coronavirus (SARS-CoV) 3a protein is crucial for its intracellular transport. *Virol. J.* **11**, 75 (2014).
- 351.Lu, W. *et al.* Severe acute respiratory syndrome-associated coronavirus 3a protein forms an ion channel

- and modulates virus release. *Proc. Natl. Acad. Sci. U. S. A.* **103**, 12540–12545 (2006).
352. Balderhaar, H. J. K. & Ungermann, C. CORVET and HOPS tethering complexes - coordinators of endosome and lysosome fusion. *J. Cell Sci.* **126**, 1307–1316 (2013).
353. Jiang, P. *et al.* The HOPS complex mediates autophagosome-lysosome fusion through interaction with syntaxin 17. *Mol. Biol. Cell* **25**, 1327–1337 (2014).
354. Dunn, L. L. *et al.* New insights into intracellular locations and functions of heme oxygenase-1. *Antioxid. Redox Signal.* **20**, 1723–1742 (2014).
355. Lee, T.-S. & Chau, L.-Y. Heme oxygenase-1 mediates the anti-inflammatory effect of interleukin-10 in mice. *Nat. Med.* **8**, 240–246 (2002).
356. Piantadosi, C. A. *et al.* Heme oxygenase-1 couples activation of mitochondrial biogenesis to anti-inflammatory cytokine expression. *J. Biol. Chem.* **286**, 16374–16385 (2011).
357. Skrzypek, K. *et al.* Interplay between heme oxygenase-1 and miR-378 affects non-small cell lung carcinoma growth, vascularization, and metastasis. *Antioxid. Redox Signal.* **19**, 644–660 (2013).
358. Chen, L.-H., Liao, C.-Y., Lai, L.-C., Tsai, M.-H. & Chuang, E. Y. Semaphorin 6A Attenuates the Migration Capability of Lung Cancer Cells via the NRF2/HMOX1 Axis. *Sci. Rep.* **9**, 13302 (2019).
359. Singh, N., Ahmad, Z., Baid, N. & Kumar, A. Host heme oxygenase-1: Friend or foe in tackling pathogens? *IUBMB Life* **70**, 869–880 (2018).
360. Kumar, A. *et al.* Heme oxygenase-1-derived carbon monoxide induces the Mycobacterium tuberculosis dormancy regulon. *J. Biol. Chem.* **283**, 18032–18039 (2008).
361. Nagasawa, M., Kanzaki, M., Iino, Y., Morishita, Y. & Kojima, I. Identification of a novel chloride channel expressed in the endoplasmic reticulum, golgi apparatus, and nucleus. *J. Biol. Chem.* **276**, 20413–20418 (2001).
362. Jia, Y., Jucius, T. J., Cook, S. A. & Ackerman, S. L. Loss of Clcc1 results in ER stress, misfolded protein accumulation, and neurodegeneration. *J. Neurosci.* **35**, 3001–3009 (2015).
363. Chu, Q. *et al.* Regulation of the ER stress response by a mitochondrial microprotein. *Nat. Commun.* **10**, 4883 (2019).

- 364.Irvin, M. R. *et al.* Genes linked to energy metabolism and immunoregulatory mechanisms are associated with subcutaneous adipose tissue distribution in HIV-infected men. *Pharmacogenet. Genomics* **21**, 798–807 (2011).
- 365.Wu, W. *et al.* The interactome of the human respiratory syncytial virus NS1 protein highlights multiple effects on host cell biology. *J. Virol.* **86**, 7777–7789 (2012).
- 366.Hubel, P. *et al.* A protein-interaction network of interferon-stimulated genes extends the innate immune system landscape. *Nat. Immunol.* **20**, 493–502 (2019).
- 367.Song, F. *et al.* Regulation and biological role of the peptide/histidine transporter SLC15A3 in Toll-like receptor-mediated inflammatory responses in macrophage. *Cell Death Dis.* **9**, 770 (2018).
- 368.Eckhardt, M. *et al.* Multiple Routes to Oncogenesis Are Promoted by the Human Papillomavirus-Host Protein Network. *Cancer Discov.* **8**, 1474–1489 (2018).
- 369.Wu, J. Q. *et al.* Transcriptional profiles in CD8+ T cells from HIV+ progressors on HAART are characterized by coordinated up-regulation of oxidative phosphorylation enzymes and interferon responses. *Virology* **380**, 124–135 (2008).
- 370.Zhou, P., Li, H., Wang, H., Wang, L.-F. & Shi, Z. Bat severe acute respiratory syndrome-like coronavirus ORF3b homologues display different interferon antagonist activities. *J. Gen. Virol.* **93**, 275–281 (2012).
- 371.Kopecky-Bromberg, S. A., Martínez-Sobrido, L., Frieman, M., Baric, R. A. & Palese, P. Severe acute respiratory syndrome coronavirus open reading frame (ORF) 3b, ORF 6, and nucleocapsid proteins function as interferon antagonists. *J. Virol.* **81**, 548–557 (2007).
- 372.Too, I. H. K., Bonne, I., Tan, E. L., Chu, J. J. H. & Alonso, S. Prohibitin plays a critical role in Enterovirus 71 neuropathogenesis. *PLoS Pathog.* **14**, e1006778 (2018).
- 373.Wai, T. *et al.* The membrane scaffold SLP2 anchors a proteolytic hub in mitochondria containing PARL and the i-AAA protease YME1L. *EMBO Rep.* **17**, 1844–1856 (2016).
- 374.Kirchhof, M. G. *et al.* Modulation of T cell activation by stomatin-like protein 2. *J. Immunol.* **181**, 1927–1936 (2008).
- 375.Zhao, J. *et al.* Severe acute respiratory syndrome coronavirus protein 6 is required for optimal replication.

J. Virol. **83**, 2368–2373 (2009).

376. Zhou, H. *et al.* The N-terminal region of severe acute respiratory syndrome coronavirus protein 6 induces membrane rearrangement and enhances virus replication. *J. Virol.* **84**, 3542–3551 (2010).
377. Calvo, E. *et al.* Severe acute respiratory syndrome coronavirus accessory proteins 6 and 9b interact in vivo. *Virus Res.* **169**, 282–288 (2012).
378. Quan, B., Seo, H.-S., Blobel, G. & Ren, Y. Vesiculoviral matrix (M) protein occupies nucleic acid binding site at nucleoporin pair (Rae1 • Nup98). *Proc. Natl. Acad. Sci. U. S. A.* **111**, 9127–9132 (2014).
379. Faria, P. A. *et al.* VSV disrupts the Rae1/mrnp41 mRNA nuclear export pathway. *Mol. Cell* **17**, 93–102 (2005).
380. Alcalá, S., Klee, M., Fernández, J., Fleischer, A. & Pimentel-Muñoz, F. X. A high-throughput screening for mammalian cell death effectors identifies the mitochondrial phosphate carrier as a regulator of cytochrome c release. *Oncogene* **27**, 44–54 (2008).
381. Nelson, C. A., Pekosz, A., Lee, C. A., Diamond, M. S. & Fremont, D. H. Structure and intracellular targeting of the SARS-coronavirus Orf7a accessory protein. *Structure* **13**, 75–85 (2005).
382. Tan, Y.-J. *et al.* Overexpression of 7a, a protein specifically encoded by the severe acute respiratory syndrome coronavirus, induces apoptosis via a caspase-dependent pathway. *J. Virol.* **78**, 14043–14047 (2004).
383. Yuan, X. *et al.* SARS coronavirus 7a protein blocks cell cycle progression at G0/G1 phase via the cyclin D3/pRb pathway. *Virology* **346**, 74–85 (2006).
384. Fielding, B. C. *et al.* Characterization of a unique group-specific protein (U122) of the severe acute respiratory syndrome coronavirus. *J. Virol.* **78**, 7311–7318 (2004).
385. Barraud, P., Banerjee, S., Mohamed, W. I., Jantsch, M. F. & Allain, F. H.-T. A bimodular nuclear localization signal assembled via an extended double-stranded RNA-binding domain acts as an RNA-sensing signal for transportin 1. *Proc. Natl. Acad. Sci. U. S. A.* **111**, E1852–61 (2014).
386. Arnold, M., Nath, A., Hauber, J. & Kehlenbach, R. H. Multiple importins function as nuclear transport receptors for the Rev protein of human immunodeficiency virus type 1. *J. Biol. Chem.* **281**, 20883–20890

- (2006).
- 387.Miyake, Y. *et al.* Influenza virus uses transportin 1 for vRNP debundling during cell entry. *Nat Microbiol* **4**, 578–586 (2019).
- 388.Gagné, B., Tremblay, N., Park, A. Y., Baril, M. & Lamarre, D. Importin β 1 targeting by hepatitis C virus NS3/4A protein restricts IRF3 and NF- κ B signaling of IFNB1 antiviral response. *Traffic* **18**, 362–377 (2017).
- 389.Zhang, W. *et al.* Extended haplotype association study in Crohn's disease identifies a novel, Ashkenazi Jewish-specific missense mutation in the NF- κ B pathway gene, HEATR3. *Genes Immun.* **14**, 310–316 (2013).
- 390.Caruso, R., Warner, N., Inohara, N. & Núñez, G. NOD1 and NOD2: signaling, host defense, and inflammatory disease. *Immunity* **41**, 898–908 (2014).
- 391.Dediego, M. L. *et al.* Pathogenicity of severe acute respiratory coronavirus deletion mutants in hACE-2 transgenic mice. *Virology* **376**, 379–389 (2008).
- 392.Lau, S. K. P. *et al.* Severe acute respiratory syndrome coronavirus-like virus in Chinese horseshoe bats. *Proc. Natl. Acad. Sci. U. S. A.* **102**, 14040–14045 (2005).
- 393.Song, H.-D. *et al.* Cross-host evolution of severe acute respiratory syndrome coronavirus in palm civet and human. *Proc. Natl. Acad. Sci. U. S. A.* **102**, 2430–2435 (2005).
- 394.Lau, S. K. P. *et al.* Severe Acute Respiratory Syndrome (SARS) Coronavirus ORF8 Protein Is Acquired from SARS-Related Coronavirus from Greater Horseshoe Bats through Recombination. *J. Virol.* **89**, 10532–10547 (2015).
- 395.Oostra, M., de Haan, C. A. M. & Rottier, P. J. M. The 29-nucleotide deletion present in human but not in animal severe acute respiratory syndrome coronaviruses disrupts the functional expression of open reading frame 8. *J. Virol.* **81**, 13876–13888 (2007).
- 396.Shi, C.-S., Nabar, N. R., Huang, N.-N. & Kehrl, J. H. SARS-Coronavirus Open Reading Frame-8b triggers intracellular stress pathways and activates NLRP3 inflammasomes. *Cell Death Discov* **5**, 101 (2019).
- 397.Le, T. M. *et al.* Expression, post-translational modification and biochemical characterization of proteins

- encoded by subgenomic mRNA8 of the severe acute respiratory syndrome coronavirus. *FEBS J.* **274**, 4211–4222 (2007).
- 398.Hosokawa, N., Kamiya, Y., Kamiya, D., Kato, K. & Nagata, K. Human OS-9, a lectin required for glycoprotein endoplasmic reticulum-associated degradation, recognizes mannose-trimmed N-glycans. *J. Biol. Chem.* **284**, 17061–17068 (2009).
- 399.Suzuki, T., Huang, C. & Fujihira, H. The cytoplasmic peptide:N-glycanase (NGLY1) - Structure, expression and cellular functions. *Gene* **577**, 1–7 (2016).
- 400.Hirao, K. *et al.* EDEM3, a soluble EDEM homolog, enhances glycoprotein endoplasmic reticulum-associated degradation and mannose trimming. *J. Biol. Chem.* **281**, 9650–9658 (2006).
- 401.Arnold, S. M. & Kaufman, R. J. The noncatalytic portion of human UDP-glucose: glycoprotein glucosyltransferase I confers UDP-glucose binding and transferase function to the catalytic domain. *J. Biol. Chem.* **278**, 43320–43328 (2003).
- 402.Riemer, J. *et al.* A luminal flavoprotein in endoplasmic reticulum-associated degradation. *Proc. Natl. Acad. Sci. U. S. A.* **106**, 14831–14836 (2009).
- 403.Saeed, M. *et al.* Role of the endoplasmic reticulum-associated degradation (ERAD) pathway in degradation of hepatitis C virus envelope proteins and production of virus particles. *J. Biol. Chem.* **286**, 37264–37273 (2011).
- 404.Davis, Z. H. *et al.* Global mapping of herpesvirus-host protein complexes reveals a transcription strategy for late genes. *Mol. Cell* **57**, 349–360 (2015).
- 405.Ye, J. *et al.* Molecular pathology in the lungs of severe acute respiratory syndrome patients. *Am. J. Pathol.* **170**, 538–545 (2007).
- 406.Staab-Weijnitz, C. A. *et al.* FK506-Binding Protein 10, a Potential Novel Drug Target for Idiopathic Pulmonary Fibrosis. *Am. J. Respir. Crit. Care Med.* **192**, 455–467 (2015).
- 407.Kim, Y.-I. *et al.* Epithelial cell-derived cytokines CST3 and GDF15 as potential therapeutics for pulmonary fibrosis. *Cell Death Dis.* **9**, 506 (2018).
- 408.Luzina, I. G. *et al.* Elevated expression of NEU1 sialidase in idiopathic pulmonary fibrosis provokes

- pulmonary collagen deposition, lymphocytosis, and fibrosis. *Am. J. Physiol. Lung Cell. Mol. Physiol.* **310**, L940–54 (2016).
409. Gurczynski, S. J. & Moore, B. B. IL-17 in the lung: the good, the bad, and the ugly. *Am. J. Physiol. Lung Cell. Mol. Physiol.* **314**, L6–L16 (2018).
410. Liu L. *et al.* [miR-21 promotes pulmonary fibrosis in rats via down-regulating the expression of ADAMTS-1]. *Xi Bao Yu Fen Zi Mian Yi Xue Za Zhi* **32**, 1636–1640 (2016).
411. Lu, J., Auduong, L., White, E. S. & Yue, X. Up-regulation of heparan sulfate 6-O-sulfation in idiopathic pulmonary fibrosis. *Am. J. Respir. Cell Mol. Biol.* **50**, 106–114 (2014).
412. Mi, S. *et al.* Blocking IL-17A promotes the resolution of pulmonary inflammation and fibrosis via TGF- β 1-dependent and -independent mechanisms. *J. Immunol.* **187**, 3003–3014 (2011).
413. Otsuka, Y., Kedersha, N. L. & Schoenberg, D. R. Identification of a cytoplasmic complex that adds a cap onto 5'-monophosphate RNA. *Mol. Cell. Biol.* **29**, 2155–2167 (2009).
414. Trotman, J. B. & Schoenberg, D. R. A recap of RNA recapping. *Wiley Interdiscip. Rev. RNA* **10**, e1504 (2019).
415. Qiu, M. *et al.* Antibody responses to individual proteins of SARS coronavirus and their neutralization activities. *Microbes Infect.* **7**, 882–889 (2005).
416. Chan, W. S. *et al.* Coronaviral hypothetical and structural proteins were found in the intestinal surface enterocytes and pneumocytes of severe acute respiratory syndrome (SARS). *Mod. Pathol.* **18**, 1432–1439 (2005).
417. Sharma, K. *et al.* SARS-CoV 9b protein diffuses into nucleus, undergoes active Crm1 mediated nucleocytoplasmic export and triggers apoptosis when retained in the nucleus. *PLoS One* **6**, e19436 (2011).
418. Shi, C.-S. *et al.* SARS-coronavirus open reading frame-9b suppresses innate immunity by targeting mitochondria and the MAVS/TRAF3/TRAF6 signalosome. *J. Immunol.* **193**, 3080–3089 (2014).
419. Liu, X.-Y., Wei, B., Shi, H.-X., Shan, Y.-F. & Wang, C. Tom70 mediates activation of interferon regulatory factor 3 on mitochondria. *Cell Res.* **20**, 994–1011 (2010).

420. Wei, B. *et al.* Tom70 mediates Sendai virus-induced apoptosis on mitochondria. *J. Virol.* **89**, 3804–3818 (2015).
421. De Snoo, M. L. *et al.* Bcl-2-associated athanogene 5 (BAG5) regulates Parkin-dependent mitophagy and cell death. *Cell Death Dis.* **10**, 907 (2019).
422. Zhang, L., Qin, Y. & Chen, M. Viral strategies for triggering and manipulating mitophagy. *Autophagy* **14**, 1665–1673 (2018).
423. Vietri, M. *et al.* Spastin and ESCRT-III coordinate mitotic spindle disassembly and nuclear envelope sealing. *Nature* **522**, 231–235 (2015).
424. Morita, E. *et al.* ESCRT-III protein requirements for HIV-1 budding. *Cell Host Microbe* **9**, 235–242 (2011).
425. Bartusch, C. & Prange, R. ESCRT Requirements for Murine Leukemia Virus Release. *Viruses* **8**, 103 (2016).
426. Gu, G. J. *et al.* Role of individual MARK isoforms in phosphorylation of tau at Ser²⁶² in Alzheimer's disease. *Neuromolecular Med.* **15**, 458–469 (2013).
427. Malikov, V. & Naghavi, M. H. Localized Phosphorylation of a Kinesin-1 Adaptor by a Capsid-Associated Kinase Regulates HIV-1 Motility and Uncoating. *Cell Rep.* **20**, 2792–2799 (2017).
428. Inberg, A. & Linial, M. Evolutional insights on uncharacterized SARS coronavirus genes. *FEBS Lett.* **577**, 159–164 (2004).
429. Shukla, A. & Hilgenfeld, R. Acquisition of new protein domains by coronaviruses: analysis of overlapping genes coding for proteins N and 9b in SARS coronavirus. *Virus Genes* **50**, 29–38 (2015).
430. Marra, M. A. *et al.* The Genome sequence of the SARS-associated coronavirus. *Science* **300**, 1399–1404 (2003).
431. Stroud, D. A. *et al.* Accessory subunits are integral for assembly and function of human mitochondrial complex I. *Nature* **538**, 123–126 (2016).
432. Dunning, C. J. R. *et al.* Human CIA30 is involved in the early assembly of mitochondrial complex I and mutations in its gene cause disease. *EMBO J.* **26**, 3227–3237 (2007).
433. Nouws, J. *et al.* Acyl-CoA dehydrogenase 9 is required for the biogenesis of oxidative phosphorylation

- complex I. *Cell Metab.* **12**, 283–294 (2010).
434. Ohishi, K., Inoue, N. & Kinoshita, T. PIG-S and PIG-T, essential for GPI anchor attachment to proteins, form a complex with GAA1 and GPI8. *EMBO J.* **20**, 4088–4098 (2001).
435. Csernok, E. *et al.* Wegener autoantigen induces maturation of dendritic cells and licenses them for Th1 priming via the protease-activated receptor-2 pathway. *Blood* **107**, 4440–4448 (2006).
436. Chiu, L.-L., Perng, D.-W., Yu, C.-H., Su, S.-N. & Chow, L.-P. Mold allergen, pen C 13, induces IL-8 expression in human airway epithelial cells by activating protease-activated receptor 1 and 2. *J. Immunol.* **178**, 5237–5244 (2007).
437. Goon Goh, F. *et al.* G-protein-dependent and -independent pathways regulate proteinase-activated receptor-2 mediated p65 NFκB serine 536 phosphorylation in human keratinocytes. *Cell. Signal.* **20**, 1267–1274 (2008).
438. Rallabhandi, P. *et al.* Analysis of proteinase-activated receptor 2 and TLR4 signal transduction: a novel paradigm for receptor cooperativity. *J. Biol. Chem.* **283**, 24314–24325 (2008).
439. Su, X., Camerer, E., Hamilton, J. R., Coughlin, S. R. & Matthay, M. A. Protease-activated receptor-2 activation induces acute lung inflammation by neuropeptide-dependent mechanisms. *J. Immunol.* **175**, 2598–2605 (2005).
440. Feld, M. *et al.* Agonists of proteinase-activated receptor-2 enhance IFN-γ-inducible effects on human monocytes: role in influenza A infection. *J. Immunol.* **180**, 6903–6910 (2008).
441. Antoniak, S. & Mackman, N. Multiple roles of the coagulation protease cascade during virus infection. *Blood* **123**, 2605–2613 (2014).
442. Allen, I. C. *et al.* NLRX1 protein attenuates inflammatory responses to infection by interfering with the RIG-I-MAVS and TRAF6-NF-κB signaling pathways. *Immunity* **34**, 854–865 (2011).
443. Cole, S. P. C. Multidrug resistance protein 1 (MRP1, ABCC1), a ‘multitasking’ ATP-binding cassette (ABC) transporter. *J. Biol. Chem.* **289**, 30880–30888 (2014).
444. Jorajuria, S., Dereuddre-Bosquet, N., Naissant-Storck, K., Dormont, D. & Clayette, P. Differential expression levels of MRP1, MRP4, and MRP5 in response to human immunodeficiency virus infection in

- human macrophages. *Antimicrob. Agents Chemother.* **48**, 1889–1891 (2004).
445. He, H. *et al.* Polarized macrophage subsets differentially express the drug efflux transporters MRP1 and BCRP, resulting in altered HIV production. *Antivir. Chem. Chemother.* **26**, 2040206617745168 (2018).
446. Weekes, M. P. *et al.* Latency-associated degradation of the MRP1 drug transporter during latent human cytomegalovirus infection. *Science* **340**, 199–202 (2013).
447. Srinivas, R. V., Middlemas, D., Flynn, P. & Fridland, A. Human immunodeficiency virus protease inhibitors serve as substrates for multidrug transporter proteins MDR1 and MRP1 but retain antiviral efficacy in cell lines expressing these transporters. *Antimicrob. Agents Chemother.* **42**, 3157–3162 (1998).
448. Young, L. C., Campling, B. G., Cole, S. P., Deeley, R. G. & Gerlach, J. H. Multidrug resistance proteins MRP3, MRP1, and MRP2 in lung cancer: correlation of protein levels with drug response and messenger RNA levels. *Clin. Cancer Res.* **7**, 1798–1804 (2001).
449. Okamura, T., Kikuchi, T., Okada, M., Wakizaka, H. & Zhang, M.-R. Imaging of activity of multidrug resistance-associated protein 1 in the lungs. *Am. J. Respir. Cell Mol. Biol.* **49**, 335–340 (2013).
450. Budulac, S. E. *et al.* Multidrug resistance-associated protein-1 (MRP1) genetic variants, MRP1 protein levels and severity of COPD. *Respir. Res.* **11**, 60 (2010).
451. Zukauskas, A. *et al.* Transporters MRP1 and MRP2 Regulate Opposing Inflammatory Signals To Control Transepithelial Neutrophil Migration during *Streptococcus pneumoniae* Lung Infection. *mSphere* **3**, (2018).
452. Odon, V., Georgana, I., Holley, J., Morata, J. & Maluquer de Motes, C. Novel Class of Viral Ankyrin Proteins Targeting the Host E3 Ubiquitin Ligase Cullin-2. *J. Virol.* **92**, (2018).
453. Westrich, J. A. *et al.* Human Papillomavirus 16 E7 Stabilizes APOBEC3A Protein by Inhibiting Cullin 2-Dependent Protein Degradation. *J. Virol.* **92**, (2018).
454. Timms, R. T. *et al.* A glycine-specific N-degron pathway mediates the quality control of protein N-myristoylation. *Science* **365**, (2019).
455. El-Gebali, S. *et al.* The Pfam protein families database in 2019. *Nucleic Acids Res.* **47**, D427–D432 (2019).
456. Melé, M., Ferreira, P. G., Reverter, F. & DeLuca, D. S. The human transcriptome across tissues and

individuals. (2015).

457. Goetz, D. H. *et al.* Substrate specificity profiling and identification of a new class of inhibitor for the major protease of the SARS coronavirus. *Biochemistry* **46**, 8744–8752 (2007).
458. Kiemer, L., Lund, O., Brunak, S. & Blom, N. Coronavirus 3CLpro proteinase cleavage sites: possible relevance to SARS virus pathology. *BMC Bioinformatics* **5**, 72 (2004).
459. Johnson, M. *et al.* NCBI BLAST: a better web interface. *Nucleic Acids Res.* **36**, W5–9 (2008).
460. Brown, N. P., Leroy, C. & Sander, C. MView: a web-compatible database search or multiple alignment viewer. *Bioinformatics* **14**, 380–381 (1998).
461. Crooks, G. E., Hon, G., Chandonia, J.-M. & Brenner, S. E. WebLogo: a sequence logo generator. *Genome Res.* **14**, 1188–1190 (2004).
462. Jensen, N. H. *et al.* N-desalkylquetiapine, a potent norepinephrine reuptake inhibitor and partial 5-HT_{1A} agonist, as a putative mediator of quetiapine's antidepressant activity. *Neuropsychopharmacology* **33**, 2303–2312 (2008).
463. Altschul, S. F. *et al.* Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res.* **25**, 3389–3402 (1997).
464. Ranwez, V., Douzery, E. J. P., Cambon, C., Chantret, N. & Delsuc, F. MACSE v2: Toolkit for the Alignment of Coding Sequences Accounting for Frameshifts and Stop Codons. *Mol. Biol. Evol.* **35**, 2582–2584 (2018).
465. Guindon, S. *et al.* New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. *Syst. Biol.* **59**, 307–321 (2010).
466. Yang, Z. PAML 4: phylogenetic analysis by maximum likelihood. *Mol. Biol. Evol.* **24**, 1586–1591 (2007).
467. Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *J. R. Stat. Soc. Series B Stat. Methodol.* **57**, 289–300 (1995).