

Price of Anarchy for Mean Field Games

René Carmona*

Christy V. Graves[†]

Zongjun Tan[‡]

July 21, 2018

Abstract

The price of anarchy, originally introduced to quantify the inefficiency of selfish behavior in routing games, is extended to mean field games. The price of anarchy is defined as the ratio of a worst case social cost computed for a mean field game equilibrium to the optimal social cost as computed by a central planner. We illustrate properties of such a price of anarchy on linear quadratic extended mean field games, for which explicit computations are possible. Various asymptotic behaviors of the price of anarchy are proved for limiting behaviors of the coefficients in the model and numerics are presented.

1 Introduction

The concept of the ‘price of anarchy’ was introduced to quantify the inefficiency of selfish behavior in finite player games [8][9][12][17][18][19]. In this report, we extend the notion of price of anarchy to mean field games (MFG). Mean field games were introduced by Lasry and Lions [13] and Caines and his collaborators [11] to describe the limiting regime of large symmetric games when the number of players, N , tends to infinity. A mean field game equilibrium characterizes the analogue of a Nash equilibrium in the $N = \infty$ regime. Thus, as in the finite player case, it is possible that the mean field game equilibrium is inefficient. In fact, in the paper of Balandat and Tomlin [2], they present a numerical example that shows that mean field game equilibria are not efficient, in general. The suboptimality of a mean field game equilibrium is also illustrated numerically for a congestion model in a paper of Achdou and Laurière [1]. More recently Cardaliaguet and Rainer gave in [4] a partial differential equation based thorough analysis of the *(in)efficiency* of the mean field game equilibria.

In this report, the goal is to define the price of anarchy in the context of mean field games, and to compute it for a class of linear quadratic mean field game models, which can be solved explicitly. In fact, we consider an even more general class of games by allowing for interaction between the players through their controls, in addition to interaction through their states. This is often referred in the literature as extended mean field game, or mean field game of control. We

*Operations Research and Financial Engineering, Princeton University, Partially supported by NSF #DMS-1716673 and ARO #W911NF-17-1-0578

[†]Program in Applied and Computational Mathematics, Princeton University, Partially supported by NSF #DMS-1515753 and NSF GRFP

[‡]Operations Research and Financial Engineering, Princeton University, Partially supported by NSF #DMS-1515753

compare the social cost of a mean field game equilibrium to the cost incurred when the players execute a strategy computed centrally.

We consider a system of N players whose private states are denoted at time t by $X_t^1, X_t^2, \dots, X_t^N$. To keep the presentations simple, we assume the state space is \mathbb{R} . We denote by μ_t^N the empirical distribution of the states, namely

$$\mu_t^N = \frac{1}{N} \sum_{i=1}^N \delta_{X_t^i}.$$

We assume that these states evolve in continuous time under the influences of controls $\alpha_t^1, \alpha_t^2, \dots, \alpha_t^N \in \mathbb{A}$, where the set of admissible controls, \mathbb{A} , will be defined later. Let ν_t^N denote the empirical measure of the controls:

$$\nu_t^N = \frac{1}{N} \sum_{i=1}^N \delta_{\alpha_t^i}.$$

We also assume that if and when interactions between these states and controls are present, they are of a mean field type, i.e. through μ_t^N and ν_t^N . The time evolution of the state for player i is given by the Itô dynamics:

$$dX_t^i = b(t, X_t^i, \mu_t^N, \alpha_t^i, \nu_t^N)dt + \sigma dW_t.$$

We work over the interval $[0, T]$ limited by a finite time horizon $T \in \mathbb{R}^+$. We assume the drift function $b : [0, T] \times \mathbb{R} \times \mathcal{P}(\mathbb{R}) \times \mathbb{A} \times \mathcal{P}(\mathbb{A}) \ni (t, x, \mu, \alpha, \nu) \rightarrow \mathbb{R}$ is Lipschitz in each of its inputs. For the sake of simplicity, we assume that the volatility, σ , is a positive constant.

Cost Functionals

We assume that we are given two functions $f : [0, T] \times \mathbb{R} \times \mathcal{P}(\mathbb{R}) \times \mathbb{A} \times \mathcal{P}(\mathbb{A}) \ni (t, x, \mu, \alpha, \nu) \rightarrow \mathbb{R}$ and $g : \mathbb{R} \times \mathcal{P}(\mathbb{R}) \ni (x, \mu) \rightarrow \mathbb{R}$ which we call running and terminal cost functions, respectively. We assume f and g are Lipschitz in each of their arguments. The goal of player i is to minimize its expected cost as given by:

$$J^i(\alpha^1, \dots, \alpha^N) = \mathbb{E} \left[\int_0^T f(t, X_t^i, \mu_t^N, \alpha_t^i, \nu_t^N) dt + g(X_T^i, \mu_T^N) \right].$$

Social Cost

We restrict ourselves to Markovian control strategies $\alpha = (\alpha_t)_{0 \leq t \leq T}$ given by feedback functions in the form $\alpha_t = \phi(t, X_t)$ and we let \mathbb{A} denote the set of such controls. If the N players use distributed Markovian control strategies of the form $\alpha_t^i = \phi(t, X_t^i)$, we define the cost (per player) to the system as the quantity $J_\phi^{(N)}$:

$$J_\phi^{(N)} = \frac{1}{N} \sum_{i=1}^N J^i(\alpha^1, \dots, \alpha^N).$$

We shall compute this social cost in the limit $N \rightarrow \infty$ when all the players use the distributed control strategies given by the same feedback function ϕ identified by solving an optimization

problem in the limit $N \rightarrow \infty$. We take the social cost to be the limit as $N \rightarrow \infty$ of $J_\phi^{(N)}$, namely

$$\begin{aligned} \lim_{N \rightarrow \infty} J_\phi^{(N)} &= \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{i=1}^N J^i(\alpha^1, \dots, \alpha^N) \\ &= \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{i=1}^N \mathbb{E} \left[\int_0^T f(t, X_t^i, \mu_t^N, \phi(t, X_t^i), \nu_t^N) dt + g(X_T^i, \mu_T^N) \right], \\ &= \lim_{N \rightarrow \infty} \mathbb{E} \left[\int_0^T \langle f(t, \cdot, \mu_{X_t}^N, \phi(t, \cdot), \nu_t^N), \mu_t^N \rangle dt + \langle g(\cdot, \mu_{X_T}^N), \mu_T^N \rangle \right], \end{aligned}$$

if we use the notation $\langle \varphi, \rho \rangle$ for the integral $\int \varphi(z) \rho(dz)$ of the function φ with respect to the measure ρ . Now if we assume that in the limit $N \rightarrow \infty$ the empirical distributions μ_t^N converge toward a measure μ_t , and thus $\nu_t^N = \frac{1}{N} \sum_{i=1}^N \delta_{\phi(t, X_t^i)}$ also converges toward a measure ν_t , then the social cost of the feedback function ϕ becomes:

$$SC(\phi) = \int_0^T \langle f(t, \cdot, \mu_t, \phi(t, \cdot), \nu_t), \mu_t \rangle dt + \langle g(\cdot, \mu_T), \mu_T \rangle$$

with the expectation, \mathbb{E} , disappearing when the limiting flows $\boldsymbol{\mu} = (\mu_t)_{0 \leq t \leq T}$ and $\boldsymbol{\nu} = (\nu_t)_{0 \leq t \leq T}$ are deterministic.

We would like to evaluate $SC(\phi)$ in the $N = \infty$ regime directly, without having to construct the deterministic measure flows $\boldsymbol{\mu}$ and $\boldsymbol{\nu}$ as limits of the finite player empirical measures. To do this, we assume that propagation of chaos holds and that the states of the N players become asymptotically independent in the limit as $N \rightarrow \infty$. We consider a representative agent whose state is given by $\mathbf{X}^\phi = (X_t^\phi)_{0 \leq t \leq T}$, the continuous time solution of the stochastic differential equation of McKean-Vlasov type:

$$dX_t^\phi = b(t, X_t^\phi, \mathcal{L}(X_t^\phi), \phi(t, X_t^\phi), \mathcal{L}(\phi(t, X_t^\phi))) dt + \sigma dW_t \quad (1)$$

controlled by ϕ . Then we can identify $\boldsymbol{\mu}$ as the law of a representative agent using the feedback function ϕ , i.e. $\mu_t = \mathcal{L}(X_t^\phi)$, and similarly, we can identify $\boldsymbol{\nu}$ as the law of the control, such that $\nu_t = \mathcal{L}(\phi(t, X_t^\phi))$. Thus, in the $N = \infty$ regime, we rewrite the social cost as

$$SC(\phi) = \int_0^T \langle f(t, \cdot, \mathcal{L}(X_t^\phi), \phi(t, \cdot), \mathcal{L}(\phi(t, X_t^\phi))), \mathcal{L}(X_t^\phi) \rangle dt + \langle g(\cdot, \mathcal{L}(X_T^\phi)), \mathcal{L}(X_T^\phi) \rangle$$

where \mathbf{X}^ϕ satisfies equation (1). For the remainder of the paper, we work in the $N = \infty$ regime. As mentioned earlier, ϕ should be identified by solving an optimal control problem. We consider two distinct problems:

- ϕ is a feedback function providing a mean field game equilibrium. We detail more precisely what is meant by ϕ providing a mean field game equilibrium in Section 1.1.
- ϕ is the feedback function minimizing the social cost $SC(\phi)$, without having to be a mean field game equilibrium, in which case we use the notation SC^{MKV} for $SC(\phi)$. This is a control problem of McKean-Vlasov type, which is detailed more precisely in Section 1.2.

The two problems are detailed more precisely in Sections 1.1 and 1.2. In Section 1.3, we define the price of anarchy based on these two problem formulations. The class of linear quadratic models is explored in Section 2, where we provide some theoretical results on the price of anarchy for this class of games, show numerical results, and detail a particular example of flocking. We conclude in Section 3.

1.1 Nash Equilibrium: Mean Field Game Formulation

The goal of this subsection is to articulate what is meant by a feedback function providing a mean field game equilibrium. To begin, we define what we call the *mean field environment*. By symmetry of the players, we suppose all of the players in the mean field game use the same feedback function, ϕ . Then the *mean field environment* specified by ϕ is characterized by $\mathcal{L}(X_t^\phi)_{0 \leq t \leq T}$ and $\mathcal{L}(\phi(t, X_t^\phi))_{0 \leq t \leq T}$ where the dynamics of $(X_t^\phi)_{0 \leq t \leq T}$ are given by equation (1). Since we search for a Nash equilibrium, we consider a representative agent who wishes to find their best response, ϕ' , to the mean field environment specified by ϕ , in which case their state is given by $\mathbf{X}^{\phi', \phi} = (X_t^{\phi', \phi})_{0 \leq t \leq T}$ solving the standard stochastic differential equation:

$$dX_t^{\phi', \phi} = b(t, X_t^{\phi', \phi}, \mathcal{L}(X_t^\phi), \phi'(t, X_t^{\phi', \phi}), \mathcal{L}(\phi(t, X_t^\phi)))dt + \sigma dW_t.$$

Consider the function:

$$\mathcal{S}(\phi', \phi) = \left[\int_0^T \langle f(t, \cdot, \mathcal{L}(X_t^\phi), \phi'(t, \cdot), \mathcal{L}(\phi(t, X_t^\phi))), \mathcal{L}(X_t^{\phi', \phi}) \rangle dt + \langle g(\cdot, \mathcal{L}(X_T^\phi)), \mathcal{L}(X_T^{\phi', \phi}) \rangle \right].$$

The best response for the representative agent in the mean field environment specified by ϕ is the feedback function minimizing this cost, namely $\phi^* = \arg \inf_{\phi'} \mathcal{S}(\phi', \phi)$. Assuming the minimizer is unique (which will be the case for the models we consider), this defines a mapping $\Phi : \phi \rightarrow \phi^*$. If there is a $\hat{\phi}$ such that $\Phi(\hat{\phi}) = \hat{\phi}$, then the players are in a mean field game equilibrium.

Thus, the search for a feedback function providing a mean field game equilibrium can be summarized as the following set of two successive steps:

1. For each feedback function $\phi : [0, T] \times \mathbb{R} \ni (t, x) \rightarrow \mathbb{R}$, solve the optimal control problem

$$\phi^* = \arg \inf_{\phi'} \mathcal{S}(\phi', \phi).$$

Define the mapping $\Phi(\phi) := \phi^*$.

2. Find a fixed point $\hat{\phi}$ of Φ such that $\Phi(\hat{\phi}) = \hat{\phi}$.

When these two steps can be taken successfully, we say that $\hat{\phi}$ provides a mean field game equilibrium. Note that $\mathbf{X}^{\hat{\phi}, \hat{\phi}} = \mathbf{X}^{\hat{\phi}}$ and therefore $\mathcal{S}(\hat{\phi}, \hat{\phi}) = SC(\hat{\phi})$ gives the social cost for the mean field game equilibrium provided by $\hat{\phi}$. Notice that there could possibly be many feedback functions providing a mean field game equilibrium. Let \mathcal{N} denote the set of all such feedback functions providing mean field game equilibria, as detailed above, i.e.

$$\mathcal{N} = \{\phi : [0, T] \times \mathbb{R} \ni (t, x) \rightarrow \mathbb{R} \mid \Phi(\phi) = \phi\}.$$

1.2 Centralized Control: Optimal Control of McKean-Vlasov Type

The goal of this subsection is to articulate how to compute the cost associated with the control problem of McKean-Vlasov type, SC^{MKV} . The central planner considers the following control problem:

$$\begin{aligned} \hat{\phi} &= \arg \inf_{\phi} SC(\phi) \\ &= \arg \inf_{\phi} \left[\int_0^T \langle f(t, \cdot, \mathcal{L}(X_t^\phi), \phi(t, \cdot), \mathcal{L}(\phi(t, X_t^\phi))), \mathcal{L}(X_t^\phi) \rangle dt + \langle g(\cdot, \mathcal{L}(X_T^\phi)), \mathcal{L}(X_T^\phi) \rangle \right]. \end{aligned}$$

Thus, the cost of the solution to the optimal control problem of McKean-Vlasov is given by

$$SC^{MKV} = SC(\hat{\phi}).$$

Remark 1. *We are not concerned with uniqueness for the control of McKean-Vlasov type problem, because $SC^{MKV} = SC(\phi_1) = SC(\phi_2)$ is still well defined even if there are two different optimal feedback functions ϕ_1 and ϕ_2 minimizing $SC(\phi)$.*

1.3 Price of Anarchy

We have described two approaches to compute the optimal feedback function ϕ . In the mean field game formulation, we require $\phi \in \mathcal{N}$, where \mathcal{N} denotes the set of feedback functions providing mean field game equilibria. In the optimal control of McKean-Vlasov type formulation, the optimal control to be adopted by all players is computed by a central planner, who optimizes the social cost function $SC(\phi)$ directly. Thus, we necessarily have:

$$SC^{MKV} \leq SC(\phi), \quad \forall \phi \in \mathcal{N}.$$

In other words, there is a ‘price of anarchy’ associated with allowing players to choose their controls selfishly. We thus define the price of anarchy (denoted PoA) as the ratio between the worst case cost for a mean field game equilibrium and the optimal cost computed by a central planner:

$$PoA = \frac{\sup_{\phi \in \mathcal{N}} SC(\phi)}{SC^{MKV}}.$$

2 Price of Anarchy for Linear Quadratic Extended Mean Field Games

The class of linear quadratic extended mean field games is a class of problems for which explicit solutions can be computed analytically, and thus, we can compute the price of anarchy explicitly. To the best of our knowledge, the case of linear quadratic extended mean field games has not been explored in the literature, as well as computing the price of anarchy for this class of games.

To begin, we need to describe in more detail the two problems that will be used to compute the price of anarchy: the linear quadratic extended mean field game, and the linear quadratic control problem of McKean-Vlasov type with dependence on the law of the control. To specify the problems, we only need to specify the drift and cost functions, b , f , and g introduced in Section 1. For the linear quadratic models, we take the drift to be linear:

$$b(t, x, \mu, \alpha, \nu) = b_1(t)x + \bar{b}_1(t)\bar{\mu} + b_2(t)\alpha + \bar{b}_2(t)\bar{\nu},$$

where $\bar{\mu}$ denotes the mean of the measure μ , namely, $\bar{\mu} = \int_{\mathbb{R}} x d\mu(x)$, and similarly for $\bar{\nu}$. We take the running and terminal costs to be quadratic:

$$\begin{aligned} f(t, x, \mu, \alpha, \nu) &= \frac{1}{2} \left(q(t)x^2 + \bar{q}(t)(x - s(t)\bar{\mu})^2 + r(t)\alpha^2 + \bar{r}(t)(\alpha - \bar{s}(t)\bar{\nu})^2 \right), \\ g(x, \mu) &= \frac{1}{2} \left(q_T x^2 + \bar{q}_T (x - s_T \bar{\mu})^2 \right). \end{aligned}$$

Remark 2. *If $\bar{b}_2(t) \equiv 0$ and $\bar{r}(t) \equiv 0$, then we have the standard mean field game or control problem of McKean-Vlasov type. (See Theorem 1 for assumptions that provide existence and uniqueness.)*

2.1 Linear Quadratic Extended Mean Field Games

To solve the linear quadratic extended mean field game (LQEMFG), we begin by considering the reduced Hamiltonian for this problem:

$$H(t, x, \bar{\mu}, \alpha, \bar{\nu}, y) = (b_1(t)x + \bar{b}_1(t)\bar{\mu} + b_2(t)\alpha + \bar{b}_2(t)\bar{\nu})y \\ + \frac{1}{2} (q(t)x^2 + \bar{q}(t)(x - s(t)\bar{\mu})^2 + r(t)\alpha^2 + \bar{r}(t)(\alpha - \bar{s}(t)\bar{\nu})^2),$$

and whenever the flows $\bar{\mu} = (\bar{\mu}_t)_{0 \leq t \leq T}$ and $\bar{\nu} = (\bar{\nu}_t)_{0 \leq t \leq T}$ are fixed, we consider for each control process $\alpha = (\alpha_t)_{0 \leq t \leq T}$ the adjoint equation:

$$dY_t = -\partial_x H(t, X_t, \bar{\mu}_t, \alpha_t, \bar{\nu}_t, Y_t)dt + Z_t dW_t \\ Y_T = \partial_x g(X_T, \mathcal{L}(X_T)).$$

According to the Pontryagin stochastic maximum principle, a sufficient condition for optimality is $\partial_\alpha H(t, X_t, \bar{\mu}_t, \hat{\alpha}_t, \bar{\nu}_t, y) = 0$. We introduce the function:

$$\hat{\alpha}(t, x, \bar{\mu}, \bar{\nu}, y) = \frac{\bar{r}(t)\bar{s}(t)\bar{\nu} - b_2(t)y}{r(t) + \bar{r}(t)},$$

and use the control $\hat{\alpha}_t = \hat{\alpha}(t, X_t, \bar{\mu}, \bar{\nu}, Y_t)$. When solving the fixed point step, we identify $\bar{\nu}_t = \mathbb{E}(\hat{\alpha}_t)$. By taking the expectation, we find:

$$\mathbb{E}(\hat{\alpha}_t) = c^{MFG}(t)\mathbb{E}(Y_t)$$

with:

$$c^{MFG}(t) = -\frac{b_2(t)}{r(t) + \bar{r}(t)(1 - \bar{s}(t))}.$$

Thus, necessarily we must have:

$$\hat{\alpha}_t = a^{MFG}(t)Y_t + b^{MFG}(t)\mathbb{E}(Y_t)$$

with:

$$a^{MFG}(t) = -\frac{b_2(t)}{r(t) + \bar{r}(t)},$$

and:

$$b^{MFG}(t) = -\frac{\bar{r}(t)\bar{s}(t)b_2(t)}{(r(t) + \bar{r}(t))(r(t) + \bar{r}(t)(1 - \bar{s}(t)))}.$$

Note that $c^{MFG}(t) = a^{MFG}(t) + b^{MFG}(t)$. The solution of the mean field game equilibrium problem is given by the solution to the FBSDE system:

$$dX_t = (b_1(t)X_t + \bar{b}_1(t)\mathbb{E}X_t + a^{MFG}(t)b_2(t)Y_t + (b^{MFG}(t)b_2(t) + c^{MFG}(t)\bar{b}_2(t))\mathbb{E}Y_t) dt + \sigma dW_t \\ dY_t = -((q(t) + \bar{q}(t))X_t - \bar{q}(t)s(t)\mathbb{E}X_t + b_1(t)Y_t) dt + Z_t dW_t \quad (2)$$

with initial condition $X_0 = \xi$, a random variable with finite mean and variance, and terminal condition $Y_T = (q_T + \bar{q}_T)X_T - \bar{q}_T s_T \mathbb{E}X_T$.

This is a linear FBSDE of McKean-Vlasov type, which can be solved explicitly under mild assumptions (or at least in the case of time-independent coefficients which we will consider later. See Appendix A). Let $\bar{\eta}_t^{MFG}$, η_t^{MFG} , \bar{x}_t^{MFG} , and v_t^{MFG} denote the solutions for this problem as described in the appendix so that $Y_t = \eta_t^{MFG} X_t + (\bar{\eta}_t^{MFG} - \eta_t^{MFG}) \bar{x}_t^{MFG}$, $\mathbb{E}(Y_t) = \bar{\eta}_t^{MFG} \bar{x}_t^{MFG}$, $\mathbb{E}(X_t) = \bar{x}_t^{MFG}$, and $Var(X_t) = v_t^{MFG}$ provide a solution to the LQEMFG problem. Then from the appendix, we have:

$$\begin{aligned} \dot{\bar{\eta}}_t^{MFG} + c^{MFG}(t)(b_2(t) + \bar{b}_2(t))(\bar{\eta}_t^{MFG})^2 + (2b_1(t) + \bar{b}_1(t))\bar{\eta}_t^{MFG} + q(t) + \bar{q}(t)(1 - s(t)) &= 0, \\ \bar{\eta}_T^{MFG} - (q_T + \bar{q}_T(1 - s_T)) &= 0, \end{aligned} \quad (3)$$

$$\begin{aligned} \dot{\eta}_t^{MFG} + a^{MFG}(t)b_2(t)(\eta_t^{MFG})^2 + 2b_1(t)\eta_t^{MFG} + q(t) + \bar{q}(t) &= 0, \\ \eta_T^{MFG} - (q_T + \bar{q}_T) &= 0, \end{aligned} \quad (4)$$

$$\begin{aligned} \dot{\bar{x}}_t^{MFG} &= (b_1(t) + \bar{b}_1(t) + c^{MFG}(t)(b_2(t) + \bar{b}_2(t))\bar{\eta}_t^{MFG})\bar{x}_t^{MFG}, \\ \bar{x}_0^{MFG} &= \mathbb{E}(\xi). \end{aligned} \quad (5)$$

And thus,

$$\bar{x}_t^{MFG} = \mathbb{E}(\xi) e^{\int_0^t (b_1(u) + \bar{b}_1(u) + c^{MFG}(u)(b_2(u) + \bar{b}_2(u))\bar{\eta}_u^{MFG}) du}, \quad (6)$$

$$v_t^{MFG} = Var(\xi) e^{\int_0^t 2(b_1(s) + a^{MFG}(s)b_2(s)\eta_s^{MFG}) ds} + \sigma^2 \int_0^t e^{2 \int_s^t (b_1(u) + a^{MFG}(u)b_2(u)\eta_u^{MFG}) du} ds. \quad (7)$$

Let $SC^{MFG} := SC(\phi)$ for the feedback function specified by this solution, i.e.

$$\phi(t, x) = a^{MFG}(t)\eta_t^{MFG}x + (a^{MFG}(t)(\bar{\eta}_t^{MFG} - \eta_t^{MFG}) + b^{MFG}(t)\bar{\eta}_t^{MFG})\bar{x}_t^{MFG}.$$

Then we can compute the social cost as described in Section 1.1:

$$\begin{aligned} SC^{MFG} &= \frac{1}{2}[(q_T + \bar{q}_T)v_T^{MFG} + (q_T + \bar{q}_T(1 - s_T)^2)(\bar{x}_T^{MFG})^2 \\ &\quad + \int_0^T (q(t) + \bar{q}(t) + (r(t) + \bar{r}(t))(a^{MFG}(t)\eta_t^{MFG})^2)v_t^{MFG} \\ &\quad + (q(t) + \bar{q}(t)(1 - s(t))^2 + (r(t) + \bar{r}(t)(1 - \bar{s}(t))^2)(c^{MFG}(t)\bar{\eta}_t^{MFG})^2)(\bar{x}_t^{MFG})^2 dt], \end{aligned}$$

where we have used the fact that:

$$\mathbb{E}(\phi(t, X_t)) = c^{MFG}(t)\bar{\eta}_t^{MFG}\bar{x}_t^{MFG},$$

and:

$$Var(\phi(t, X_t)) = (a^{MFG}(t)\eta_t^{MFG})^2 v_t^{MFG}.$$

2.2 Linear Quadratic Control of McKean-Vlasov Type Involving the Law of the Control

To solve the linear quadratic optimal control problem of McKean-Vlasov type involving the law of the control (LQEMKV), we begin with the reduced Hamiltonian, which is the same as in the LQEMFG problem:

$$\begin{aligned} H(t, x, \bar{\mu}, \alpha, \bar{\nu}, y) &= (b_1(t)x + \bar{b}_1(t)\bar{\mu} + b_2(t)\alpha + \bar{b}_2(t)\bar{\nu})y \\ &\quad + \frac{1}{2} (q(t)x^2 + \bar{q}(t)(x - s(t)\bar{\mu})^2 + r(t)\alpha^2 + \bar{r}(t)(\alpha - \bar{s}(t)\bar{\nu})^2). \end{aligned}$$

Since we require $\bar{\nu}_t$ to be equal to $\mathbb{E}(\alpha_t)$ throughout the optimization, it is not sufficient to minimize the Hamiltonian with respect to the α input alone in order to guarantee optimality. A sufficient condition for control problems of McKean-Vlasov type involving the law of the control is derived in [5]. Since we consider a Hamiltonian that depends on the means of $\bar{\mu}$ and $\bar{\nu}$ instead of the full distributions, the sufficient condition is the following: $\hat{\alpha}(t, X_t, \bar{\mu}, \bar{\nu}, Y_t)$ should satisfy:

$$\partial_\alpha H(t, X_t, \mathbb{E}(X_t), \hat{\alpha}_t, \mathbb{E}(\hat{\alpha}_t), Y_t) + \tilde{\mathbb{E}} \left[\partial_{\bar{\nu}} H(t, \tilde{X}_t, \mathbb{E}(X_t), \hat{\alpha}_t, \mathbb{E}(\hat{\alpha}_t), \tilde{Y}_t) \right] = 0,$$

where the adjoint equation is given by:

$$\begin{aligned} dY_t &= - \left[\partial_x H(t, X_t, \bar{\mu}_t, \alpha_t, \bar{\nu}_t, Y_t) + \tilde{\mathbb{E}} \left[\partial_{\bar{\mu}} H(t, \tilde{X}_t, \bar{\mu}_t, \tilde{\alpha}_t, \bar{\nu}_t, \tilde{Y}_t) \right] \right] dt + Z_t dW_t \\ Y_T &= \partial_x g(X_T, \mathcal{L}(X_T)) + \tilde{\mathbb{E}} \left[\partial_{\bar{\mu}} g(\tilde{X}_T, \mathcal{L}(X_T))(X_T) \right], \end{aligned}$$

and $(\tilde{X}, \tilde{Y}, \tilde{\alpha})$ denotes an independent copy of (X, Y, α) . In the present LQ case, the sufficient condition can be used to solve for:

$$\hat{\alpha}_t = a^{MKV}(t)Y_t + b^{MKV}(t)\mathbb{E}(Y_t),$$

and:

$$\mathbb{E}(\hat{\alpha}_t) = c^{MKV}(t)\mathbb{E}(Y_t),$$

with:

$$\begin{aligned} a^{MKV}(t) &= -\frac{b_2(t)}{r(t) + \bar{r}(t)} \\ b^{MKV}(t) &= -\frac{1}{r(t) + \bar{r}(t)} \left(\bar{b}_2(t) - \frac{\bar{r}(t)\bar{s}(t)(\bar{s}(t) - 2)(b_2(t) + \bar{b}_2(t))}{r(t) + \bar{r}(t)(1 - \bar{s}(t))^2} \right) \\ c^{MKV}(t) &= -\frac{b_2(t) + \bar{b}_2(t)}{r(t) + \bar{r}(t)(1 - \bar{s}(t))^2}. \end{aligned}$$

So the solution of the optimal control problem of McKean-Vlasov type is given by the solution to the FBSDE system:

$$\begin{aligned} dX_t &= (b_1(t)X_t + \bar{b}_1(t)\mathbb{E}X_t + a^{MKV}(t)b_2(t)Y_t + (b^{MKV}(t)b_2(t) + c^{MKV}(t)\bar{b}_2(t))\mathbb{E}Y_t) dt + \sigma dW_t \\ dY_t &= -((q(t) + \bar{q}(t))X_t + s(t)\bar{q}(t)(s(t) - 2)\mathbb{E}X_t + b_1(t)Y_t + \bar{b}_1(t)\mathbb{E}Y_t) dt + Z_t dW_t \end{aligned} \tag{8}$$

with initial condition $X_0 = \xi$, and terminal condition $Y_T = (q_T + \bar{q}_T)X_T + s_T\bar{q}_T(s_T - 2)\mathbb{E}X_T$.

As in the previous section, this is a linear FBSDE of McKean-Vlasov type, which can be solved explicitly under mild assumptions (or at least in the case of time-independent coefficients which we will consider later. See Appendix A). Let $\bar{\eta}_t^{MKV}$, η_t^{MKV} , \bar{x}_t^{MKV} , and v_t^{MKV} so that $Y_t = \eta_t^{MKV}X_t + (\bar{\eta}_t^{MKV} - \eta_t^{MKV})\bar{x}_t^{MKV}$, $\mathbb{E}(Y_t) = \bar{\eta}_t^{MKV}\bar{x}_t^{MKV}$, $\mathbb{E}(X_t) = \bar{x}_t^{MKV}$, and $Var(X_t) = v_t^{MKV}$ provide a solution to the LQEMKV problem. Then from the appendix, we have:

$$\begin{aligned} \dot{\bar{\eta}}_t^{MKV} + c^{MKV}(t)(b_2(t) + \bar{b}_2(t))(\bar{\eta}_t^{MKV})^2 + 2(b_1(t) + \bar{b}_1(t))\bar{\eta}_t^{MKV} + q(t) + \bar{q}(t)(1 - s(t))^2 &= 0, \\ \bar{\eta}_T^{MKV} - (q_T + \bar{q}_T(1 - s_T))^2 &= 0, \end{aligned} \tag{9}$$

$$\begin{aligned}\dot{\eta}_t^{MKV} + a^{MKV}(t)b_2(t)(\eta_t^{MKV})^2 + 2b_1(t)\eta_t^{MKV} + q(t) + \bar{q}(t) &= 0, \\ \eta_T^{MKV} - (q_T + \bar{q}_T) &= 0,\end{aligned}\tag{10}$$

$$\begin{aligned}\dot{\bar{x}}_t^{MKV} &= (b_1(t) + \bar{b}_1(t) + c^{MKV}(t)(b_2(t) + \bar{b}_2(t))\bar{\eta}_t^{MKV})\bar{x}_t^{MKV}, \\ \bar{x}_0^{MKV} &= \mathbb{E}(\xi).\end{aligned}\tag{11}$$

And thus,

$$\bar{x}_t^{MKV} = \mathbb{E}(\xi)e^{\int_0^t (b_1(u) + \bar{b}_1(u) + c^{MKV}(u)(b_2(u) + \bar{b}_2(u))\bar{\eta}_u^{MKV})du},\tag{12}$$

$$v_t^{MKV} = Var(\xi)e^{\int_0^t 2(b_1(s) + a^{MKV}(s)b_2(s)\eta_s^{MKV})ds} + \sigma^2 \int_0^t e^{2\int_s^t (b_1(u) + a^{MKV}(u)b_2(u)\eta_u^{MKV})du} ds.\tag{13}$$

Then $SC^{MKV} = SC(\phi)$ where ϕ is the feedback function specified by this solution, i.e.

$$\phi(t, x) = a^{MKV}(t)\eta_t^{MKV}x + (a^{MKV}(t)(\bar{\eta}_t^{MKV} - \eta_t^{MKV}) + b^{MKV}(t)\bar{\eta}_t^{MKV})\bar{x}_t^{MKV}.$$

Then we can compute the social cost, denoted SC^{MKV} , as described in Section 1.2:

$$\begin{aligned}SC^{MKV} &= \frac{1}{2}[(q_T + \bar{q}_T)v_T^{MKV} + (q_T + \bar{q}_T(1 - s_T)^2)(\bar{x}_T^{MKV})^2 \\ &\quad + \int_0^T (q(t) + \bar{q}(t) + (r(t) + \bar{r}(t))(a^{MKV}(t)\eta_t^{MKV})^2)v_t^{MKV}dt \\ &\quad + \int_0^T (q(t) + \bar{q}(t)(1 - s(t))^2 + (r(t) + \bar{r}(t)(1 - \bar{s}(t))^2)(c^{MKV}(t)\bar{\eta}_t^{MKV})^2)(\bar{x}_t^{MKV})^2dt],\end{aligned}$$

where we have used the fact that:

$$\mathbb{E}(\phi(t, X_t)) = c^{MKV}(t)\bar{\eta}_t^{MKV}\bar{x}_t^{MKV},$$

and:

$$Var(\phi(t, X_t)) = (a^{MKV}(t)\eta_t^{MKV})^2v_t^{MKV}.$$

2.3 Theoretical Results

For the remainder of the paper, we assume the coefficients are independent of time and nonnegative:

$$(b_1(t), \bar{b}_1(t), b_2(t), \bar{b}_2(t), q(t), \bar{q}(t), r(t), \bar{r}(t), s(t), \bar{s}(t)) = (b_1, \bar{b}_1, b_2, \bar{b}_2, q, \bar{q}, r, \bar{r}, s, \bar{s}) \in (\mathbb{R}^+)^{10}$$

and therefore,

$$\begin{aligned}(a^{MFG}(t), b^{MFG}(t), c^{MFG}(t)) &= (a^{MFG}, b^{MFG}, c^{MFG}) \\ (a^{MKV}(t), b^{MKV}(t), c^{MKV}(t)) &= (a^{MKV}, b^{MKV}, c^{MKV}).\end{aligned}$$

Theorem 1. *Assume the following:*

$$\begin{array}{llll} b_2 & > & 0 & b_2 + \bar{b}_2 & > & 0 \\ r + \bar{r} & > & 0 & r + \bar{r}(1 - \bar{s}) & > & 0 & r + \bar{r}(1 - \bar{s})^2 & > & 0 \\ q + \bar{q} & > & 0 & q + \bar{q}(1 - s) & > & 0 & q + \bar{q}(1 - s)^2 & > & 0 \\ q_T + \bar{q}_T & \geq & 0 & q_T + \bar{q}_T(1 - s_T) & \geq & 0 & q_T + \bar{q}_T(1 - s_T)^2 & \geq & 0 \end{array}\tag{14}$$

then there exists a unique solution to the LQEMFG problem, and there exists a unique solution to the LQEMKV problem. And therefore, $PoA = \frac{SC^{MFG}}{SC^{MKV}}$ where $SC^{MFG} := SC(\phi)$ for ϕ given by the explicit solution constructed in Appendix A.

Remark 3. Note that existence in Theorem 1 follows from the explicit construction in Appendix A, because the above conditions provide existence to the solutions of the Riccati equations. Uniqueness comes from the connection between LQEMFG or LQEMKV and deterministic LQ optimal control. (See Section 3.5.1 in [7]).

To compute the price of anarchy, it is useful to make the following observations:

$$\begin{aligned} a^{MFG} &= a^{MKV} =: a \\ \eta_t^{MFG} &= \eta_t^{MKV} =: \eta_t \\ v_t^{MFG} &= v_t^{MKV} =: v_t \end{aligned}$$

Proposition 1. Assuming (14), if furthermore,

$$\begin{aligned} (s\bar{q}(s-1) + \bar{b}_1\bar{\eta}_t^{MKV})\bar{x}_t^{MKV} &= 0, \quad \forall t \in [0, T] \\ \bar{\eta}_t^{MKV}\bar{x}_t^{MKV} [(b^{MFG} - b^{MKV})b_2 + (c^{MFG} - c^{MKV})\bar{b}_2] &= 0, \quad \forall t \in [0, T] \\ s_T\bar{q}_T(s_T-1)\bar{x}_T^{MKV} &= 0, \end{aligned}$$

then $PoA = 1$.

Proof. Comparing the FBSDE systems (2) and (8), the result is clear. \square

Corollary 1. Assuming (14), if furthermore, $\bar{b}_1 = 0$, $s\bar{q}(s-1) = 0$, and $s_T\bar{q}_T(s_T-1) = 0$ and at least one of the following holds: $b_2 = \bar{b}_2 = 0$ or $\frac{b_2(r+\bar{r}(1-\bar{s})^2)}{(b_2+\bar{b}_2)(r+\bar{r}(1-\bar{s}))} = 1$ then $PoA = 1$.

Remark 4. The only result similar to Proposition 1 that we are aware of is Remark 6.1 in [16].

Using the above observations, we can rewrite:

$$\begin{aligned} SC^{MFG} &= \frac{1}{2}(q_T + \bar{q}_T)v_T + \frac{1}{2}(q_T + \bar{q}_T(1-s_T)^2)(\bar{x}_T^{MFG})^2 + \frac{1}{2}\int_0^T (q + \bar{q} + (r + \bar{r})(a\eta_t)^2)v_t dt \\ &\quad + \frac{1}{2}\int_0^T [q + \bar{q}(1-s)^2 + (r + \bar{r}(1-\bar{s})^2)(c^{MFG}\bar{\eta}_t^{MFG})^2](\bar{x}_t^{MFG})^2 dt, \end{aligned} \quad (15)$$

and:

$$\begin{aligned} SC^{MKV} &= \frac{1}{2}(q_T + \bar{q}_T)v_T + \frac{1}{2}(q_T + \bar{q}_T(1-s_T)^2)(\bar{x}_T^{MKV})^2 + \frac{1}{2}\int_0^T (q + \bar{q} + (r + \bar{r})(a\eta_t)^2)v_t dt \\ &\quad + \frac{1}{2}\int_0^T [q + \bar{q}(1-s)^2 + (r + \bar{r}(1-\bar{s})^2)(c^{MKV}\bar{\eta}_t^{MKV})^2](\bar{x}_t^{MKV})^2 dt. \end{aligned} \quad (16)$$

In the following, we intend to simplify the explicit solutions (15) and (16) for the social costs in the LQEMFG and LQEMKV problems. First, consider the quantity $\int_0^T (\bar{\eta}_t^{MFG})^2(\bar{x}_t^{MFG})^2 dt$. Using equation (3), we have:

$$\begin{aligned} &\int_0^T (\bar{\eta}_t^{MFG})^2(\bar{x}_t^{MFG})^2 dt \\ &= -\frac{1}{c^{MFG}(b_2 + \bar{b}_2)} \left[\int_0^T \dot{\bar{\eta}}_t^{MFG}(\bar{x}_t^{MFG})^2 dt + \int_0^T ((2b_1 + \bar{b}_1)\bar{\eta}_t^{MFG} + (q + \bar{q}(1-s))) (\bar{x}_t^{MFG})^2 dt \right] \end{aligned}$$

Then by integration by parts:

$$= -\frac{1}{c^{MFG}(b_2 + \bar{b}_2)} \left[\bar{\eta}_T^{MFG} (\bar{x}_T^{MFG})^2 - \bar{\eta}_0^{MFG} (\bar{x}_0^{MFG})^2 - 2 \int_0^T \bar{\eta}_t^{MFG} \bar{x}_t^{MFG} \dot{\bar{x}}_t^{MFG} dt \right. \\ \left. + \int_0^T ((2b_1 + \bar{b}_1) \bar{\eta}_t^{MFG} + (q + \bar{q}(1-s)) (\bar{x}_t^{MFG})^2) dt \right]$$

Then using equation (5):

$$= 2 \int_0^T (\bar{\eta}_t^{MFG})^2 (\bar{x}_t^{MFG})^2 dt - \frac{1}{c^{MFG}(b_2 + \bar{b}_2)} \left[\bar{\eta}_T^{MFG} (\bar{x}_T^{MFG})^2 - \bar{\eta}_0^{MFG} (\mathbb{E}(\xi))^2 \right. \\ \left. - \int_0^T (\bar{b}_1 \bar{\eta}_t^{MFG} - (q + \bar{q}(1-s))) (\bar{x}_t^{MFG})^2 dt \right].$$

Finally, we solve:

$$\int_0^T (\bar{\eta}_t^{MFG})^2 (\bar{x}_t^{MFG})^2 dt = \frac{1}{c^{MFG}(b_2 + \bar{b}_2)} \left[\bar{\eta}_T^{MFG} (\bar{x}_T^{MFG})^2 - \bar{\eta}_0^{MFG} (\mathbb{E}(\xi))^2 \right. \\ \left. + \int_0^T (-\bar{b}_1 \bar{\eta}_t^{MFG} + (q + \bar{q}(1-s))) (\bar{x}_t^{MFG})^2 dt \right].$$

If we denote:

$$\lambda := \frac{c^{MFG}}{c^{MKV}} = \frac{b_2}{b_2 + \bar{b}_2} \cdot \frac{r + \bar{r}(1-s)^2}{r + \bar{r}(1-\bar{s})},$$

$$h_{var} := \frac{1}{2} \int_0^T (q + \bar{q} + (r + \bar{r})(a\eta_t)^2) v_t dt + \frac{1}{2} (q_T + \bar{q}_T) v_T,$$

and use the terminal condition for $\bar{\eta}_T^{MFG}$, then equation (15) can be rewritten as:

$$SC^{MFG} = h_{var} + \frac{1}{2} \int_0^T [\bar{b}_1 \lambda \bar{\eta}_t^{MFG} + (q + \bar{q}(1-s)^2) - \lambda(q + \bar{q}(1-s))] (\bar{x}_t^{MFG})^2 dt, \\ + \frac{1}{2} \lambda (\bar{\eta}_0^{MFG} (\mathbb{E}(\xi))^2 - (q_T + \bar{q}_T(1-s_T)) (\bar{x}_T^{MFG})^2) + \frac{1}{2} (q_T + \bar{q}_T(1-s_T)^2) (\bar{x}_T^{MFG})^2. \quad (17)$$

Repeating the calculation for $\int_0^T (\bar{\eta}_t^{MFG})^2 (\bar{x}_t^{MFG})^2 dt$ (using equations (9) and (11)), we arrive at:

$$\int_0^T (\bar{\eta}_t^{MKV})^2 (\bar{x}_t^{MKV})^2 dt = \frac{1}{c^{MKV}(b_2 + \bar{b}_2)} \left[\bar{\eta}_T^{MKV} (\bar{x}_T^{MKV})^2 - \bar{\eta}_0^{MKV} (\mathbb{E}(\xi))^2 \right. \\ \left. + \int_0^T (q + \bar{q}(1-s)^2) (\bar{x}_t^{MKV})^2 dt \right],$$

Using the terminal condition for $\bar{\eta}_T^{MKV}$, equation (16) can be rewritten as:

$$SC^{MKV} = h_{var} + \frac{1}{2} \bar{\eta}_0^{MKV} (\mathbb{E}(\xi))^2. \quad (18)$$

Let's denote the (weighted) difference between the solutions of the Riccati equations associated with $\bar{\eta}_t^{MFG}$ and $\bar{\eta}_t^{MKV}$ by:

$$\Delta\bar{\eta}_t = \lambda\bar{\eta}_t^{MFG} - \bar{\eta}_t^{MKV}. \quad (19)$$

Proposition 2. *Under assumption (14), the difference in the social costs in the LQEMFG and LQEMKV problems can be represented by*

$$\Delta SC := SC^{MFG} - SC^{MKV} = \frac{1}{2} \cdot \frac{(b_2 + \bar{b}_2)^2}{r + \bar{r}(1 - \bar{s})^2} \int_0^T (\Delta\bar{\eta}_t \cdot \bar{x}_t^{MFG})^2 dt.$$

Proof. The solutions $\bar{\eta}_t^{MFG}$ and $\bar{\eta}_t^{MKV}$ for the Riccati equations (3) and (9), respectively, are well defined under assumption (14) (see Appendix A). We notice that $\Delta\bar{\eta}_t$ defined in (19) satisfies the following linear first-order differential equation:

$$\frac{d(\Delta\bar{\eta}_t)}{dt} = \gamma_t \Delta\bar{\eta}_t + \beta_t, \quad \Delta\bar{\eta}_T = \lambda\bar{\eta}_T^{MFG} - \bar{\eta}_T^{MKV}$$

with coefficients:

$$\begin{cases} \gamma_t &= -2b_1 - 2\bar{b}_1 + \frac{(b_2 + \bar{b}_2)^2}{r + \bar{r}(1 - \bar{s})^2} (\lambda\bar{\eta}_t^{MFG} + \bar{\eta}_t^{MKV}), \\ \beta_t &= \bar{b}_1 \lambda \bar{\eta}_t^{MFG} + (q + \bar{q}(1 - s)^2) - \lambda(q + \bar{q}(1 - s)). \end{cases}$$

Since $q_T + \bar{q}_T(1 - s_T) = \bar{\eta}_T^{MFG}$, $q_T + \bar{q}_T(1 - s_T)^2 = \bar{\eta}_T^{MKV}$ and $\lambda\bar{\eta}_0^{MFG} - \bar{\eta}_0^{MKV} = \Delta\bar{\eta}_0$, we deduce from equations (17) and (18) that:

$$\begin{aligned} SC^{MFG} - SC^{MKV} &= \frac{1}{2} \left[\Delta\bar{\eta}_0 (\mathbb{E}(\xi))^2 - \Delta\bar{\eta}_T (\bar{x}_T^{MFG})^2 + \int_0^T \beta_t (\bar{x}_t^{MFG})^2 dt \right] \\ &= \frac{1}{2} \int_0^T \left[-\frac{d(\Delta\bar{\eta}_t (\bar{x}_t^{MFG})^2)}{dt} + \left(\frac{d(\Delta\bar{\eta}_t)}{dt} - \gamma_t \Delta\bar{\eta}_t \right) (\bar{x}_t^{MFG})^2 \right] dt \\ &= \frac{1}{2} \int_0^T \left[-2\Delta\bar{\eta}_t \bar{x}_t^{MFG} \dot{\bar{x}}_t^{MFG} - \gamma_t \Delta\bar{\eta}_t (\bar{x}_t^{MFG})^2 \right] dt \\ &= \frac{1}{2} \int_0^T \Delta\bar{\eta}_t (\bar{x}_t^{MFG})^2 \left[-2 \left(b_1 + \bar{b}_1 - \frac{(b_2 + \bar{b}_2)^2}{r + \bar{r}(1 - \bar{s})^2} \lambda \bar{\eta}_t^{MFG} \right) - \gamma_t \right] dt \\ &= \frac{1}{2} \cdot \frac{(b_2 + \bar{b}_2)^2}{r + \bar{r}(1 - \bar{s})^2} \int_0^T (\Delta\bar{\eta}_t \cdot \bar{x}_t^{MFG})^2 dt, \end{aligned}$$

where we use equation (5) for the fourth equality. □

Remark 5. *We can see directly from Proposition 2 that the social cost in the LQEMFG problem is larger than (or possibly equal to) the social cost in the LQEMKV problem. This result is consistent with the definition of the price of anarchy in Section 1.3.*

Therefore, under assumption (14), the price of anarchy for the LQ model is given by

$$PoA = 1 + \frac{\Delta SC}{SC^{MKV}} = 1 + \frac{\frac{(b_2 + \bar{b}_2)^2}{r + \bar{r}(1 - \bar{s})^2} \int_0^T (\Delta \bar{\eta}_t \cdot \bar{x}_t^{MFG})^2 dt}{\int_0^T \left[q + \bar{q} + \frac{b_2^2}{r + \bar{r}} \eta_t^2 \right] v_t dt + (q_T + \bar{q}_T) v_T + \bar{\eta}_0^{MKV} (\mathbb{E}(\xi))^2}. \quad (20)$$

Corollary 2. *Assuming (14), if the initial condition ξ is such that $\mathbb{E}(\xi) = 0$, then $PoA = 1$.*

Proof. By equation (6), $\mathbb{E}(\xi) = 0$ implies $\bar{x}_t^{MFG} = 0$, $\forall t \in [0, T]$. Therefore by Proposition 2, $\Delta SC = 0$. From equation (7), which we recall is equivalent to equation (13), $v_t > 0$ which implies $h_{var} > 0$. Therefore by equation (18), $SC^{MKV} > 0$. We conclude that $PoA = 1$. \square

We study in the following the variation of PoA by letting one of the coefficients tend to 0 or to ∞ . It will be useful for us to note here the scalar Riccati equations associated with $u_t := \lambda \bar{\eta}_t^{MFG}$, $w_t := \bar{\eta}_t^{MKV}$ and η_t :

$$\dot{u}_t - 2A^u u_t - B u_t^2 + C^u = 0 \quad u_T = D^u \quad (21)$$

$$\dot{w}_t - 2A^w w_t - B w_t^2 + C^w = 0 \quad w_T = D^w \quad (22)$$

$$\dot{\eta}_t - 2A^\eta \eta_t - B^\eta \eta_t^2 + C^\eta = 0 \quad \eta_T = D^\eta \quad (23)$$

where

$$\begin{aligned} A^u &= -\left(b_1 + \frac{\bar{b}_1}{2}\right) & A^w &= -(b_1 + \bar{b}_1) & A^\eta &= -b_1 \\ B^u &= \frac{(b_2 + \bar{b}_2)^2}{r + \bar{r}(1 - \bar{s})^2} & B^w &= \frac{(b_2 + \bar{b}_2)^2}{r + \bar{r}(1 - \bar{s})^2} & B^\eta &= \frac{b_2^2}{r + \bar{r}} \\ C^u &= \lambda(q + \bar{q}(1 - s)) & C^w &= q + \bar{q}(1 - s)^2 & C^\eta &= q + \bar{q} \\ D^u &= \lambda(q_T + \bar{q}_T(1 - s_T)) & D^w &= q_T + \bar{q}_T(1 - s_T)^2 & D^\eta &= q_T + \bar{q}_T \end{aligned}$$

If $B^u \neq 0$, $B^u D^u \geq 0$ and $B^u C^u > 0$, we have (see equation (40) in Appendix A) the existence and uniqueness for u_t which can be expressed by:

$$u_t = \frac{(C^u + D^u \delta_u^+) - (C^u + D^u \delta_u^-) e^{-(\delta_u^+ - \delta_u^-)(T-t)}}{(-\delta_u^- + D^u B^u) + (\delta_u^+ - D^u B^u) e^{-(\delta_u^+ - \delta_u^-)(T-t)}} \quad (24)$$

where $\delta_u^\pm = -A^u \pm \sqrt{(A^u)^2 + B^u C^u}$. Note that under the above conditions, we have $\delta_u^- < 0 < \delta_u^+$ and thus, $u_t \geq 0$ for all $t \in [0, T]$. We have analogous expressions for w_t and η_t , which are also nonnegative, in terms of δ_w^\pm and δ_η^\pm , respectively. Note that $B^u = B^w =: B$, and the above conditions on the parameters B , C^u , D^u , C^w , D^w , B^η , C^η , and D^η are exactly the conditions in assumption (14).

To make the following computations easier to follow, we repeat equations (20), (6), and (7), which we recall is equivalent to equation (13), using the above notations:

$$PoA = 1 + \frac{\Delta SC}{SC^{MKV}} = 1 + \frac{B \int_0^T (u_t - w_t)^2 \cdot (\bar{x}_t^{MFG})^2 dt}{\int_0^T [q + \bar{q} + B^\eta \eta_t^2] v_t dt + (q_T + \bar{q}_T) v_T + w_0 (\mathbb{E}(\xi))^2}, \quad (25)$$

$$\bar{x}_t^{MFG} = \mathbb{E}(\xi) e^{\int_0^t (b_1 + \bar{b}_1 - B u_s) ds}, \quad (26)$$

$$v_t = Var(\xi) e^{\int_0^t 2(b_1 - B^\eta \eta_s) ds} + \sigma^2 \int_0^t e^{2 \int_s^t (b_1 - B^\eta \eta_u) du} ds. \quad (27)$$

Proposition 3. *Assuming (14), and assuming $b_1 > 0$,*

$$\lim_{r \rightarrow \infty} PoA = 1 \quad \text{and} \quad \lim_{\bar{r} \rightarrow \infty} PoA = 1.$$

Proof. First, we consider $r \rightarrow \infty$. For every given $r > 0$, we have existence and uniqueness of the solutions u_t^r , w_t^r and η_t^r to the scalar Riccati equations (21)-(23). Note that we have added the superscript r to emphasize the dependence on this parameter.

When $r \rightarrow \infty$, we have:

$$\lambda^r \longrightarrow \lambda^{r \rightarrow \infty} := \frac{b_2}{b_2 + \bar{b}_2},$$

and

$$\begin{aligned} B^r &\longrightarrow 0, \quad B^{\eta, r} \longrightarrow 0, \\ C^{u, r} &\longrightarrow C_u^{r \rightarrow \infty} := \lambda^{r \rightarrow \infty} (q + \bar{q}(1 - s)), \quad D^{u, r} \rightarrow D_u^{r \rightarrow \infty} := \lambda^{r \rightarrow \infty} (q_T + \bar{q}_T(1 - s_T)). \end{aligned}$$

Let $u^{r \rightarrow \infty} : [0, T] \rightarrow \mathbb{R}$ be the solution to the linear first-order differential equation:

$$(u_t^{r \rightarrow \infty})' - 2A^u u_t^{r \rightarrow \infty} + C_u^{r \rightarrow \infty} = 0, \quad u_T^{r \rightarrow \infty} = D_u^{r \rightarrow \infty}. \quad (28)$$

Then we have:

$$u_t^{r \rightarrow \infty} = \left(D_u^{r \rightarrow \infty} - \frac{C_u^{r \rightarrow \infty}}{2A^u} \right) e^{-2A^u(T-t)} + \frac{C_u^{r \rightarrow \infty}}{2A^u}.$$

It is easy to show that for every time $t \in [0, T]$, $\lim_{r \rightarrow \infty} u_t^r = u_t^{r \rightarrow \infty}$, and thus, $\lim_{r \rightarrow \infty} B^r u_t^r = 0$. Next, our goal is to bound the u_t^r uniformly over $t \in [0, T]$ for large enough r .

Note that $A^u < 0$, B^r , $C^{u, r}$, $\lambda^{r \rightarrow \infty}$, $C_u^{r \rightarrow \infty} > 0$, $D^{u, r}$, $D_u^{r \rightarrow \infty} \geq 0$, and $\delta_u^{-, r} < 0 < \delta_u^{+, r}$. Let $\epsilon > 0$. Then there exists $r^* > 0$ such that $\max_{r > r^*} \{B^r, C^{u, r}, D^{u, r}\} \leq \max\{C_u^{r \rightarrow \infty}, D_u^{r \rightarrow \infty}\} + \epsilon =: c_1$. We have the following two inequalities for $r > r^*$ and $\forall t \in [0, T]$:

$$\begin{aligned} &\left| (C^{u, r} + D^{u, r} \delta_u^{+, r}) - (C^{u, r} + D^{u, r} \delta_u^{-, r}) e^{-(\delta_u^{+, r} - \delta_u^{-, r})(T-t)} \right| \\ &= C^{u, r} (1 - e^{-(\delta_u^{+, r} - \delta_u^{-, r})(T-t)}) + D^{u, r} (\delta_u^{+, r} - \delta_u^{-, r}) e^{-(\delta_u^{+, r} - \delta_u^{-, r})(T-t)} \\ &\leq C^{u, r} + D^{u, r} (\delta_u^{+, r} - \delta_u^{-, r}) \\ &= C^{u, r} + 2D^{u, r} \sqrt{(A^u)^2 + B^r C^{u, r}} \\ &\leq c_1 + 2c_1 \sqrt{(A^u)^2 + c_1^2} \\ &< \infty, \end{aligned}$$

and

$$\begin{aligned}
& \left| (-\delta_u^{-,r} + D^{u,r} B^r) + (\delta_u^{+,r} - D^{u,r} B^r) e^{-(\delta_u^{+,r} - \delta_u^{-,r})(T-t)} \right| \\
&= D^{u,r} B^r (1 - e^{-(\delta_u^{+,r} - \delta_u^{-,r})(T-t)}) + \delta_u^{+,r} e^{-(\delta_u^{+,r} - \delta_u^{-,r})(T-t)} - \delta_u^{-,r} \\
&\geq \delta_u^{+,r} e^{-2\sqrt{(A^u)^2 + B^r C^{u,r}}(T-t)} \\
&= (-A^u + \sqrt{(A^u)^2 + B^r C^{u,r}}) e^{-2\sqrt{(A^u)^2 + B^r C^{u,r}}(T-t)} \\
&\geq (-A^u + |A_u|) e^{-2T\sqrt{(A^u)^2 + c_1^2}} \\
&\geq -2A^u e^{-2T\sqrt{(A^u)^2 + c_1^2}} \\
&> 0.
\end{aligned}$$

Thus, we can deduce that for $r > r^*$, for every $t \in [0, T]$:

$$\begin{aligned}
|u_t^r| &= \frac{\left| (C^{u,r} + D^{u,r} \delta_u^{+,r}) - (C^{u,r} + D^{u,r} \delta_u^{-,r}) e^{-(\delta_u^{+,r} - \delta_u^{-,r})(T-t)} \right|}{\left| (-\delta_u^{-,r} + D^{u,r} B^r) + (\delta_u^{+,r} - D^{u,r} B^r) e^{-(\delta_u^{+,r} - \delta_u^{-,r})(T-t)} \right|} \\
&\leq \frac{c_1 + 2c_1 \sqrt{(A^u)^2 + c_1^2}}{-2A^u e^{-2T\sqrt{(A^u)^2 + c_1^2}}} \\
&=: c_2.
\end{aligned}$$

From equation (26) and by the bounded convergence theorem, we have for every $t \in [0, T]$:

$$\lim_{r \rightarrow \infty} \bar{x}_t^{MFG,r} = \mathbb{E}(\xi) e^{(b_1 + \bar{b}_1)t} =: \bar{x}_t^{MFG,r \rightarrow \infty}.$$

Moreover, $\bar{x}_t^{MFG,r}$ is uniformly bounded over $r > r^*$ and $t \in [0, T]$. From the nonnegativity of u_t , we have:

$$\left| \bar{x}_t^{MFG,r} \right| \leq |\mathbb{E}(\xi)| e^{(b_1 + \bar{b}_1)T}, \quad \forall t \in [0, T], \quad r > r^*.$$

Similarly, the solutions w_t^r and η_t^r of equations (22) and (23) converge for each time $t \in [0, T]$ towards the solutions $w_t^{r \rightarrow \infty}$ and $\eta_t^{r \rightarrow \infty}$ of some linear first-order differential equations. Namely, for every $t \in [0, T]$,

$$\lim_{r \rightarrow \infty} w_t^r = w_t^{r \rightarrow \infty}, \quad \lim_{r \rightarrow \infty} \eta_t^r = \eta_t^{r \rightarrow \infty},$$

and the functions w_t^r and η_t^r are uniformly bounded over $t \in [0, T]$ and large r .

Again, by the bounded convergence theorem we have for every $t \in [0, T]$:

$$\lim_{r \rightarrow \infty} \int_0^T (u_t^r - w_t^r)^2 (\bar{x}_t^{MFG,r})^2 dt = \int_0^T (u_t^{r \rightarrow \infty} - w_t^{r \rightarrow \infty})^2 (\bar{x}_t^{MFG,r \rightarrow \infty})^2 dt < \infty,$$

and thus,

$$\lim_{r \rightarrow \infty} \Delta SC^r = \lim_{r \rightarrow \infty} \frac{1}{2} \cdot B^r \int_0^T (u_t^r - w_t^r)^2 (\bar{x}_t^{MFG,r})^2 dt = 0.$$

From equation (27) and by the bounded convergence theorem, we have for each $t \in [0, T]$:

$$\lim_{r \rightarrow \infty} v_t^r = \text{Var}(\xi) e^{2b_1 t} + \sigma^2 \int_0^t e^{2b_1(t-s)} ds =: v_t^{r \rightarrow \infty}.$$

The variance function v_t^r is also uniformly bounded over $t \in [0, T]$ for large r . $w_0^{r \rightarrow \infty} \geq 0$ and $v_t^{r \rightarrow \infty} > 0$ for $t > 0$. Hence,

$$\lim_{r \rightarrow \infty} SC^{MKV,r} = \frac{1}{2} \left(\int_0^T (q + \bar{q}) v_t^{r \rightarrow \infty} dt + (q_T + \bar{q}_T) v_T^{r \rightarrow \infty} + w_0^{r \rightarrow \infty} (\mathbb{E}(\xi))^2 \right) > 0.$$

Therefore, from Proposition 2 and equation (25), we have $\lim_{r \rightarrow \infty} PoA^r = 1$. Idem, by replacing $\lambda^{r \rightarrow \infty}$ with $\lambda^{\bar{r} \rightarrow \infty} := \frac{b_2}{b_2 + \bar{b}_2} (1 - \bar{s})$, we obtain $\lim_{\bar{r} \rightarrow \infty} PoA^{\bar{r}} = 1$. □

Proposition 4. *Assuming (14), if initial condition ξ satisfies $\mathbb{E}(\xi) \neq 0$, and if furthermore, $\bar{b}_2 = 0$, then:*

$$\lim_{b_2 \rightarrow 0} PoA = 1,$$

whereas if $\bar{b}_2 > 0$, then:

$$\lim_{b_2 \rightarrow 0} PoA =: PoA^{0,b_2} > 1.$$

$$\lim_{b_2 \rightarrow \infty} PoA = 1.$$

Proof. First, we consider $b_2 \rightarrow 0$. If $\bar{b}_2 = 0$, we can use the same technique in Proposition 3 to show that $\lim_{b_2 \rightarrow 0} PoA = 1$. Now assume $\bar{b}_2 > 0$. As $b_2 \rightarrow 0$, we have $\lambda \rightarrow 0$ and

$$B \rightarrow \frac{\bar{b}_2^2}{r + \bar{r}(1 - \bar{s})^2} =: B^{0,b_2} > 0, \quad B^\eta \rightarrow 0, \quad C^u \rightarrow 0, \quad D^u \rightarrow 0.$$

We also have $\delta_u^\pm \rightarrow -A^u \pm |A^u|$ and therefore, $\delta_u^+ - \delta_u^- \rightarrow 2|A^u|$. Thus, from equation (24) we deduce that for every fixed time $t \in [0, T]$:

$$u_t \xrightarrow{b_2 \rightarrow 0} 0.$$

From equation (6), we have $\bar{x}_t^{MFG} = \mathbb{E}(\xi) e^{\int_0^t (b_1 + \bar{b}_1 - B u_s) ds}$, and therefore, $\bar{x}_t^{MFG} \rightarrow \mathbb{E}(\xi) \exp((b_1 + \bar{b}_1)t) =: \bar{x}_t^{MFG,0,b_2}$, which is bounded over $[0, T]$. Moreover, since $B^{0,b_2} > 0, B^{0,b_2} C^w > 0, B^{0,b_2} D^w \geq 0$, we still have the unique solution for w_t^{0,b_2} when $b_2 \rightarrow 0$.

Hence, from Proposition 2, the bounded convergence theorem, and the assumption on ξ , we deduce

$$\begin{aligned} \lim_{b_2 \rightarrow 0} \Delta SC &= \frac{1}{2} B^{0,b_2} \int_0^T (w_t^{0,b_2} \cdot \bar{x}_t^{MFG,0,b_2})^2 dt, \\ 0 &< \lim_{b_2 \rightarrow 0} \Delta SC < \infty, \end{aligned}$$

Since $B^\eta \rightarrow 0$, using the same argument shown in Proposition 3, we get:

$$\begin{aligned} \text{when } A^\eta \neq 0, \quad \eta_t &\rightarrow \eta_t^{0,b_2} := \left(D^\eta - \frac{C^\eta}{2A^\eta} \right) e^{-2A^\eta(T-t)} + \frac{C^\eta}{2A^\eta}, \\ \text{when } A^\eta = 0, \quad \eta_t &\rightarrow \eta_t^{0,b_2} := D^\eta - C^\eta(T-t), \end{aligned}$$

which are bounded over $[0, T]$. Thus, the variance function v_t converges to some positive bounded function $v_t^{0,b_2} := Var(\xi)e^{2b_1t} + \sigma^2 \int_0^t e^{2b_1(t-s)} ds > 0$ for every given time $t \in [0, T]$. Using the fact that $w_t^{0,b_2} \not\equiv 0$ and together with equation (18), we have $0 < \lim_{b_2 \rightarrow 0} SC^{MKV} < \infty$.

Therefore, combined with equation (25), we obtain

$$\lim_{b_2 \rightarrow 0} PoA = 1 + \frac{\lim_{b_2 \rightarrow 0} \Delta SC}{\lim_{b_2 \rightarrow 0} SC^{MKV}} =: PoA^{0,b_2} > 1.$$

Now we consider the case when $b_2 \rightarrow \infty$. It can be seen later in the proof that both SC^{MFG} and SC^{MKV} tend towards 0. In order to study the limit of PoA , the idea becomes to estimate the order of magnitude in terms of b_2 for the quantity ΔSC and SC^{MKV} .

When $b_2 \rightarrow \infty$, we have

$$\begin{aligned} \lambda^{b_2} &\rightarrow \frac{r+\bar{r}(1-\bar{s})^2}{r+\bar{r}(1-\bar{s})} =: \lambda^{b_2 \rightarrow \infty} \\ B^{b_2} &\rightarrow \infty, \quad B^{\eta,b_2} \rightarrow \infty \\ C^{u,b_2} &\rightarrow \lambda^{b_2 \rightarrow \infty}(q + \bar{q}(1-s)) =: C^{u,b_2 \rightarrow \infty}, \quad D^{u,b_2} \rightarrow \lambda^{b_2 \rightarrow \infty}(q_T + \bar{q}_T(1-s_T)) =: D^{u,b_2 \rightarrow \infty} \\ A^u, A^w, C^w, D^w, A^\eta, C^\eta, D^\eta &\text{ are independent of } b_2 \end{aligned}$$

Moreover

$$\begin{aligned} \frac{\delta_u^{\pm,b_2}}{b_2 + \bar{b}_2} &= \frac{A^u}{b_2 + \bar{b}_2} \pm \sqrt{\frac{C^{u,b_2}}{r + \bar{r}(1-\bar{s})^2} + \frac{(A^u)^2}{(b_2 + \bar{b}_2)^2}} \\ &\xrightarrow{b_2 \rightarrow \infty} \pm \sqrt{\frac{q + \bar{q}(1-s)}{r + \bar{r}(1-\bar{s})}} =: \pm c_{\delta_u^{b_2 \rightarrow \infty}} \end{aligned}$$

Thus we have $\delta_u^{+,b_2} - \delta_u^{-,b_2} \xrightarrow{b_2 \rightarrow \infty} +\infty$ and for any $t \in [0, T]$,

$$e^{-(\delta_u^{+,b_2} - \delta_u^{-,b_2})(T-t)} \xrightarrow{b_2 \rightarrow \infty} 0.$$

From the explicit formula (24) of $u_t^{b_2}$, we can compute the limit of $(b_2 + \bar{b}_2)u_t^{b_2}$ for any $t \in [0, T]$ by dividing the numerator with $(b_2 + \bar{b}_2)$ and the denominator with $(b_2 + \bar{b}_2)^2$, namely

$$\begin{aligned} (b_2 + \bar{b}_2)u_t^{b_2} &= \frac{\left(\frac{C^{u,b_2}}{b_2 + \bar{b}_2} + D^{u,b_2} \cdot \frac{\delta_u^{+,b_2}}{b_2 + \bar{b}_2} \right) - \left(\frac{C^{u,b_2}}{b_2 + \bar{b}_2} + D^{u,b_2} \cdot \frac{\delta_u^{-,b_2}}{b_2 + \bar{b}_2} \right) e^{-(\delta_u^{+,b_2} - \delta_u^{-,b_2})(T-t)}}{\left(\frac{-\delta_u^{-,b_2}}{(b_2 + \bar{b}_2)^2} + \frac{D^{u,b_2}}{r + \bar{r}(1-\bar{s})^2} \right) + \left(\frac{\delta_u^{+,b_2}}{(b_2 + \bar{b}_2)^2} - \frac{D^{u,b_2}}{r + \bar{r}(1-\bar{s})^2} \right) e^{-(\delta_u^{+,b_2} - \delta_u^{-,b_2})(T-t)}} \\ &\xrightarrow{b_2 \rightarrow \infty} (r + \bar{r}(1-\bar{s})^2) \cdot c_{\delta_u^{b_2 \rightarrow \infty}} =: c_3 \end{aligned}$$

Idem, we can have for any $t \in [0, T]$,

$$(b_2 + \bar{b}_2)w_t^{b_2} \xrightarrow{b_2 \rightarrow \infty} (r + \bar{r}(1-\bar{s})^2) \cdot c_{\delta_w^{b_2 \rightarrow \infty}} =: c_4, \quad \text{with} \quad c_{\delta_w^{b_2 \rightarrow \infty}} = \sqrt{\frac{q + \bar{q}(1-s)^2}{r + \bar{r}(1-\bar{s})^2}};$$

and for any $t \in [0, T)$,

$$b_2 \eta_t^{b_2} \xrightarrow{b_2 \rightarrow \infty} (r + \bar{r}) c_{\delta_\eta^{b_2 \rightarrow \infty}} =: c_5, \quad \text{with} \quad c_{\delta_\eta^{b_2 \rightarrow \infty}} = \sqrt{\frac{q + \bar{q}}{r + \bar{r}}}.$$

The next step is to provide an estimation of the order of magnitude of ΔSC and SC^{MKV} . For all $t \in [0, T)$, we denote $h_u(b_2, t) = (b_2 + \bar{b}_2)u_t^{b_2}$, $h_w(b_2, t) = (b_2 + \bar{b}_2)w_t^{b_2}$ and $h_\eta(b_2, t) = b_2 \eta_t^{b_2}$.

Since $u_t^{b_2}$, $w_t^{b_2}$ and \bar{x}_t^{MFG, b_2} are continuous functions on $[0, T]$, we deduce that the functions $t \mapsto \int_0^t u_s^{b_2} ds$ and $t \mapsto \int_0^t (u_s^{b_2} - w_s^{b_2})^2 (\bar{x}_s^{MFG, b_2})^2 ds$ are continuous on $[0, T]$ and differentiable on $(0, T)$. Then with equation (26), we can apply the Mean Value Theorem so that there exists two fixed times $0 < t_2 < t_1 < T$ for which

$$\begin{aligned} \Delta SC &= \frac{1}{2} B^{b_2} \int_0^T (u_t^{b_2} - w_t^{b_2})^2 (\bar{x}_t^{MFG, b_2})^2 dt \\ &= \frac{1}{2} B^{b_2} (u_{t_1}^{b_2} - w_{t_1}^{b_2})^2 (\bar{x}_{t_1}^{MFG, b_2})^2 \cdot T \\ &= \frac{T B^{b_2}}{2} \cdot (u_{t_1}^{b_2} - w_{t_1}^{b_2})^2 \cdot \mathbb{E}[\xi]^2 \exp \left(2(b_1 + \bar{b}_1)t_1 - 2 \int_0^{t_1} B^{b_2} u_s^{b_2} ds \right) \\ &= \kappa_1 \left[(b_2 + \bar{b}_2)(u_{t_1}^{b_2} - w_{t_1}^{b_2}) \right]^2 \cdot \exp \left(-2B^{b_2} u_{t_2}^{b_2} \cdot t_1 \right) \\ &= \kappa_1 [h_u(b_2, t_1) - h_w(b_2, t_1)]^2 \cdot e^{-\kappa_2 (b_2 + \bar{b}_2) h_u(b_2, t_2)}, \end{aligned} \quad (29)$$

where $\kappa_1 = \frac{T \mathbb{E}[\xi]^2 e^{2(b_1 + \bar{b}_1)t_1}}{2(r + \bar{r}(1 - \bar{s})^2)} > 0$ and $\kappa_2 = \frac{2t_1}{r + \bar{r}(1 - \bar{s})^2} > 0$ are two positive constants independent of b_2 .

Similarly, with equation (27) of the variance v_t , we know that the functions $t \mapsto v_t$ and $t \mapsto \int_0^t (q + \bar{q} + B^{\eta, b_2} (\eta_s^{b_2})^2) v_s^{b_2} ds$ are continuous on $[0, T]$ and differentiable on $(0, T)$. Then the Mean Value Theorem yields that there exists a time $t_3 \in (0, T)$ so that

$$\int_0^T (q + \bar{q} + B^{\eta, b_2} (\eta_s^{b_2})^2) v_s^{b_2} ds = \left(q + \bar{q} + \frac{1}{r + \bar{r}} h_\eta(b_2, t_3)^2 \right) v_{t_3}^{b_2} \cdot T, \quad (30)$$

and there exists 3 time points t_4, t_5, t_6 satisfying $0 < t_4 < T$ and $0 < t_5 < t_6 < T$ so that

$$\begin{aligned} v_T^{b_2} &= Var(\xi) \exp \left(2 \int_0^T (b_1 - B^{\eta, b_2} \eta_s^{b_2}) ds \right) + \sigma^2 \int_0^T \exp \left(2 \int_s^T (b_1 - B^{\eta, b_2} \eta_u^{b_2}) du \right) ds \\ &= Var(\xi) e^{2b_1 T} \exp \left(-2B^{\eta, b_2} \eta_{t_4}^{b_2} \cdot T \right) + \sigma^2 \exp \left(2 \int_{t_5}^T (b_1 - B^{\eta, b_2} \eta_u^{b_2}) du \right) \cdot T \\ &= Var(\xi) e^{2b_1 T} e^{-\frac{2T}{r + \bar{r}} b_2 h_\eta(b_2, t_4)} + \sigma^2 T e^{2b_1(T - t_5)} \exp \left(-2B^{\eta, b_2} \eta_{t_6}^{b_2} (T - t_5) \right) \\ &= \kappa_3 e^{-\kappa_4 b_2 h_\eta(b_2, t_4)} + \kappa_5 e^{-\kappa_6 b_2 h_\eta(b_2, t_6)} \end{aligned} \quad (31)$$

where $\kappa_3 = Var(\xi) e^{2b_1 T}$, $\kappa_4 = \frac{2T}{r + \bar{r}}$, $\kappa_5 = \sigma^2 T e^{2b_1(T - t_5)}$, $\kappa_6 = \frac{2(T - t_5)}{r + \bar{r}}$ are four positive constants independent of b_2 .

With equation (29) and (30), we can multiply ΔSC and SC^{MKV} by $(b_2 + \bar{b}_2)$ and then deduce that

$$\begin{aligned} \frac{\Delta SC}{SC^{MKV}} &= \frac{\kappa_1 [h_u(b_2, t_1) - h_w(b_2, t_1)]^2 \cdot e^{-\kappa_2(b_2 + \bar{b}_2)h_u(b_2, t_2)}}{\left((q + \bar{q})T + \frac{T}{r + \bar{r}} h_\eta(b_2, t_3)^2 \right) v_{t_3}^{b_2} + (q_T + \bar{q}_T) v_T^{b_2} + \frac{1}{b_2 + \bar{b}_2} (\mathbb{E}(\xi))^2 h_w(b_2, 0)} \\ &= \frac{l_{u,v}(b_2) \cdot (b_2 + \bar{b}_2) e^{-\kappa_2(b_2 + \bar{b}_2)h_u(b_2, t_2)}}{l_\eta(b_2) \cdot b_2 v_{t_3}^{b_2} + l_1(b_2) \cdot b_2 v_T^{b_2} + (\mathbb{E}(\xi))^2 h_w(b_2, 0)} \end{aligned} \quad (32)$$

where $l_{u,v}(b_2) = \kappa_1 [h_u(b_2, t_1) - h_w(b_2, t_1)]^2$, $l_\eta(b_2) = \left((q + \bar{q})T + \frac{T}{r + \bar{r}} h_\eta(b_2, t_3)^2 \right) \frac{b_2 + \bar{b}_2}{b_2}$ and $l_1(b_2) = (q_T + \bar{q}_T) \frac{b_2 + \bar{b}_2}{b_2}$. With the limits of $h_u(b_2, t)$, $h_w(b_2, t)$ and $h_\eta(b_2, t)$ for any fixed $t \in [0, T)$, it is easy to check that

$$l_{u,v}(b_2, t_3) \xrightarrow{b_2 \rightarrow \infty} \kappa_1(c_3 - c_4)^2, \quad l_\eta(b_2) \xrightarrow{b_2 \rightarrow \infty} (q + \bar{q} + \frac{c_5^2}{r + \bar{r}})T, \quad l_1(b_2) \xrightarrow{b_2 \rightarrow \infty} q_T + \bar{q}_T.$$

Finally, we will show that for any fixed $t \in [0, T)$ and $\kappa > 0$, we have

$$(b_2 + \bar{b}_2) e^{-\kappa(b_2 + \bar{b}_2)h_u(b_2, t)} \xrightarrow{b_2 \rightarrow \infty} 0 \quad \text{and} \quad b_2 e^{-\kappa b_2 h_\eta(b_2, t)} \xrightarrow{b_2 \rightarrow \infty} 0. \quad (33)$$

Therefore, the numerator of equation (32) tends towards 0 whilst the denominator tends towards $(\mathbb{E}(\xi))^2 c_4 > 0$ because equation (31) gives raise to

$$b_2 v_T^{b_2} = \kappa_3 \cdot b_2 e^{-\kappa_4 b_2 h_\eta(b_2, t_4)} + \kappa_5 \cdot b_2 e^{-\kappa_6 b_2 h_\eta(b_2, t_6)} \xrightarrow{b_2 \rightarrow \infty} 0,$$

and so $b_2 v_{t_3}^{b_2} \xrightarrow{b_2 \rightarrow \infty} 0$. Together with the limits of $l_{u,v}(b_2)$, $l_\eta(b_2)$ and $l_1(b_2)$, we have

$$\frac{\Delta SC}{SC^{MKV}} \xrightarrow{b_2 \rightarrow \infty} 0.$$

Take $(b_2 + \bar{b}_2) e^{-\kappa(b_2 + \bar{b}_2)h_u(b_2, t)}$ as an example. For any $t \in [0, T)$, we previously showed that $\lim_{b_2 \rightarrow \infty} h(b_2, t) = c_3 > 0$. Let $\epsilon_1 > 0$ such that $c_3 - 2\epsilon_1 > 0$. Then there exists $b_2^* > 0$ so that for all $b_2 \geq b_2^*$ we have $|h_u(b_2, t) - c_3| \leq \epsilon_1$. Thus $h_u(b_2, t) - c_2 + 2\epsilon_1 > \epsilon_1$ and

$$\begin{aligned} (b_2 + \bar{b}_2) e^{-\kappa(b_2 + \bar{b}_2)h_u(b_2, t)} &= e^{-\kappa(b_2 + \bar{b}_2)(h_u(b_2, t) - c_3 + 2\epsilon_1)} \cdot e^{-\kappa(c_3 - 2\epsilon_1)(b_2 + \bar{b}_2) + \ln(b_2 + \bar{b}_2)} \\ &\leq e^{-\kappa(b_2^* + \bar{b}_2)\epsilon_1} \cdot \exp(-\kappa(c_3 - 2\epsilon_1)(b_2 + \bar{b}_2) + \ln(b_2 + \bar{b}_2)). \end{aligned}$$

Since the first term in the right hand side of the above inequality is a constant depending only on ϵ_1 , and the second term tends towards 0 when $b_2 \rightarrow \infty$, then we must have

$$(b_2 + \bar{b}_2) e^{-\kappa(b_2 + \bar{b}_2)h_u(b_2, t)} \xrightarrow{b_2 \rightarrow \infty} 0, \quad \forall 0 \leq t < T.$$

Thus, the limits in (33) hold true and following the previous analysis we conclude that

$$\lim_{b_2 \rightarrow \infty} PoA = 1.$$

□

Proposition 5. Assuming (14), the initial condition ξ satisfies $\mathbb{E}(\xi) \neq 0$, and assuming $\bar{b}_1 > 0$, $q_T + \bar{q}_T(1 - s_T) > 0$, and $q_T + \bar{q}_T(1 - s_T)^2 > 0$ then

$$\lim_{\bar{b}_2 \rightarrow 0} PoA =: PoA^{\bar{b}_2 \rightarrow 0} > 1 \quad \text{and} \quad \lim_{\bar{b}_2 \rightarrow \infty} PoA = 1.$$

Proof. First, we consider $\bar{b}_2 \rightarrow 0$. For every given $\bar{b}_2 > 0$, we have existence and uniqueness of the solutions $u_t^{\bar{b}_2}$, $w_t^{\bar{b}_2}$ and $\eta_t^{\bar{b}_2}$ to the scalar Riccati equations (21)-(23). Note that, as before, we have added the superscript \bar{b}_2 to emphasize the dependence on this parameter.

When $\bar{b}_2 \rightarrow 0$, we have:

$$\lambda^{\bar{b}_2} \rightarrow \lambda^{\bar{b}_2 \rightarrow 0} := \frac{r + \bar{r}(1 - \bar{s})^2}{r + \bar{r}(1 - \bar{s})},$$

and

$$B^{\bar{b}_2} \rightarrow \frac{b_2^2}{r + \bar{r}(1 - \bar{s})^2} =: B^{\bar{b}_2 \rightarrow 0} > 0, \quad C^{u, \bar{b}_2} \rightarrow \lambda^{\bar{b}_2 \rightarrow 0}(q + \bar{q}(1 - s)) =: C_u^{\bar{b}_2 \rightarrow 0} > 0, \\ D^{u, \bar{b}_2} \rightarrow \lambda^{\bar{b}_2 \rightarrow 0}(q_T + \bar{q}_T(1 - s_T)) =: D_u^{\bar{b}_2 \rightarrow 0} \geq 0.$$

Let $u^{\bar{b}_2 \rightarrow 0} : [0, T] \rightarrow \mathbb{R}$ be the solution to the limiting Riccati equation:

$$\left(u_t^{\bar{b}_2 \rightarrow 0}\right)' - 2A^u u_t^{\bar{b}_2 \rightarrow 0} + B^{\bar{b}_2 \rightarrow 0}(u_t^{\bar{b}_2 \rightarrow 0})^2 + C_u^{\bar{b}_2 \rightarrow 0} = 0, \quad u_T^{\bar{b}_2 \rightarrow 0} = D_u^{\bar{b}_2 \rightarrow 0}, \quad (34)$$

which we recall has an explicit solution. It is easy to show that for every time $t \in [0, T]$, $\lim_{\bar{b}_2 \rightarrow 0} u_t^{\bar{b}_2} = u_t^{\bar{b}_2 \rightarrow 0}$. Next, our goal is to bound $u_t^{\bar{b}_2}$ uniformly over $t \in [0, T]$ for small enough \bar{b}_2 , following the methodology of the proof of Proposition 3. Because $\bar{b}_1 > 0$, we have $A^u < 0$. For any $\epsilon > 0$, there exists a $\bar{b}_2^* > 0$ such that $\max_{\bar{b}_2 < \bar{b}_2^*} \{B^{\bar{b}_2}, C^{u, \bar{b}_2}, D^{u, \bar{b}_2}\} < \max\{B^{\bar{b}_2 \rightarrow 0}, C_u^{\bar{b}_2 \rightarrow 0}, D_u^{\bar{b}_2 \rightarrow 0}\} + \epsilon =: c_1$. Thus, for all $\bar{b}_2 < \bar{b}_2^*$ and for every $t \in [0, T]$:

$$\left|u_t^{\bar{b}_2}\right| \leq \frac{c_1 + 2c_1 \sqrt{(A^u)^2 + c_1^2}}{-2A^u e^{-2T} \sqrt{(A^u)^2 + c_1^2}} \\ =: c_2.$$

Similarly, for every time $t \in [0, T]$, $\lim_{\bar{b}_2 \rightarrow 0} w_t^{\bar{b}_2} = w_t^{\bar{b}_2 \rightarrow 0}$, and $w^{\bar{b}_2}$ is uniformly bounded over $[0, T]$ for small enough \bar{b}_2 .

From equation (26), the assumption on ξ , and by the bounded convergence theorem, we have for every $t \in [0, T]$:

$$\lim_{\bar{b}_2 \rightarrow 0} \bar{x}_t^{MFG, \bar{b}_2} = \mathbb{E}(\xi) e^{\int_0^t (b_1 + \bar{b}_1 - B^{\bar{b}_2 \rightarrow 0} u_s^{\bar{b}_2 \rightarrow 0}) ds} =: \bar{x}_t^{MFG, \bar{b}_2 \rightarrow 0} > 0.$$

Moreover, \bar{x}^{MFG, \bar{b}_2} is uniformly bounded for all $\bar{b}_2 < \bar{b}_2^*$ and for all $t \in [0, T]$. From the nonnegativity of u_t , we have:

$$\left|\bar{x}_t^{MFG, \bar{b}_2}\right| \leq |\mathbb{E}(\xi)| e^{(b_1 + \bar{b}_1)T}, \quad \forall t \in [0, T], \quad \forall \bar{b}_2 < \bar{b}_2^*.$$

By our assumption on \bar{b}_1 , we have $A^u \neq A^w$ and thus, $u_t^{\bar{b}_2 \rightarrow 0} \neq w_t^{\bar{b}_2 \rightarrow 0}$ on a set of positive Lebesgue measure. Thus, from Proposition 2, and by the bounded convergence theorem, we deduce:

$$\lim_{\bar{b}_2 \rightarrow 0} \Delta SC^{\bar{b}_2} = \frac{1}{2} B^{\bar{b}_2 \rightarrow 0} \int_0^T \left((u_t^{\bar{b}_2 \rightarrow 0} - w_t^{\bar{b}_2 \rightarrow 0})^2 \left(\bar{x}_t^{MFG, \bar{b}_2 \rightarrow 0}\right)^2 dt > 0.$$

Meanwhile, η_t does not depend on \bar{b}_2 , and therefore the variance v_t also does not depend on \bar{b}_2 . It is easy to check that $0 < v_t < \infty$ for $t > 0$ and $0 \leq w_0^{\bar{b}_2 \rightarrow 0} < \infty$, and thus,

$$0 < \lim_{\bar{b}_2 \rightarrow 0} SC^{MKV, \bar{b}_2} = \frac{1}{2} \int_0^T [q + \bar{q} + B^\eta \eta_t^2] v_t dt + \frac{1}{2} (q_T + \bar{q}_T) v_T + \frac{1}{2} w_0^{\bar{b}_2 \rightarrow 0} (\mathbb{E}(\xi))^2 < \infty.$$

Hence, we deduce:

$$\lim_{\bar{b}_2 \rightarrow 0} PoA^{\bar{b}_2} =: PoA^{\bar{b}_2 \rightarrow 0} > 1.$$

Now, for the case when $\bar{b}_2 \rightarrow \infty$, we have $\lambda^{\bar{b}_2} \rightarrow 0$ and $B^{\bar{b}_2} \rightarrow \infty$, $C^{u, \bar{b}_2} \rightarrow 0$, $D^{u, \bar{b}_2} \rightarrow 0$. It can be shown that:

$$\lim_{\bar{b}_2 \rightarrow \infty} C^{u, \bar{b}_2} \sqrt{b_2 + \bar{b}_2} = 0,$$

and

$$\lim_{\bar{b}_2 \rightarrow \infty} \frac{\delta_u^{\pm, \bar{b}_2}}{\sqrt{b_2 + \bar{b}_2}} = \pm \sqrt{\frac{b_2(q + \bar{q}(1-s))}{r + \bar{r}(1-s)}} =: \pm \delta^{\bar{b}_2 \rightarrow \infty},$$

and therefore,

$$\lim_{\bar{b}_2 \rightarrow \infty} \delta_u^{\pm, \bar{b}_2} = \pm \infty,$$

$$\lim_{\bar{b}_2 \rightarrow \infty} \frac{\delta_u^{\pm, \bar{b}_2}}{b_2 + \bar{b}_2} = 0.$$

Also, note from the assumptions $q_T + \bar{q}_T(1-s_T) > 0$ and $q_T + \bar{q}_T(1-s_T)^2 > 0$, we have $D^u > 0$ and $D^w > 0$. From equation (24), for $t \in [0, T]$ we can deduce:

$$\begin{aligned} (b_2 + \bar{b}_2)^{\frac{3}{2}} \cdot u_t^{\bar{b}_2} &= \frac{\left(C^{u, \bar{b}_2} \sqrt{b_2 + \bar{b}_2} + \zeta \cdot \frac{\delta_u^{+, \bar{b}_2}}{\sqrt{b_2 + \bar{b}_2}} \right) - \left(C^{u, \bar{b}_2} \sqrt{b_2 + \bar{b}_2} + \zeta \cdot \frac{\delta_u^{-, \bar{b}_2}}{\sqrt{b_2 + \bar{b}_2}} \right) e^{-(\delta_u^{+, \bar{b}_2} - \delta_u^{-, \bar{b}_2})(T-t)}}{\left(-\frac{\delta_u^{-, \bar{b}_2}}{b_2 + \bar{b}_2} + \frac{b_2(q_T + \bar{q}_T(1-s_T))}{r + \bar{r}(1-s)} \right) + \left(\frac{\delta_u^{+, \bar{b}_2}}{b_2 + \bar{b}_2} - \frac{b_2(q_T + \bar{q}_T(1-s_T))}{r + \bar{r}(1-s)} \right) e^{-(\delta_u^{+, \bar{b}_2} - \delta_u^{-, \bar{b}_2})(T-t)}}} \\ &\xrightarrow{\bar{b}_2 \rightarrow \infty} \delta^{\bar{b}_2 \rightarrow \infty} (r + \bar{r}(1-s)^2) =: c_3, \end{aligned}$$

where $\zeta = \frac{b_2(r + \bar{r}(1-s)^2)}{r + \bar{r}(1-s)} (q_T + \bar{q}_T(1-s_T))$.

Since $u_t^{\bar{b}_2}$, $w_t^{\bar{b}_2}$ and $\bar{x}_t^{MFG, \bar{b}_2}$ are continuous over $[0, T]$, then the functions $t \mapsto \int_0^t u_s^{\bar{b}_2} ds$ and $t \mapsto \int_0^t (u_s^{\bar{b}_2} - w_s^{\bar{b}_2})^2 (\bar{x}_s^{MFG, \bar{b}_2})^2 ds$ are continuous over $[0, T]$ and differentiable over $(0, T)$. Then with the equation (26) we can apply the Mean Value theorem and deduce that there exists two fixed times $0 < t_2 < t_1 < T$ such that:

$$\begin{aligned} \Delta SC &= \frac{1}{2} B^{\bar{b}_2} \int_0^T (u_t^{\bar{b}_2} - w_t^{\bar{b}_2})^2 (\bar{x}_t^{MFG, \bar{b}_2})^2 dt \\ &= \frac{1}{2} B^{\bar{b}_2} (u_{t_1}^{\bar{b}_2} - w_{t_1}^{\bar{b}_2})^2 (\bar{x}_{t_1}^{MFG, \bar{b}_2})^2 \cdot T \\ &= \frac{TB^{\bar{b}_2}}{2} \cdot (u_{t_1}^{\bar{b}_2} - w_{t_1}^{\bar{b}_2})^2 \cdot \mathbb{E}[\xi]^2 \exp \left(2(b_1 + \bar{b}_1)t_1 - 2 \int_0^{t_1} B^{\bar{b}_2} u_s^{\bar{b}_2} ds \right) \\ &= \kappa_1 (u_{t_1}^{\bar{b}_2} - w_{t_1}^{\bar{b}_2})^2 \cdot \left[(b_2 + \bar{b}_2) \exp \left(-B^{\bar{b}_2} u_{t_2}^{\bar{b}_2} \cdot t_1 \right) \right]^2 \\ &= \kappa_1 (u_{t_1}^{\bar{b}_2} - w_{t_1}^{\bar{b}_2})^2 \cdot \left((b_2 + \bar{b}_2) e^{-\kappa(b_2 + \bar{b}_2)^{\frac{1}{2}} h(\bar{b}_2)} \right)^2, \end{aligned} \tag{35}$$

where $\kappa = \frac{t_1}{r+\bar{r}(1-\bar{s})^2} > 0$ and $\kappa_1 = \frac{T\mathbb{E}[\xi]^2 e^{2(b_1+\bar{b}_1)t_1}}{2(r+\bar{r}(1-\bar{s})^2)} > 0$ are two positive constants independent of \bar{b}_2 , and $h(\bar{b}_2) = (b_2 + \bar{b}_2)^{\frac{3}{2}} u_{t_2}^{\bar{b}_2}$.

Our next step is to show that $(b_2 + \bar{b}_2)e^{-\kappa(b_2+\bar{b}_2)^{\frac{1}{2}}h(\bar{b}_2)}$ tends towards 0 when $\bar{b}_2 \rightarrow \infty$. We previously showed that $\lim_{\bar{b}_2 \rightarrow \infty} h(\bar{b}_2) = c_3 > 0$. Let $\epsilon_1 > 0$ such that $c_3 - 2\epsilon_1 > 0$. Then there exists $\bar{b}_2^{*,1} > 0$ so that for all $\bar{b}_2 > \bar{b}_2^{*,1}$ we have $|h(\bar{b}_2) - c_3| < \epsilon_1$. Then $h(\bar{b}_2) - c_3 + 2\epsilon_1 > \epsilon_1$ and

$$\begin{aligned} (b_2 + \bar{b}_2)e^{-\kappa(b_2+\bar{b}_2)^{1/2}h(\bar{b}_2)} &= e^{-\kappa(b_2+\bar{b}_2)^{1/2}(h(\bar{b}_2)-c_3+2\epsilon_1)} \cdot e^{-\kappa(c_3-2\epsilon_1)(b_2+\bar{b}_2)^{1/2}+\ln(b_2+\bar{b}_2)} \\ &\leq e^{-\kappa(b_2+\bar{b}_2^{*,1})^{1/2}\epsilon_1} \cdot e^{-\kappa(c_3-2\epsilon_1)(b_2+\bar{b}_2)^{\frac{1}{2}}+\ln(b_2+\bar{b}_2)} \\ &= c_5 e^{-\kappa(c_3-2\epsilon_1)(b_2+\bar{b}_2)^{1/2}+\ln(b_2+\bar{b}_2)} \\ &\xrightarrow{\bar{b}_2 \rightarrow \infty} 0. \end{aligned}$$

Namely, for any $\epsilon > 0$, there exists a $\bar{b}_2^{*,2} \geq \bar{b}_2^{*,1}$, such that for all $\bar{b}_2 > \bar{b}_2^{*,2}$ we have:

$$|(b_2 + \bar{b}_2)e^{-\kappa(b_2+\bar{b}_2)^{\frac{1}{2}}h(\bar{b}_2)}| < \epsilon. \quad (36)$$

Now, in order to show that ΔSC tends to 0, we need to bound the terms $u_{t_1}^{\bar{b}_2}$ and $w_{t_1}^{\bar{b}_2}$ for large enough \bar{b}_2 . Given $\epsilon > 0$, there exists a $\bar{b}_2^{*,3} \geq 1$ such that for any $\bar{b}_2 > \bar{b}_2^{*,3}$ we have $(b_2 + \bar{b}_2)^{\frac{3}{2}} \left| u_{t_1}^{\bar{b}_2} \right| \leq c_3 + \epsilon$. Thus, for any $\bar{b}_2 > \bar{b}_2^{*,3} \geq 1$ we deduce:

$$\left| u_{t_1}^{\bar{b}_2} \right| \leq \frac{c_3 + \epsilon}{(b_2 + \bar{b}_2)^{\frac{3}{2}}} \leq \frac{c_3 + \epsilon}{(b_2 + \bar{b}_2^{*,3})^{\frac{3}{2}}} \leq c_3 + \epsilon.$$

Idem, we can show that for $t_1 \in [0, T]$:

$$\begin{aligned} (b_2 + \bar{b}_2)w_{t_1}^{\bar{b}_2} &= \frac{\left(\frac{C^w}{b_2+b_2} + D^w \cdot \frac{\delta_w^{+, \bar{b}_2}}{b_2+b_2} \right) - \left(\frac{C^w}{b_2+b_2} + D^w \cdot \frac{\delta_w^{-, \bar{b}_2}}{b_2+b_2} \right) e^{-(\delta_w^{+, \bar{b}_2} - \delta_w^{-, \bar{b}_2})(T-t_1)}}{\left(-\frac{\delta_w^{-, \bar{b}_2}}{(b_2+b_2)^2} + \frac{D^w}{r+\bar{r}(1-\bar{s})^2} \right) + \left(\frac{\delta_w^{+, \bar{b}_2}}{(b_2+b_2)^2} - \frac{D^w}{r+\bar{r}(1-\bar{s})^2} \right) e^{-(\delta_w^{+, \bar{b}_2} - \delta_w^{-, \bar{b}_2})(T-t_1)}}} \\ &\xrightarrow{\bar{b}_2 \rightarrow \infty} \sqrt{(r + \bar{r}(1 - \bar{s})^2)C^w} =: c_4 > 0, \end{aligned}$$

where we have used the fact that $\lim_{\bar{b}_2 \rightarrow \infty} \frac{\delta_w^{\pm, \bar{b}_2}}{b_2+b_2} = \sqrt{\frac{C^w}{r+\bar{r}(1-\bar{s})^2}}$. Thus, $\left| w_{t_1}^{\bar{b}_2} \right|$ can also be bounded by $c_4 + \epsilon$ for large enough \bar{b}_2 .

Hence, together with equation (35) and (36), we obtain that for any $\epsilon > 0$, there exists a \bar{b}_2^* large enough so that for all $\bar{b}_2 > \bar{b}_2^*$ we have

$$\Delta SC \leq \kappa_1(c_3 + c_4 + 2\epsilon)^2 \epsilon^2.$$

Thus, by taking $\epsilon \rightarrow 0$, we have:

$$\lim_{\bar{b}_2 \rightarrow \infty} \Delta SC = 0.$$

Moreover, recall that η_t and v_t are invariant with respect to \bar{b}_2 and $0 < v_t < \infty$ for $t > 0$. Clearly we also have $0 \leq w_0^{\bar{b}_2 \rightarrow \infty} < \infty$. Thus,

$$0 < \lim_{\bar{b}_2 \rightarrow \infty} SC^{MKV, \bar{b}_2} < \infty.$$

Therefore, we conclude that:

$$\lim_{\bar{b}_2 \rightarrow \infty} PoA^{\bar{b}_2} = 1.$$

□

Proposition 6. *Assuming (14), the initial condition ξ satisfies $\mathbb{E}(\xi) \neq 0$, $Var(\xi) > 0$, and $\lambda \neq 1$, then*

$$\begin{aligned} \lim_{b_1 \rightarrow 0} PoA &=: PoA^{0,b_1} > 1 & \text{and} & \quad \lim_{\bar{b}_1 \rightarrow 0} PoA &=: PoA^{0,\bar{b}_1} > 1, \\ \lim_{b_1 \rightarrow \infty} PoA &= 1 & \text{and} & \quad \lim_{\bar{b}_1 \rightarrow \infty} PoA = \infty. \end{aligned}$$

Proof. First, we consider $b_1 \rightarrow 0$. We have $A^{u,b_1} \rightarrow A^{u,b_1 \rightarrow 0}$, $\delta_u^{+,b_1} \rightarrow \delta_u^{+,b_1 \rightarrow 0} > 0$, and $\delta_u^{-,b_1} \rightarrow \delta_u^{-,b_1 \rightarrow 0} < 0$, and similarly for A^{w,b_1} , A^{η,b_1} , δ_w^{\pm,b_1} , and δ_η^{\pm,b_1} . Clearly we have for each $t \in [0, T]$, $\lim_{b_1 \rightarrow 0} u_t^{b_1} =: u_t^{b_1 \rightarrow 0}$, $\lim_{b_1 \rightarrow 0} w_t^{b_1} =: w_t^{b_1 \rightarrow 0}$, and $\lim_{b_1 \rightarrow 0} \eta_t^{b_1} =: \eta_t^{b_1 \rightarrow 0}$. Next, we show that the three sequences are uniformly bounded. Let $0 < \epsilon < -\delta_u^{-,b_1 \rightarrow 0}$. There exists a b_1^* such that $|\delta_u^{+,b_1} - \delta_u^{+,b_1 \rightarrow 0}| < \epsilon$ and $|\delta_u^{-,b_1} - \delta_u^{-,b_1 \rightarrow 0}| < \epsilon$ for each $b_1 < b_1^*$. Then for each $b_1 < b_1^*$ and $t \in [0, T]$, we have:

$$\begin{aligned} |u_t^{b_1}| &\leq \frac{C^u + D^u (\delta_u^{+,b_1} - \delta_u^{-,b_1})}{-\delta_u^{-,b_1}} \\ &\leq \frac{C^u + D^u (\delta_u^{+,b_1 \rightarrow 0} - \delta_u^{-,b_1 \rightarrow 0} + 2\epsilon)}{-\delta_u^{-,b_1 \rightarrow 0} - \epsilon} \\ &=: c, \end{aligned}$$

and similarly for $|w_t^{b_1}|$ and $|\eta_t^{b_1}|$. By the assumption on λ , $u_t^{b_1 \rightarrow 0} \neq w_t^{b_1 \rightarrow 0}$ on a set of positive Lebesgue measure.

From equation (26), the assumption on ξ , and by the bounded convergence theorem, we have for every $t \in [0, T]$:

$$\lim_{b_1 \rightarrow 0} \bar{x}_t^{MFG,b_1} = \mathbb{E}(\xi) e^{\int_0^t (\bar{b}_1 - B u_s^{b_1 \rightarrow 0}) ds} =: \bar{x}_t^{MFG,b_1 \rightarrow 0} > 0.$$

Moreover, $\bar{x}_t^{MFG,b_1 \rightarrow 0}$ is uniformly bounded over $b_1 < b_1^*$ and $t \in [0, T]$, i.e.

$$|\bar{x}_t^{MFG,b_1}| \leq |\mathbb{E}(\xi)| e^{\bar{b}_1 T}, \quad \forall t \in [0, T], \quad b_1 < b_1^*.$$

Therefore, by Proposition 2, and the bounded convergence theorem,

$$0 < \lim_{b_1 \rightarrow 0} \Delta SC^{b_1} < \infty.$$

By the bounded convergence theorem, $\lim_{b_1 \rightarrow 0} v_t^{b_1} =: v_t^{b_1 \rightarrow 0}$, which is bounded over $t \in [0, T]$, and thus:

$$0 < \lim_{b_1 \rightarrow 0} SC^{MKV,b_1} < \infty.$$

Therefore, $\lim_{b_1 \rightarrow 0} PoA^{b_1} =: PoA^{b_1 \rightarrow 0} > 1$. The proof can be repeated to show $\lim_{\bar{b}_1 \rightarrow 0} PoA^{\bar{b}_1} =: PoA^{\bar{b}_1 \rightarrow 0} > 1$.

Now we consider $b_1 \rightarrow \infty$. For each $b_1 > 0$, we have the following bounds over $t \in [0, T]$:

$$\begin{aligned} |u_t^{b_1}| &\leq \frac{C^u + D^u(\delta_u^{+,b_1} - \delta_u^{-,b_1})}{-\delta_u^{-,b_1}} =: h_u(b_1) \\ |w_t^{b_1}| &\leq \frac{C^w + D^w(\delta_w^{+,b_1} - \delta_w^{-,b_1})}{-\delta_w^{-,b_1}} =: h_w(b_1) \\ |\eta_t^{b_1}| &\leq \frac{C^\eta + D^\eta(\delta_\eta^{+,b_1} - \delta_\eta^{-,b_1})}{-\delta_\eta^{-,b_1}} =: h_\eta(b_1) \\ |\bar{x}_t^{MFG,b_1}| &\leq |\mathbb{E}(\xi)| e^{T(b_1 + \bar{b}_1)} =: h_{\bar{x}}(b_1). \end{aligned}$$

Thus, from Proposition 2, we have:

$$\Delta SC^{b_1} \leq \frac{1}{2} BT (h_u(b_1) + h_w(b_1))^2 (h_{\bar{x}}(b_1))^2.$$

We also have the following lower bound:

$$\begin{aligned} SC^{MKV,b_1} &\geq \frac{1}{2} (q_T + \bar{q}_T) v_T^{b_1} \\ &\geq (q_T + \bar{q}_T) Var(\xi) e^{2T(b_1 - B^\eta h_\eta(b_1))}. \end{aligned}$$

Let $h(b_1) = \max\{h_u(b_1), h_w(b_1), h_\eta(b_1)\}$. It is easy to check that $\lim_{b_1 \rightarrow \infty} h(b_1) = \infty$. Combining the above bounds and using the assumption $Var(\xi) > 0$, we have:

$$\begin{aligned} \frac{\Delta SC^{b_1}}{SC^{MKV,b_1}} &\leq \frac{2BT(h(b_1))^2(h_{\bar{x}}(b_1))^2}{\frac{1}{2}(q_T + \bar{q}_T)Var(\xi)e^{2T(b_1 - B^\eta h(b_1))}} \\ &= \frac{4BT(h(b_1))^2(|\mathbb{E}(\xi)| e^{T(b_1 + \bar{b}_1)})^2}{(q_T + \bar{q}_T)Var(\xi)e^{2T(b_1 - B^\eta h(b_1))}} \\ &= \frac{4BT e^{2T(\bar{b}_1)} (\mathbb{E}(\xi))^2}{(q_T + \bar{q}_T)Var(\xi)} \cdot \frac{(h(b_1))^2}{e^{-2TB^\eta h(b_1)}} \\ &\xrightarrow{b_1 \rightarrow \infty} 0. \end{aligned}$$

Therefore,

$$\lim_{b_1 \rightarrow \infty} \frac{\Delta SC^{b_1}}{SC^{MKV,b_1}} = 0,$$

and thus, $\lim_{b_1 \rightarrow \infty} PoA^{b_1} = 1$. ToDo: last case. □

2.4 Numerical Results

The solutions to the problems we have considered are given by the formulas derived in Appendix A, which are explicit up to evaluating integrals. Using the simple rectangle rule to evaluate integrals,

we numerically compute the price of anarchy when the coefficients are time-independent, nonnegative, and satisfy assumption (14). In particular, when we allow for full interaction (i.e. through the states and the controls), we choose the following default values:

$$\begin{aligned}\xi &\equiv 1, \quad T = 1 \\ b_1 &= 1, \quad \bar{b}_1 = 1, \quad b_2 = 1, \quad \bar{b}_2 = 1, \quad \sigma = 1 \\ q &= 1, \quad \bar{q} = 1, \quad s = 0.5, \quad r = 1, \quad \bar{r} = 1, \quad \bar{s} = 0.5 \\ q_T &= 1, \quad \bar{q}_T = 1, \quad s_T = 0.5.\end{aligned}$$

Unless otherwise stated, the parameters stay at these default values. For results involving only interaction through the states, we set $\bar{b}_2 = 0$ and $\bar{r} = 0$. For results involving only interaction through the controls, we set $\bar{b}_1 = 0$, $\bar{q} = 0$, and $\bar{q}_T = 0$. Figures 1-5 show the price of anarchy as we vary one parameter at a time for each of three cases: full interaction (i.e. through the states and the controls), interaction only through the states, and interaction only through the controls.




Figure 1: PoA as we vary b_1 (left) and \bar{b}_1 (right).




Figure 2: PoA as we vary b_2 (left) and \bar{b}_2 (right).

The results show various limiting behaviors, such as some of the cases proved in the previous section. In Figure 1, we note that Proposition 6 is confirmed. For all three cases, as $b_1 \rightarrow 0$, we see that $PoA \rightarrow PoA^{b_1 \rightarrow 0} > 1$ and as $\bar{b}_1 \rightarrow 0$, we see that $PoA \rightarrow PoA^{\bar{b}_1 \rightarrow 0} > 1$. We also




Figure 3: PoA as we vary q (left) and q_T (right).




Figure 4: PoA as we vary \bar{q} (left) and \bar{q}_T (right).




Figure 5: PoA as we vary r (left) and \bar{r} (right).

see that $PoA \rightarrow 1$ as $b_1 \rightarrow \infty$ and $PoA \rightarrow \infty$ as $\bar{b}_1 \rightarrow \infty$. Proposition 4 is confirmed in Figure 2. When there is only interaction through the states, then $\bar{b}_2 = 0$ and we see that $PoA \rightarrow 1$ as $b_2 \rightarrow 0$. When there is full interaction or only interaction through the controls, then $\bar{b}_2 \neq 0$ and we see that $PoA \rightarrow PoA^{b_2 \rightarrow 0} > 1$ as $b_2 \rightarrow 0$. For all three cases, we note that the condition $\frac{r+\bar{r}(1-\bar{s})^2}{r+\bar{r}(1-\bar{s})} = \frac{q+\bar{q}(1-s)^2}{q+\bar{q}(1-s)}$ is satisfied, and thus, $PoA \rightarrow 1$ as $b_2 \rightarrow \infty$. For Proposition 5, Figure 2

confirms that $PoA \rightarrow PoA^{\bar{b}_2 \rightarrow 0} > 1$ as $\bar{b}_2 \rightarrow 0$ and $PoA \rightarrow 1$ as $\bar{b}_2 \rightarrow \infty$. Figure 5 confirms as in Proposition 3 that $PoA \rightarrow 1$ as $r \rightarrow \infty$ or $\bar{r} \rightarrow \infty$.

2.5 A Particular Example: Flocking

Mean field game models of flocking have been proposed in the literature by Nourian et al [14][15]. Here we consider a slightly different formulation as described in Section 3.6.1 of the book [7]. A representative bird in the flock controls their velocity, X_t , through the drift:

$$b(t, x, \mu, \alpha, \nu) = \alpha.$$

They choose the control with two goals in mind: to minimize their kinetic energy put into their control, and to align their velocity with the average velocity of the group. Thus, they consider the cost functions given by:

$$\begin{aligned} f(t, x, \mu, \alpha, \nu) &= \frac{1}{2} (\bar{q}|x - s\bar{\mu}|^2 + \alpha^2), \\ g(x, \mu) &= 0. \end{aligned}$$

In our general linear quadratic formulation, this is equivalent to taking $b_1 = 0$, $\bar{b}_1 = 0$, $b_2 = 1$, $\bar{b}_2 = 0$, $q = 0$, $s = 1$, $r = 1$, $\bar{r} = 0$, and $\bar{s} = 0$. Note that for these values of the parameters, the assumptions of Theorem 1 and Corollary 1 are satisfied. Therefore, $PoA = 1$. In fact, if we took the state space to be \mathbb{R}^d instead of \mathbb{R} , the result would still hold.

3 Conclusion

We defined the price of anarchy (PoA) in the context of extended mean field games as the ratio of the worst case social cost when the players are in a mean field game equilibrium to the social cost as computed by a central planner. Since the central planner does not require that the players be in a mean field game equilibrium, the central planner will realize a social cost that is no worse than that of a mean field game equilibrium. Thus, $PoA \geq 1$.

We computed the price of anarchy for linear quadratic extended mean field games, for which explicit computations are possible. We identify a large class of models for which $PoA = 1$ (see Proposition 1 and Corollary 1), as well as some limiting cases where $PoA \rightarrow 1$ as certain parameters tend to zero or to infinity. The numerics support our theoretical results.

4 Acknowledgments

The authors would like to thank the organizers of CEMRACS 2017 for putting together such a successful summer school.

References

- [1] Y. Achdou and M. Laurière. On the system of partial differential equations arising in mean field type control. *arXiv preprint arXiv:1503.05044*, 2015.

- [2] M. Balandat and C. J. Tomlin. On efficiency in mean field differential games. In *American Control Conference (ACC), 2013*, pages 2527–2532. IEEE, 2013.
- [3] A. Bensoussan, J. Frehse, and P. Yam. *Mean field games and mean field type control theory*, volume 101. Springer, 2013.
- [4] P. Cardaliaguet and C. Rainer. On the (in)efficiency of MFG equilibria. Technical report, 2018.
- [5] R. Carmona, B. Acciaio, and J. Backhoff Veraguas. Generalized McKean-Vlasov (Mean Field) Control: a stochastic maximum principle and a transport perspective. Technical report, 2018.
- [6] R. Carmona and F. Delarue. Probabilistic analysis of mean-field games. *SIAM Journal on Control and Optimization*, 51(4):2705–2734, 2013.
- [7] R. Carmona and F. Delarue. *Probabilistic Theory of Mean Field Games with Applications I*. Springer, 2018.
- [8] G. Christodoulou and E. Koutsoupias. On the price of anarchy and stability of correlated equilibria of linear congestion games. In *European Symposium on Algorithms*, pages 59–70. Springer, 2005.
- [9] G. Christodoulou and E. Koutsoupias. The price of anarchy of finite congestion games. In *Proceedings of the thirty-seventh annual ACM symposium on Theory of computing*, pages 67–73. ACM, 2005.
- [10] M. Huang, P. E. Caines, and R. P. Malhamé. Social optima in mean field LQG control: centralized and decentralized strategies. *IEEE Transactions on Automatic Control*, 57(7):1736–1751, 2012.
- [11] M. Huang, R. P. Malhamé, and P. E. Caines. Large population stochastic dynamic games: closed-loop McKean-Vlasov systems and the Nash certainty equivalence principle. *Communications in Information & Systems*, 6(3):221–252, 2006.
- [12] E. Koutsoupias and C. Papadimitriou. Worst-case equilibria. In *Annual Symposium on Theoretical Aspects of Computer Science*, pages 404–413. Springer, 1999.
- [13] J. M. Lasry and P. L. Lions. Mean field games. *Japanese Journal of Mathematics*, 2(1):229–260, 2007.
- [14] M. Nourian, P. E. Caines, and R. P. Malhamé. Synthesis of Cucker-Smale type flocking via mean field stochastic control theory: Nash equilibria. In *Communication, Control, and Computing (Allerton), 2010 48th Annual Allerton Conference on*, pages 814–819. IEEE, 2010.
- [15] M. Nourian, P. E. Caines, and R. P. Malhamé. Mean field analysis of controlled Cucker-Smale type flocking: Linear analysis and perturbation equations. *IFAC Proceedings Volumes*, 44(1):4471–4476, 2011.
- [16] M. Nourian, P. E. Caines, R. P. Malhamé, and M. Huang. Nash, social and centralized solutions to consensus problems via mean field control theory. *IEEE Transactions on Automatic Control*, 58(3):639–653, 2013.

- [17] T. Roughgarden. Intrinsic robustness of the price of anarchy. In *Proceedings of the forty-first annual ACM symposium on Theory of computing*, pages 513–522. ACM, 2009.
- [18] T. Roughgarden and É. Tardos. How bad is selfish routing? *Journal of the ACM (JACM)*, 49(2):236–259, 2002.
- [19] Q. Zhu and T. Başar. Price of anarchy and price of information in N-person linear-quadratic differential games. In *American Control Conference (ACC), 2010*, pages 762–767. IEEE, 2010.

Appendices

A Solving Linear FBSDEs of McKean-Vlasov Type

Consider a linear FBSDE system of McKean-Vlasov type:

$$\begin{aligned}
dX_t &= (a_t^x X_t + a_t^{\bar{x}} \mathbb{E}X_t + a_t^y Y_t + a_t^{\bar{y}} \mathbb{E}Y_t) dt + \sigma dW_t \\
X_0 &= \xi \\
dY_t &= (b_t^x X_t + b_t^{\bar{x}} \mathbb{E}X_t + b_t^y Y_t + b_t^{\bar{y}} \mathbb{E}Y_t) dt + Z_t dW_t \\
Y_T &= c^x X_T + c^{\bar{x}} \mathbb{E}X_T.
\end{aligned} \tag{37}$$

For the LQEMFG model considered in Section 2.1, the FBSDE system in equation (2) is of the form of equation (37) if we set:

$$\begin{aligned}
a_t^x &= b_1(t), \quad a_t^{\bar{x}} = \bar{b}_1(t), \quad a_t^y = a^{MFG}(t)b_2(t), \quad a_t^{\bar{y}} = b^{MFG}(t)b_2(t) + c^{MFG}(t)\bar{b}_2(t) \\
b_t^x &= -(q(t) + \bar{q}(t)), \quad b_t^{\bar{x}} = \bar{q}(t)s(t), \quad b_t^y = -b_1(t), \quad b_t^{\bar{y}} = 0 \\
c_t^x &= q_T + \bar{q}_T, \quad c_t^{\bar{x}} = -\bar{q}_T s_T.
\end{aligned}$$

For the LQEMKV model considered in Section 2.2, the FBSDE system in equation (8) is of the form of equation (37) if we set:

$$\begin{aligned}
a_t^x &= b_1(t), \quad a_t^{\bar{x}} = \bar{b}_1(t), \quad a_t^y = a^{MKV}(t)b_2(t), \quad a_t^{\bar{y}} = b^{MKV}(t)b_2(t) + c^{MKV}(t)\bar{b}_2(t) \\
b_t^x &= -(q(t) + \bar{q}(t)), \quad b_t^{\bar{x}} = -s(t)\bar{q}(t)(s(t) - 2), \quad b_t^y = -b_1(t), \quad b_t^{\bar{y}} = -\bar{b}_1 \\
c_t^x &= q_T + \bar{q}_T, \quad c_t^{\bar{x}} = s_T \bar{q}_T (s_T - 2).
\end{aligned}$$

Now we return to the general FBSDE system (37). By taking expectations in equation (37), and letting \bar{x}_t and \bar{y}_t denote $\mathbb{E}X_t$ and $\mathbb{E}Y_t$, respectively, we get:

$$\begin{aligned}
\dot{\bar{x}}_t &= (a_t^x + a_t^{\bar{x}})\bar{x}_t + (a_t^y + a_t^{\bar{y}})\bar{y}_t \\
\bar{x}_0 &= \mathbb{E}(\xi) \\
\dot{\bar{y}}_t &= (b_t^x + b_t^{\bar{x}})\bar{x}_t + (b_t^y + b_t^{\bar{y}})\bar{y}_t \\
\bar{y}_T &= (c^x + c^{\bar{x}})\bar{x}_T
\end{aligned} \tag{38}$$

where the dot is the standard ODE notation for a derivative. We then make the ansatz $\bar{y}_t = \bar{\eta}_t \bar{x}_t + \bar{\chi}_t$ for deterministic functions $[0, T] \ni t \mapsto \bar{\eta}_t \in \mathbb{R}$ and $[0, T] \ni t \mapsto \bar{\chi}_t \in \mathbb{R}$. By plugging in the ansatz,

the system in equation (38) is equivalent to the ODE system:

$$\begin{aligned}\dot{\bar{\eta}}_t + (a_t^y + a_t^{\bar{y}})\bar{\eta}_t^2 + (a_t^x + a_t^{\bar{x}} - b_t^y - b_t^{\bar{y}})\bar{\eta}_t - b_t^x - b_t^{\bar{x}} &= 0 \\ \bar{\eta}_T - c^x - c^{\bar{x}} &= 0 \\ \dot{\bar{\chi}}_t + (\bar{\eta}_t(a_t^y + a_t^{\bar{y}}) - b_t^y - b_t^{\bar{y}})\bar{\chi}_t &= 0 \\ \bar{\chi}_T &= 0.\end{aligned}$$

The first equation is a Riccati equation. Note that $\bar{\chi}_t$ solves a first order homogeneous linear equation. Thus $\bar{\chi}_t = 0, \forall t \in [0, T]$. Once the equation for $\bar{\eta}_t$ is solved, we can compute \bar{x}_t by solving the linear ODE:

$$\begin{aligned}\dot{\bar{x}}_t &= (a_t^x + a_t^{\bar{x}} + (a_t^y + a_t^{\bar{y}})\bar{\eta}_t)\bar{x}_t \\ \bar{x}_0 &= \mathbb{E}(\xi)\end{aligned}$$

and thus,

$$\bar{x}_t = \mathbb{E}(\xi) e^{\int_0^t (a_u^x + a_u^{\bar{x}} + (a_u^y + a_u^{\bar{y}})\bar{\eta}_u) du}.$$

Once we have computed $(\bar{x}_t)_{0 \leq t \leq T}$, we can rewrite the original FBSDE system:

$$\begin{aligned}dX_t &= (a_t^x X_t + a_t^y Y_t + a_t^0) dt + \sigma dW_t \\ X_0 &= \xi \\ dY_t &= (b_t^x X_t + b_t^y Y_t + b_t^0) dt + Z_t dW_t \\ Y_T &= c^x X_T + c^0\end{aligned}$$

with:

$$\begin{aligned}a_t^0 &= (a_t^{\bar{x}} + a_t^{\bar{y}}\bar{\eta}_t)\bar{x}_t \\ b_t^0 &= (b_t^{\bar{x}} + b_t^{\bar{y}}\bar{\eta}_t)\bar{x}_t \\ c^0 &= c^{\bar{x}}\bar{x}_T.\end{aligned}$$

Now we make the ansatz: $Y_t = \eta_t X_t + \chi_t$, which reduces the problem to the ODE system:

$$\begin{aligned}\dot{\eta}_t + a_t^y \eta_t^2 + (a_t^x - b_t^y)\eta_t - b_t^x &= 0 \\ \eta_T &= c^x, \\ \dot{\chi}_t + (-b_t^y + a_t^y \eta_t)\chi_t + a_t^0 \eta_t - b_t^0 &= 0, \\ \chi_T &= c^0, \\ Z_t &= \sigma \eta_t.\end{aligned}$$

Again, the first equation is a Riccati equation. Note that it is not necessary to solve for χ_t because of the relationship:

$$\bar{\eta}_t \bar{x}_t = \bar{y}_t = \mathbb{E}(Y_t) = \mathbb{E}(\eta_t X_t + \chi_t) = \eta_t \bar{x}_t + \chi_t.$$

Thus,

$$\chi_t = (\bar{\eta}_t - \eta_t)\bar{x}_t.$$

In summary, the solution to the linear FBSDE of McKean-Vlasov type is reduced to solving linear ODEs and Riccati equations. It will also be useful to compute $Var(X_t)$, which we denote by v_t . After we have solved the above equations, we have

$$\begin{aligned} dX_t &= ((a_t^x + a_t^y \eta_t)X_t + a_t^y \chi_t + a_t^0) dt + \sigma dW_t \\ X_0 &= \xi. \end{aligned}$$

Thus,

$$v_t = Var(X_t) = Var(\xi) e^{\int_0^t 2(a_s^x + a_s^y \eta_s) ds} + \sigma^2 \int_0^t e^{2 \int_s^t (a_u^x + a_u^y \eta_u) du} ds.$$

In the case where the coefficients are time-independent, the Riccati equations for $\bar{\eta}_t$ and η_t can be solved explicitly.

Scalar Riccati Equation

If the scalar Riccati equation

$$\dot{\rho}_t - B\rho_t^2 - 2A\rho_t + C = 0$$

with terminal condition $\rho_T = D$ satisfies:

$$B \neq 0, BD \geq 0, BC > 0, \quad (39)$$

then it has a unique solution:

$$\rho_t = \frac{(C + D\delta^+) - (C + D\delta^-)e^{-(\delta^+ - \delta^-)(T-t)}}{(-\delta^- + DB) + (\delta^+ - DB)e^{-(\delta^+ - \delta^-)(T-t)}} \quad (40)$$

where $\delta^\pm = -A \pm \sqrt{(A)^2 + BC}$.

Furthermore, if $B \rightarrow 0$ and $A \neq 0$, we can deduce that the limiting solution of the scalar Riccati equation coincides with the linear first-order differential equation:

$$\dot{\rho}_t - 2A\rho_t + C = 0$$

with terminal condition $\rho_T = D$, namely:

$$\rho_t = \left(D - \frac{C}{2A}\right) e^{-2A(T-t)} + \frac{C}{2A}.$$

If $B \rightarrow 0$ and $A = 0$, the limiting solution of the scalar Riccati equation coincides with the linear first-order differential equation:

$$\dot{\rho}_t + C = 0$$

with terminal condition $\rho_T = D$, namely:

$$\rho_t = D + C(T - t).$$

Hence, returning to the linear FBSDE (37), for $\bar{\eta}_t$, we use:

$$\begin{cases} A = -\frac{1}{2}(a^x + a^{\bar{x}} - b^y - b^{\bar{y}}) \\ B = -(a^y + a^{\bar{y}}) \\ C = -(b^x + b^{\bar{x}}) \\ D = c^x + c^{\bar{x}} \end{cases}$$

The conditions (39) are satisfied if:

$$\left\{ \begin{array}{l} -(a^y + a^{\bar{y}}) > 0 \\ -(b^x + b^{\bar{x}}) > 0 \\ c^x + c^{\bar{x}} \geq 0 \end{array} \right.$$

For η_t , we use:

$$\left\{ \begin{array}{l} A = -\frac{1}{2}(a^x - b^y) \\ B = -a^y \\ C = -b^x \\ D = c^x. \end{array} \right.$$

The conditions (39) are satisfied if:

$$\left\{ \begin{array}{l} -a^y > 0 \\ -b^x > 0 \\ c^x \geq 0. \end{array} \right.$$

Returning to the LGEMFG and LGEMKV problems, if we assume the coefficients are nonnegative, we see that these conditions are exactly assumption (14).