

ANNEX I

SUMMARY OF PRODUCT CHARACTERISTICS

▼ This medicinal product is subject to additional monitoring. This will allow quick identification of new safety information. Healthcare professionals are asked to report any suspected adverse reactions. See section 4.8 for how to report adverse reactions.

1. NAME OF THE MEDICINAL PRODUCT

TECVAYLI 10 mg/mL solution for injection
TECVAYLI 90 mg/mL solution for injection

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

TECVAYLI 10 mg/mL solution for injection
One 3 mL vial contains 30 mg of teclistamab (10 mg/mL).

TECVAYLI 90 mg/mL solution for injection
One 1.7 mL vial contains 153 mg of teclistamab (90 mg/mL).

Teclistamab is a humanised immunoglobulin G4-proline, alanine, alanine (IgG4-PAA) bispecific antibody directed against the B cell maturation antigen (BCMA) and CD3 receptors, produced in a mammalian cell line (Chinese hamster ovary [CHO]) using recombinant DNA technology.

For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Solution for injection (injection).

The solution is colourless to light yellow, with a pH of 5.2 and osmolarity of approximately 296 mOsm/L (10 mg/mL solution for injection), and approximately 357 mOsm/L (90 mg/mL solution for injection).

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

TECVAYLI is indicated as monotherapy for the treatment of adult patients with relapsed and refractory multiple myeloma, who have received at least three prior therapies, including an immunomodulatory agent, a proteasome inhibitor, and an anti-CD38 antibody and have demonstrated disease progression on the last therapy.

4.2 Posology and method of administration

Treatment with TECVAYLI should be initiated and supervised by physicians experienced in the treatment of multiple myeloma.

TECVAYLI should be administered by a healthcare professional with adequately trained medical personnel and appropriate medical equipment to manage severe reactions, including cytokine release syndrome (CRS) (see section 4.4).

Posology

Pre-treatment medicinal products should be administered prior to each dose of TECVAYLI in the step-up dosing schedule (see below).

TECVAYLI step-up dosing schedule should not be administered in patients with active infection (see Table 3 and section 4.4).

Recommended dosing schedule

The recommended dosing schedule for TECVAYLI is provided in Table 1. The recommended doses of TECVAYLI are 1.5 mg/kg by subcutaneous injection (SC) weekly, preceded by step-up doses of 0.06 mg/kg and 0.3 mg/kg.

Treatment with TECVAYLI should be initiated according to the step-up dosing schedule in Table 1 to reduce the incidence and severity of cytokine release syndrome. Due to the risk of cytokine release syndrome, patients should be instructed to remain within proximity of a healthcare facility, and monitored for signs and symptoms daily for 48 hours after administration of all doses within the TECVAYLI step-up dosing schedule (see section 4.4).

Failure to follow the recommended doses or dosing schedule for initiation of therapy, or re-initiation of therapy after dose delays, may result in increased frequency and severity of adverse reactions related to mechanism of action, particularly cytokine release syndrome (see section 4.4).

Table 1: TECVAYLI dosing schedule

Dosing schedule	Day	Dose ^a	
Step-up dosing schedule ^e	Day 1	Step-up dose 1	0.06 mg/kg single dose
	Day 3 ^b	Step-up dose 2	0.3 mg/kg single dose
	Day 5 ^c	First maintenance dose	1.5 mg/kg single dose
Weekly dosing schedule ^e	One week after first maintenance dose and weekly thereafter ^d	Subsequent maintenance doses	1.5 mg/kg once weekly

^a Dose is based on actual body weight and should be administered subcutaneously.

^b Step-up dose 2 may be given between 2 to 7 days after Step-up dose 1.

^c First maintenance dose may be given between 2 to 7 days after Step-up dose 2. This is the first full treatment dose (1.5 mg/kg).

^d Maintain a minimum of five days between weekly maintenance doses.

^e See Table 2 for recommendations on restarting TECVAYLI after dose delays.

Duration of treatment

Patients should be treated with TECVAYLI until disease progression or unacceptable toxicity.

Pre-treatment medicinal products

The following pre-treatment medicinal products must be administered 1 to 3 hours before each dose of the TECVAYLI step-up dosing schedule (see Table 1) to reduce the risk of cytokine release syndrome (see sections 4.4 and 4.8).

- Corticosteroid (oral or intravenous dexamethasone 16 mg)
- Antihistamine (oral or intravenous diphenhydramine 50 mg, or equivalent)
- Antipyretics (oral or intravenous acetaminophen 650 to 1 000 mg, or equivalent)

Administration of pre-treatment medicinal products may also be required prior to administration of subsequent doses of TECVAYLI for the following patients:

- Patients who repeat doses within the TECVAYLI step-up dosing schedule due to dose delays (Table 2), or
- Patients who experienced CRS following the previous dose (Table 3).

Prevention of herpes zoster reactivation

Prior to starting treatment with TECVAYLI, antiviral prophylaxis should be considered for the prevention of herpes zoster virus reactivation, per local institutional guidelines.

Restarting TECVAYLI after dose delay

If a dose of TECVAYLI is delayed, therapy should be restarted based on the recommendations listed in Table 2 and TECVAYLI resumed according to the dosing schedule (see Table 1). Pre-treatment medicinal products should be administered as indicated in Table 2. Patients should be monitored accordingly (see section 4.2).

Table 2: Recommendations for restarting therapy with TECVAYLI after dose delay

Last dose administered	Duration of delay from the last dose administered	Action
Step-up dose 1	More than 7 days	Restart TECVAYLI step-up dosing schedule at Step-up dose 1 (0.06 mg/kg) ^a .
Step-up dose 2	8 days to 28 days	Repeat Step-up dose 2 (0.3 mg/kg) ^a and continue TECVAYLI step-up dosing schedule.
	More than 28 days	Restart TECVAYLI step-up dosing schedule at Step-up dose 1 (0.06 mg/kg) ^a .
Any maintenance doses	8 days to 28 days	Continue TECVAYLI dosing schedule at maintenance dose (1.5 mg/kg) ^a .
	more than 28 days	Restart TECVAYLI step-up dosing schedule at Step-up dose 1 (0.06 mg/kg) ^a .

^a Pre-treatment medicinal products should be administered prior to TECVAYLI dose and patients monitored accordingly.

Dose modifications

Treatment with TECVAYLI should be initiated according to the step-up dosing schedule in Table 1.

Dose reductions of TECVAYLI are not recommended.

Dose delays may be required to manage toxicities related to TECVAYLI (see section 4.4). Recommendations on restarting TECVAYLI after a dose delay are provided in Table 2.

Recommended actions after adverse reactions following administration of TECVAYLI are listed in Table 3.

Table 3: Recommended actions taken after adverse reactions following administration of TECVAYLI

Adverse reactions	Grade	Actions
Cytokine release syndrome ^a (see section 4.4)	Grade 1 <ul style="list-style-type: none"> • Temperature $\geq 38^{\circ}\text{C}^{\text{b}}$ 	<ul style="list-style-type: none"> • Withhold TECVAYLI until adverse reaction resolves. • See Table 4 for management of cytokine release syndrome. • Administer pre-treatment medicinal products prior to next dose of TECVAYLI.
	Grade 2 <ul style="list-style-type: none"> • Temperature $\geq 38^{\circ}\text{C}^{\text{b}}$ with either: <ul style="list-style-type: none"> • Hypotension responsive to fluids and not requiring vasopressors, or • Oxygen requirement of low-flow nasal cannula^c or blow-by 	<ul style="list-style-type: none"> • Withhold TECVAYLI until adverse reaction resolves. • See Table 4 for management of cytokine release syndrome. • Administer pre-treatment medicinal products prior to next dose of TECVAYLI. • Monitor patient daily for 48 hours following the next dose of TECVAYLI. Instruct patients to remain within proximity of a healthcare facility during daily monitoring.
	Grade 3 (Duration: less than 48 hours) <ul style="list-style-type: none"> • Temperature $\geq 38^{\circ}\text{C}^{\text{b}}$ with either: <ul style="list-style-type: none"> • Hypotension requiring one vasopressor with or without vasopressin, or • Oxygen requirement of high-flow nasal cannula^c, facemask, non-rebreather mask, or Venturi mask 	<ul style="list-style-type: none"> • Monitor patient daily for 48 hours following the next dose of TECVAYLI. Instruct patients to remain within proximity of a healthcare facility during daily monitoring.
	Grade 3 (Recurrent or duration: more than 48 hours) <ul style="list-style-type: none"> • Temperature $\geq 38^{\circ}\text{C}^{\text{b}}$ with either: <ul style="list-style-type: none"> • Hypotension requiring one vasopressor with or without vasopressin, or • Oxygen requirement of high-flow nasal cannula^c, facemask, non-rebreather mask, or Venturi mask. 	<ul style="list-style-type: none"> • Permanently discontinue therapy with TECVAYLI. • See Table 4 for management of cytokine release syndrome.
	Grade 4 <ul style="list-style-type: none"> • Temperature $\geq 38^{\circ}\text{C}^{\text{b}}$ with either: <ul style="list-style-type: none"> • Hypotension requiring multiple vasopressors (excluding vasopressin), or • Oxygen requirement of positive pressure (e.g., continuous positive airway pressure [CPAP], bilevel positive airway pressure [BiPAP], intubation, and mechanical ventilation). 	

Immune effector cell-associated neurotoxicity syndrome (ICANS) ^d (see section 4.4)	Grade 1	<ul style="list-style-type: none"> • Withhold TECVAYLI until adverse reaction resolves. • See Table 5 for management of immune effector cell-associated neurotoxicity syndrome.
	Grade 2 Grade 3 (First occurrence)	<ul style="list-style-type: none"> • Withhold TECVAYLI until adverse reaction resolves. • See Table 5 for management of immune effector cell-associated neurotoxicity syndrome. • Monitor patient daily for 48 hours following the next dose of TECVAYLI. Instruct patients to remain within proximity of a healthcare facility during daily monitoring.
	Grade 3 (Recurrent) Grade 4	<ul style="list-style-type: none"> • Permanently discontinue therapy with TECVAYLI. • See Table 5 for management of immune effector cell-associated neurotoxicity syndrome.
Infections (see section 4.4)	All Grades	<ul style="list-style-type: none"> • Do not administer TECVAYLI step-up dosing schedule in patients with active infection. TECVAYLI step-up dosing schedule may proceed upon resolution of active infection.
	Grade 3 Grade 4	<ul style="list-style-type: none"> • Withhold subsequent maintenance doses of TECVAYLI (i.e., doses administered after TECVAYLI step-up dosing schedule) until infection improves to Grade 2 or better.
Haematologic toxicities (see sections 4.4 and 4.8)	Absolute neutrophil count less than $0.5 \times 10^9/L$	<ul style="list-style-type: none"> • Withhold TECVAYLI until absolute neutrophil count is $0.5 \times 10^9/L$ or higher.
	Febrile neutropenia	<ul style="list-style-type: none"> • Withhold TECVAYLI until absolute neutrophil count is $1.0 \times 10^9/L$ or higher, and fever resolves.
	Haemoglobin less than 8 g/dL	<ul style="list-style-type: none"> • Withhold TECVAYLI until haemoglobin is 8 g/dL or higher.
	Platelet count less than 25 000/ μL Platelet count between 25 000/ μL and 50 000/ μL with bleeding	<ul style="list-style-type: none"> • Withhold TECVAYLI until platelet count is 25 000/μL or higher and no evidence of bleeding.
Other adverse reactions (see section 4.8) ^e	Grade 3 Grade 4	<ul style="list-style-type: none"> • Withhold TECVAYLI until adverse reaction improves to Grade 2 or better.

-
- ^a Based on American Society for Transplantation and Cellular Therapy (ASTCT) grading for CRS (Lee et al 2019).
- ^b Attributed to CRS. Fever may not always be present concurrently with hypotension or hypoxia as it may be masked by interventions such as antipyretics or anticytokine therapy (e.g., tocilizumab or corticosteroids).
- ^c Low-flow nasal cannula is ≤ 6 L/min, and high-flow nasal cannula is > 6 L/min.
- ^d Based on ASTCT grading for ICANS.
- ^e Based on National Cancer Institute Common Terminology Criteria for Adverse Events (NCI-CTCAE), Version 4.03.

Special populations

Paediatric population

There is no relevant use of TECVAYLI in the paediatric population for the treatment of multiple myeloma.

Elderly (65 years of age and older)

No dosage adjustment is necessary (see section 5.2).

Renal impairment

No dosage adjustment is recommended for patients with mild or moderate renal impairment (see section 5.2).

Hepatic impairment

No dosage adjustment is recommended for patients with mild hepatic impairment (see section 5.2).

Method of administration

TECVAYLI is for subcutaneous injection only.

For instructions on handling of the medicinal product before administration, see section 6.6.

4.3 Contraindications

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

4.4 Special warnings and precautions for use

Traceability

In order to improve the traceability of biological medicinal products, the name and the batch number of the administered product should be clearly recorded.

Cytokine release syndrome (CRS)

Cytokine release syndrome, including life-threatening or fatal reactions, may occur in patients receiving TECVAYLI.

Clinical signs and symptoms of CRS may include but are not limited to fever, hypoxia, chills, hypotension, tachycardia, headache, and elevated liver enzymes. Potentially life-threatening complications of CRS may include cardiac dysfunction, adult respiratory distress syndrome, neurologic toxicity, renal and/or hepatic failure, and disseminated intravascular coagulation (DIC).

Treatment should be initiated with TECVAYLI according to the step-up dosing schedule to reduce risk of CRS. Pre-treatment medicinal products (corticosteroids, antihistamine and antipyretics) should

be administered prior to each dose of the TECVAYLI step-up dosing schedule to reduce risk of CRS (see section 4.2).

The following patients should be instructed to remain within proximity of a healthcare facility and monitored daily for 48 hours:

- If the patient has received any dose within the TECVAYLI step-up dosing schedule (for CRS).
- If the patient has received TECVAYLI after experiencing Grade 2 or higher CRS.

Patients who experience CRS following their previous dose should be administered pre-treatment medicinal products prior to the next dose of TECVAYLI.

Patients should be counselled to seek medical attention should signs or symptoms of CRS occur. At the first sign of CRS, patients should be immediately evaluated for hospitalisation. Treatment with supportive care, tocilizumab and/or corticosteroids should be instituted, based on severity as indicated in Table 4 below. The use of myeloid growth factors, particularly granulocyte macrophage-colony stimulating factor (GM-CSF), has the potential to worsen CRS symptoms and should be avoided during CRS. Treatment with TECVAYLI should be withheld until CRS resolves as indicated in Table 3 (see section 4.2).

Management of cytokine release syndrome

CRS should be identified based on clinical presentation. Patients should be evaluated and treated for other causes of fever, hypoxia, and hypotension.

If CRS is suspected, TECVAYLI should be withheld until the adverse reaction resolves (see Table 3). CRS should be managed according to the recommendations in Table 4. Supportive care for CRS (including but not limited to anti-pyretic agents, intravenous fluid support, vasopressors, supplemental oxygen, etc.) should be administered as appropriate. Laboratory testing to monitor for disseminated intravascular coagulation (DIC), haematology parameters, as well as pulmonary, cardiac, renal, and hepatic function should be considered.

Table 4: Recommendations for management of cytokine release syndrome with tocilizumab and corticosteroids

Grade ^e	Presenting symptoms	Tocilizumab ^a	Corticosteroids ^b
Grade 1	Temperature $\geq 38^{\circ}\text{C}^{\text{c}}$	May be considered	Not applicable
Grade 2	Temperature $\geq 38^{\circ}\text{C}^{\text{c}}$ with either: <ul style="list-style-type: none"> • Hypotension responsive to fluids and not requiring vasopressors, or • Oxygen requirement of low-flow nasal cannula^d or blow-by 	<p>Administer tocilizumab^b 8 mg/kg intravenously over 1 hour (not to exceed 800 mg).</p> <p>Repeat tocilizumab every 8 hours as needed, if not responsive to intravenous fluids or increasing supplemental oxygen.</p> <p>Limit to a maximum of 3 doses in a 24-hour period; maximum total of 4 doses.</p>	<p>If no improvement within 24 hours of starting tocilizumab, administer methylprednisolone 1 mg/kg intravenously twice daily, or dexamethasone 10 mg intravenously every 6 hours.</p> <p>Continue corticosteroid use until the event is Grade 1 or less, then taper over 3 days.</p>

Grade 3	Temperature $\geq 38^{\circ}\text{C}^{\text{c}}$ with either: <ul style="list-style-type: none"> • Hypotension requiring one vasopressor with or without vasopressin, or • Oxygen requirement of high-flow nasal cannula^d, facemask, non-rebreather mask, or Venturi mask 	Administer tocilizumab 8 mg/kg intravenously over 1 hour (not to exceed 800 mg). Repeat tocilizumab every 8 hours as needed, if not responsive to intravenous fluids or increasing supplemental oxygen. Limit to a maximum of 3 doses in a 24-hour period; maximum total of 4 doses.	If no improvement, administer methylprednisolone 1 mg/kg intravenously twice daily, or dexamethasone 10 mg intravenously every 6 hours. Continue corticosteroid use until the event is Grade 1 or less, then taper over 3 days.
Grade 4	Temperature $\geq 38^{\circ}\text{C}^{\text{c}}$ with either: <ul style="list-style-type: none"> • Hypotension requiring multiple vasopressors (excluding vasopressin), or • Oxygen requirement of positive pressure (e.g., continuous positive airway pressure [CPAP], bilevel positive airway pressure [BiPAP], intubation, and mechanical ventilation) 	Administer tocilizumab 8 mg/kg intravenously over 1 hour (not to exceed 800 mg). Repeat tocilizumab every 8 hours as needed if not responsive to intravenous fluids or increasing supplemental oxygen. Limit to a maximum of 3 doses in a 24-hour period; maximum total of 4 doses.	As above, or administer methylprednisolone 1 000 mg intravenously per day for 3 days, per physician discretion. If no improvement or if condition worsens, consider alternate immunosuppressants ^b .

^a Refer to tocilizumab prescribing information for details.

^b Treat unresponsive CRS per institutional guidelines.

^c Attributed to CRS. Fever may not always be present concurrently with hypotension or hypoxia as it may be masked by interventions such as antipyretics or anticytokine therapy (e.g., tocilizumab or corticosteroids).

^d Low-flow nasal cannula is ≤ 6 L/min, and high-flow nasal cannula is >6 L/min.

^e Based on ASTCT grading for CRS (Lee et al 2019).

Neurologic toxicities

Serious or life-threatening neurologic toxicities, including Immune Effector Cell-Associated Neurotoxicity Syndrome (ICANS) may occur following treatment with TECVAYLI.

Patients should be monitored for signs or symptoms of neurologic toxicities during treatment and treated promptly.

Patients should be counselled to seek medical attention should signs or symptoms of neurologic toxicity occur. At the first sign of neurologic toxicity, including ICANS, patients should be immediately evaluated and treated based on severity. Patients who experience Grade 2 or higher ICANS or first occurrence of Grade 3 ICANS with the previous dose of TECVAYLI should be instructed to remain within proximity of a healthcare facility and monitored for signs and symptoms daily for 48 hours.

For ICANS and other neurologic toxicities, treatment with TECVAYLI should be withheld as indicated in Table 3 (see section 4.2).

Due to the potential for ICANS, patients should be advised not to drive or operate heavy machinery during the TECVAYLI step-up dosing schedule and for 48 hours after completing the TECVAYLI step-up dosing schedule and in the event of new onset of any neurological symptoms (see section 4.7).

Management of neurologic toxicities

At the first sign of neurologic toxicity, including ICANS, neurology evaluation should be considered. Other causes of neurologic symptoms should be ruled out. TECVAYLI should be withheld until adverse reaction resolves (see Table 3). Intensive care and supportive therapy should be provided for severe or life-threatening neurologic toxicities. General management for neurologic toxicity (e.g., ICANS with or without concurrent CRS) is summarised in Table 5.

Table 5: Guidelines for management of immune effector cells-associated neurotoxicity syndrome (ICANS)

Grade	Presenting symptoms ^a	Concurrent CRS	No Concurrent CRS
Grade 1	ICE score 7-9 ^b Or, depressed level of consciousness ^c : awakens spontaneously.	Management of CRS per Table 4.	Monitor neurologic symptoms and consider neurology consultation and evaluation, per physician discretion.
		Monitor neurologic symptoms and consider neurology consultation and evaluation, per physician discretion. Consider non-sedating, anti-seizure medicinal products (e.g., levetiracetam) for seizure prophylaxis.	
Grade 2	ICE score 3-6 ^b Or, depressed level of consciousness ^c : awakens to voice.	Administer tocilizumab per Table 4 for management of CRS. If no improvement after starting tocilizumab, administer dexamethasone ^d 10 mg intravenously every 6 hours if not already taking other corticosteroids. Continue dexamethasone use until resolution to Grade 1 or less, then taper.	Administer dexamethasone ^d 10 mg intravenously every 6 hours. Continue dexamethasone use until resolution to Grade 1 or less, then taper.
		Consider non-sedating, anti-seizure medicinal products (e.g., levetiracetam) for seizure prophylaxis. Consider neurology consultation and other specialists for further evaluation, as needed.	
Grade 3	ICE score 0-2 ^b Or, depressed level of consciousness ^c : awakens only to tactile stimulus, or seizures ^c , either: • any clinical seizure, focal or generalised	Administer tocilizumab per Table 4 for management of CRS. In addition, administer dexamethasone ^d 10 mg intravenously with the first dose of tocilizumab, and repeat dose every 6 hours. Continue dexamethasone use until resolution to Grade 1 or less, then taper.	Administer dexamethasone ^d 10 mg intravenously every 6 hours. Continue dexamethasone use until resolution to Grade 1 or less, then taper.

	<p>that resolves rapidly, or</p> <ul style="list-style-type: none"> • non-convulsive seizures on electroencephalogram (EEG) that resolve with intervention, or <p>raised intracranial pressure: focal/local oedema on neuroimaging^c.</p>	<p>Consider non-sedating, anti-seizure medicinal products (e.g., levetiracetam) for seizure prophylaxis. Consider neurology consultation and other specialists for further evaluation, as needed.</p>	
Grade 4	<p>ICE score 0^b</p> <p>Or, depressed level of consciousness either:</p> <ul style="list-style-type: none"> • patient is unarousable or requires vigorous or repetitive tactile stimuli to arouse, or • stupor or coma, or <p>seizures^c, either:</p> <ul style="list-style-type: none"> • life-threatening prolonged seizure (>5 minutes), or • repetitive clinical or electrical seizures without return to baseline in between, or <p>motor findings^c:</p> <ul style="list-style-type: none"> • deep focal motor weakness such as hemiparesis or paraparesis, or <p>raised intracranial pressure / cerebral oedema^c, with signs/symptoms such as:</p> <ul style="list-style-type: none"> • diffuse cerebral oedema on neuroimaging, or • decerebrate or decorticate posturing, or • cranial nerve VI palsy, or • papilloedema, or • cushing's triad 	<p>Administer tocilizumab per Table 4 for management of CRS.</p> <p>As above, or consider administration of methylprednisolone 1 000 mg per day intravenously with first dose of tocilizumab, and continue methylprednisolone 1 000 mg per day intravenously for 2 or more days.</p>	<p>As above, or consider administration of methylprednisolone 1 000 mg per day intravenously for 3 days; if improves, then manage as above.</p>
		<p>Consider non-sedating, anti-seizure medicinal products (e.g., levetiracetam) for seizure prophylaxis. Consider neurology consultation and other specialists for further evaluation, as needed. In case of raised intracranial pressure/cerebral oedema, refer to institutional guidelines for management.</p>	

-
- ^a Management is determined by the most severe event, not attributable to any other cause.
- ^b If patient is arousable and able to perform Immune Effector Cell-Associated Encephalopathy (ICE) Assessment, assess: **Orientation** (oriented to year, month, city, hospital = 4 points); **Naming** (name 3 objects, e.g., point to clock, pen, button = 3 points); **Following Commands** (e.g., “show me 2 fingers” or “close your eyes and stick out your tongue” = 1 point); **Writing** (ability to write a standard sentence = 1 point; and **Attention** (count backwards from 100 by ten = 1 point). If patient is unarousable and unable to perform ICE Assessment (Grade 4 ICANS) = 0 points.
- ^c Attributable to no other cause.
- ^d All references to dexamethasone administration are dexamethasone or equivalent

Infections

Severe, life-threatening, or fatal infections have been reported in patients receiving TECVAYLI (see section 4.8). New or reactivated viral infections occurred during therapy with TECVAYLI. Progressive multifocal leukoencephalopathy (PML) has also occurred during therapy with TECVAYLI.

Patients should be monitored for signs and symptoms of infection prior to and during treatment with TECVAYLI and treated appropriately. Prophylactic antimicrobials should be administered according to local institutional guidelines.

TECVAYLI step-up dosing schedule should not be administered in patients with active infection. For subsequent doses, TECVAYLI should be withheld as indicated in Table 3 (see section 4.2).

Hepatitis B virus reactivation

Hepatitis B virus reactivation can occur in patients treated with medicinal products directed against B cells, and in some cases, may result in fulminant hepatitis, hepatic failure, and death.

Patients with evidence of positive HBV serology should be monitored for clinical and laboratory signs of HBV reactivation while receiving TECVAYLI, and for at least six months following the end of TECVAYLI treatment.

In patients who develop reactivation of HBV while on TECVAYLI, treatment with TECVAYLI should be withheld as indicated in Table 3 and manage per local institutional guidelines (see section 4.2).

Hypogammaglobulinaemia

Hypogammaglobulinaemia has been reported in patients receiving TECVAYLI (see section 4.8).

Immunoglobulin levels should be monitored during treatment with TECVAYLI. Intravenous or subcutaneous immunoglobulin therapy was used to treat hypogammaglobulinemia in 39% of patients. Patients should be treated according to local institutional guidelines, including infection precautions, antibiotic or antiviral prophylaxis, and administration of immunoglobulin replacement.

Vaccines

Immune response to vaccines may be reduced when taking TECVAYLI.

The safety of immunisation with live viral vaccines during or following TECVAYLI treatment has not been studied. Vaccination with live virus vaccines is not recommended for at least 4 weeks prior to the start of treatment, during treatment and least 4 weeks after treatment.

Neutropenia

Neutropenia and febrile neutropenia have been reported in patients who received TECVAYLI (see section 4.8).

Complete blood cell counts should be monitored at baseline and periodically during treatment. Supportive care should be provided per local institutional guidelines.

Patients with neutropenia should be monitored for signs of infection.

Treatment with TECVAYLI should be withheld as indicated in Table 3 (see section 4.2).

Excipients

This medicinal product contains less than 1 mmol (23 mg) sodium per dose, that is to say essentially 'sodium-free'.

4.5 Interaction with other medicinal products and other forms of interaction

No interaction studies have been performed with TECVAYLI.

The initial release of cytokines associated with the start of TECVAYLI treatment could suppress CYP450 enzymes. The highest risk of interaction is expected to be from initiation of TECVAYLI step-up schedule up to 7 days after the first maintenance dose or during a CRS event. During this time period, toxicity or medicinal product concentrations (e.g., cyclosporine) should be monitored in patients who are receiving concomitant CYP450 substrates with a narrow therapeutic index. The dose of the concomitant medicinal product should be adjusted as needed.

4.6 Fertility, pregnancy and lactation

Women of child-bearing potential/Contraception in males and females

Pregnancy status for females of child-bearing potential should be verified prior to starting treatment with TECVAYLI.

Women of child-bearing potential should use effective contraception during treatment and for 3 months after the final dose of TECVAYLI. In clinical studies, male patients with a female partner of child-bearing potential used effective contraception during treatment and for three months after the last dose of teclistamab.

Pregnancy

There are no available data on the use of teclistamab in pregnant women or animal data to assess the risk of teclistamab in pregnancy. Human IgG is known to cross the placenta after the first trimester of pregnancy. Therefore, teclistamab, a humanised IgG4-based antibody, has the potential to be transmitted from the mother to the developing foetus. TECVAYLI is not recommended for women who are pregnant. TECVAYLI is associated with hypogammaglobulinaemia, therefore, assessment of immunoglobulin levels in newborns of mothers treated with TECVAYLI should be considered.

Breast-feeding

It is not known whether teclistamab is excreted in human or animal milk, affects breast-fed infants or affects milk production. Because of the potential for serious adverse reactions in breast-fed infants from TECVAYLI, patients should be advised not to breast-feed during treatment with TECVAYLI and for at least three months after the last dose.

Fertility

There are no data on the effect of teclistamab on fertility. Effects of teclistamab on male and female fertility have not been evaluated in animal studies.

4.7 Effects on ability to drive and use machines

TECVAYLI has major influence on the ability to drive and use machines.

Due to the potential for ICANS, patients receiving TECVAYLI are at risk of depressed level of consciousness (see section 4.8). Patients should be instructed to avoid driving and operating heavy or potentially dangerous machinery during and for 48 hours after completion of TECVAYLI step-up dosing schedule and in the event of new onset of any neurological symptoms (Table 1) (see section 4.2 and section 4.4).

4.8 Undesirable effects

The most frequent adverse reactions of any grade in patients were hypogammaglobulinaemia (75%), cytokine release syndrome (72%), neutropenia (71%), anaemia (55%), musculoskeletal pain (52%), fatigue (41%), thrombocytopenia (40%), injection site reaction (38%), upper respiratory tract infection (37%), lymphopenia (35%), diarrhoea (28%), pneumonia (28%), nausea (27%), pyrexia (27%), headache (24%), cough (24%), constipation (21%) and pain (21%).

Serious adverse reactions were reported in 65% patients who received TECVAYLI, including pneumonia (16%), COVID-19 (15%), cytokine release syndrome (8%), sepsis (7%), pyrexia (5%), musculoskeletal pain (5%), acute kidney injury (4.8%), diarrhoea (3.0%), cellulitis (2.4%), hypoxia (2.4%), febrile neutropenia (2.4%), and encephalopathy (2.4%).

Tabulated list of adverse reactions

The safety data of TECVAYLI was evaluated in MajesTEC-1, which included 165 adult patients with multiple myeloma who received the recommended dosing regimen of TECVAYLI as monotherapy. The median duration of TECVAYLI treatment was 8.5 (Range: 0.2 to 24.4) months.

Table 6 summarises adverse reactions reported in patients who received TECVAYLI. The safety data of TECVAYLI was also evaluated in the all treated population (N=302) with no additional adverse reactions identified.

Adverse reactions observed during clinical studies are listed below by frequency category. Frequency categories are defined as follows: very common ($\geq 1/10$); common ($\geq 1/100$ to $< 1/10$); uncommon ($\geq 1/1\ 000$ to $< 1/100$); rare ($\geq 1/10\ 000$ to $< 1/1\ 000$); very rare ($< 1/10\ 000$) and not known (frequency cannot be estimated from the available data).

Within each frequency grouping, adverse reactions are presented in order of decreasing seriousness.

Table 6: Adverse reactions in patients with multiple myeloma treated with TECVAYLI in MajesTEC-1 at the recommended dose for monotherapy use

System Organ Class	Adverse Reaction	Frequency (All grades)	N=165	
			n (%)	
			Any Grade	Grade 3 or 4
Infections and infestations	Pneumonia ¹	Very common	46 (28%)	32 (19%)
	Sepsis ²	Common	13 (7.9%)	11 (6.7%)
	COVID-19 ³	Very common	30 (18%)	20 (12%)
	Upper respiratory tract infection ⁴	Very common	61 (37%)	4 (2.4%)
	Cellulitis	Common	7 (4.2%)	5 (3.0%)
Blood and lymphatic system disorders	Neutropenia	Very common	117 (71%)	106 (64%)
	Febrile neutropenia	Common	6 (3.6%)	5 (3.0%)

	Thrombocytopenia	Very common	66 (40%)	35 (21%)
	Lymphopenia	Very common	57 (35%)	54 (33%)
	Anaemia ⁵	Very common	90 (55%)	61 (37%)
	Leukopenia	Very common	29 (18%)	12 (7.3%)
	Hypofibrinogenaemia	Common	16 (9.7%)	2 (1.2%)
	Immune system disorders			
	Cytokine release syndrome	Very common	119 (72%)	1 (0.6%)
	Hypogammaglobulinaemia ⁶	Very common	123 (75%)	3 (1.8%)

Metabolism and nutrition disorders	Hyperamylasaemia	Common	6 (3.6%)	4 (2.4%)
	Hyperkalaemia	Common	8 (4.8%)	2 (1.2%)
	Hypercalcaemia	Very common	19 (12%)	5 (3.0%)
	Hyponatraemia	Common	13 (7.9%)	8 (4.8%)
	Hypokalaemia	Very common	23 (14%)	8 (4.8%)
	Hypocalcaemia	Common	12 (7.3%)	0
	Hypophosphataemia	Very common	20 (12%)	10 (6.1%)
	Hypoalbuminaemia	Common	4 (2.4%)	1 (0.6%)
	Hypomagnesaemia	Very common	22 (13%)	0
	Decreased appetite	Very common	20 (12%)	1 (0.6%)
Nervous system disorders	Immune effector cell-associated neurotoxicity syndrome	Common	5 (3.0%)	0
	Encephalopathy ⁷	Common	16 (9.7%)	0
	Neuropathy peripheral ⁸	Very common	26 (16%)	1 (0.6%)
	Headache	Very common	39 (24%)	1 (0.6%)
Vascular disorders	Hemorrhage ⁹	Very common	20 (12%)	5 (3.0%)
	Hypertension ¹⁰	Very common	21 (13%)	9 (5.5%)
Respiratory, thoracic and mediastinal disorders	Hypoxia	Common	16 (9.7%)	6 (3.6%)
	Dyspnoea ¹¹	Very common	22 (13%)	3 (1.8%)
	Cough ¹²	Very common	39 (24%)	0
Gastrointestinal disorders	Diarrhoea	Very common	47 (28%)	6 (3.6%)
	Vomiting	Very common	21 (13%)	1 (0.6%)
	Nausea	Very common	45 (27%)	1 (0.6%)
	Constipation	Very common	34 (21%)	0
Musculoskeletal and connective tissue disorders	Musculoskeletal pain ¹³	Very common	85 (52%)	14 (8.5%)
General disorders and administration site conditions	Pyrexia	Very common	45 (27%)	1 (0.6%)
	Injection site reaction ¹⁴	Very common	62 (38%)	1 (0.6%)
	Pain ¹⁵	Very common	34 (21%)	3 (1.8%)
	Oedema ¹⁶	Very common	23 (14%)	0
	Fatigue ¹⁷	Very common	67 (41%)	5 (3.0%)

Investigations	Blood creatinine increased	Common	9 (5.5%)	0
	Transaminase elevation ¹⁸	Common	16 (9.7%)	4 (2.4%)
	Lipase increased	Common	10 (6.1%)	2 (1.2%)
	Blood alkaline phosphatase increased	Very common	18 (11%)	3 (1.8%)
	Gamma-glutamyltransferase increased	Common	16 (9.7%)	5 (3.0%)
	Activated partial thromboplastin time prolonged	Common	13 (7.9%)	2 (1.2%)
	International normalised ratio increased	Common	10 (6.1%)	2 (1.2%)

Adverse events are coded using MedDRA Version 24.0.

Note: The output includes the diagnosis of CRS and ICANS; the symptoms of CRS or ICANS are excluded.

- ¹ Pneumonia includes Enterobacter pneumonia, lower respiratory tract infection, lower respiratory tract infection viral, Metapneumovirus pneumonia, Pneumocystis jirovecii pneumonia, pneumonia, Pneumonia adenoviral, Pneumonia bacterial, Pneumonia klebsiella, Pneumonia moraxella, Pneumonia pneumococcal, Pneumonia pseudomonal, Pneumonia respiratory syncytial viral, Pneumonia staphylococcal and Pneumonia viral.
- ² Sepsis includes bacteraemia, Meningococcal sepsis, neutropenic sepsis, Pseudomonal bacteraemia, Pseudomonal sepsis, sepsis and Staphylococcal bacteraemia.
- ³ COVID-19 includes asymptomatic COVID-19 and COVID-19.
- ⁴ Upper respiratory tract infection includes bronchitis, nasopharyngitis, pharyngitis, respiratory tract infection, respiratory tract infection bacterial, rhinitis, rhinovirus infection, sinusitis, tracheitis, upper respiratory tract infection and viral upper respiratory tract infection.
- ⁵ Anaemia includes anaemia, iron deficiency and iron deficiency anaemia.
- ⁶ Hypogammaglobulinaemia includes patients with adverse events of hypogammaglobulinaemia, hypoglobulinaemia, immunoglobulins decreased, and/or patients with laboratory IgG levels below 500 mg/dL following treatment with teclistamab.
- ⁷ Encephalopathy includes confusional state, depressed level of consciousness, lethargy, memory impairment and somnolence.
- ⁸ Neuropathy peripheral includes dysaesthesia, hypoaesthesia, hypoaesthesia oral, neuralgia, paraesthesia, paraesthesia oral, peripheral sensory neuropathy and sciatica.
- ⁹ Hemorrhage includes conjunctival hemorrhage, epistaxis, haematoma, haematuria, haemoperitoneum, haemorrhoidal hemorrhage, lower gastrointestinal hemorrhage, melaena, mouth hemorrhage and subdural haematoma.
- ¹⁰ Hypertension includes essential hypertension and hypertension.
- ¹¹ Dyspnoea includes acute respiratory failure, dyspnoea and dyspnoea exertional.
- ¹² Cough includes allergic cough, cough, productive cough and upper-airway cough syndrome.
- ¹³ Musculoskeletal pain includes arthralgia, back pain, bone pain, musculoskeletal chest pain, musculoskeletal pain, myalgia, neck pain and pain in extremity.
- ¹⁴ Injection site reaction includes injection site bruising, injection site cellulitis, injection site discomfort, injection site erythema, injection site haematoma, injection site induration, injection site inflammation, injection site oedema, injection site pruritus, injection site rash, injection site reaction and injection site swelling.
- ¹⁵ Pain includes ear pain, flank pain, groin pain, non-cardiac chest pain, oropharyngeal pain, pain, pain in jaw, toothache and tumour pain.
- ¹⁶ Oedema includes face oedema, fluid overload, oedema peripheral and peripheral swelling.
- ¹⁷ Fatigue includes asthenia, fatigue and malaise
- ¹⁸ Transaminase elevation includes alanine aminotransferase increased and aspartate aminotransferase increased.

Description of selected adverse reactions

Cytokine release syndrome

In MajesTEC-1 (N=165), CRS was reported in 72% of patients following treatment with TECVAYLI. One-third (33%) of patients experienced more than one CRS event. Most patients experienced CRS following Step-up Dose 1 (44%), Step-up Dose 2 (35%), or the initial maintenance dose (24%). Less than 3% of patients developed first occurrence of CRS following subsequent doses of TECVAYLI. CRS events were Grade 1 (50%) and Grade 2 (21%) or Grade 3 (0.6%). The median time to onset of CRS was 2 (Range: 1 to 6) days after the most recent dose, with a median duration of 2 (Range: 1 to 9) days.

The most frequent signs and symptoms associated with CRS were fever (72%), hypoxia (13%), chills (12%), hypotension (12%), sinus tachycardia (7%), headache (7%), and elevated liver enzymes (aspartate aminotransferase and alanine aminotransferase elevation) (3.6% each).

In MajesTEC-1, tocilizumab, corticosteroids and tocilizumab in combination with corticosteroids were used to treat CRS in 32%, 11% and 3% of CRS events, respectively.

Neurologic toxicities

In MajesTEC-1 (N=165), neurologic toxicity events were reported in 15% of patients receiving TECVAYLI. Neurologic toxicity events were Grade 1 (8.5%), Grade 2 (5.5%), or Grade 4 (<1%). The most frequently reported neurologic toxicity event was headache (8%).

ICANS was reported in 3% of patients receiving TECVAYLI at the recommended dose. The most frequent clinical manifestation of ICANS reported were confusional state (1.2%) and dysgraphia (1.2%). The onset of neurologic toxicity can be concurrent with CRS, following resolution of CRS, or in the absence of CRS. Seven of nine ICANS events (78%) were concurrent with CRS (during or within 7 days of CRS resolution). The median time to onset of ICANS was 4 (Range: 2 to 5) days after the most recent dose, with a median duration of 3 (Range: 1 to 20) days.

Immunogenicity

Patients treated with subcutaneous teclistamab monotherapy (N=238) in MajesTEC-1 were evaluated for antibodies to teclistamab using an electrochemiluminescence-based immunoassay. One subject (0.4%) developed neutralising antibodies to teclistamab of low-titre.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system listed in Appendix V.

4.9 Overdose

Symptoms and signs

The maximum tolerated dose of teclistamab has not been determined. In clinical studies, doses of up to 6 mg/kg have been administered.

Treatment

In the event of an overdose, the patient should be monitored for any signs or symptoms of adverse reactions, and appropriate symptomatic treatment should be instituted immediately.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: {group}, ATC code: not yet assigned

Mechanism of action

Teclistamab is a full-size, IgG4-PAA bispecific antibody that targets the CD3 receptor expressed on the surface of T cells and B cell maturation antigen (BCMA), which is expressed on the surface of

malignant multiple myeloma B-lineage cells, as well as late-stage B cells and plasma cells. With its dual binding sites, teclistamab is able to draw CD3⁺ T cells in close proximity to BCMA⁺ cells, resulting in T cell activation and subsequent lysis and death of BCMA⁺ cells, which is mediated by secreted perforin and various granzymes stored in the secretory vesicles of cytotoxic T cells. This effect occurs without regard to T cell receptor specificity or reliance on major histocompatibility complex (MHC) Class 1 molecules on the surface of antigen presenting cells.

Pharmacodynamic effects

Within the first month of treatment, activation of T-cells, redistribution of T-cells, reduction of B-cells and induction of serum cytokines were observed.

Within one month of treatment with teclistamab, the majority of responders had reduction in soluble BCMA, and a greater reduction in soluble BCMA was observed in subjects with deeper responses to teclistamab.

Clinical efficacy and safety

The efficacy of TECVAYLI monotherapy was evaluated in patients with relapsed or refractory multiple myeloma in a single-arm, open-label, multi-centre, Phase 1/2 study (MajesTEC-1). The study included patients who had previously received at least three prior therapies, including a proteasome inhibitor, an immunomodulatory agent, and an anti-CD38 monoclonal antibody. The study excluded patients who experienced stroke or seizure within the past 6 months, and patients with Eastern Cooperative Oncology Group performance score (ECOG PS) ≥ 2 , plasma cell leukaemia, known active CNS involvement or exhibited clinical signs of meningeal involvement of multiple myeloma, or active or documented history of autoimmune disease with the exception of vitiligo, Type 1 diabetes and prior autoimmune thyroiditis.

Patients received initial step-up doses of 0.06 mg/kg and 0.3 mg/kg of TECVAYLI administered subcutaneously, followed by the maintenance dose of TECVAYLI 1.5 mg/kg, administered subcutaneously once weekly thereafter, until disease progression or unacceptable toxicity (see section 4.2). The median duration between Step-up Dose 1 and Step-up Dose 2 was 2.9 (Range: 2-7) days. The median duration between Step-up Dose 2 and the initial maintenance dose was 3.1 (Range: 2-9) days. Patients were hospitalised for monitoring for at least 48 hours after administration of each dose of the TECVAYLI Step-up dosing schedule.

The efficacy population included 165 patients. The median age was 64 (Range: 33-84) years with 15% of subjects ≥ 75 years of age; 58% were male; 81% were White, 13% were Black, 2% were Asian. The International Staging System (ISS) at study entry was 52% in Stage I, 35% in Stage II and 12% in Stage III. High-risk cytogenetics (presence of del(17p), t(4;14) or t(14;16)) were present in 26% of patients. Seventeen percent of patients had extramedullary plasmacytomas.

The median time since initial diagnosis of multiple myeloma to enrolment was 6 (Range: 0.8-22.7) years. The median number of prior therapies was 5 (Range: 2-14), with 23% of patients who received 3 prior therapies. Eighty-two percent of patients received prior autologous stem cell transplantation, and 4.8% of patients received prior allogeneic transplantation. Seventy-eight percent of patients were triple-class refractory (refractory to proteasome inhibitor, an immunomodulatory agent and an anti-CD38 monoclonal antibody).

Efficacy results were based on overall response rate, as determined by the Independent Review Committee (IRC) assessment, using International Myeloma Working Group (IMWG) 2016 criteria (see Table 7).

Table 7: Efficacy results for MajesTEC-1

	All Treated (N=165)
Overall response rate (ORR: sCR, CR, VGPR, PR) n(%)	104 (63.0%)
95% CI (%)	(55.2%, 70.4%)
Stringent complete response (sCR)	54 (32.7%)
Complete response (CR)	11 (6.7%)
Very good partial response (VGPR)	32 (19.4%)
Partial response (PR)	7 (4.2%)
Duration of Response (DOR) (months)	
Number of Responders	104
DOR (Months): Median (95% CI)	18.4 (14.9, NE) ¹
Time to First Response (months)	
Number of responders	104
Median	1.2
Range	(0.2; 5.5)
MRD negativity rate² in all treated patients, n (%) [N=165]	44 (26.7%)
95% CI (%)	(20.1%, 34.1%)
MRD negativity rate^{2,3} in patients achieving CR or sCR, n (%) [N=65]	30 (46.2%)
95% CI (%)	(33.7%, 59.0%)

¹ NE=not estimable

² MRD-negativity rate is defined as the proportion of participants who achieved MRD negative status (at 10^{-5}) at any timepoint after initial dose, and prior to progressive disease (PD) or subsequent anti-myeloma therapy.

³ Only MRD assessments (10^{-5} testing threshold) within 3 months of achieving CR/sCR until death/progression/subsequent therapy (exclusive) are considered.

Paediatric population

The European Medicines Agency has waived the obligation to submit the results of studies with TECVAYLI in all subsets of the paediatric population in multiple myeloma (see section 4.2 for information on paediatric use).

5.2 Pharmacokinetic properties

Teclistamab exhibited approximately dose-proportional pharmacokinetics following subcutaneous administration across a dose range of 0.08 mg/kg to 3 mg/kg (0.05 to 2.0 times the recommended dose). The mean accumulation ratio following subcutaneous weekly dosing of teclistamab at steady state (based on the 7th weekly maintenance dose), was 2.71- and 3.05-fold for C_{max} and AUC_{tau} , respectively. The mean bioavailability following teclistamab subcutaneous administration was 69%, relative to intravenous dosing.

Pharmacokinetic parameters of teclistamab following the 1st and 7th recommended maintenance dose of 1.5 mg/kg are shown in Table 8.

Table 8: Pharmacokinetic parameters of teclistamab following the at first and seventh recommended maintenance dose (1.5 mg/kg) in patients with relapsed or refractory multiple myeloma in MajesTEC-1

Pharmacokinetic Parameters	1st maintenance dose of 1.5 mg/kg	7th maintenance dose of 1.5 mg/kg (steady-state)
T_{max} (hours)	72.0 (45.8 – 193) (n=40)	48.9 (0.0 – 166) (n=15)
C_{max} (µg/mL)	8.74 ± 3.65 (n=40)	25.3 ± 11.1 (n=15)
C_{trough} (µg/mL)	7.67 ± 3.52 (n=38)	22.1 ± 10.9 (n=27)
AUC_{tau} (µg·h/mL)	1 169 ± 481 (n=38)	3 905 ± 1 748 (n=13)

T_{\max} = Time to reach the C_{\max} ; C_{\max} = Maximum observed serum teclistamab concentration; C_{trough} = Observed serum teclistamab concentration prior to next dose; AUC_{tau} = Area under the concentration-time curve over the weekly dosing interval. Data are presented as mean \pm standard deviation, except for T_{\max} which is presented as median (minimum, maximum).

Distribution

Based on the population pharmacokinetic model, mean volume of distribution was 4.13 L (48.8% CV (coefficient of variation)) for the central compartment, and 1.34 L for the peripheral compartment.

Excretion

Teclistamab exhibited both time-independent and time-dependent clearance. Based on the population pharmacokinetic model, the mean time-independent clearance of teclistamab is 0.449 L/day (53.6% CV), with the median of time-dependent clearance contributing approximately 43% of the total clearance at baseline and decreasing rapidly thereafter to less than 10% after Week 8.

Based on non-compartmental analysis, the mean half-life (SD) was 3.8 (1.7) days (individual values ranging up to 8.8 days) following the first treatment intravenous dose of teclistamab.

Population pharmacokinetic analysis (based on MajesTEC-1) showed that soluble BCMA did not impact teclistamab serum concentrations.

Special populations

The pharmacokinetics of TECVAYLI in paediatric patients aged 17 years and younger have not been investigated.

Results of population pharmacokinetic analyses indicate that age (24 to 84 years) and sex did not influence the pharmacokinetics of teclistamab.

Renal impairment

No formal studies of TECVAYLI in patients with renal impairment have been conducted.

Results of population pharmacokinetic analyses indicate that mild renal impairment ($60 \text{ mL/min/1.73 m}^2 \leq \text{estimated glomerular filtration rate (eGFR)} < 90 \text{ mL/min/1.73 m}^2$) or moderate renal impairment ($30 \text{ mL/min/1.73 m}^2 \leq \text{eGFR} < 60 \text{ mL/min/1.73 m}^2$) did not significantly influence the pharmacokinetics of teclistamab. Limited data are available from patients with severe renal impairment.

Hepatic impairment

No formal studies of TECVAYLI in patients with hepatic impairment have been conducted.

Results of population pharmacokinetic analyses indicate that mild hepatic impairment (total bilirubin >1 to 1.5 times upper limit of normal (ULN) and any aspartate aminotransferase (AST), or total bilirubin $\leq \text{ULN}$ and $\text{AST} > \text{ULN}$) did not significantly influence the pharmacokinetics of teclistamab. No data are available in patients with moderate and severe hepatic impairment.

5.3 Preclinical safety data

Carcinogenicity and mutagenicity

No animal studies have been performed to assess the carcinogenic or genotoxic potential of teclistamab.

Reproductive toxicology and fertility

No animal studies have been conducted to evaluate the effects of teclistamab on reproduction and foetal development. In the 5-week repeat-dose toxicity study in cynomolgus monkeys, there were no notable effects in the male and female reproductive organs at doses up to 30 mg/kg/week (approximately 22 times the maximum recommended human dose, based on AUC exposure) intravenously for five weeks.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

EDTA disodium salt dihydrate
Glacial acetic acid
Polysorbate 20 (E432)
Sodium acetate trihydrate
Sucrose
Water for injections

6.2 Incompatibilities

In the absence of compatibility studies, this medicinal product must not be mixed with other medicinal products.

6.3 Shelf life

Unopened vial

18 months

Prepared syringe

The prepared syringes should be administered immediately. If immediate administration is not possible, in-use storage times of the prepared syringe should be no longer than 20 hours at 2 °C - 8 °C or ambient temperature (15 °C – 30 °C). Discard after 20 hours if not used.

6.4 Special precautions for storage

Store in a refrigerator (2 °C - 8 °C).
Do not freeze.
Store in the original carton in order to protect from light.

6.5 Nature and contents of container

3 mL solution for injection in a Type 1 glass vial with an elastomeric closure, and aluminium seal with a flip-off button containing 30 mg of teclistamab (10 mg/mL).
Pack size of 1 vial.

1.7 mL solution for injection in a Type 1 glass vial with an elastomeric closure, and aluminium seal with a flip-off button containing 153 mg of teclistamab (90 mg/mL).
Pack size of 1 vial.

6.6 Special precautions for disposal and other handling

It is very important that the instructions for preparation and administration provided in this section are strictly followed to minimise potential dosing errors with TECVAYLI 10 mg/mL and TECVAYLI 90 mg/mL vials.

TECVAYLI should be administered via subcutaneous injection only. Do not administer TECVAYLI intravenously.

TECVAYLI should be administered by a healthcare professional with adequately trained medical personnel and appropriate medical equipment to manage severe reactions, including cytokine release syndrome (see section 4.4).

TECVAYLI 10 mg/mL and TECVAYLI 90 mg/mL vials are for single use only.

TECVAYLI vials of different concentrations should not be combined to achieve maintenance dose.

Aseptic technique should be used to prepare and administer TECVAYLI.

Any unused medicinal product or waste material should be disposed in accordance with local requirements.

Preparation of TECVAYLI

- Verify the prescribed dose for each TECVAYLI injection. To minimise errors, use the following tables to prepare TECVAYLI injection.
 - Use Table 9 to determine the total dose, injection volume and number of vials required, based on patient's actual body weight for Step-up dose 1 using TECVAYLI 10 mg/mL vial.

Table 9: Injection volumes of TECVAYLI (10 mg/mL) for Step-up dose 1 (0.06 mg/kg)

Step-Up dose 1 (0.06 mg/kg)	Body weight (kg)	Total dose (mg)	Volume of injection (mL)	Number of vials (1 vial=3 mL)
	35-39	2.2	0.22	1
	40-44	2.5	0.25	1
	45-49	2.8	0.28	1
	50-59	3.3	0.33	1
	60-69	3.9	0.39	1
	70-79	4.5	0.45	1
	80-89	5.1	0.51	1
	90-99	5.7	0.57	1
	100-109	6.3	0.63	1
	110-119	6.9	0.69	1
	120-129	7.5	0.75	1
	130-139	8.1	0.81	1
	140-149	8.7	0.87	1
	150-160	9.3	0.93	1

- Use Table 10 to determine the total dose, injection volume and number of vials required based on patient's actual body weight for Step-up dose 2 using TECVAYLI 10 mg/mL vial.

Table 10: Injection volumes of TECVAYLI (10 mg/mL) for Step-up dose 2 (0.3 mg/kg)

	Body weight (kg)	Total dose (mg)	Volume of injection (mL)	Number of vials (1 vial=3 mL)
Step-up dose 2 (0.3 mg/kg)	35-39	11	1.1	1
	40-44	13	1.3	1
	45-49	14	1.4	1
	50-59	16	1.6	1
	60-69	19	1.9	1
	70-79	22	2.2	1
	80-89	25	2.5	1
	90-99	28	2.8	1
	100-109	31	3.1	2
	110-119	34	3.4	2
	120-129	37	3.7	2
	130-139	40	4.0	2
	140-149	43	4.3	2
	150-160	47	4.7	2

- Use Table 11 to determine the total dose, injection volume and number of vials required based on patient's actual body weight for the maintenance dose using TECVAYLI 90 mg/mL vial.

Table 11: Injection volumes of TECVAYLI (90 mg/mL) for maintenance dose (1.5 mg/kg)

	Body weight (kg)	Total dose (mg)	Volume of injection (mL)	Number of vials (1 vial=1.7 mL)
Maintenance dose (1.5 mg/kg)	35-39	56	0.62	1
	40-44	63	0.70	1
	45-49	70	0.78	1
	50-59	82	0.91	1
	60-69	99	1.1	1
	70-79	108	1.2	1
	80-89	126	1.4	1
	90-99	144	1.6	1
	100-109	153	1.7	1
	110-119	171	1.9	2
	120-129	189	2.1	2
	130-139	198	2.2	2
	140-149	216	2.4	2
	150-160	234	2.6	2

- Remove the appropriate TECVAYLI vial from refrigerated storage (2 °C – 8 °C) and equilibrate to ambient temperature (15 °C – 30 °C), as needed, for at least 15 minutes. Do not warm TECVAYLI in any other way.
- Once equilibrated, gently swirl the vial for approximately 10 seconds to mix. Do not shake.
- Withdraw the required injection volume of TECVAYLI from the vial(s) into an appropriately sized syringe using a transfer needle.
 - Each injection volume should not exceed 2.0 mL. Divide doses requiring greater than 2.0 mL equally into multiple syringes.
- TECVAYLI is compatible with stainless steel injection needles and polypropylene and polycarbonate syringe material.
- Replace the transfer needle with an appropriately sized needle for injection.
- Visually inspect TECVAYLI for particulate matter and discolouration prior to administration. Do not use if the solution is discoloured, or cloudy, or if foreign particles are present.
 - TECVAYLI solution for injection is colourless to light yellow.

Administration of TECVAYLI

- Inject the required volume of TECVAYLI into the subcutaneous tissue of the abdomen (preferred injection site). Alternatively, TECVAYLI may be injected into the subcutaneous tissue at other sites (e.g., thigh). If multiple injections are required, TECVAYLI injections should be at least 2 cm apart.
- Do not inject into tattoos or scars or areas where the skin is red, bruised, tender, hard or not intact.

7. MARKETING AUTHORISATION HOLDER

Janssen-Cilag International NV
Turnhoutseweg 30
B-2340 Beerse
Belgium

8. MARKETING AUTHORISATION NUMBER(S)

EU/1/22/1675/001 (10 mg/ml)
EU/1/22/1675/002 (90 mg/ml)

9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

10. DATE OF REVISION OF THE TEXT

Detailed information on this medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu>.

ANNEX II

- A. MANUFACTURER OF THE BIOLOGICAL ACTIVE SUBSTANCE AND MANUFACTURER RESPONSIBLE FOR BATCH RELEASE**
- B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**
- C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**
- D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**
- E. SPECIFIC OBLIGATION TO COMPLETE POST-AUTHORISATION MEASURES FOR THE CONDITIONAL MARKETING AUTHORISATION**

**A. MANUFACTURER OF THE BIOLOGICAL ACTIVE SUBSTANCE AND
MANUFACTURER RESPONSIBLE FOR BATCH RELEASE**

Name and address of the manufacturer of the biological active substance

Janssen Sciences Ireland UC
Barnahely,
Ringaskiddy, Co. Cork
Ireland

Name and address of the manufacturer responsible for batch release

Janssen Biologics B.V.
Einsteinweg 101
2333 CB Leiden
The Netherlands

B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE

Medicinal product subject to restricted medical prescription (see Annex I: Summary of Product Characteristics, section 4.2).

**C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING
AUTHORISATION**

- **Periodic safety update reports (PSURs)**

The requirements for submission of PSURs for this medicinal product are set out in the list of Union reference dates (EURD list) provided for under Article 107c(7) of Directive 2001/83/EC and any subsequent updates published on the European medicines web-portal.

The marketing authorisation holder (MAH) shall submit the first PSUR for this product within 6 months following authorisation.

**D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND
EFFECTIVE USE OF THE MEDICINAL PRODUCT**

- **Risk management plan (RMP)**

The marketing authorisation holder (MAH) shall perform the required pharmacovigilance activities and interventions detailed in the agreed RMP presented in Module 1.8.2 of the marketing authorisation and any agreed subsequent updates of the RMP.

An updated RMP should be submitted:

- At the request of the European Medicines Agency;
- Whenever the risk management system is modified, especially as the result of new information being received that may lead to a significant change to the benefit/risk profile or as the result of an important (pharmacovigilance or risk minimisation) milestone being reached.

- **Additional risk minimisation measures**

The MAH shall ensure that in each Member State where TECVAYLI is marketed, all patients/carers who are expected to use teclistamab have access to/are provided with the Patient Card which will

inform and explain to patients the risks of CRS. The Patient Card also includes a warning message for healthcare professionals treating the patient that the patient is receiving teclistamab.

The Patient Card will contain the following key messages:

- A description of the key signs and symptoms of CRS
- A description of when to seek urgent attention from the healthcare provider or seek emergency help, should signs and symptoms of CRS present themselves
- The prescribing physician's contact details

E. SPECIFIC OBLIGATION TO COMPLETE POST-AUTHORISATION MEASURES FOR THE CONDITIONAL MARKETING AUTHORISATION

This being a conditional marketing authorisation and pursuant to Article 14-a of Regulation (EC) No 726/2004, the MAH shall complete, within the stated timeframe, the following measures:

Description	Due date
In order to confirm the efficacy and safety of Teclistamab indicated as monotherapy for the treatment of adult patients with relapsed and refractory multiple myeloma, who have received at least three prior therapies, including an immunomodulatory agent, a proteasome inhibitor, and an anti-CD38 antibody, and have demonstrated disease progression on the last therapy, the MAH shall submit the results of study 64007957MMY3001, a Phase 3 Randomised Study Comparing Teclistamab in Combination with Daratumumab SC versus Daratumumab SC, Pomalidomide, and Dexamethasone (DPd) or Daratumumab SC, Bortezomib, and Dexamethasone (DVd) in Participants with Relapsed or Refractory Multiple Myeloma	March 2028
In order to further characterise the duration of response and long-term safety in subjects with multiple myeloma who have been previously treated with ≥ 3 prior lines of therapy, including an immunomodulatory agent, a PI and anti-CD38 antibody, the MAH shall submit the final study report of 64007957MMY1001, a Phase 1/2, First-in-Human, Open-Label, Dose Escalation Study of Teclistamab, a Humanised BCMA x CD3 Bispecific Antibody, in Subjects with Relapsed or Refractory Multiple Myeloma	December 2028

ANNEX III
LABELLING AND PACKAGE LEAFLET

A. LABELLING

PARTICULARS TO APPEAR ON THE OUTER PACKAGING**CARTON****1. NAME OF THE MEDICINAL PRODUCT**

TECVAYLI 10 mg/mL solution for injection
teclistamab

2. STATEMENT OF ACTIVE SUBSTANCE(S)

One 3 mL vial contains 30 mg of teclistamab (10 mg/mL)

3. LIST OF EXCIPIENTS

Excipients: EDTA disodium salt dihydrate, glacial acetic acid, polysorbate 20, sodium acetate trihydrate, sucrose, water for injections.

4. PHARMACEUTICAL FORM AND CONTENTS

Solution for injection
1 vial, 30 mg/3 mL
Step-up dose

5. METHOD AND ROUTE(S) OF ADMINISTRATION

Read the package leaflet before use.
For subcutaneous use only.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY

Do not shake.

8. EXPIRY DATE

EXP

9. SPECIAL STORAGE CONDITIONS

Store in a refrigerator.
Do not freeze.

Store in the original carton in order to protect from light.

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Janssen-Cilag International NV
Turnhoutseweg 30
B-2340 Beerse
Belgium

12. MARKETING AUTHORISATION NUMBER(S)

EU/1/22/1675/001

13. BATCH NUMBER

Lot

14. GENERAL CLASSIFICATION FOR SUPPLY

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

Justification for not including Braille accepted.

17. UNIQUE IDENTIFIER – 2D BARCODE

2D barcode carrying the unique identifier included.

18. UNIQUE IDENTIFIER - HUMAN READABLE DATA

PC
SN
NN

MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS VIAL LABEL
--

1. NAME OF THE MEDICINAL PRODUCT AND ROUTE(S) OF ADMINISTRATION
--

TECVAYLI 10 mg/mL injection
teclistamab
teclistamab
SC

2. METHOD OF ADMINISTRATION

3. EXPIRY DATE

EXP

4. BATCH NUMBER

Lot

5. CONTENTS BY WEIGHT, BY VOLUME OR BY UNIT
--

30 mg/3 mL

6. OTHER

PARTICULARS TO APPEAR ON THE OUTER PACKAGING**CARTON****1. NAME OF THE MEDICINAL PRODUCT**

TECVAYLI 90 mg/mL solution for injection
teclistamab

2. STATEMENT OF ACTIVE SUBSTANCE(S)

One 1.7 mL vial contains 153 mg of teclistamab (90 mg/mL).

3. LIST OF EXCIPIENTS

Excipients: EDTA disodium salt dihydrate, glacial acetic acid, polysorbate 20, sodium acetate trihydrate, sucrose, water for injections.

4. PHARMACEUTICAL FORM AND CONTENTS

Solution for injection
1 vial, 153 mg/1.7 mL
Maintenance dose

5. METHOD AND ROUTE(S) OF ADMINISTRATION

Read the package leaflet before use.
For subcutaneous use only.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY

Do not shake.

8. EXPIRY DATE

EXP

9. SPECIAL STORAGE CONDITIONS

Store in a refrigerator.
Do not freeze.

Store in the original carton in order to protect from light.

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Janssen-Cilag International NV
Turnhoutseweg 30
B-2340 Beerse
Belgium

12. MARKETING AUTHORISATION NUMBER(S)

EU/1/22/1675/002

13. BATCH NUMBER

Lot

14. GENERAL CLASSIFICATION FOR SUPPLY

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

Justification for not including Braille accepted.

17. UNIQUE IDENTIFIER – 2D BARCODE

2D barcode carrying the unique identifier included.

18. UNIQUE IDENTIFIER - HUMAN READABLE DATA

PC
SN
NN

MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS VIAL LABEL
--

1. NAME OF THE MEDICINAL PRODUCT AND ROUTE(S) OF ADMINISTRATION
--

TECVAYLI 90 mg/mL injection
teclistamab
teclistamab
SC

2. METHOD OF ADMINISTRATION

3. EXPIRY DATE

EXP

4. BATCH NUMBER

Lot

5. CONTENTS BY WEIGHT, BY VOLUME OR BY UNIT
--

153 mg/1.7 mL

6. OTHER

B. PACKAGE LEAFLET

Package leaflet: Information for the patient

TECVAYLI 10 mg/mL solution for injection TECVAYLI 90 mg/mL solution for injection

teclistamab

▼ This medicine is subject to additional monitoring. This will allow quick identification of new safety information. You can help by reporting any side effects you may get. See the end of section 4 for how to report side effects.

Read all of this leaflet carefully before you are given this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or nurse.
- If you get any side effects, talk to your doctor or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What TECVAYLI is and what it is used for
2. What you need to know before you are given TECVAYLI
3. How TECVAYLI is given
4. Possible side effects
5. How to store TECVAYLI
6. Contents of the pack and other information

1. What TECVAYLI is and what it is used for

TECVAYLI is a cancer medicine that contains the active substance ‘teclistamab’ and is used to treat adults with a type of cancer of the bone marrow called multiple myeloma. It is used for patients who have had at least three other kinds of treatment that have not worked or have stopped working.

How TECVAYLI works

TECVAYLI is an antibody, a type of protein which has been designed to recognise and attach to specific targets in your body. TECVAYLI targets B cell maturation antigen (BCMA), which is found on multiple myeloma cancer cells, and cluster of differentiation 3 (CD3), which is found on so-called T cells of your immune system. This medicine works by attaching to these cells and bringing them together, so that your immune system can destroy the multiple myeloma cancer cells.

2. What you need to know before you are given TECVAYLI

You must not be given TECVAYLI if you are allergic to teclistamab, or any of the other ingredients of this medicine (listed in section 6).

If you are not sure if you are allergic, talk to your doctor or nurse before you are given TECVAYLI.

Warnings and precautions

Talk to your doctor or nurse before you are given TECVAYLI if you have had a stroke or seizure within the past 6 months.

TECVAYLI and vaccines

Talk to your doctor or nurse before you are given TECVAYLI if you have had a recent vaccination or are going to have a vaccination.

You should not receive live vaccines from four weeks before until four weeks after you are treated with TECVAYLI.

Tests and checks

Before you are given TECVAYLI, your doctor will check your blood counts for signs of infection. If you have any infection, it will be treated before you start TECVAYLI. Your doctor will also check if you are pregnant or breast-feeding.

During treatment with TECVAYLI, your doctor will monitor you for side effects. Your doctor will regularly check your blood counts, as the number of blood cells and other blood components may decrease.

Look out for serious side effects.

Tell your doctor or nurse right away if you experience any of the following:

- Signs of a condition known as ‘cytokine release syndrome’ (CRS). Cytokine release syndrome is a serious immune reaction with symptoms such as fever, chills, nausea, headache, fast heartbeat, feeling dizzy, and difficulty breathing.
- Effects on your nervous system. Symptoms include feeling confused, feeling less alert, or having difficulty writing. Some of these may be signs of a serious immune reaction called ‘immune effector cell-associated neurotoxicity syndrome’ (ICANS).
- Signs and symptoms of an infection.

Tell your doctor or nurse if you notice any signs of the above.

Children and adolescents

Do not give TECVAYLI to children or young people below 18 years of age, because it is not known how this medicine will affect them.

Other medicines and TECVAYLI

Tell your doctor or nurse if you are taking, have recently taken, or might take any other medicines. This includes medicines you can get without a prescription and herbal medicines.

Pregnancy and breast-feeding

It is not known if TECVAYLI affects an unborn baby or if it passes into breast milk.

Pregnancy-information for women

Tell your doctor or nurse before you are given TECVAYLI if you are pregnant, think you might be pregnant or are planning to have a baby.

If you become pregnant while being treated with this medicine, tell your doctor or nurse straight away.

Pregnancy-information for men

If your partner becomes pregnant while you are taking this medicine, tell your doctor straight away.

Contraception

If you or your partner could become pregnant, you must use effective contraception during treatment and for 3 months after stopping treatment with TECVAYLI.

Breast-feeding

You and your doctor will decide if the benefit of breast-feeding is greater than the risk to your baby. If you and your doctor decide to stop taking this medicine, you should not breast-feed for 3 months after stopping treatment.

Driving and using machines

Some people may feel tired, dizzy, or confused while taking TECVAYLI. Do not drive, use tools, operate heavy machinery, or do things that could pose a danger to yourself until at least 48 hours after receiving your third dose of TECVAYLI, or as instructed by your doctor.

TECVAYLI contains sodium

TECVAYLI contains less than 1 mmol sodium (23 mg) per dose, that is to say essentially 'sodium-free'.

3. How TECVAYLI is given**How much is given**

Your doctor will determine your dose of TECVAYLI. The dose will depend on your body weight. The first two doses will be lower.

TECVAYLI is given as follows:

- You will receive 0.06 mg for each kilogram of bodyweight for your first dose.
- You will receive 0.3 mg per kilogram of bodyweight as your second dose 2-7 days later.
- You will then receive a 'Maintenance dose' of 1.5 mg per kilogram of bodyweight 2-7 days after your second dose.
- You will then continue receiving a 'Maintenance dose' once a week as long as you are getting benefit from TECVAYLI.

Your doctor will monitor you for side effects after each of your first three doses. They will do this for 2 days after each dose.

You should stay close to a healthcare facility after the first three doses in case you have side effects.

How the medicine is given

TECVAYLI will be given to you by a doctor or nurse as an injection under your skin ('subcutaneous' injection). It is given in the stomach area (abdomen) or thigh.

Other medicines given during treatment with TECVAYLI

You will be given medicines 1-3 hours before each of your first three doses of TECVAYLI, which help to lower the chance of side effects, such as cytokine release syndrome. These may include:

- medicines to reduce the risk of an allergic reaction (antihistamines)
- medicines to reduce the risk of inflammation (corticosteroids)
- medicines to reduce the risk of fever (such as paracetamol)

You may also be given these medicines for later doses of TECVAYLI based on any symptoms you have.

You may also be given additional medicines based on any symptoms you experience or your medical history.

If you are given more TECVAYLI than you should

This medicine will be given by your doctor or nurse, and it is unlikely that you will receive too much. In the event that you are given too much (an overdose), your doctor will check you for side effects.

If you forget your appointment to have TECVAYLI

It is very important to go to all your appointments. If you miss an appointment, make another one as soon as possible.

If you have any further questions on the use of this medicine, ask your doctor or nurse.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Serious side effects

Get medical help straight away if you get any of the following serious side effects, which may be severe and can be fatal.

Very common (may affect more than 1 in 10 people):

- serious immune reaction ('cytokine release syndrome') that may cause fever, chills, nausea, headache, fast heart beat, feeling dizzy, and difficulty breathing
- low level of antibodies called 'immunoglobulins' in the blood (hypogammaglobulinaemia), which may make infections more likely
- low levels of a type of white blood cells (neutropenia)
- infection, which may include fever, chills, shivering, cough, shortness of breath, rapid breathing and rapid pulse

Common (may affect up to 1 in 10 people):

- Effects on your nervous system. These may be signs of a serious immune reaction called 'immune effector cell associated neurotoxicity syndrome' (ICANS). Some of the symptoms are:
 - feeling confused
 - feeling less alert
 - having difficulty writing

Tell your doctor right away if you notice any of the above-listed serious side effects.

Other side effects

Other side effects are listed below. Tell your doctor or nurse if you get any of these side effects.

Very common (may affect more than 1 in 10 people):

- lung infection (pneumonia)
- COVID-19 infection caused by a virus called coronavirus (SARS-CoV-2)
- infected nose, sinuses or throat (upper respiratory tract infection)
- low levels of red blood cells (anaemia)
- low levels of blood platelets (cells that help blood to clot; thrombocytopaenia)
- low number of white blood cells (leukopenia)
- low levels of a type of white blood cells (lymphopenia)
- low level of 'phosphate', 'magnesium' or 'potassium' in the blood (hypophosphataemia, hypomagnesaemia or hypokalaemia)
- increased level of 'calcium' (hypercalcaemia)
- increased 'alkaline phosphatase' in the blood
- decreased appetite
- feeling sick (nausea), diarrhoea, constipation, vomiting
- headache
- nerve damage that may cause tingling, numbness, pain or loss of pain sensation
- high blood pressure (hypertension)
- bleeding, which can be severe (haemorrhage)
- cough
- being short of breath (dyspnea)
- fever
- feeling very tired
- pain or muscle aches
- swollen hands, ankles or feet (oedema)
- skin reactions at or near the injection site, including redness of the skin, itching, swelling, pain, bruising, rash, bleeding

Common (may affect up to 1 in 10 people)

- severe infection throughout the body (sepsis)
- skin infection causing redness (cellulitis)

- low number of a type of white blood cell with a fever (febrile neutropenia)
- low levels of 'fibrinogen,' a type of protein in the blood, making it more difficult to form clots
- change in brain function (encephalopathy)
- low level of 'calcium' or 'sodium' in the blood (hypocalcaemia or hyponatremia)
- high level of 'potassium' in the blood (hyperkalemia)
- low level of 'albumin' in the blood (hypoalbuminaemia)
- low level of oxygen in the blood (hypoxia)
- increased level of 'gamma-glutamyltransferase' in the blood
- increased level of liver enzymes 'transaminases' in the blood
- increased level of 'creatinine' in the blood
- increased level of 'amylase' in the blood (hyperamylasemia)
- increased level of 'lipase' in the blood (hyperlipasaemia)
- blood tests may show it takes longer for blood to clot (INR increased and PTT prolongation)

Reporting of side effects

If you get any side effects, talk to your doctor or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via [the national reporting system listed in Appendix V](#). By reporting side effects, you can help provide more information on the safety of this medicine.

5. How to store TECVAYLI

TECVAYLI will be stored at the hospital or clinic by your doctor.

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton and vial label after "EXP". The expiry date refers to the last day of that month.

Store in a refrigerator (2 °C - 8 °C). Do not freeze.

Store in the original carton in order to protect from light.

Medicines should not be disposed of via wastewater or household waste. Your healthcare professional will throw away any medicines that are no longer being used. These measures will help protect the environment.

6. Contents of the pack and other information

What TECVAYLI contains

- The active substance is teclistamab. TECVAYLI comes in two different strengths:
 - 10 mg/mL - one 3 mL vial contains 30 mg teclistamab
 - 90 mg/mL - one 1.7 mL vial contains 153 mg teclistamab
- The other ingredients are EDTA disodium salt dihydrate, glacial acetic acid, polysorbate 20, sodium acetate trihydrate, sucrose, water for injections (see "TECVAYLI contains sodium" in section 2).

What TECVAYLI looks like and contents of the pack

TECVAYLI is a solution for injection (injection) and is a colourless to light yellow liquid. TECVAYLI is supplied as a carton pack containing 1 glass vial.

Marketing Authorisation Holder

Janssen-Cilag International NV

Turnhoutseweg 30
B-2340 Beerse
Belgium

Manufacturer

Janssen Biologics B.V.
Einsteinweg 101
2333 CB Leiden
The Netherlands

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder:

België/Belgique/Belgien

Janssen-Cilag NV
Tel/Tél: +32 14 64 94 11
janssen@jacbe.jnj.com

Lietuva

UAB "JOHNSON & JOHNSON"
Tel: +370 5 278 68 88
lt@its.jnj.com

България

„Джонсън & Джонсън България” ЕООД
Тел.: +359 2 489 94 00
jjsafety@its.jnj.com

Luxembourg/Luxemburg

Janssen-Cilag NV
Tél/Tel: +32 14 64 94 11
janssen@jacbe.jnj.com

Česká republika

Janssen-Cilag s.r.o.
Tel: +420 227 012 227

Magyarország

Janssen-Cilag Kft.
Tel.: +36 1 884 2858
janssenhu@its.jnj.com

Danmark

Janssen-Cilag A/S
Tlf: +45 4594 8282
jacdk@its.jnj.com

Malta

AM MANGION LTD
Tel: +356 2397 6000

Deutschland

Janssen-Cilag GmbH
Tel: +49 2137 955 955
jancil@its.jnj.com

Nederland

Janssen-Cilag B.V.
Tel: +31 76 711 1111
janssen@jacnl.jnj.com

Eesti

UAB "JOHNSON & JOHNSON" Eesti filiaal
Tel: +372 617 7410
ee@its.jnj.com

Norge

Janssen-Cilag AS
Tlf: +47 24 12 65 00
jacno@its.jnj.com

Ελλάδα

Janssen-Cilag Φαρμακευτική Α.Ε.Β.Ε.
Τηλ: +30 210 80 90 000

Österreich

Janssen-Cilag Pharma GmbH
Tel: +43 1 610 300

España

Janssen-Cilag, S.A.
Tel: +34 91 722 81 00
contacto@its.jnj.com

Polska

Janssen-Cilag Polska Sp. z o.o.
Tel.: +48 22 237 60 00

France

Janssen-Cilag
Tél: 0 800 25 50 75 / +33 1 55 00 40 03
medisource@its.jnj.com

Portugal

Janssen-Cilag Farmacêutica, Lda.
Tel: +351 214 368 600

Hrvatska

Johnson & Johnson S.E. d.o.o.
Tel: +385 1 6610 700
jjsafety@JNJCR.JNJ.com

Ireland

Janssen Sciences Ireland UC
Tel: +353 1 800 709 122

Ísland

Janssen-Cilag AB
c/o Vistor hf.
Sími: +354 535 7000
janssen@vistor.is

Italia

Janssen-Cilag SpA
Tel: 800.688.777 / +39 02 2510 1
janssenita@its.jnj.com

Κύπρος

Βαρνάβας Χατζηπαναγής Ατδ
Τηλ: +357 22 207 700

Latvija

UAB "JOHNSON & JOHNSON" filiāle Latvijā
Tel: +371 678 93561
lv@its.jnj.com

România

Johnson & Johnson România SRL
Tel: +40 21 207 1800

Slovenija

Johnson & Johnson d.o.o.
Tel: +386 1 401 18 00
Janssen_safety_slo@its.jnj.com

Slovenská republika

Johnson & Johnson, s.r.o.
Tel: +421 232 408 400

Suomi/Finland

Janssen-Cilag Oy
Puh/Tel: +358 207 531 300
jacfi@its.jnj.com

Sverige

Janssen-Cilag AB
Tfn: +46 8 626 50 00
jacse@its.jnj.com

United Kingdom (Northern Ireland)

Janssen Sciences Ireland UC
Tel: +44 1 494 567 444

This leaflet was last revised in

This medicine has been given 'conditional approval'. This means that there is more evidence to come about this medicine.

The European Medicines Agency will review new information on this medicine at least every year and this leaflet will be updated as necessary.

Other sources of information

Detailed information on this medicine is available on the European Medicines Agency web site:
<http://www.ema.europa.eu>.

This leaflet is available in all EU/EEA languages on the European Medicines Agency website.

The following information is intended for healthcare professionals only:

It is very important to that the instructions for preparation and administration provided in this section are strictly followed to minimise potential dosing errors with TECVAYLI 10 mg/mL and TECVAYLI 90 mg/mL vials.

TECVAYLI should be administered via subcutaneous injection only. Do not administer TECVAYLI intravenously.

TECVAYLI should be administered by a healthcare professional with adequately trained medical personnel and appropriate medical equipment to manage severe reactions, including cytokine release syndrome.

TECVAYLI 10 mg/mL and TECVAYLI 90 mg/mL vials are for single use only.

TECVAYLI vials of different strengths should not be combined to achieve maintenance dose.

Aseptic technique should be used to prepare and administer TECVAYLI.

Any unused medicinal product or waste material should be disposed in accordance with local requirements.

Preparation of TECVAYLI

- Verify the prescribed dose for each TECVAYLI injection. To minimise errors, use the following tables to prepare TECVAYLI injection.
 - Use Table 1 to determine total dose, injection volume and number of vials required based on patient's actual body weight for Step-up dose 1 using TECVAYLI 10 mg/mL vial.

Table 1: Injection volumes of TECVAYLI (10 mg/mL) for Step-up dose 1 (0.06 mg/kg)

Step-Up dose 1 (0.06 mg/kg)	Body weight (kg)	Total dose (mg)	Volume of injection (mL)	Number of vials (1 vial=3 mL)
	35-39	2.2	0.22	1
	40-44	2.5	0.25	1
	45-49	2.8	0.28	1
	50-59	3.3	0.33	1
	60-69	3.9	0.39	1
	70-79	4.5	0.45	1
	80-89	5.1	0.51	1
	90-99	5.7	0.57	1
	100-109	6.3	0.63	1
	110-119	6.9	0.69	1
	120-129	7.5	0.75	1
	130-139	8.1	0.81	1
	140-149	8.7	0.87	1
	150-160	9.3	0.93	1

- Use Table 2 to determine total dose, injection volume and number of vials required based on patient's actual body weight for Step-up dose 2 using TECVAYLI 10 mg/mL vial.

Table 2: Injection volumes of TECVAYLI (10 mg/mL) for Step-up dose 2 (0.3 mg/kg)

Step-up dose 2 (0.3 mg/kg)	Body weight (kg)	Total dose (mg)	Volume of injection (mL)	Number of vials (1 vial=3 mL)
---------------------------------------	-----------------------------	----------------------------	-------------------------------------	--

35-39	11	1.1	1
40-44	13	1.3	1
45-49	14	1.4	1
50-59	16	1.6	1
60-69	19	1.9	1
70-79	22	2.2	1
80-89	25	2.5	1
90-99	28	2.8	1
100-109	31	3.1	2
110-119	34	3.4	2
120-129	37	3.7	2
130-139	40	4.0	2
140-149	43	4.3	2
150-160	47	4.7	2

- Use Table 3 to determine total dose, injection volume and number of vials required based on patient's actual body weight for the maintenance dose using TECVAYLI 90 mg/mL vial.

Table 3: Injection volumes of TECVAYLI (90 mg/mL) for maintenance dose (1.5 mg/kg)

	Body weight (kg)	Total dose (mg)	Volume of injection (mL)	Number of vials (1 vial=1.7 mL)
Maintenance dose (1.5 mg/kg)	35-39	56	0.62	1
	40-44	63	0.70	1
	45-49	70	0.78	1
	50-59	82	0.91	1
	60-69	99	1.1	1
	70-79	108	1.2	1
	80-89	126	1.4	1
	90-99	144	1.6	1
	100-109	153	1.7	1
	110-119	171	1.9	2
	120-129	189	2.1	2
	130-139	198	2.2	2
	140-149	216	2.4	2
	150-160	234	2.6	2

- Remove the appropriate strength TECVAYLI vial from refrigerated storage (2 °C–8 °C) and equilibrate to ambient temperature (15 °C – 30 °C), as needed, for at least 15 minutes. Do not warm TECVAYLI in any other way.
- Once equilibrated, gently swirl the vial for approximately 10 seconds to mix. Do not shake.
- Withdraw the required injection volume of TECVAYLI from the vial(s) into an appropriately sized syringe using a transfer needle.
 - Each injection volume should not exceed 2.0 mL. Divide doses requiring greater than 2.0 mL equally into multiple syringes.
- TECVAYLI is compatible with stainless steel needles, polypropylene and polycarbonate syringe material.
- Replace the transfer needle with an appropriately sized needle for injection.
- Visually inspect TECVAYLI for particulate matter and discolouration prior to administration. Do not use if the solution is discoloured, or cloudy, or if foreign particles are present.
 - TECVAYLI solution for injection is colourless to light yellow.

Administration of TECVAYLI

- Inject the required volume of TECVAYLI into the subcutaneous tissue of the abdomen (preferred injection site). Alternatively, TECVAYLI may be injected into the subcutaneous

tissue of the thigh. If multiple injections are required, TECVAYLI injections should be at least 2 cm apart.

- Do not inject into tattoos or scars or areas where the skin is red, bruised, tender, hard or not intact.

Traceability

In order to improve the traceability of biological medicinal products, the name and the batch number of the administered product should be clearly recorded.

ANNEX IV

CONCLUSIONS ON THE GRANTING OF THE CONDITIONAL MARKETING AUTHORISATION PRESENTED BY THE EUROPEAN MEDICINES AGENCY

Conclusions presented by the European Medicines Agency on:

- **Conditional marketing authorisation**

The CHMP having considered the application is of the opinion that the risk-benefit balance is favourable to recommend the granting of the conditional marketing authorisation as further explained in the European Public Assessment Report.